

12. A. de Rujula, S. L. Glashow, H. D. Politzer, S. B. Treiman, F. Wilczek and A. Zee, Princeton preprint (1974) (to be published).
13. C. H. Albright and C. Jarlskog, CERN-TH-1828 (1974).
14. We change the last term in Eq. (3.8c) of Ref. (12) to $17.5(1-x)^7$; this has the effect of flattening F_2 as $x \rightarrow 0$.
15. D. Gross, Phys. Rev. Letters 32 1071 (1974).
See also A. de Rujula, Phys. Rev. Letters 32 1143 (1974);
D. Gross and S. B. Treiman, Phys. Rev. Letters 1145 (1974).
16. Several authors have discussed the large ω limit in a manner similar to Ref. (12) but outside the context of asymptotically free theory:
V. N. Gribov and L. N. Lipatov, Yad. Fis. 15 781 (1972) [Sov. J. Nucl. Phys. 15, 438 (1972)] Eqns. 93 and ff) For further references see Ref. (12).
17. The calculations in this section were done with W. Caswell and R. Shrock (unpublished). M. Calvo (Princeton University thesis, unpublished) has independently analyzed corrections to other parton model sum rules as well as the Callan-Gross relation.
18. K. Wilson, Phys. Rev 179, 1499 (1969); C. Callan, Phys. Rev. D5 3203 (1972); K. Symanzik, Commun. Math. Phys. 23 491 (1971).
19. We thank S. Adler for emphasizing this question.
20. M. S. Chanowitz and S. D. Drell, Phys. Rev. Letters 30 807 (1973); V. Barger, Phys. Letters 49B 43 (1974).