

“STANDARD”

RADIUM
PREPARATIONS



PERMANENT

“Standard” Radium Solution for Drinking

—

Each bottle contains two micrograms radium element in 60 cc. aqua dist.

—

Maximum-equilibrium constant of radium emanation, 5400 mache units.



PERMANENT

“Standard” Radium Solution for Intravenous Use.

—

In Ampules of 2 cc. N. P. S. containing 5, 10, 25, 50, or 100 micrograms radium element.



PERMANENT RADIO-ACTIVITY

“Standard” Radium Compress

—

A means of applying radium locally for the relief of pain.

—

A flexible pad of standardized, guaranteed radium element content.

INDICATIONS

Subacute and Chronic Joint and Muscular Conditions.
High Blood Pressure. Nephritis.
The Simple and Pernicious Anemias.

“The value of radium is unquestionably established in chronic and subacute arthritis of all kinds (luetic and tuberculous excepted) acute, subacute and chronic joint and muscular rheumatism (so called) in gout, sciatica, neuralgia, polyneuritis, lumbago and the lancinating pain of tabes.”—Rowntree and Baetjer, Journal A. M. A. Oct. 18, 1913.

<p>For Descriptive and Clinical Literature Address.</p>	<p>RADIUM CHEMICAL COMPANY</p>	<p>Chicago C. W. Hanford, M. D., 749 1st Nat'l Bank Bldg.</p>
<p>New York C. Everett Field, M. D., 50 E. 41st St.</p>	<p>Boston Samuel Delano, M. D., 29 Newbury St.</p>	<p>San Francisco Fred I. Lackenbach Biologic Depot 908 Butler Bldg</p>
	<p>PITTSBURGH</p>	

FIGURE 3 This advertisement for radium preparations to be used internally and externally appeared in the June 1916 issue of the journal *Radium*.