

- ¹⁹ There may be a very serious problem with treating all these matrix elements on the same footing. Some of our operators have the form $\bar{p}'p'\bar{\lambda}n$. According to quark model ideas, and the empirical success of Zweig's rule for heavy quarks, we might expect the matrix elements of such operators between light hadrons to be very small. The reader should keep this caveat in mind in assessing the importance of these operators for weak nonleptonic decays.
- ²⁰ K. M. Smith et al., Nucl. Phys. B91, 45 (1975) and references therein.
- ²¹ S. Nussinov and J. Rosner, Phys. Rev. Lett. 23, 1264 (1969); J. C. Pati and C. H. Woo, Phys. Rev. D3, 2920 (1971); See also R. L. Kingsley, Phys. Letters 40B, 387 (1972).
- ²² M. Suzuki, Phys. Rev. Lett. 15, 986 (1965); H. Sugawara, Phys. Rev. Lett. 15, 870, 997 (ϵ) (1965); Y. Hara and Y. Nambu, Phys. Rev. Lett. 16, 875 (1966); C. Callan and S. Treiman, Phys. Rev. Lett. 16, 153 (1966).
- ²³ G. Altarelli, N. Cabibbo and L. Maiani, Nucl. Phys. B88, 285 (1975);
R. L. Kingsley, et al., Ref. 12.
- ²⁴ A. Benvenuti, et al., Phys. Rev. Lett. 34, 419 (1975).
- ²⁵ F. Wilczek, Ref. 3.
- ²⁶ J. -E. Augustin, et al., Phys. Rev. Lett. 34, 764 (1975).
- ²⁷ M. Gaillard, B. W. Lee and J. L. Rosner, Rev. Mod. Phys. 47, 277 (1975); See also Sec. III.