

closing of the 5f shell. The divalent ion is at least 1.3 V more stable than the monovalent.

This work was performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore Laboratory under contract No. W-7405-ENG-48.

REFERENCES

1. I. Zvara, V. Z. Belov, V. P. Domanov, B. L. Zhookov, T. Ryott, Z. Chubener, M. R. Shalaevski, Joint Institute for Nuclear Research, Dubna, USSR, Preprint P6-10334 (in Russian) 1976.
2. L. Brewer, J. Opt. Soc. Amer. 61, 1101 (1971).
3. E. K. Hulet, R. W. Lougheed, J. D. Brady, R. E. Stone, and M. S. Coops, Science 158, 486 (1967).
4. G. R. Choppin, B. G. Harvey, and S. G. Thompson, J. Inorg. Nucl. Chem. 2, 66 (1955).
5. E. P. Horwitz and C. A. A. Bloomquist, Inorg. Nucl. Chem. Lett. 5, 753 (1969).
6. J. Mály and B. B. Cunningham, J. Inorg. Nucl. Chem. Lett. 3, 455 (1967).
7. J. Mály, J. Inorg. Nucl. Chem. 31, 741 (1969).
8. K. Samhoun, F. David, R. L. Hahn, G. D. O'Kelley, J. R. Tarrant, and D. E. Hobart, J. Inorg. Nucl. Chem. 41, 1749 (1979).
9. F. David, K. Samhoun, E. K. Hulet, P. A. Baisden, R. Dougan, J. H. Landrum, R. W. Lougheed, J. F. Wild, and G. D. O'Kelley, to be published.
10. L. J. Nugent, J. Inorg. Nucl. Chem. 37, 1967 (1975).
11. N. B. Mikheev, V. I. Spitsyn, A. N. Kamenskaya, I. A. Rumer, B. A. Gvozdev, N. A. Rosenkevich, and L. N. Auerman, Dokl. Akad. Nauk SSSR 208, 1146 (1973).
12. E. K. Hulet, R. W. Lougheed, P. A. Baisden, J. H. Landrum, J. F. Wild, and R. F. D. Lundqvist, J. Inorg. Nucl. Chem. 41, 1743 (1979). ✓