

Experiments Utilizing ICRF Heating on TFTR

J.R. Wilson, J.C. Hosea, M.G. Bell, M. Bitter, R. Boivin, E.D. Fredrickson,
G.J. Greene, G.W. Hammett, K.W. Hill, D.J. Hoffman*, H. Hsuan,
M. Hughest†, A.C. Janos, D.L. Jassby, F.C. Jobes, D.W. Johnson,
C.K. Phillips, D.K. Mansfield, K.M. McGuire, S.S. Medley, D. Mueller,
Y. Nagayama‡, M. Ono, D.K. Owens, H.K. Park, M. Phillipst,
A.T. Ramsey, G.L. Schmidt, S.D. Scott, J.E. Stevens, B.C. Stratton,
E. Synakowski, G. Taylor, M. Ulrickson, K.L. Wong, M.C. Zarnstorff,
S.J. Zweben

Princeton Plasma Physics Laboratory

Princeton, New Jersey 08543

*Oak Ridge National Laboratory

Oak Ridge, Tennessee 08731

†Grumman Corporation

Princeton, New Jersey 08540

‡University of Tokyo

Tokyo, Japan

MASTER

ABSTRACT

A variety of experiments have been performed on the TFTR tokamak utilizing ICRF heating. Of special interest has been the insight into plasma performance gained by utilizing a different heating scheme other than the usual NBI. Utilizing ICRF heating