GA–A23155

A COMPARISON OF PLASMA PERFORMANCE BETWEEN SINGLE-NULL AND DOUBLE-NULL CONFIGURATIONS DURING ELMING H–MODE

by

T.W. PETRIE, M.E. FENSTERMACHER,* S.L. ALLEN,* T.N. CARLSTROM, P. GOHIL, R.J. GROEBNER, C.M. GREENFIELD, A.W. HYATT, C.J. LASNIER,* R.J. LA HAYE, A.W. LEONARD, M.A. MAHDAVI, T.H. OSBORNE, G.D. PORTER,* T.L. RHODES,† D.M. THOMAS, J.G. WATKINS,‡ W.P. WEST, N.S. WOLF,* and The DIII–D TEAM

This is a preprint of a paper presented at the Twenty-Sixth European Physical Society Conference on Controlled Fusion and Plasma Physics, June 14–18, 1999 in Maastricht, The Netherlands, and to be published in *The Proceedings.*

*Lawrence Livermore National Laboratory, Livermore, California.
†University of California, Los Angeles, California.
‡Sandia National Laboratories, Albuquerque, New Mexico.

Work supported by
the U.S. Department of Energy
under Contract Nos. DE-AC03-99ER54463, W-7405-ENG-48,
DE-AC04-94AL85000 and Grant No. DE-FG03-86ER53225

GA PROJECT 30033
JULY 1999

GENERAL ATOMICS