

Existence of Core Localized Toroidicity-induced Alfvén Eigenmode

G. Y. Fu

Princeton Plasma Physics Laboratory, Princeton University

P. O. Box 451

Princeton, New Jersey 08543, U.S.A.

(December 9, 1994)

Abstract

The core-localized toroidicity-induced Alfvén eigenmode (TAE) is shown to exist at finite plasma pressure due to finite aspect ratio effects in tokamak plasma. The new critical beta for the existence of the TAE mode is given by $\alpha \approx 3\epsilon + 2s^2$, where $\epsilon = r/R$ is the inverse aspect ratio, s is the magnetic shear and $\alpha = -Rq^2 d\beta/dr$ is the normalized pressure gradient. In contrast, previous critical α is given by $\alpha \approx s^2$. In the limit of $s \ll \sqrt{r/R}$, the new critical α is greatly enhanced by the finite aspect ratio effects.

52.35.Bj, 52.55Fa, 52.55.Pi

MASTER

Typeset using REVTeX

The toroidicity-induced Alfvén eigenmode [1,2] (TAE) exists in a tokamak plasma due to toroidal mode coupling and finite magnetic shear. The mode frequency is located inside a continuum gap induced by toroidicity. Recently, it has been shown [3–5] that there is a critical pressure gradient above which the TAE mode no longer exists. In the limit of small shear, the critical value is given by $\alpha = \alpha_c = s^2/(1 + s)$, where $s = rdq/qdr$ is the magnetic shear, $\alpha = -Rq^2d\beta/dr$ represents the product of magnetic field curvature and plasma pressure gradient with R being the major radius, r being the plasma radius, q being the safety factor and β being the plasma toroidal beta. This result implies that the TAE mode should not exist in the core of tokamak plasma where the values of α and s are comparable and small for typical parameters. In particular, $\alpha/s \gg 1$ asymptotically when r goes to zero. However, recent numerical results [6] indicate that a TAE mode, which is localized at a single gap, does exist in the core of the tokamak plasma, where $\epsilon = r/R \sim 0.1$ and $s \sim \alpha \sim \epsilon$. We call this type of TAE mode the core-localized TAE mode. These core-localized modes are particularly susceptible to destabilization by fusion alpha particles in a tokamak plasma since the density profile of the alpha particles is sharply peaked at the center of plasma [6].

In this work, we will show that the effects of finite aspect ratio change the critical α qualitatively using the high- n ballooning mode equation. The previous results [3,4], based on the standard $s - \alpha$ model ballooning mode equation [7], is only valid in the limit of $\epsilon \ll s^2$. We will show that our new critical α reduces to $\alpha_c = 3\epsilon$ in the limit of $s \ll \sqrt{\epsilon} \ll 1$. In the following, we will derive an analytic expression for critical α by taking into account the effects of finite aspect ratio.

We start with the following ballooning mode equation:

$$\begin{aligned} \frac{\partial}{\partial \theta} [G_1(\theta) + G_2(\theta)h^2(\theta)] \frac{\partial}{\partial \theta} \Phi + \Omega^2 [G_1(\theta) + G_2(\theta)h^2(\theta)] (1 + 4\epsilon \cos \theta) \Phi \\ + \alpha [\cos \theta + h(\theta) \sin \theta] \Phi = 0 \end{aligned} \quad (1)$$

where $h(\theta) = s\theta - \bar{\alpha} \sin \theta$, $G_1(\theta) = 1 - 2(\epsilon + \Delta') \cos \theta$, $G_2(\theta) = 1 + 2\Delta' \cos \theta$, $\bar{\alpha} =$

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, make any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

$\alpha + 2\epsilon - 2(1 - s)\Delta'$, $\Delta' = \epsilon(1/4 + \beta_p)$ is the radial derivative of the Shafranov shift, and Ω is the mode frequency normalized to the Alfvén frequency. The poloidal beta is defined as $\beta_p = 8\pi(\langle p \rangle - p)/B_p^2$ where $\langle \rangle$ denotes a volume average.

Equation 1 is derived for a large aspect ratio, low-beta tokamak plasma by using the Shafranov shifted circle flux coordinates [8]. We note that the previous work [3,4] neglects the ϵ term in $G_1(\theta)$ and $G_2(\theta)$ and assumes $\alpha \gg \epsilon$. In this limit, the Eq. (1) reduces to the standard $s - \alpha$ model ballooning equation [7]:

$$\begin{aligned} \frac{\partial}{\partial \theta} [1 + h^2(\theta)] \frac{\partial}{\partial \theta} \Phi + \Omega^2 [1 + h^2(\theta)] (1 + 4\epsilon \cos \theta) \Phi \\ + \alpha [\cos \theta + h(\theta) \sin \theta] \Phi = 0. \end{aligned} \quad (2)$$

Comparing Eqs. (1) and (2), the new terms in Eq. (1), of order $O(\epsilon)$, come from the intrinsic toroidicity and pressure-induced Shafranov shift. We will show below that these ϵ terms in G_1 and G_2 can not be neglected when ϵ is comparable or larger than s^2 .

To make analytic progress, we make the following transformation $\Phi \rightarrow (1/\sqrt{F})\Phi$ with $F = G_1(\theta) + G_2(\theta)h^2(\theta)$, the Eq. (1) then becomes

$$\frac{\partial^2}{\partial \theta^2} \Phi + \left[\Omega^2(1 + 4\epsilon \cos \theta) - \frac{F''}{2F} + \frac{(F')^2}{4F^2} + \frac{\alpha(\cos(\theta) + h(\theta) \sin \theta)}{F} \right] \Phi = 0 \quad (3)$$

where the prime denotes the derivative with respect to θ . Assuming $\epsilon \ll 1$ and $s \sim \alpha \sim O(\epsilon)$, we expand Eq. (3) to the second order of ϵ . The resulting equation is given by

$$\frac{\partial^2}{\partial \theta^2} \Phi + \Omega^2(1 + 4\hat{\epsilon} \cos \theta) \Phi + \left[\frac{H_1(\theta)}{1 + s^2\theta^2} - \frac{H_2(\theta)}{(1 + s^2\theta^2)^2} \right] \Phi = 0 \quad (4)$$

where the functions H_1 and H_2 are defined as

$$H_1(\theta) = (\alpha - \delta) \cos \theta - 2\alpha\Delta' \cos^2 \theta + (2\delta\Delta' + \alpha\bar{\alpha} - \bar{\alpha}^2) \sin^2 \theta \quad (5)$$

$$H_2(\theta) = (s - \bar{\alpha} \cos \theta)^2 + 2\delta(\delta - \alpha) \cos^2 \theta - (\delta^2 + 2\bar{\alpha}^2 - 2\alpha\bar{\alpha}) \sin^2 \theta \quad (6)$$

with $\hat{\epsilon} \approx \epsilon + \Delta'$ and $\delta = \epsilon + 2\Delta'$.

Following Fu and Cheng [3], a dispersion relation for the TAE mode frequency can be derived by solving Eq. (4) asymptotically. The result is given by

$$\sqrt{\frac{\Omega^2 - \Omega_-^2}{\Omega_+^2 - \Omega^2}} = \frac{\pi}{4s}(1 + \epsilon + s)(\alpha_c - \alpha) \quad (7)$$

where Ω_- and Ω_+ are the lower bound and the upper bound of the Alfvén continuum gap respectively. The critical α for the existence of TAE mode, α_c , is given by

$$\alpha_c = \frac{(1 + \epsilon)(\epsilon + 2\Delta') + s^2 - 2(\epsilon - \Delta')s}{1 + \epsilon + s}. \quad (8)$$

In the limit of $\epsilon \rightarrow 0$ and $\Delta' \rightarrow 0$, the critical value given in Eq. (8) reduces to the previous results [3–5]. It should be noted that Δ' is dependent on the poloidal beta which is proportional to α . This implicit dependence can be made explicit in the core of plasma where we can expand the plasma pressure near $r = 0$. Thus, we have following relations near the center of plasma for typical pressure profile of form $p \propto (1 - r^2)^b$ with b being a constant: $\alpha = 4\epsilon\beta_p$ and $\bar{\alpha} = \delta = 3\epsilon/2 + \alpha/2$. The critical α then reduces to

$$\alpha_c = \frac{3\epsilon(1 + \epsilon) + 2s^2 - 3\epsilon s}{1 + \epsilon + s} \quad (9)$$

It is instructive to note that in the limit of zero shear, the critical alpha is simply given by $\alpha_c = 3\epsilon$ which corresponds to $\beta_{pc} = 3/4$. Thus, the finite aspect ratio effects greatly enhance the critical beta for small shear. It is also interesting to note that in the limit of $\epsilon \rightarrow 0$, the critical α is a factor two larger than the previous results [3–5]. This factor of two comes from the pressure-induced Shafranov shift.

The critical α , given by Eq. (9), has been confirmed by solving Eq.(1) numerically. Figure 1 shows that the analytic results for the critical α (solid line) agree very well with the numerical results (solid dots). For comparison, we have shown in Fig. 2 both the previous results (solid line marked with $r/R = 0$) and our new results (solid line marked with $r/R = 0.1$). As a reference, the ballooning mode first stability boundary is also shown (dotted line). We observe that the effects of finite aspect ratio makes the critical α much

closer to the first stability boundary for small value of shear. (It was shown previously [3] that the critical α for TAE mode approaches the first stability boundary asymptotically for large values of shear). Although our results are based on shifted-circle model equilibria, we conjecture that the critical α for TAE mode is also close to the ballooning first stability limit in general equilibria, and that the TAE can generally exist as long as the tokamak plasma is ballooning stable.

Finally, the analytic critical α in the limit of $s^2 \ll \epsilon$, $\alpha_c = 3\epsilon$, has been confirmed in both magnitude and scaling by numerical calculations using a global stability code NOVA [6].

In conclusion, we have shown that the core-localized TAE mode exists at finite plasma pressure in tokamaks due to finite aspect ratio effects. The critical beta for the existence of TAE mode in the core region of plasma is given by $\beta_p = 3/4$ in the limit of $s \ll \sqrt{r/R} \ll 1$.

The author gratefully acknowledges stimulating discussion with Drs. H. L. Berk, C. Z. Cheng and J. W. Van Dam. This work is supported by the U.S. Department of Energy under Contract No. DE-AC02-76-CHO-3073.

REFERENCES

- [1] C. Z. Cheng and M. S. Chance, *Phys. Fluids* **29**, 3695 (1986).
- [2] C. Z. Cheng, L. Chen, and M. S. Chance, *Ann. Phys. NY* **161**, 21 (1984).
- [3] G. Y. Fu and C. Z. Cheng, *Phys. Fluids* **B2**, 985 (1990).
- [4] F. Zonca and L. Chen, *Bull. Am. Phys. Soc.* **37**, 1436 (1992).
- [5] J. W. Van Dam and H. L. Berk, *Bull. Am. Phys. Soc.* **38**, 1944 (1993).
- [6] G. Y. Fu et al., to be submitted to *Phys. Rev. Lett.*
- [7] J. W. Connor, R. J. Hastie and J. B. Taylor, *Phys. Rev. Lett.* **40**, 396 (1978).
- [8] J. D. Meiss and R. D. Hazeltine, *Phys. Fluids* **B2**, 2563 (1990).

FIGURES

FIG. 1. The critical α value, α_c , as a function of shear s obtained with $\epsilon = 0.1$ near the center of plasma.

FIG. 2. The critical α for the TAE as a function of shear s from both the previous results (solid line marked with $r/R = 0$) and the new results (solid line marked with $r/R = 0.1$). The ballooning first stability boundary is also shown (dotted line).




Fig. 1




Fig. 2

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA
 Prof. R.C. Cross, Univ. of Sydney, AUSTRALIA
 Plasma Research Lab., Australian Nat. Univ., AUSTRALIA
 Prof. I.R. Jones, Flinders Univ, AUSTRALIA
 Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA
 Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA
 Prof. M. Goossens, Astronomisch Instituut, BELGIUM
 Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM
 Commission-Europeen, DG. XII-Fusion Prog., BELGIUM
 Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM
 Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL
 Prof. Dr. I.C. Nascimento, Instituto Fisica, Sao Paulo, BRAZIL
 Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL
 Documents Office, Atomic Energy of Canada Ltd., CANADA
 Ms. M. Morin, CCFM/Tokamak de Varennes, CANADA
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
 Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA
 Prof. J. Teichmann, Univ. of Montreal, CANADA
 Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA
 Prof. T.W. Johnston, INRS-Energie, CANADA
 Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA
 Dr. C.R. James,, Univ. of Alberta, CANADA
 Dr. P. Lukác, Komenského Univerzita, CZECHO-SLOVAKIA
 The Librarian, Culham Laboratory, ENGLAND
 Library, R61, Rutherford Appleton Laboratory, ENGLAND
 Mrs. S.A. Hutchinson, JET Library, ENGLAND
 Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS
 P. Mähönen, Univ. of Helsinki, FINLAND
 Prof. M.N. Bussac, Ecole Polytechnique,, FRANCE
 C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE
 J. Radet, CEN/CADARACHE - Bat 506, FRANCE
 Prof. E. Economou, Univ. of Crete, GREECE
 Ms. C. Rinni, Univ. of Ioannina, GREECE
 Preprint Library, Hungarian Academy of Sci., HUNGARY
 Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA
 Dr. P. Kaw, Inst. for Plasma Research, INDIA
 Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL
 Librarian, International Center for Theo Physics, ITALY
 Miss C. De Palo, Associazione EURATOM-ENEA , ITALY
 Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY
 Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY
 Dr. H. Yamato, Toshiba Res & Devel Center, JAPAN
 Prof. I. Kawakami, Hiroshima Univ., JAPAN
 Prof. K. Nishikawa, Hiroshima Univ., JAPAN
 Librarian, Naka Fusion Research Establishment, JAERI, JAPAN
 Director, Japan Atomic Energy Research Inst., JAPAN
 Prof. S. Itoh, Kyushu Univ., JAPAN
 Research Info. Ctr., National Instit. for Fusion Science, JAPAN
 Prof. S. Tanaka, Kyoto Univ., JAPAN
 Library, Kyoto Univ., JAPAN
 Prof. N. Inoue, Univ. of Tokyo, JAPAN
 Secretary, Plasma Section, Electrotechnical Lab., JAPAN
 Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
 Dr. G.S. Lee, Korea Basic Sci. Ctr., KOREA
 J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
 D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
 Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
 Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
 Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
 Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
 Z. Li, S.W. Inst Physics, PEOPLE'S REPUBLIC OF CHINA
 Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
 Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
 Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
 Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
 Dr. L. Stenflo, Univ. of UMEA, SWEDEN
 Library, Royal Inst. of Technology, SWEDEN
 Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
 Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
 Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
 Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
 Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. Apparatus, USSR
 Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
 Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
 Librarian, The Ukr.SSR Academy of Sciences, USSR
 Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
 Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
 Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
 Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
 Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
 Librarian, Max-Planck-Institut, W. GERMANY