

49/5-13-92 g5 ②

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2836
UC-427

PPPL-2836

THIRD HARMONIC GENERATION WITH
ULTRA-HIGH INTENSITY LASER PULSES


BY

J.M. RAX AND N.J. FISCH

April, 1992

PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY



NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
Prices available from (615) 576-8401.

Available to the public from the:

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
703-487-4650

THIRD HARMONIC GENERATION WITH ULTRA-HIGH INTENSITY LASER PULSES

J.M. Rax* and N.J. Fisch

Princeton Plasma Physics Laboratory, Princeton University, Princeton NJ 08540

Abstract

When an intense, plane-polarized, laser pulse interacts with a plasma, the relativistic nonlinearities induce a third harmonic polarization. A phase-locked growth of a third harmonic wave can take place, but the difference between the nonlinear dispersion of the pump and driven waves leads to a rapid unlocking, resulting in a saturation. What becomes third harmonic amplitude oscillations are identified here, and the nonlinear phase velocity and the renormalized electron mass due to plasmon screening are calculated. A simple phase-matching scheme, based on a resonant density modulation, is then proposed and analyzed.

*Permanent address: Association EURATOM-CEA, CEN Cad. 13108 Saint-Paul-lez-Durance, FRANCE.

MASTER

Recent advances in pulse compression ¹ now make possible the exploration of laser-plasma interactions, at fluxes above 10^{18} W/cm². Harmonic generation is among the important new processes that take place at such high laser intensities, and has been recently investigated ² and identified as a promising candidate for a coherent light source at very short wavelength.

The nonlinear orbit of an electron in an intense, plane-polarized, laser pulse can be the current source of two different processes. (i) Spontaneous harmonic Compton scattering: ³ this process is incoherent, so that the emitted power scales as the density, i.e., as the square of the plasma frequency ω_p^2 . (ii) Collective forward harmonic Compton scattering: in a cold plasma the phases of the currents are fixed by the pump, so that a coherent harmonic wave, in phase with the nonlinear currents, can grow or decay. The efficiency of this phase-locked coherent harmonic generation was recently studied, ² and was shown to scale as ω_p^8 .

The main issue of harmonic generation in condensed and gaseous media is the phase velocity mismatch between the pump and the harmonic waves. ⁴ This is also the case for harmonic generation in a plasma, and the efficiency of the conversion of power to high harmonics is dramatically sensitive to this mismatch. What happens then, as we demonstrate, is that the harmonic wave does not really grow at all; rather, there are amplitude oscillations at a saturated level, scaling with ω_p^2 . Also, we demonstrate that, by modulating the density, linear growth can be accomplished with an efficiency scaling ω_p^4 or $\omega_p^{8/3}$.

Consider an intense pulse, such that the plasma period, ω_p^{-1} , is shorter than the pulse duration, $\delta\omega^{-1}$. In this regime, each electron is displaced in the direction of the pulse as the pulse passes it by. Then, after a transient response, a nonlinear oscillation, driven by the wave, and modulated by the plasma collective effects, is set up. To analyze this nonlinear response, we use a Lagrangian description of the plasma, rather than a Eulerian one. This method has proved to be powerful in studying the generation of beat waves ⁵ and plasma wakes. ⁶

The nonlinearity parameter of an intense electromagnetic wave, with vector potential A , is eA/mc , where c is the velocity of light, and $-e$ and m are respectively the electron charge and

mass. For ultra-intense waves, $eA/mc \geq 1$, and the electron quiver velocity becomes relativistic, so that the polarization currents saturate at the value $\epsilon_0 \omega_p^2 mc/e$. On the other hand, the displacement currents increase with A , as $\epsilon_0 \omega^2 A$, where $\omega/2\pi$ is the wave frequency. Because of the saturation of the polarization currents, the wave dynamics is dominated by nonlinearity, and ω_p^2/ω^2 can be used as an expansion parameter, with all orders in eA/mc kept. This is to be contrasted with the usual nonrelativistic plasma electrodynamics, which is essentially an expansion in eA/mc . The large value of the pump field causes electrons to respond with an effective mass, $^7 M$, so that, it turns out that a density expansion scheme is valid if $\omega_p^2/(\omega^2 M) \approx \omega_p^2/(\omega^2 eA/mc)$ is a small parameter. In fact, this parameter is small over a very broad range.

Intermediate, but important, results are the calculations of the nonlinear phase velocity of an intense laser wave and the electron renormalized effective mass, due to plasmon inertia.

To overcome the problems that we identify, we propose and analyze two phase-matching schemes, based on a resonant density modulation. In the following, except in the final part, we will use $e=m=c=\omega=1$.

Consider an intense, plane-polarized, laser wave propagating along the z axis:

$$A(z,t) = A(z,t) \cos[t - z + \phi(t)] \mathbf{e}_x, \quad (1)$$

where $\phi(t)$ is a slowly varying phase [$d\phi/dt = O(\omega_p^2)$], which accounts for the the nonlinear dispersion of the phase velocity, and where $A(z,t)$ is a slowly varying envelope, whose dynamics is insignificant to the problem, provided that $\partial A/\partial z < A$, $\partial A/\partial t < A$, and $\delta\omega < \omega_p$. Under such conditions, when an electron enters the pulse, it behaves essentially as in an infinite wave. The power transfer from the pump to the harmonic wave is diminished primarily by the phase-velocity mismatch, and, to a much lesser extent, by the group-velocity mismatch. This latter mismatch accounts for the imperfect overlapping of the two pulses, and it takes place on a far longer time scale. The effect of the finite length of the envelope, will be evaluated at the end of this letter.

Each electron is described by its unperturbed position z_0 , and follows a Lagrangian orbit, $h(z_0, t) = z(t) - z_0$, $x(z_0, t)$, about its rest position. To lowest order in ω_p^2 , when all collective plasma effects are neglected, the electrons perform the well known "figure-8" motion: ⁸

$$x = \frac{A}{M} \sin[M\tau(t, z_0) + \phi], \quad h = \frac{A^2}{8M^2} \sin[2M\tau(t, z_0) + 2\phi], \quad t = z_0 + M\tau + \frac{A^2}{8M^2} \sin[2M\tau + 2\phi]. \quad (2)$$

The proper time, τ , can be obtained from the last equation, which is a Kepler equation: $M\tau(t, z_0) = t - z_0 + \sum_{n=1}^{\infty} J_n[-nA^2/4M^2] n^{-1} \sin[2n(t - z_0) + 2n\phi]$, where the J_n are the ordinary Bessel functions of order n . The effective mass of the electron in an intense wave ⁷ is $M = \sqrt{1 + A^2/2}$.

The sum of all the Lagrangian currents, $-(dx/dt)\delta[z - z_0 - h(t, z_0)]$, gives the Eulerian current, which is the source term of the Maxwell's equations. With the Lorentz gauge we obtain

$$\frac{\partial^2 \mathbf{A}}{\partial z^2} - \frac{\partial^2 \mathbf{A}}{\partial t^2} = \omega_p^2 \int dz_0 \delta[z - z_0 - h(t, z_0)] \frac{\mathbf{A}(z, t)}{\gamma(z_0, t)} = \omega_p^2 \frac{\mathbf{A}}{\gamma(1 + \partial h / \partial z_0)}, \quad (3)$$

where we have introduced the relativistic energy, $\gamma^2 = 1 + (dh/d\tau)^2 + (dx/d\tau)^2$, and used the conservation of the transverse canonical momentum, $\gamma dx/dt = A$. After some algebra, we can rewrite, $\gamma(1 + \partial h / \partial z_0) = \gamma - p$, where p is the longitudinal momentum, $p = dh/d\tau$. The zero-order orbit, described by Eq. (2), gives $\gamma - p = M$, with the result that, although the microscopic Lagrangian currents contain the various harmonics of ω , the Eulerian current, to this order, contains only the fundamental.

The relativistic nonlinearity does manifest itself through an effective plasma frequency, ω_p^2/M . The nonlinear dispersion, described by the slowly varying phase, ϕ , is then easily calculated with Eqs. (1-3), to get $d\phi/dt = \omega_p^2/2M$. The nonlinear phase velocity of the pump, V^* , can be written as $V^* = 1 + \omega_p^2/2M$.

Because of the cancellation between the relativistic velocity anharmonicity, and the relativistic density oscillations, $\gamma(1 + \partial h / \partial z_0) = M$, harmonic generation occurs only at the order ω_p^4 , i.e., at the order ω_p^2 for the Lagrangian orbits. In this order, the Coulomb interactions, responsible for the plasma collective effects, enter the Lorentz equations. The transverse x dynamics remains

unaltered, but, on applying the Gauss theorem to the perturbed density, one finds an additional restoring force,⁹ proportional to the density, ω_p^2 , and the displacement, h .

$$\frac{dh}{d\tau} = p, \quad \frac{dp}{d\tau} = -\frac{A^2}{2} \sin[2(t-z)+2\phi] - \omega_p^2 \gamma h, \quad (4a)$$

$$\frac{dt}{d\tau} = \gamma, \quad \frac{d\gamma}{d\tau} = -\frac{A^2}{2} \sin[2(t-z)+2\phi] \left[1 + \frac{\omega_p^2}{2M}\right] - \omega_p^2 p h. \quad (4b)$$

These equations describe a perturbed nonlinear oscillator. To implement an ω_p^2 expansion scheme, we must be careful to avoid secular terms.¹⁰ On the basis of the unperturbed solution, Eq. (2), we seek a first order solution of the form:

$$h = \frac{A^2}{8M^{*2}} \sin[2M^*\tau+2\phi] + O[\omega_p^2], \quad t = z_0 + M^*\tau + \frac{A^2}{8M^{*2}} \sin[2M^*\tau+2\phi] + O[\omega_p^2]. \quad (5)$$

The plasma effects add up higher order harmonic terms, $O[\omega_p^2]$, and renormalize the nonlinear fundamental frequency M , to give a new effective mass, $M^*=M + O[\omega_p^2]$, dressed by plasmons.

Solving Eq. (4) with Eq. (5) leads to:

$$\gamma - p = M^* + \omega_p^2 \frac{A^2}{16M^{*2}} \cos[2M^*\tau+2\phi], \quad t - z = M^* \tau + \omega_p^2 \frac{A^2}{32M^{*3}} \sin[2M^*\tau+2\phi]. \quad (6)$$

The dressed effective mass is then obtained by demanding that there be no secular drift along the z direction. It is convenient to use $\gamma^2 - p^2 = 1 + (A \cos[t - z + \phi])^2$, and, after some algebra, we obtain:

$$M^* = M - \omega_p^2 \frac{A^4}{64 M^4}. \quad (7)$$

This last result is not specific to the problem of harmonic generation, and is, in fact, quite general. Equation (7) is the effective mass of an electron in an intense wave, when Coulomb collective interactions are taken into account to first order in ω_p^2 . The plasma collective effects decrease the bare effective mass M , because the collective forces are restoring forces, $-\omega_p^2 h$, which oppose the driving fast oscillations, and screen A .

On the basis of Eqs. (3) and (6), we see that the transverse Eulerian polarization current has a third harmonic component, of order ω_p^4 , which is able to excite a third harmonic transverse wave,

$$\mathbf{a}(z,t) = a(t) \cos[3(t-z)+\phi(t)] \mathbf{e}_x, \quad (8)$$

with the amplitude, a , and phase, ϕ , evolving on the slow time scale of the problem, i.e. at most $O(\omega_p^2)$. To order ω_p^4 , and provided that $a < A$, the coupled Maxwell's equations are:

$$\frac{\partial^2 A}{\partial z^2} - \frac{\partial^2 A}{\partial t^2} = \frac{\omega_p^2}{M^*} A - \frac{\omega_p^4 A^2}{32M^{*4}} A, \quad \frac{\partial^2 a}{\partial z^2} - \frac{\partial^2 a}{\partial t^2} = \frac{\omega_p^2}{M^*} a - \frac{\omega_p^4 A^2}{32M^{*4}} A \cos[3(t-z)+3\phi(t)] e_x. \quad (9)$$

The first term, on the right hand side of the second equation, describes the reactive dispersion due to the polarization currents, but, with dressed electrons, whose inertia is given by Eq. (7). The second term drives the harmonic generation. On the basis of the first equation, one can calculate the next order plasma correction to the nonlinear dispersion of A , $2d\phi/dt = \omega_p^2/M^* - \omega_p^4 A^2/32M^{*4}$; thus, to this order, the nonlinear phase velocity of the pump is

$$V^* = 1 + \frac{\omega_p^2}{2M} - \frac{\omega_p^4 A^2}{64M^4} + \frac{\omega_p^4 A^4}{256M^6}. \quad (10)$$

The equations for the slowly varying amplitude and phase, a and ϕ , take a form similar to the Rosenbluth-Liu⁵ equations for beat wave generation, up to a dephasing term,

$$\frac{da}{dt} = -\omega_p^4 \frac{A^3}{192M^4} \sin(\theta) \quad (11 a)$$

$$\frac{d\theta}{dt} = -\frac{4\omega_p^2}{3M} - \omega_p^4 \frac{A^3}{192M^4} \frac{\cos(\theta)}{a} \quad (11 b)$$

where $\theta(t) = \phi(t) - 3\phi(t)$. The tendency to phase-lock at large a , due to the second term on the right hand side of Eq. (11b), is cancelled by the phase-velocity mismatch, described by the first term. No phase-locking occurs, so that instead of growing linearly with time, the amplitude oscillates.

Figure 1 depicts the phase portrait of this dynamical system, Eq. (11), with various orbits arranged around the elliptic points, $[a = \pm \omega_p^2 A^3 / 256M^3, \theta = n\pi]$. These equations are a Pfaffian system, and the first integral is: $I = a^2 + \omega_p^2 \frac{A^3}{128M^3} a \cos(\theta)$. Two classes of orbits are easily identified: circulating one ($I > 0$, for large $|a|$) and trapped one ($I < 0$, for small $|a|$). The equation of both types of orbits is: $[Ut] = -[\text{Arcsin} \frac{2UI - U^2 a^2 + 2V^2}{2V\sqrt{V^2 + 2UI}}]$, where $U = -4\omega_p^2/3M$, and $V = -\omega_p^4 A^3 / 192M^4$. If the amplitude is large enough, the $I > 0$ orbits can be approximated by $a = [-V/U] \cos(Ut)$.

The previous poor conversion to third harmonic can be improved upon through a resonant density modulation, in order to detrap the $I=0$ orbit, as one proceed to show.

Imagine a one dimensional plasma media, with alternating, along z , high and low density sections. The laser pulse will induce harmonic generation in the active high density sections, but the interaction with the low density plasma will result only in a reactive phase shift between the pump and the harmonic. Choosing the width of the reactive low density sections to compensate the phase mismatch due to the active one, now restores linear growth. The bold line on Fig. 2 displays the amplitude-phase orbit in such a configuration. After s steps, one can reach $a \approx s\omega_p^2 A^3/128M^3$. Thus, the power conversion efficiency scales as, $P_3/P_1 \approx 10^{-3}s^2(eA/mc)^\alpha(\omega_p/\omega)^4$, where the α exponent is 4, if $A < 1$, 0, if $A \approx 1$, and -2, if $A > 1$.

Strong density modulations might be set up by, for example, the laser ablation of a multiple-layered media, or a nonlinear wave. However, such a strong resonant modulation, may not be necessary, it would be more convenient to use a low frequency, small amplitude, density wave in an homogeneous plasma, such as, for example, a long wavelength ion-acoustic wave.

Actually, even a weak modulation has a dramatic effect which can be studied within the framework of the previous model. Consider the density modulation, $\delta n/n = \epsilon \sin(\Omega t)$, with $\epsilon < 1$, and $\Omega < \delta\omega < \omega_p < \omega$, so that the Eq. (11) becomes:

$$\frac{da}{dt} = -\frac{\omega_p^4 A^3}{192M^4} [1 + 2\epsilon \sin(\Omega t)] \sin(\theta) \quad (12 a)$$

$$\frac{d\theta}{dt} = -\frac{4\omega_p^2}{3M} [1 + \epsilon \sin(\Omega t)] - \frac{\omega_p^4 A^3}{192M^4} [1 + 2\epsilon \sin(\Omega t)] \frac{\cos(\theta)}{a}. \quad (12 b)$$

As soon as the amplitude becomes larger than $\omega_p^2/256$, the second term on the right hand side of Eq. (12b) is negligible compared to the first one, and the amplitude equation becomes:

$$\frac{da}{dt} = -\frac{\omega_p^4 A^3}{192M^4} [1 + 2\epsilon \sin(\Omega t)] \sum_N J_N(-4\epsilon\omega_p^2/3M\Omega) \sin \left[\frac{4\omega_p^2}{3M} t + N\Omega t + N\frac{\pi}{2} \right]. \quad (13)$$

The amplitude is driven by a sum of oscillating terms, and, if one of these oscillations is resonant, it induces a secular linear growth. When N is odd the resonance condition is

$$N\Omega + 4\omega_p^2/3M = 0. \quad (14)$$

The associated resonant term quickly dominates the other bounded components, and we can average out the oscillating part of a , to study the secular part of the third harmonic amplitude, $\langle a \rangle$. Taking the asymptotic expansion of $J_N(N\epsilon)$ in the resulting equation, $d\langle a \rangle/dt = J_N(N\epsilon)\omega_p^4 A^3/192M^4$, we obtain

$$\langle a \rangle = 6 \cdot 10^{-3} A^3 M^{-11/3} \omega_p^{10/3} \epsilon^{-1/3} \Omega^{1/3} \text{Ai}[1.52 M^{-2/3} \omega_p^{4/3} \epsilon^{-1/3} \Omega^{-2/3}(1-\epsilon)] t. \quad (15)$$

Ai is the Airy function, and one can take the typical value $\epsilon^{-1/3} \text{Ai} \approx 0.1$. The efficiency scaling becomes $P_3/P_1 \approx 10^{-5} (eA/mc)^\alpha (\omega_p/\omega)^{20/3} (\Omega/\omega)^{2/3} (\omega t)^2$. The exponent α is now $-10/3$ if $A > 1$.

So far, we have solved a nonlinear initial value problem, i.e., an infinite plane wave, $A(t)$, is turned on adiabatically in an infinite plasma, and we have studied the associated response, $a(t)$. This solution is relevant to the corresponding initial boundary value problem, i.e., the study of the $a(z,t)$ response to an $A(z,t)$ pump, provided that the wave-packet overlapping problem, due to group velocity mismatch, takes place on a long time scale. A departure from overlapping appears after a time $c\delta\omega^{-1}/2\omega(\partial v_g/\partial\omega) \approx \delta\omega^{-1}\omega^2/\omega_p^2$. This time is to be compared with the time needed to complete one generation cycle $\approx \omega/\omega_p^2$. Since $\omega \gg \delta\omega$ for a wave packet, the overlapping mismatch comes into play on a time scale far longer than all the other processes.

Nevertheless, this overlapping does limit the maximum number of step in the strong modulation scheme, as well as the time in Eq. (15) ($t_{\max} = \delta\omega^{-1}\omega^2/\omega_p^2$).

To assess more carefully the potential of phase-matching with a small amplitude wave, consider an ion acoustic wave. The quasineutral, low frequency, long-wavelength dispersion relation for a wave with $\delta n/n = \epsilon \sin(\Omega_s t - K_s z)$ is $\Omega_s = K_s \sqrt{T/m_i}$, T is the electron temperature and m_i is the ion mass. Under typical laboratory conditions, $\Omega_s < \omega_{pi} < \delta\omega < \omega_{pe} < \omega$, the pulse length is smaller than the ion acoustic wavelength, so that, Eq. (12) apply, provided that we

use the effective modulation frequency, $\Omega = \Omega_s v_g / \sqrt{T/m_i}$, seen by the pulse. We do not need to know the exact expression of v_g , the nonlinear group velocity of the pulse, because $v_g = 1 + O[(\omega_p/\omega)^2]$ and we are working to the lowest relevant order in ω_p/ω . Thus the equivalent frequency to be used in Eq. (12) is $\Omega = \Omega_s / \sqrt{T/m_i}$. Taking the group velocity mismatch as the ultimate limitation, and considering the regime $A=1$, we obtain: $P_3/P_1 \approx 10^{-2} (T/1.eV)^{1/3} (\omega_p/\omega)^{8/3} (\Omega_s/\omega)^{2/3} (\omega/\delta\omega)^2$.

To summarize, we have set up, discussed, and solved, the equations for relativistic third harmonic generation in a plasma within the framework of a Lagrangian description. The important problem of phase mismatch has been identified and addressed. The nonlinear phase velocity, the renormalized electron mass, and the conversion efficiency have been calculated. Two simple modulation schemes, to overcome saturation, have been proposed and analyzed.

Using plasma for third harmonic generation has advantages over using other nonlinear condensed, or gaseous media. Plasma do not suffer material breakdown at high intensities, and can convert radiation over a very broad range of frequencies.

Acknowledgments: We gratefully acknowledge useful discussions with E. Valeo, P. Sprangle, J. Dawson, T. Johnston, and W. Mori. This work was supported by the U.S Department of Energy under contract number DE-AC02-76-CHO3073.

References

- ¹M. Pessot, J. Squier, G. Mourou and D.J. Harter, *Opt. Lett.* **14**, 797 (1987). M. Pessot, J. Squier, P. Bado, G. Mourou and D.J. Harter, *IEEE J. Quantum Electron.* **QE-25**, 61 (1989).
- ²P. Sprangle, E. Esarey and A. Ting, *Phys. Rev. Lett.* **64**, 2011 (1990). P. Sprangle, E. Esarey and A. Ting, *Phys. Rev. A* **41**, 4463 (1990). A. Ting, E. Esarey, and P. Sprangle, *Phys. Fluids. B* **2**, 1390 (1990).
- ³E.S. Sarachik and G.T. Schappert, *Phys. Rev. D* **10**, 2738 (1970). J.E. Gunn, and J.P. Ostriker, *Astro. J.*, **165**, 523 (1971). R.E. Waltz and O.P. Manley *Phys. Fluids* **21**, 808 (1978).
- ⁴N. Bloembergen, *Nonlinear Optics* (Benjamin, New York, 1965).
- ⁵M.N. Rosenbluth and C.S. Liu, *Phys. Rev. Lett.* **29**, 701 (1972).
- ⁶J.M. Rax and N.J. Fisch, *Phys. Fluids B* **4** (to appear, 1992).
- ⁷T.W. Kibble, *Phys. Rev.* **138**, 740 (1965). J.H. Eberly, and H.R. Reiss *Phys. Rev.* **145**, 1035 (1966). T.W. Kibble, *Phys. Rev.* **150**, 1060 (1966).
- ⁸L.D. Landau and E. M. Lifshitz, *The Classical Theory of Fields* (Pergamon, Oxford, 1975).
- ⁹J.M. Dawson, *Phys. Rev.* **113**, 383 (1958). R.C. Davidson and P.P. Schram *Nuclear Fusion*, **8**, 183 (1968).
- ¹⁰N. Minorsky, *Non Linear Oscillations* (Van Nostrand, Princeton, 1962).

Figure captions

Fig. 1: Phase portrait of Eq. (11), $eA/mc=1$, $(\omega_p/\omega)^2=0.1$, starting from the background noise, near $[a=0, \theta=(2n+1)\pi/2]$, the third harmonic wave describes the separatrix orbit, $I=0$, and the amplitude oscillates between the values $\pm\omega_p^2 A^3/128M^3$, the $I>0$ open orbits oscillate within the range $\omega_p^2 A^3/128M^3$.

Fig. 2: Amplitude-phase orbit (bold line) in a resonant stack of high and low density plasma layers, the horizontal straight lines, which are, in fact, a very small amplitude oscillations, correspond to passive phase shifting. The length of the active high density sections is to be of the order of an odd multiple of $3M\pi/8\omega_p^2$.




Figure 1




Figure 2

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA
 Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA
 Plasma Research Lab., Australian Nat. Univ., AUSTRALIA
 Prof. I.R. Jones, Flinders Univ, AUSTRALIA
 Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA
 Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA
 Prof. M. Goossens, Astronomisch Instituut, BELGIUM
 Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM
 Commission-Européenne, DG. XII-Fusion Prog., BELGIUM
 Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM
 Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL
 Instituto Nacional De Pesquisas Especiais-INPE, BRAZIL
 Documents Office, Atomic Energy of Canada Ltd., CANADA
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
 Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA
 Prof. J. Teichmann, Univ. of Montreal, CANADA
 Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA
 Prof. T.W. Johnston, INRS-Energie, CANADA
 Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA
 Dr. C.R. James, Univ. of Alberta, CANADA
 Dr. P. Lukáč, Komenského Univerzita, CZECHO-SLOVAKIA
 The Librarian, Culham Laboratory, ENGLAND
 Library, R61, Rutherford Appleton Laboratory, ENGLAND
 Mrs. S.A. Hutchinson, JET Library, ENGLAND
 Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS
 P. Mähönen, Univ. of Helsinki, FINLAND
 Prof. M.N. Bussac, Ecole Polytechnique, FRANCE
 C. Moutet, Lab. de Physique des Milieux Ionisés, FRANCE
 J. Radet, CEN/CADARACHE - Bat 506, FRANCE
 Prof. E. Economou, Univ. of Crete, GREECE
 Ms. C. Rinni, Univ. of Ioannina, GREECE
 Dr. T. Mui, Academy Bibliographic Ser., HONG KONG
 Preprint Library, Hungarian Academy of Sci., HUNGARY
 Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA
 Dr. P. Kaw, Inst. for Plasma Research, INDIA
 Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL
 Librarian, International Center for Theoretical Physics, ITALY
 Miss C. De Palo, Associazione EURATOM-ENEA, ITALY
 Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY
 Prof. G. Rostagni, Istituto Gas Ionizzati Del Cnr, ITALY
 Dr. H. Yamato, Toshiba Res & Devel Center, JAPAN
 Prof. I. Kawakami, Hiroshima Univ., JAPAN
 Prof. K. Nishikawa, Hiroshima Univ., JAPAN
 Director, Japan Atomic Energy Research Inst, JAPAN
 Prof. S. Itoh, Kyushu Univ., JAPAN
 Research Info. Ctr., National Inst. for Fusion Science, JAPAN
 Prof. S. Tanaka, Kyoto Univ., JAPAN
 Library, Kyoto Univ., JAPAN
 Prof. N. Inoue, Univ. of Tokyo, JAPAN
 Secretary, Plasma Section, Electrotechnical Lab., JAPAN
 S. Mori, Technical Advisor, JAERI, JAPAN
 Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
 J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
 D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
 Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
 Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
 Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
 Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
 Z. Li, S.W. Inst Physics, PEOPLE'S REPUBLIC OF CHINA
 Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
 Dr. O. Petrus, AL I GUZA Univ., ROMANIA
 Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
 Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
 Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
 Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
 L. Stenflo, Univ. of UMEA, SWEDEN
 Librarian, Royal Inst. of Technology, SWEDEN
 Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
 Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
 Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
 Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
 Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. Apparatus, USSR
 Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
 Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
 Librarian, The Ukr.SSR Academy of Sciences, USSR
 Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
 Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
 Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
 Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
 Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
 Librarian, Max-Planck-Institut, W. GERMANY
 Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA

END

**DATE
FILMED**

6/12/92

