



ORNL/RASA-91/20

**OAK RIDGE  
NATIONAL  
LABORATORY**

**MARTIN MARIETTA**

**Radiological Survey Results at  
13 Congress Street,  
Beverly, Massachusetts  
(VB003)**

**R. D. Foley  
M. S. Uziel**

**MANAGED BY  
MARTIN MARIETTA ENERGY SYSTEMS, INC.  
FOR THE UNITED STATES  
DEPARTMENT OF ENERGY**

HEALTH AND SAFETY RESEARCH DIVISION

Environmental Restoration and Waste Management Non-Defense Programs  
(Activity No. EX 20 20 01 0; ADS3170000)

**Radiological Survey Results at 13 Congress Street,  
Beverly, Massachusetts (VB003)**

R. D. Foley and M. S. Uziel

Date Issued - July 1992

Investigation Team

R. E. Swaja - Measurement Applications and Development Manager  
W. D. Cottrell - FUSRAP Project Director  
R. D. Foley - Survey Team Leader

Survey Team Members

J. F. Allred	M. E. Murray
A. C. Butler*	V. P. Patania
R. D. Foley	D. E. Rice
D. Mackenzie†	D. A. Rose
W. H. Shinpaugh*	

\*D. R. Stone and Associates, Inc.

†H&R Technical Associates

Work performed by the  
MEASUREMENT APPLICATIONS AND DEVELOPMENT GROUP

Prepared by the  
OAK RIDGE NATIONAL LABORATORY  
Oak Ridge, Tennessee 37831-6285  
managed by  
MARTIN MARIETTA ENERGY SYSTEMS, INC.  
for the  
U. S. DEPARTMENT OF ENERGY  
under contract DE-AC05-84OR21400

## CONTENTS

LIST OF FIGURES .....	v
LIST OF TABLES .....	vii
ACKNOWLEDGMENTS .....	ix
ABSTRACT .....	xi
INTRODUCTION .....	1
SURVEY METHODS .....	2
SURFACE RADIATION MEASUREMENTS .....	2
SOIL SAMPLING AND ANALYSES .....	2
SURVEY RESULTS .....	3
SURFACE RADIATION MEASUREMENTS .....	3
SOIL SAMPLES .....	3
SIGNIFICANCE OF FINDINGS .....	4
REFERENCES .....	4

## LIST OF FIGURES

1	Diagram showing general location of the former Ventron site .....	5
2	Diagram showing location of 13 Congress Street, Beverly, Massachusetts, in relation to the former Ventron site .....	6
3	Diagram of the property located at 13 Congress Street, Beverly, Massachusetts .....	7
4	View looking north at the house at 13 Congress Street, Beverly, Massachusetts .....	8
5	View looking south at the house at 13 Congress Street, Beverly, Massachusetts .....	8
6	Diagram showing surface gamma exposure rates and soil sampling locations at 13 Congress Street, Beverly, Massachusetts .....	9

## LIST OF TABLES

1	Applicable guidelines for protection against radiation .....	10
2	Background radiation levels and concentrations of selected radionuclides in soil in the Beverly, Massachusetts, area .....	11
3	Concentrations of radionuclides in soil samples from 13 Congress Street, Beverly, Massachusetts .....	12

## ACKNOWLEDGMENTS

This project was sponsored by the Office of Environmental Restoration, U.S. Department of Energy, under contract DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc. The authors wish to acknowledge the contributions of R. F. Carrier, W. D. Cottrell, C. A. Johnson, V. P. Patania, and D. A. Roberts of the Measurement Applications and Development (MAD) Group, Oak Ridge National Laboratory, for participation in the analyses, editing, and reporting of data for this survey. The authors also wish to thank other members of the MAD Group: J. F. Allred and D. A. Rose for sample preparation and T. R. Stewart for computer graphics.

## ABSTRACT

At the request of the U.S. Department of Energy (DOE), a team from Oak Ridge National Laboratory conducted a radiological survey at 13 Congress Street, Beverly, Massachusetts. The survey was performed in May 1991. The purpose of the survey was to determine if uranium from work performed under government contract at the former Ventron facility had migrated off-site to neighboring areas. The survey included a surface gamma scan, a beta-gamma scan of paved areas, and the collection of soil samples for radionuclide analyses.

Results of the survey demonstrated no radionuclide concentrations or radiation measurements in excess of the DOE Formerly Utilized Sites Remedial Action Program guidelines.

# Radiological Survey Results at 13 Congress Street, Beverly, Massachusetts (VB003)\*

## INTRODUCTION

The Metal Hydrides Corporation in Beverly, Massachusetts (which became the Ventron Corporation in 1965), was one of many companies performing work during the 1940s associated with the development of nuclear energy for defense-related projects under contract to the Manhattan Engineer District (MED) and the Atomic Energy Commission (AEC). Operations conducted under government contract at such sites included the procurement, storage, and processing of uranium oxides, salts, and metals, and the subsequent machining of these products. As a result of activities involving these materials, equipment, buildings, and land at some of the sites became radiologically contaminated with small amounts of material resulting in low levels of contamination on the properties. At contract termination, release limits and decontamination operations were typically applied in conformance with standards currently deemed adequate for purposes of health and environmental protection. Subsequent to original assessments and the release of these facilities, new research and information have resulted in the development of more stringent guidelines for release of such facilities for unrestricted use. Furthermore, in some instances, documentation is limited or nonexistent, and conditions at a specific site may be unknown. It is the policy of the U.S. Department of Energy (DOE) to verify that radiological conditions at such facilities comply with existing guidelines.<sup>1</sup> The Formerly Utilized Sites Remedial Action Program (FUSRAP) was established in 1974 to assist in assessment and cleanup activities at these sites.

The radiological survey detailed in this report was performed under the FUSRAP program and is one of several surveys conducted in May 1991 on properties in the vicinity of the former Ventron facility by members of the Oak Ridge National Laboratory (ORNL) at the request of DOE. The city of Beverly lies on Beverly Harbor ~15 miles northeast of the central Boston area. The former Ventron facility, now owned by Morton International, is located at the confluence of the Bass and Danvers rivers on Congress Street near the Beverly-Salem bridge (Fig. 1, p. 5).<sup>2</sup>

From 1942 to 1948 the Metal Hydrides Corporation (later to become the Ventron facility) converted uranium oxide to uranium metal powder at the facility under contract to MED in support of the war effort. As better methods for production of uranium metal were developed, Metal Hydrides shifted its operations toward recovering uranium from scrap uranium and turnings from the slug fabrication plant at Hanford, Washington.<sup>1</sup> Contracts between Metal Hydrides and the government were completed in 1954.

---

\*The survey was performed by members of the Measurement Applications and Development Group of the Health and Safety Research Division at Oak Ridge National Laboratory under DOE contract DE-AC05-84OR21400.

Following a radiological screening survey at the site in 1977, a comprehensive survey was performed in 1982.<sup>2</sup> In 1987 DOE contractors removed the uranium-contaminated roof from a Ventron building, which had begun to leak. Radioactive materials remaining on the site do not pose a health hazard under present-use conditions but could cause radiation exposure to workers if excavation or major renovation took place on the property. DOE plans a complete characterization study of the site in 1992 and the initiation of remedial action soon thereafter.

The surveys of the property reported in this document and of other surrounding properties are part of DOE's continuing effort to assess the former Ventron site and plan for remedial action. The objective of the surveys was to determine if uranium from plant operations had migrated off-site to neighboring areas including Beverly Harbor and, if so, to what degree. The location of this vicinity property in relation to the former Ventron site is shown in Fig. 2 (p. 6). The property at 13 Congress Street includes a house and a small building to the north (Fig. 3, p. 7). Most of the property surface is covered with grass or asphalt with concrete bordering the south, east, and northeast sides of the house. Other features include a stone and concrete wall just north of the house, a rock wall at the northern boundary of the property, and a stone retaining wall along part of the western property boundary. The radiological survey included measurements of radiation levels over the surface of the property and analysis of soil samples for the presence of radionuclides.

## SURVEY METHODS

A comprehensive description of the survey methods and instrumentation used in this survey is given in *Procedures Manual for the ORNL Radiological Survey Activities (RSA) Program*, ORNL/TM-8600 (April 1987).<sup>3</sup>

### SURFACE RADIATION MEASUREMENTS

Gamma radiation levels were determined using a portable NaI gamma scintillation meter. Because NaI gamma scintillators are energy dependent, measurements of gamma radiation levels in counts per minute are normalized to pressurized ionization chamber (PIC) measurements to estimate gamma exposure rates in  $\mu\text{R/h}$ . Using a Geiger-Mueller pancake detector, beta-gamma radiation levels in cpm were measured over selected paved and structural surfaces and then converted to mrad/h.

### SOIL SAMPLING AND ANALYSES

Surface and subsurface soil samples were systematically collected over the property in a pattern sufficient to obtain a characterization of the radionuclide content of the soil. All soil samples were analyzed to determine concentrations of  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{226}\text{Ra}$ .

## SURVEY RESULTS

DOE guidelines are summarized in Table 1 (p. 10). Typical background radiation levels for the Beverly, Massachusetts, area are presented in Table 2 (p. 11). These data are provided for comparison with survey results presented in this section. All direct measurement results presented in this report are gross readings; background radiation levels have not been subtracted. Similarly, background concentrations have not been subtracted from radionuclide concentrations measured in soil samples.

Photographs taken in May 1991 of the property at 13 Congress Street, Beverly, Massachusetts, are shown in Figs. 4 and 5 (p. 8).

### SURFACE RADIATION MEASUREMENTS

Results of the surface gamma scan over the property surface are shown in Fig. 6 (p. 9). Surface gamma exposure rates averaged 8 and 9  $\mu\text{R/h}$  over the areas covered with grass and asphalt and 10  $\mu\text{R/h}$  near the stone retaining wall along the western boundary of the property. Blocks of granite bordering the south edge of the asphalt sidewalk (see Fig. 4, p. 8) also measured 10  $\mu\text{R/h}$ . These values are comparable to typical background measurements in the Beverly, Massachusetts, area (6 to 9  $\mu\text{R/h}$ , Table 2, p. 11). Slightly higher exposure rates are frequently associated with structural materials such as granite or brick, which inherently contain slightly elevated concentrations of naturally occurring radionuclides.

Beta-gamma dose rates ranging from 0.04 to 0.06 mrad/h were measured over selected hard surfaces on the property. All measurements were comparable to background levels measured in the vicinity.

### SOIL SAMPLES

Soil sample locations are shown in Fig. 6 (p. 9), and results of analyses are listed in Table 3 (p. 12). Concentrations of  $^{226}\text{Ra}$  and  $^{232}\text{Th}$  in surface soil (0-15 cm) ranged from 0.77 to 1.4 pCi/g and from 0.75 to 1.4 pCi/g, respectively. Concentrations of  $^{226}\text{Ra}$  in subsurface soil (15-30 cm) ranged from 0.67 to 1.6;  $^{232}\text{Th}$  ranged from 0.65 to 1.6 pCi/g in subsurface soil. These values are slightly above typical background levels in the Beverly area (Table 2, p. 11) but well below DOE guidelines (Table 1, p. 10). Uranium-238 concentrations in surface soil ranged from 2.8 to 11 pCi/g. Subsurface  $^{238}\text{U}$  concentrations ranged from 1.3 to 5.3 pCi/g. Uranium-238 levels in most of the soil samples from 13 Congress Street were above typical background levels in the Beverly area (Table 2, p. 11); however, all samples were well below guidelines of 35 to 40 pCi/g that have been applied at other FUSRAP sites (Table 1, p. 10).

## SIGNIFICANCE OF FINDINGS

The results of the radiological survey at 13 Congress Street, Beverly, Massachusetts, demonstrated no radionuclide concentrations or radiation measurements above established DOE guidelines.

## REFERENCES

1. U. S. Department of Energy, *A Background Report for the Formerly Utilized Manhattan Engineer District/Atomic Energy Commission Sites Program*, DOE/EV-0097, September 1980.
2. W. D. Cottrell and R. F. Carrier, *Results of the Radiological Survey at the Ventron Site, Beverly, Massachusetts*, ORNL/TM-10053, Martin Marietta Energy Systems, Inc., Oak Ridge Natl. Lab., May 1988.
3. T. E. Myrick, B. A. Berven, W. D. Cottrell, W. A. Goldsmith, and F. F. Haywood, *Procedures Manual for the ORNL Radiological Survey Activities (RASA) Program*, ORNL/TM-8600, Martin Marietta Energy Systems, Inc., Oak Ridge Natl. Lab., April 1987.




Fig. 1. Diagram showing general location of the former Ventron site.






Fig. 2. Diagram showing location of 13 Congress Street, Beverly, Massachusetts, in relation to the former Ventrone site.




**Fig. 3. Diagram of the property located at 13 Congress Street, Beverly, Massachusetts.**



**Fig. 4. View looking north at the house at 13 Congress Street, Beverly, Massachusetts.**



**Fig. 5. View looking south at the house at 13 Congress Street, Beverly, Massachusetts.**



**Fig. 6. Diagram showing surface gamma exposure rates and soil sampling locations at 13 Congress Street, Beverly, Massachusetts.**

**Table 1. Applicable guidelines for protection against radiation**  
(Limits for uncontrolled areas)

Mode of exposure	Exposure conditions	Guideline value
Gamma radiation	Indoor gamma radiation level (above background)	20 $\mu\text{R}/\text{h}^a$
Radionuclide concentrations in soil (generic)	Maximum permissible concentration of the following radionuclides in soil above background levels, averaged over a 100-m <sup>2</sup> area <sup>226</sup> Ra <sup>232</sup> Th <sup>230</sup> Th	5 pCi/g averaged over the first 15 cm of soil below the surface; 15 pCi/g when averaged over 15-cm-thick soil layers more than 15 cm below the surface
Derived concentrations	<sup>238</sup> U	Site specific <sup>b</sup>
Guideline for non-homogeneous contamination (used in addition to the 100-m <sup>2</sup> guideline) <sup>c</sup>	Applicable to locations with an area $\leq 25$ m <sup>2</sup> , with significantly elevated concentrations of radionuclides ("hot spots")	$G_A = G_i(100/A)^{1/2}$ , where $G_A$ = guideline for "hot spot" of area (A) $G_i$ = guideline averaged over a 100-m <sup>2</sup> area

<sup>a</sup>The 20  $\mu\text{R}/\text{h}$  shall comply with the basic dose limit (100 mrem/year) when an appropriate-use scenario is considered.

<sup>b</sup>DOE guidelines for uranium are derived on a site-specific basis. Guidelines of 35–40 pCi/g have been applied at other FUSRAP sites. *Source*: J. L. Marley and R. F. Carrier, *Results of the Radiological Survey at 4 Elmhurst Avenue, Colonie, New York (AL219)*, ORNL/RASA-87/117, Martin Marietta Energy Systems, Inc., Oak Ridge Natl. Lab., February 1988; B. A. Berven et al., *Radiological Survey of the Former Kellex Research Facility, Jersey City, New Jersey*, DOE/EV-0005/29, ORNL-5734, Martin Marietta Energy Systems, Inc., Oak Ridge Natl. Lab., February 1982.

<sup>c</sup>DOE guidelines specify that every reasonable effort shall be made to identify and to remove any source that has a concentration exceeding 30 times the guideline value, irrespective of area (adapted from *Revised Guidelines for Residual Radioactive Material at FUSRAP and Remote SFMP Sites*, April 1987).

*Sources*: Adapted from U.S. Department of Energy, DOE Order 5400.5, April 1990, and U.S. Department of Energy, *Guidelines for Residual Radioactive Material at Formerly Utilized Sites Remedial Action Program and Remote Surplus Facilities Management Program Sites*, Rev. 2, March 1987.

**Table 2. Background radiation levels and concentrations of selected radionuclides in soil in the Beverly, Massachusetts, area**

Type of radiation measurement or sample	Radiation level or radionuclide concentration	
	Range	Average
Gamma exposure rate at ground surface ( $\mu\text{R/h}$ ) <sup>a</sup>	6-9	7
Concentration of radionuclides in soil (pCi/g) <sup>a</sup>		
<sup>238</sup> U	0.69-1.05	0.89
<sup>226</sup> Ra	0.70-0.94	0.84
<sup>232</sup> Th	0.76-0.94	0.84

<sup>a</sup>Values obtained from three locations in the Beverly area.

**Table 3. Concentrations of radionuclides in soil samples from  
13 Congress Street, Beverly, Massachusetts**

Sample <sup>a</sup>	Depth (cm)	Radionuclide concentration (pCi/g) <sup>b</sup>		
		<sup>226</sup> Ra	<sup>232</sup> Th	<sup>238</sup> U
<i>Systematic samples<sup>c</sup></i>				
S1A	0-15	1.4 ±0.04	1.4 ±0.07	11 ±0.8
S1B	15-30	0.87±0.03	0.95±0.06	5.3 ±1
S2A	0-15	1.0 ±0.05	0.96±0.08	8.6 ±1
S2B	15-30	0.96±0.02	0.94±0.04	4.6 ±1
S3A	0-15	0.77±0.02	0.75±0.04	2.8 ±0.8
S3B	15-30	0.67±0.01	0.65±0.03	1.3 ±0.5
S4A	0-15	0.83±0.04	0.79±0.07	5.8 ±0.6
S4B	15-30	0.94±0.02	1.0 ±0.03	1.9 ±0.6
S5A	0-15	1.1 ±0.04	1.3 ±0.06	6.5 ±2
S5B	15-30	1.6 ±0.03	1.6 ±0.04	2.6 ±0.6

<sup>a</sup>Sample locations are shown on Fig. 6.

<sup>b</sup>Indicated counting error is at the 95% confidence level ( $\pm 2\sigma$ ).

<sup>c</sup>Systematic samples are taken at locations irrespective of gamma exposure rates.

### INTERNAL DISTRIBUTION

- |                       |                               |
|-----------------------|-------------------------------|
| 1. B. A. Berven       | 23. R. E. Rodriguez           |
| 2. K. J. Brown        | 24. P. S. Rohwer              |
| 3. R. F. Carrier      | 25-27. R. E. Swaja            |
| 4-9. W. D. Cottrell   | 28-30. M. S. Uziel            |
| 10-15. R. D. Foley    | 31. J. K. Williams            |
| 16. R. O. Hultgren    | 32-37. MAD Records Center     |
| 17. C. A. Johnson     | 38. Central Research Library  |
| 18. S. V. Kaye        | 39. ORNL Technical Lib., Y-12 |
| 19. A. P. Malinauskas | 40-41. Laboratory Records     |
| 20. M. E. Murray      | 42. Laboratory Records – RC   |
| 21. P. T. Owen        | 43. ORNL Patent Section       |
| 22. D. A. Roberts     |                               |

### EXTERNAL DISTRIBUTION

44. J. D. Berger, Oak Ridge Associated Universities, E/SH Division, Environmental Survey and Site Assessment Program, P.O. Box 117, Oak Ridge, TN 37831-0117
45. J. J. Fiore, Director, Office of Eastern Area Programs, Office of Environmental Restoration, U.S. Department of Energy, (EM-42), Trevion II, Washington, DC 20585
- 46-48. G. K. Hovey, Bechtel National, Inc., FUSRAP Department, Oak Ridge Corporate Center, 151 Lafayette Drive, P.O. Box 350, Oak Ridge, TN 37831-0350
49. L. K. Price, Director, Former Sites Restoration Division, Oak Ridge Field Office, U.S. Department of Energy, P.O. Box 2001, Oak Ridge, TN 37831-8723
50. J. W. Wagoner II, Director, Division of Off-Site Programs, Office of Eastern Area Programs, Office of Environmental Restoration, U.S. Department of Energy, (EM-421), Trevion II, Washington, DC 20585
- 51-55. W. A. Williams, Designation and Certification Manager, Division of Off-Site Programs, Office of Eastern Area Programs, Office of Environmental Restoration, U.S. Department of Energy, (EM-421), Trevion II, Washington, DC 20585
56. C. D. Young, Office of Technical Services, Roy F. Weston, Inc., 12800 Middlebrook Road, Suite 207, Germantown, MD 20874
- 57-61. Office of Scientific and Technical Information, U.S. Department of Energy, P.O. Box 62, Oak Ridge, TN 37831

**END**

**DATE  
FILMED**

**9/16/92**

