

MULTI-DEVICE SCALING OF NEOCLASSICAL TEARING MODE ONSET WITH BETA

by

R.J. LA HAYE, R.J. BUTTERY,* S. GUENTER,†
G.T.A. HUYSMANS,‡ and H.R. WILSON*

This is a preprint of a paper presented at the Twenty-Sixth European Physical Society Conference on Controlled Fusion and Plasma Physics, June 14-18, 1999 in Maastricht, The Netherlands, and to be published in *The Proceedings*.

*Euratom/UKAEA Fusion Association Culham Science Centre, Abingdon, United Kingdom.

†MPI f. Plasmaphysik, Garching, Federal Republic of Germany.

‡JET Joint Undertaking, Abingdon, United Kingdom, presently at CEA, Cadarache, France.

Work supported by
the U.S. Department of Energy
under Contract No. DE-AC03-99ER54463,
the United Kingdom Dept of Trade & Industry, and EURATOM

GA PROJECT 30033
JULY 1999



DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, make any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Multi-Device Scaling of Neoclassical Tearing Mode Onset with Beta

R.J. La Haye, R.J. Buttery,¹ S. Guenter,² G.T.A. Huysmans,³ and H.R. Wilson¹

General Atomics, P.O. Box 85608, San Diego, California 92186-5608

¹Euratom/UKAEA Fusion Association Culham Science Centre, Abingdon, OX14 3DB, UK.

²MPI f. Plasmaphysik, Garching, Federal Republic of Germany

³Jet Joint Undertaking, Abingdon, United Kingdom, presently at CEA, Cadarache, France

The islands from tearing modes driven unstable and sustained by helically perturbed neoclassical bootstrap current at high beta often provide the practical limit to long-pulse, high confinement tokamak operation [1,2]. The discharges studied are ELMy H-mode single-null divertor (SND) at $q_{95} \geq 3$. Periodic sawteeth with $m/n=1/1$ and $2/2$ are observed to induce $m/n=3/2$ neoclassical tearing modes (NTMs) in the tokamaks Asdex-Upgrade [3], DIII-D [4] and JET [5]; confinement can drop by up to 30%, constituting a "soft" beta limit. Data for the onset of these modes was obtained by slowly raising beta on a time scale longer than the sawteeth period and observing the beta value at onset. Comparison of the measured critical beta to a model for the critical beta is made in terms of dimensionless parameters. This modeling is then used for extrapolation/prediction to a reactor-grade tokamak.

For a classically stable tearing mode, $\Delta' < 0$, the perturbed neoclassical bootstrap current, proportional to beta poloidal β_{θ} , can induce destabilization if β_{θ} exceeds a critical value [1,2]

$$\beta_{\theta c} \approx \frac{(-\Delta' r)(w_s/r) / (\epsilon^{1/2} L_q / L_p)}{\frac{w_s^2}{w_s^2 + w_d^2} - \frac{w_{pol}^2}{w_s^2}} \quad (1)$$

Here w_s is the "seed" island width which must exceed a "threshold" island width w_{th} , i.e. the NTM will grow if $w_s > w_{th}$ and $\beta_{\theta} > \beta_{\theta c}$. Experiments indicate [3-6] that $w_{th} \approx w_{pol}$, the predicted polarization/inertial threshold, which scales [7] as $w_{pol}/r \propto \rho_{i*} g^{1/2}(\epsilon, \nu)$. Here g is a function of the collisionality $\nu = \nu_i / \epsilon \omega_{e*}$, with $g = 1$ at $\nu \ll 1$ and $g \approx \epsilon^{-3/2} \gg 1$ for $\nu \gg 1$. The smaller relative threshold w_d/r predicted due to incomplete flattening of the pressure within an island, scales as $(\chi_{\perp} / \chi_{\parallel})^{1/4} \propto (\chi_{BOHM} / C_s w^{-1})^{1/4} \propto \rho_{i*}^{1/3}$ and is evaluated to have $w_d^2 \ll w_{pol}^2$.

Scaling or extrapolation to a reactor grade tokamak also requires understanding of how the relative seed island w_s/r scales. For example, if w_s/r decreases faster than w_{pol}/r decreases, with ρ_{i*} for example, then $\beta_{\theta c}$ can increase as the denominator in Eq. (1) becomes smaller. On the other hand, $\beta_{\theta c}$ decreases $\propto w_{pol}/r$ if $w_{pol}^2/w_s^2 < 1$ is constant, i.e. $\beta_{\theta c} \propto \rho_{i*} g^{1/2}(\nu, \epsilon)$, an unfavorable scaling for a reactor grade tokamak which will have low ρ_{i*} . The relative seed island scaling is taken as $w_s/r \propto \beta_{\theta}^{\gamma} S^{-\alpha} \propto \rho_{i*}^{3\alpha} \nu^{\alpha}$ (with $\gamma = \alpha/2$ for simplicity and proper limit at zero beta and α determined from fits to the experimental data) where $S \propto \beta^{1/2} / \nu \rho_{i*}^3$ is the magnetic Reynold's number; this allows for the effects of geometrically coupled perturbations of the periodic sawtooth instability to the $q=3/2$ surface [8]. Note that in this form, $w_s/w_{pol} \propto \rho_{i*}^{3\alpha-1}$, at constant collisionality, and the eventual stabilization of a reactor grade tokamak depends on whether $\alpha \geq 1/3$. For $\alpha = 1/3$, $\beta_{\theta c}/\rho_{i*}$ is a function of collisionality only and thus $\beta_{\theta c}$ decreases with lower ρ_{i*} . However, if $\alpha > 1/3$, $\beta_{\theta c}/\rho_{i*}$ will eventually tend to infinity, i.e.

indicate stability, at low enough ρ_{i*} , high S so that $w_s^2 < w_{pol}^2$ and the seed island has become too small to excite the instability even though the threshold island width has also become small.

Data from Asdex-Upgrade (AUG), DIII-D and JET at the onset of sawtooth induced $m/n=3/2$ neoclassical tearing mode is plotted in Fig. 1(a) and (b) with all values evaluated at the $q=3/2$ surface on the outboard midplane and $\beta_N = \beta(\%)/(I/aB)$ the global normalized beta. Figure 1(a) shows that a single separable power law in $v = v_i/\epsilon\omega_{e*}$ does not fit multiple-devices and that α is thus $\neq 1/3$. Note the similarity in the v scaling (positive power) of the "medium-sized" tokamaks AUG and DIII-D and the opposite v scaling (negative power) of the "large-sized" JET. JET being larger and, typically, of lower ρ_{i*} and higher S may be entering the regime where, with $\alpha > 1/3$, w_s is decreasing faster than w_{pol} with lower ρ_{i*} . Figure 1(b) shows the dependence of $\beta_N/f(v)$ normalized by $1/3$ [crossing of curves value in Fig 1(a)] versus ρ_{i*} where the three separate fitted $f(v)$ from Fig. 1(a) are used. The linear dependence on ρ_{i*} is a strong indication of the validity of the polarization/inertial threshold model and the value of multi-device dimensionless comparisons for extending the data range, here in ρ_{i*} .

As β_N/ρ_{i*} is not a function of v alone, $\alpha \neq 1/3$ and the complete dimensionless model from Eq. (1) is fitted to the AUG/DIII-D/JET data as

$$\frac{\beta_{NC}}{C_0\rho_{i*}} = \frac{C_1\rho_{i*}^{3\alpha-1}v^\alpha}{\frac{1}{1+C_4\rho_{i*}^{2/3-6\alpha}v^{-2\alpha}} - \frac{(1+C_2v)/(1+C_3v)}{C_1\rho_{i*}^{2/3-6\alpha}v^{2\alpha}}} \quad (2)$$

with fitted constants C_0, \dots, C_4 and α . Common values are assumed for all three tokamaks. Note that: (1) $C_0 \propto -\Delta'r$ and is assumed constant for all devices and parameter regimes; (2) C_1 and α give the seed island amplitude and scaling; (3) C_2 and C_3 determine the "knee" in the transition from low to high collisionality regimes of the polarization threshold; and (4) C_4 gives the amplitude of the transport threshold, nearly negligible for existing devices but retained as potentially significant for low ρ_{i*} reactor-grade tokamaks.




Fig. 1. (a) Critical β_N normalized by ρ_{i*} versus collisionality v . (b) Critical β_N normalized by the separate fitted functions of v [from Fig. 1(a)] versus ρ_{i*} . ■ JET, ● AUG, ▲ DIII-D.

In fitting, the constants are adjusted to minimize the sum of the square of the difference in the fitted critical β_N and the experimental critical β_N . A test is made to ensure that all data points are on the "unstable" branch of Eq. (1) and not on the "saturated" branch. The unstable branch occurs for seed islands less than a minimum value w_{\min} for which $\beta_{\theta c}$ in Eq. (1) is a minimum, i.e.,

$$w_{\min} \approx (3w_{\text{pol}}^2 + 3w_d^2)^{1/2}, \quad (3a)$$

assuming $w_{\min}^2 \gg w_d^2$ and

$$\beta_{\theta c, \min} \approx -\Delta' \frac{3\sqrt{3}}{2} (w_{\text{pol}}^2 + w_d^2)^{1/2} / (\epsilon^{1/2} L_q / L_p). \quad (3b)$$

This is readily adapted for Eq. (2). The best fit contour plot of the combined data of the critical β_N in $\rho_{i^*} - v_i/\epsilon\omega_{e^*}$ space is shown in Fig. 2. Note that at fixed $\rho_{i^*} = 7.0 \times 10^{-3}$, for example, the horizontal dashed line typical of existing data, the minimum in β_N with v reproduces the change in sign of $d\beta_N/dv$ from AUG/DIII-D to JET. High critical β_N occurs at either: (1) high ρ_{i^*} -high v (larger w_{pol} , upper right of Fig. 2, AUG/DIII-D) or (2) low ρ_{i^*} and/or low v — high S (smaller w_s , lower left of Fig. 2, JET). At fixed v , the vertical dashed line in Fig. 2, β_{NC} decreases with ρ_{i^*} until a minimum is predicted to be reached then rises rapidly with yet lower ρ_{i^*} as $w_s^2/w_{\text{pol}}^2 \rightarrow 1$. The nominal ITER/EDA operating point ($\beta_N = 2.5$, $\rho_{i^*} = 1.2 \times 10^{-3}$ and $v_i/\epsilon\omega_{e^*} = 0.13$ at Greenwald density $G = 1.35$) is predicted to be just stable with $w_s < w_{\text{threshold}}$ and the sawteeth would be unable to excite the 3/2 neoclassical




Fig. 2. Contour plot of best fit of combined data to the NTM dimensionless scaling model. The minimum in β_N (\times) with v at a fixed ρ_{i^*} is shown along with the predicted minimum with ρ_{i^*} at a fixed v (\bullet). The nominal operating point for the ITER/EDA (\star) is just in the stable region.

tearing mode. Thus, the model suggests that a reactor-grade tokamak could operate in a regime where stability improves with decreasing normalized ion gyroradius, a favorable result.

Future work is to: (1) include data from other devices, particularly the high field Alcator-C-Mod and the large JT-60U and (2) to directly confirm the $w_s/r \propto S^{-\alpha}$ scaling. Data from Alcator C-Mod at higher field and smaller size but with similar ρ_{i*} and v as existing devices would lend confidence in the dimensionless scaling approach. Data from the larger size JT-60U could access lower ρ_{i*} . The directly measured scaling of w_s/r is needed to refine and confirm the model.

This is a report of work supported in part by the U.S. Department of Energy under Contract No. DE-AC03-99ER54463 and the United Kingdom Department of Trade and Industry and EURATOM.

References

- [1] O. Sauter, *et al.*, Phys. Plasmas **4** (1997) 1654.
- [2] C.C. Hegna, Phys. Plasmas **5** (1998) 1767.
- [3] S. Guenter, *et al.*, Nucl. Fusion **38** (1998) 1431.
- [4] R.J. La Haye and O. Sauter, Nucl. Fusion **38** (1998) 987.
- [5] G.T.A. Huysmans, *et al.*, presented at 17th IAEA Fusion Energy Conference, October 19–24, 1998, Yokohama, Japan, to be published in proceedings.
- [6] D. Gates, *et al.*, Nucl. Fusion **37** (1997) 1593.
- [7] H.R. Wilson, *et al.*, Phys. Plasmas **3** (1996) 248.
- [8] C.C. Hegna, *et al.*, Phys. Plasmas **6** (1999) 130.