

13
2-15-95 JZ (1)

PNL-10400
~~SECRET~~
DRAFT
UC-2010

Identification of Contaminants of Concern

Columbia River Comprehensive Impact Assessment

B. A. Napier
N. C. Batishko
D. A. Heise-Craff
M. F. Jarvis
S. F. Snyder

January 1995

Prepared for the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830

Pacific Northwest Laboratory
Richland, Washington 99352

MASTER

~~RESTRICTION~~ DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor Battelle Memorial Institute, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or Battelle Memorial Institute. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PACIFIC NORTHWEST LABORATORY
operated by
BATTELLE MEMORIAL INSTITUTE
for the
UNITED STATES DEPARTMENT OF ENERGY
under Contract DE-AC06-76RLO 1830

Printed in the United States of America

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831;
prices available from (615) 576-8401. FTS 626-8401.

Available to the public from the National Technical Information Service,
U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.



The contents of this report were printed on recycled paper

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.



Department of Energy

Richland Operations Office
P.O. Box 550
Richland, Washington 99352

FEB - 6 1995

Those on Attached List:

SOLICITATION OF COMMENTS ON DRAFT IDENTIFICATION OF CONTAMINANTS OF CONCERN REPORT

The U.S. Department of Energy, Richland Operations Office (RL), has negotiated an agreement with the State of Washington Department of Ecology (Ecology) and the U.S. Environmental Protection Agency (EPA) to perform a comprehensive impact assessment of current and residual Hanford-derived contaminants to the Columbia River for remedial decisions at the Hanford Site. Associated current human health and environmental impacts will be assessed. This process will utilize an ecosystem approach for guiding remedial decisions. 100-Area, 200-Area, and 300-Area operable units will continue to assess contaminant sources and remediation. The Columbia River Comprehensive Impact Assessment (CRCIA) will address all Columbia River contaminants, risk assessments, and remediation. If unacceptable levels of human health or environmental risk are found, appropriate remedial actions will be initiated consistent with the National Contingency Plan and the Hanford Federal Facility Agreement and Consent Order (Tri-Party Agreement) through the Hanford Past Practice Strategy. Remedial decisions resulting from the CRCIA will consider impacts to the environment and natural resources from alternative remedial options as part of the remedial decision process.

The initial CRCIA effort was the development of a "compendium" of existing data on Columbia River contamination (a bibliography of data sources has been developed in support of this effort).

The next step is to define the "contaminants of concern" for this study. A contaminants of concern report has been produced which documents an initial review, from a risk perspective, of historical data concerning current or potential contamination in the Columbia River. Sampling data were examined for over 600 chemical and radioactive contaminants. A screening analysis was performed to identify those substances present in such quantities that they may pose a meaningful human or ecological risk. The substances identified will require a more detailed analysis to assess their impact on humans or the river ecosystem.

The next document to be produced will be an identification of the "species of concern" for this study. It is the intention of RL and PNL to solicit early input, from those interested, concurrently with the review and comment period of the enclosed document.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Addressees

-2-

FEB - 6 1995

RL is pleased to provide you a copy of the draft subject document for your review. This document has not undergone review by RL, or the regulatory agencies (Ecology and EPA); therefore, there is ample opportunity to provide meaningful input prior to its finalization. Please provide comments to Mr. Randy Brich at this address by March 10, 1995. Mr. Brich may be reached at (509) 376-9031. RL and its contractor for this study, Pacific Northwest Laboratory (PNL), are pleased to meet with those interested; please let Mr. Brich know if you desire to do so.

Sincerely,



Julie K. Erickson, Director
River Sites Restoration Division

RSD:RFB

Enclosure

cc w/encl:
R. F. Stanley, Ecology

Addressees - Letter dated FEB - 6 1995

<u>No. of Copies</u>		<u>No. of Copies</u>
	Offsite	
12	DOE Office of Scientific and Technical Information	F. R. Cook 2552 Harris Richland, WA 99352
	M. Bauer Confederated Tribes and Bands of the Yakama Indian Nations P.O. Box 151 Toppenish, WA 98948	J. P. Corley 2213 Torbett Richland, WA 99352
	S. Benz 600 S. Kent #71 Kennewick, WA 99336	2 DNR Hanford Projects Attn: J. R. Wilkinson Confederated Tribes of the Umatilla Indian Nation P.O. Box 638 Pendleton, OR 97801
	L. Block U.S. Fish and Wildlife Service Moses Lake Field Office P.O. Box 1157 Moses Lake, WA 98837	G. deBruler Columbia River United P.O. Box 667 Bingen, WA 98605
	D. Bradshaw Audobon Society of Portland 5151 Northwest Cornell Road Portland, OR 97210	C. Denniston Greenpeace 11815 - 20th SW Seattle, WA 98146
	N. Buske Search Technical Services Star Route Box 17 Davenport, WA 99112	B. Drost USGS Water Resources Division Washington District 1201 Pacific Avenue Suite 600 Tacoma, WA 98402
	D. Cellarius Sierra Club Northwest Office 1516 Melrose Avenue Seattle, WA 98122	D. Dunning Oregon Dept. of Energy 625 Marion Street NE Salem, OR 97310
	C. Cline State of Washington Department of Ecology P.O. Box 47600 Olympia, WA 98504-7600	3 J. Erickson Division of Radiation Protection Washington State Dept. of Health P.O. Box 47827 Olympia, WA 98504-7827

No. of
CopiesNo. of
Copies

M. Fox, President
Hanford Family
P.O. Box 1462
Richland, WA 99352

R. Gardiner
614 Sherman Drive
The Dalles, OR 97058

N. J. Germond
League of Women Voters
224 Iron Mountain Boulevard
Lake Oswego, OR 97034

D. Maez
4214 W. John Day Place
Kennewick, WA 99336

C. Mebane
National Oceanic and Atmospheric
Administration
c/o EPA Seattle Office
1200 6th Avenue, HW-113
Seattle, WA 98101

J. Monteith
Oregon Natural Resources Council
1161 Lincoln Street
Eugene, OR 97401

D. Nichols
WNP-1
Building 52
Richland, WA 99352

R. Patt
Water Resources Department
158 - 12th Street N.E.
Salem, OR 97310-0210

G. Pollet
Heart of America Northwest
1305 4th Avenue, 208
Seattle, WA 98101

2 D. Powauke
Nez Perce Tribe
P.O. Box 365
Lapwai, ID 83540

S. M. Alexander
State of Washington
Department of Ecology
1315 W. Fourth Avenue
Kennewick, WA 99336-6018

L. Stembridge
HEAL
1408 West Broadway Avenue
Spokane, WA 99201-1902

C. Stephan
1520 Torthay Court
Richland, WA 99352

3 D. Stewart-Smith
Oregon Department of Energy
625 Marion Street NE
Salem, OR 97310

J. P. Thomas
HHIN Resource Center
1719 Smith Tower
506 Second Avenue
Seattle, WA 98104

Onsite


5 **State of Washington
Department of Ecology**

D. P. Holland (4) B5-18
J. W. Yokel B5-18

5 **U.S. Department of Energy
Richland Operations Office**

DOE Public Reading Room H2-53

<u>No. of Copies</u>		<u>No. of Copies</u>	
4	Westinghouse Hanford Company	G. L. Harvey	K2-02
		P. C. Hays	K9-41
	Administrative Record (3)	D. A. Heise-Craff	K3-54
	A. S. Carlson	L. R. Huesties	K6-61
		M. F. Jarvis	K6-55
6	U.S. Environmental Protection Agency	A. D. Maughan	K6-52
		T. B. Miley	K5-12
	L. E. Gadbois (5)	L. F. Morasch	K6-62
	D. R. Sherwood	B. A. Napier (5)	K3-54
		D. B. Shipler	K9-13
		S. F. Snyder	K3-54
37	Pacific Northwest Laboratory	W. L. Templeton	K9-13
		W. H. Walters	K9-33
	N. C. Batishko	L. K. Wenrich (5)	K3-70
	S. D. Cannon	B. K. Wise	K9-04
	R. L. Dirkes	Records Center (2)	K3-70
	P. W. Eslinger	Publishing Coordination	K1-06
	S. L. Friant	Technical Library (5)	P8-55
	M. S. Hanson		



The environmental quality of the Columbia River is of special interest to the public, government, and tribal governments as a source of drinking water, for crop irrigation, as ecological habitat, and for recreation. The following actions have been taken to encourage public involvement in the CRCIA Project:

- PNL has an open door policy for this project. Non-PNL individuals can visit the laboratory, interact with scientists, and observe work in progress.
- Data and documents used in the CRCIA Project are being made available to all interested parties.
- Public meetings are being conducted to obtain input to the development of work scope and technical approaches as well as to review data and work progress.

Abstract

The Columbia River Comprehensive Impact Assessment (CRCIA) Project is conducted for the U.S. Department of Energy by the Pacific Northwest Laboratory. The CRCIA Project will evaluate the current human and ecological risks from the Columbia River attributable to past and present activities on the Hanford Site. To perform a comprehensive assessment, the contaminants released from the Hanford Site must be identified. This report identifies the contaminants released and identifies those that should be considered in detailed risk analyses.



Summary

Introduction

The Columbia River Comprehensive Impact Assessment (CRCIA) Project is conducted for the U.S. Department of Energy by the Pacific Northwest Laboratory (PNL). The CRCIA Project will evaluate the current human and ecological risks from the Columbia River attributable to past and present activities on the Hanford Site. To perform a comprehensive assessment, the contaminants released from the Hanford Site must be identified. This report identifies the contaminants released and identifies those that will be considered in detailed risk analyses.

Scope of Work

The CRCIA Project is primarily concerned with the current risks from contaminants of Hanford origin. Therefore, the most recent sampling data (from 1980 through 1994) were used to estimate the source term (amount and types of radionuclides and chemicals released to the environment from Hanford facilities) for the risk calculations. For this study, the focus is on the Columbia River water, sediment, soil, and groundwater within 150 meters (500 feet) of the Columbia River, which means a spatial focus on the Hanford 100, 300, and 1100 Areas. A multi-stage screening process was developed to prioritize these various contaminants in terms of human health risk and ecosystem risk. Each stage of the process identifies contaminants of interest to the project, based on the potential for human and ecological risk. The combined results of the total screening then compose the total list of concern.

In addition to radiological and chemical contaminants, the potential for radiation doses arising from discrete radioactive particles in the river sediment or from direct irradiation from near-river Hanford facilities is also addressed.

Although the primary concern is the current status of the Columbia River, additional consideration is given to the potential impact of contaminants currently known to be in the Hanford Site groundwater. Consideration is not given to the potential impact of contaminants that are not presently in the groundwater but which may be in soils or facilities away from the Columbia River.

Technical Approach

The first step in the approach was to collect a comprehensive list of potential contaminants. This list was prepared by examining published data, reports, and contaminant databases. The review of the available data indicated that concentrations of various radionuclides, carcinogenic chemicals, and hazardous chemicals had been measured in surface water (Columbia River, springs, and seeps), groundwater, river sediment, and near-river soil. A multi-stage screening process was developed to prioritize these various contaminants in terms of human health risk and ecosystem risk. Each stage of the process identifies contaminants of interest. The combined results of the entire screening process then compose the total list of contaminants of concern. The following screening processes were used.

Initial Screening: Initial screening eliminated the contaminants on the list that showed no detectable levels of activity or concentration.

Radionuclide Screening: Radionuclide screening is based on a scenario of exposure to an individual. The exposure includes external exposure, consumption of untreated river water, consumption of freshwater fish, and consumption of small amounts of sediment. Internal risks are estimated using the U.S. Environmental Protection Agency (EPA) indicator for ingestion, called a slope factor (EPA 1994a). This indicator represents the risk of cancer to an individual from sources other than natural background radiation per unit (e.g., picocurie) of radioactive material taken into the body. Similarly, external exposure to contaminated sediment is addressed by assuming the parameters associated with the EPA slope factor for external exposure are appropriate (EPA 1994a).

Carcinogenic Chemical Screening: The individual exposure scenario for carcinogens in river water are the same as those for radionuclides, except there is no factor for external exposure because there is no external risk from chemicals.

Toxic Chemical Screening: For hazardous, but noncarcinogenic, chemicals, the screening is based on a ratio of the estimated daily intake to the EPA chronic oral reference dose (EPA 1994a). The chronic oral reference dose is the safe dose level EPA established for specific chemicals. In other words, the chemicals in the individual exposure scenario are investigated to screen out those that are ingested in amounts below the EPA's safe levels. The exposure scenario is the same as for the radionuclides or carcinogens.

Ambient Water Quality Criteria Screening: For aquatic plants and animals (biota), the measured or surrogate (estimated) concentration of the contaminant in water is compared with the applicable EPA water quality criterion (EPA 1992). The ambient water quality criteria are those concentrations of chemicals identified by EPA as safe and protective of aquatic life.

Aquatic Biota Toxicity Screening: Limited data were available that identify the concentrations of certain chemicals that result in toxic effects to aquatic life. Where possible, the threshold concentration for fresh water at which any effect was noted was used. Where not possible, the lowest concentration lethal to 50 percent (called LC50) of small, freshwater fish (e.g., guppies, mosquito fish, rainbow trout) was used (EPA 1985). To relate these lethal effects to less significant effects, the screening used a value of 1 percent of the LC50. For a few analytes (substances for which an analysis is made) for which fish data were not available, test results for crayfish or insects were used as a surrogate.

Background Screening: During the screening process, a few radionuclides and chemicals had measurements determined to be within their respective naturally occurring background levels. Because concentrations were not above naturally occurring background, the following contaminants were eliminated from further consideration: the radionuclides beryllium-7 and potassium-40; the chemicals barium, bismuth, boron, chlorine, fluorine, lithium, silicon, silver, sulfide, titanium, vanadium, and zirconium.

Nonhazardous Screening: The screening process identified several materials as nonhazardous under environmental conditions (EPA 1991; EPA 1989). These contaminants eliminated from further consideration are aluminum, calcium, iron, magnesium, potassium, and sodium.

All of the screenings require an estimate of the contaminant's concentration in river water. Only the direct river water measurements provide this information. When direct measurements of river water were not available, surrogate water concentration was estimated. To estimate surrogate concentrations in water, certain assumptions were used.

Groundwater Contamination: Groundwater adjacent to the Columbia River can flow into the river, and Columbia River water can flow into the groundwater, depending on river flow. Therefore, concentrations of contaminants in groundwater near the river are difficult to predict, and concentrations measured near the shore differ from those measured further inland. Raymond et al. (1976) and Cline et al. (1985) report an estimated flow rate of 100 cubic feet per second (cfs) over the entire Hanford Reach. For conservatism (i.e., to provide an estimate of the resulting concentration in the river that, if incorrect, would err on the high side), the value of 100 cfs was adopted for the screening. In effect, this implies that the entire groundwater that flows from beneath Hanford to the Columbia River is contaminated to the maximum level measured.

River Sediment: Sediment within the river is both a reservoir of contaminants and a source of contamination of the river water, as the material is dissolved into or carried away by the river. An equilibrium ratio of 1:100,000 was used (i.e., the concentration of the contaminant in the sediment is assumed to be 100,000 times higher than in the Columbia River waters). This assumption is based on a limited number of samples and an empirical equation (Napier et al. 1988, p. 4.82).

Near-River Soil: Contaminants in Hanford waste sites or other sites adjacent to the Columbia River (e.g., operating facilities, spills, etc.) may pose a threat of future contamination of the river. For the purpose of screening, all contaminants are assumed to be environmentally mobile and potentially dissolvable in groundwater. Based on this assumption, the surrogate groundwater contamination is assumed to have the same concentration of contaminants as the soil. The total area of industrial activity comprises approximately 6 percent of the Hanford Site (Dirkes et al. 1994, p. 5). Because it is unreasonable to assume that all of Hanford soil is contaminated to the maximum concentration measured, an effective area of 1 percent is assumed. This means that the study assumed that 1 percent of Hanford soil is contaminated to the same extent as the highest amounts measured in Hanford soil.

Results

Analyses for more than 600 different radionuclides and chemicals have been performed on Hanford-related environmental samples. A large number of these potential contaminants have never been detected in the Hanford/Columbia River environments. Screening on the basis of potential impact on human health or the health of Columbia River ecosystems has been performed for the roughly 100 radionuclides and chemicals that have been detected in environmental samples. Several different types of screenings were employed. The results were consistent in that the same materials were identified numerous times by the various screenings. Application of the screenings for contaminants within 150 meters (500 feet) of the Columbia River yields a list of 20 contaminants of concern, plus direct irradiation. These contaminants are given in the first column of Table S.1.

Table S.1. List of Identified Contaminants of Concern^(a)

In Columbia River, Groundwater, ^(b) Sediment, and Soil	Groundwater Plumes Away from the Columbia River ^(c)	Continued Public Interest
Antimony Arochlor 1248 (PCB) Arsenic Cesium-134 Cesium-137 Chlordane Chromium ^(d) Cobalt-60/particles Copper Diesel Fuel Europium-152 Europium-154 Lead Manganese Mercury Nitrate/nitrite ^(d) Phosphate Silver Chloride Strontium-90 Zinc	Carbon Tetrachloride Fluoride	Chloroform Cyanide Iodine-129 Plutonium-239/240 Technetium-99 Trichloroethylene Tritium (Hydrogen-3) Uranium
(a) Direct irradiation is also identified as being of concern. (b) Hanford groundwater within 150 meters (500 feet) of the Columbia River. (c) Hanford groundwater farther than 150 meters (500 feet) from the Columbia River. (d) These contaminants are also of concern in groundwater plumes away from the Columbia River but are not repeated in that list to avoid duplication.		

Existing Hanford groundwater contamination farther than 150 meters (500 feet) (see Table 3.3) from the Columbia River was also addressed. The contaminants identified by the screening process do not appear to be currently entering the river but have the potential to do so within 10 to 200 years (Freshley and Graham 1988). Two contaminants (chromium and nitrate) in Hanford groundwater away from the river are already included in this study because they are in or near the river. Only carbon tetrachloride and fluoride were added to the list as a result of the study of groundwater away from the river. Carbon tetrachloride and fluoride have not yet been found in the river.

Although the screenings did not indicate a potential risk, several potential or existing contaminants are of particularly high public interest (third column in Table S.1). Essentially all of these are the object of ongoing evaluation by the Surface Environmental Surveillance Project (SESP) conducted by PNL at Hanford. The CRCIA Project should remain current on SESP activities and include SESP results in all project reports (see Section 8.0).

Each of the identified contaminants can be considered to have resulted from past plutonium-production operations at Hanford. The radionuclides on the list generally represent those identified with river water or Hanford Reach sediment. The radionuclides resulted from activation of materials in the old production reactors. It is likely that the cesium isotopes are related to global fallout (Dirkes et al. 1994). Most of the metals identified from Hanford groundwater or sediment can be related to various Hanford operations in the 100 Areas. The polychlorinated biphenyl (PCB), Arochlor 1248, is used in equipment and the insecticide, Chlordane, has been used at Hanford facilities, but both are still essentially associated with soil near the river. The nitrate groundwater plumes result from past Hanford operations in the 100 and 200 Areas.

The identification of the radionuclides and chemicals as being of concern to the CRCIA Project does not imply that each or all of these compounds is necessarily a prominent problem for the river or those who live downstream. The screening and selection process described in this report is a conservative (cautious) process designed to focus the resources of the project on those contaminants with potential risk.

Glossary

100 Areas - site of the Hanford production reactors, which include B, C, D, DR, F, H, KE, KW, and N reactors.

200 Areas - site of the Hanford chemical separations plants, which include the bismuth phosphate process plants (B and T Plants), plutonium uranium extraction plant (A Plant/PUREX), and reduction and oxidation plant (S Plant/REDOX).

300 Area - site of research, development, and fuel-fabrication operations.

400 Area - site of the Fast Flux Test Facility.

600 Area - all land within the Hanford Site not occupied by the 100, 200, 300, 400, 1100, or 3000 Areas.

1100 Area - site of the warehousing, vehicle maintenance, and transportation operations center.

3000 Area - site of engineering, construction, and research and development activities.

analytes - substances for which an analysis is made.

bioconcentration factor - ratio between the radionuclide concentration in biota and the radionuclide concentration in the water in which the biota live and feed.

biota - plants and animals.

carcinogenic (chemicals) - having the property of enhancing the possibility of contracting cancer later in life following exposure.

CERCLA - *Comprehensive Environmental Response, Compensation, and Liability Act of 1980.*

Ci - abbreviation for curie.

concentration - amount of a specified substance (e.g., a radioactive element) in a unit amount of another substance (e.g., river water, milk).

conceptual model - any representation of a biological or mechanical process.

CRCIA - Columbia River Comprehensive Impact Assessment.

curie - unit of radioactivity corresponding to 3.7×10^{10} (37 billion) disintegrations per second (abbreviated Ci), 1 curie = 3.7×10^{10} becquerel.

DOE - U.S. Department of Energy.

Ecology - Washington State Department of Ecology.

EIS - environmental impact statement.

EPA - U.S. Environmental Protection Agency.

exposure - process of coming into contact with environmental materials.

internal exposure - contact with materials taken into the body through inhalation or ingestion.

external exposure - contact with materials on the outside of the body, as from submersion in water or immersion in air.

gross beta - total activity of beta-emitting radionuclides that are not distinguished separately by instrumentation or radiochemical analyses.

half-life - time required for an initial number of radioactive atoms to be reduced to half that number by radiological transformations.

Hanford Reach - stretch of the Columbia River downstream of Priest Rapids Dam and upstream of the confluence of the Yakima and Columbia Rivers.

hazardous (chemicals) - having the property of being toxic, at some level of exposure. Generally used to differentiate from carcinogenic.

HEIS - Hanford Environmental Information System. An electronic database that consolidates the data gathered during environmental monitoring and restoration of the Hanford Site.

HWMA - *Washington State Hazardous Waste Management Act of 1976.*

IRIS - Integrated Risk Information System, an EPA database that provides data on chronic health hazards (reference dose values), carcinogenicity (unit risk factors or slope factors), EPA regulatory actions, supplementary data, and a bibliography for each listed chemical.

irradiation - exposure of an object to ionizing radiation.

isotope - one of two or more atoms having the same atomic number but different mass.

LFI - limited field investigation conducted as part of Tri-Party Agreement activities to identify those Hanford waste sites that are recommended to remain as candidates for interim remedial measures.

MEPAS - Multimedia Environmental Pollutant Assessment System, a computer code that can be used to estimate the transport and fate of environmental pollutants.

model - conceptual representation of a physical/biological process. The representation may be graphical or a set of mathematical equations that simulate the process being modeled. See also conceptual model.

natural uranium - naturally occurring mixture of uranium (0.7 percent uranium-235 and 99.3 percent uranium-238).

NPL - national priorities list.

operable unit - term used to identify specific areas designated for cleanup.

PCB - polychlorinated biphenyl.

picocurie - one-millionth of a millionth curie (10^{-12}).

plume - definitive volume of air, water, or soil containing contaminants released from a contaminant source.

PNL - Pacific Northwest Laboratory.

production reactor - facility (B, C, D, DR, F, H, KE, KW, or N reactors) in which uranium or other fuel was irradiated with neutrons to produce radioactive materials. Used primarily at Hanford to produce plutonium for weapons; used also for research. Synonymous with "reactor."

radioactivity - spontaneous emission of radiation (alpha, beta, gamma rays, and/or neutrons) by some isotopes as they transform into other isotopes.

radionuclide - radioactive isotope of an element.

RCRA - *Resource Conservation and Recovery Act of 1976.*

reactor - see production reactor.

reference dose - EPA's estimate of the smallest daily intake of a hazardous material that first leads to deleterious health effects.

RI/FS - remedial investigation/feasibility study.

SARA - *Superfund Amendments and Reauthorization Act.*

seeps - very small springs of groundwater.

SESP - Surface Environmental Surveillance Project.

slope factor - EPA's value which represents the lifetime excess cancer risk per unit of intake.

source term - amount of radioactivity (curies) of a radionuclide or amount of a chemical released to the environment from a facility over a given time.

springs - source of water issuing from the ground.

SST - single-shell tank.

stack - tall chimney that was the primary release point of exhaust air from a reactor or separations plant building.

surrogate (measurement) - estimated substitute measurement used when actual measurements not available.

TPA - Tri-Party Agreement (officially, *Hanford Federal Facility Agreement and Consent Order*).

TSD - treatment, storage, and disposal facilities or units at Hanford.

TWRS - tank waste remediation system.

UST - underground storage tank.

VOC - volatile organic compounds.

Contents

Preface	iii
Abstract	v
Summary	vii
Glossary	xiii
1.0 Introduction	1.1
1.1 Background	1.1
1.2 Purpose	1.3
1.3 Scope	1.4
1.4 Preview of Report	1.4
2.0 Data Sources	2.1
2.1 General References	2.1
2.2 Hanford Environmental Information System	2.6
2.3 Remedial Investigation/Feasibility Studies	2.6
2.4 Hanford Site Environmental Reports	2.7
2.5 Limited Field Investigations	2.7
2.6 Discrete Radioactive Particles and Other Direct Exposure Sources	2.9
2.7 National Environmental Policy Act (NEPA) Documents	2.11
3.0 Composite List of Identified Radionuclides and Chemicals	3.1
3.1 Risk-Based Standards Database	3.1
3.2 Environmental Sampling Data Reports	3.1
3.3 Detected Analytes	3.2

4.0	Screening Approach	4.1
4.1	Screening Equations	4.1
4.1.1	Radionuclide Screening	4.1
4.1.2	Carcinogenic Chemical Screening	4.2
4.1.3	Toxic Chemical Screening	4.3
4.1.4	Ambient Water Quality Criteria Screening	4.3
4.1.5	Aquatic Biota Toxicity Screening	4.3
4.2	Estimation of Contaminant Concentrations in River Water	4.4
4.2.1	Radionuclides	4.4
4.2.2	Chemicals	4.6
4.3	Screening Results	4.7
4.3.1	River Water Sample Screening	4.7
4.3.2	Groundwater Sample Screening	4.8
4.3.3	River Sediment Sample Screening	4.8
4.3.4	Near-River Soil Sample Screening	4.9
4.4	Use of Suspect Measurements	4.10
5.0	Discrete Radioactive Particles	5.1
6.0	Direct Irradiation from Hanford Facilities	6.1
7.0	Potential Future Groundwater Sources	7.1
7.1	Existing Groundwater Plumes	7.1
7.2	Potential Future Groundwater Sources	7.3
8.0	Materials of Additional Public Interest	8.1
9.0	Conclusions	9.1

10.0 Perspective	10.1
11.0 References	11.1
Appendix A - Complete List of Analytes Evaluated at Hanford	A.1
Appendix B - Parameter Values Used in Screening Analyses	B.1
Appendix C - Complete Numerical Results	C.1

Figures

1.1 Map of the Hanford Site	1.2
7.1 Nitrate Plume	7.2

Tables

S.1 List of Identified Contaminants of Concern	x
3.1 Maximum Detected Concentrations in the Columbia River and Groundwater in the Hanford Site 100, 300, and 1100 Areas Near the Columbia River, 1980-1994	3.3
3.2 Maximum Detected Concentrations in Soil and Sediment in the Hanford Site 100, 300, and 1100 Areas, 1980-1994	3.5
3.3 Maximum Detected Concentrations in Groundwater in the Hanford Site 100, 200, and 600 Areas Away from the Columbia River, 1980-1994	3.7
4.1 Contaminants of Potential Interest Identified via Screening of Columbia River Samples	4.8
4.2 Contaminants of Potential Interest Identified via Screening of Groundwater Near the Columbia River	4.9
4.3 Contaminants of Potential Interest Identified via Screening of Columbia River Sediment Samples	4.9
4.4 Contaminants of Potential Interest Identified via Screening of Soil Near the Columbia River	4.10

7.1	Contaminants of Potential Interest Identified via Screening of Groundwater Away from the Columbia River	7.3
9.1	List of Identified Contaminants of Concern	9.2
A.1	Radionuclide and Chemical Activity/Concentrations in the Columbia River and Groundwater Near the Columbia River	A.2
A.2	Radionuclide and Chemical Activity/Concentrations in Soil and Sediment	A.11
A.3	Maximum Detected Concentrations in Groundwater in the Hanford Site 100, 200, and 600 Areas Away from the Columbia River, 1980-1994	A.21
B.1	Parameters Used to Screen the Columbia River and Groundwater Near the Columbia River	B.2
B.2	Parameters Used to Screen Soil and Sediment	B.4
B.3	Parameters Used to Screen Groundwater Away from the Columbia River	B.7
C.1	Results for the Columbia River and Groundwater Near the Columbia River	C.2
C.2	Results for Soil and Sediment	C.5
C.3	Results for Groundwater Away from the Columbia River	C.8

1.0 Introduction

Pacific Northwest Laboratory (PNL) is conducting a comprehensive assessment of the Columbia River. The purpose of the Columbia River Comprehensive Impact Assessment (CRCIA) Project is to evaluate the current human and ecological risk from radioactive and other hazardous materials in the Columbia River as a result of past and present activities at the Hanford Site near Richland, Washington. Many thousands of radionuclides and hazardous chemicals^(a) have been generated or used at Hanford over the past five decades, only some of which may be of current concern for human or ecological risk. The intent of this report is to focus the resources of the project on the contaminants of greatest concern.


1.1 Background

The Hanford Site in southcentral Washington State was acquired by the federal government in 1943 and was dedicated for many years to the production of plutonium for national defense and the management of resulting wastes. The production of nuclear materials for weapons ended at Hanford in 1987. With the shutdown of the production facilities, missions were diversified to include research and development in the areas of energy, waste management, and environmental restoration.

The Hanford Site is about 1,450 square kilometers (560 square miles) of semi-arid shrub-steppe located just north of the confluence of the Yakima River with the Columbia River (Figure 1.1). Approximately 6 percent of the Hanford Site has been used for operations in the following areas:

- 100-B/C, 100-D, 100-F, 100-H, 100-K, and 100-N Areas, which lie along the Columbia River in the northern portion of the Hanford Site, are the sites of the nine Hanford plutonium production reactors (now shut down)
- 200-East and 200-West Areas, which lie in the center of the Hanford Site, are the sites of the chemical reprocessing facilities and low-level- and high-level-waste management facilities
- 300 Area, near the southern border of the Hanford Site, is the site used for nuclear fuel manufacturing and research facilities
- 400 Area, between the 200 and 300 Areas, is the site of the Fast Flux Test Facility
- 1100 Area and 3000 Area, a corridor northwest of the city of Richland, are sites used for warehousing, vehicle maintenance, transportation operations center, construction, engineering, and research and development activities.

(a) In this report, organic chemicals, inorganic chemicals, ions, elements, and other chemical compounds are simply referred to as chemicals.



S9501004.1

Figure 1.1. Map of the Hanford Site

Fifty-one miles of the Columbia River, known as the Hanford Reach, flows through or borders the Hanford Site. The Hanford Reach is roughly from Priest Rapids Dam to the confluence of the Yakima River with the Columbia River. This stretch of the river offers a unique example of the river and riparian (riverside) ecologies that characterized the Columbia Basin ecosystem prior to construction of hydroelectric dams on the river. The Hanford Reach comprises the last unimpounded stretch of the Columbia River in the United States. Nearly 60 percent of the Columbia River's native wild stock of fall chinook salmon spawn in the reach (National Parks Service 1992). River water is used downstream from the Hanford Site by Washington and Oregon residents for drinking water, agriculture, industry, transportation, and recreation. The riverbanks and islands provide habitat for several species of threatened or endangered plants (e.g., Columbia milkvetch and Hoover's desert parsley) and animals (e.g., bald eagles) (National Parks Service 1992).

Plutonium production operations in the 100 Areas historically have resulted in releases of contaminants directly to the Columbia River and left extensive contamination in some areas of the surface soil, subsurface soil, and groundwater. Contamination reaches the river through groundwater seepage.

Facilities in the 200 Areas were built to process irradiated fuel from the production reactors. The subsequent operation of these facilities resulted in the storage, disposal, and some releases of radioactive and nonradioactive wastes to the environment. Contamination exists in the surface, subsurface, and groundwater in the 200 Areas. Contaminated groundwater has moved out of the operating areas into areas adjoining the operating areas.

The 300 Area is the site of former reactor fuel processing activities. The 300 Area is also the location of nuclear research and development facilities serving the Hanford Site. Wastes in the 300 Area have resulted from the fuel fabrication process and various research activities. Contamination exists in the surface, subsurface, and groundwater.

The 1100 Area just north of Richland serves as the warehousing, vehicle maintenance, and transportation operations center for the Hanford Site. Wastes present result primarily from disposal of batteries, paints and solvents, and antifreeze. Immediately adjacent to the 1100 Area is the 3000 Area, home of Hanford Site engineering, construction, and research and development activities. Minor chemical contamination from paints, solvents, and related activities is also present here.

The 600 Area is defined to include all land within the Hanford Site not occupied by the 100, 200, 300, 400, 1100, and 3000 Areas. Land uses within the 600 Area include a 41-hectare (100-acre) tract subleased from the state of Washington for the disposal of commercial low-level nuclear waste and nuclear power facilities operated by the Washington Public Power Supply System. Most contamination in the 600 Area reaches the Columbia River by groundwater.

1.2 Purpose

This report documents an initial review of the abundance of historical data concerning contamination, current or potential, of the Columbia River. The initial review focuses on the availability of key data for particular contaminants at specific locations in specific media. The result is a list of

contaminants of concern for current human or ecological risk. The list will help focus the effects of health risk assessments because the contaminants on this list are those with the highest risk levels.

The list of contaminants of concern will also be used to help define future sampling requirements to obtain current data for use in the CRCIA Project.

1.3 Scope

This study is primarily concerned with the current risks from contaminants of Hanford origin. Therefore, the most recent sampling data are used to provide the applicable source term for the risk calculations. For this study, the focus is on the Columbia River water, sediment, soil, and groundwater within 150 meters (500 feet) of the Columbia River, which means a spatial focus on the Hanford 100, 300, and 1100 Areas. A multi-stage screening process was developed to prioritize these various sources in terms of human health risk and ecosystem risk. Each stage of the process identifies pollutants of interest. The combined results of the total screening then compose the total list of concern.

The potential is also addressed for radiation doses arising from discrete radioactive particles in the river sediment or from direct irradiation from near-river Hanford facilities.

Although the primary concern is the current status of the Columbia River, additional consideration is given to the potential for future impact by contaminants currently present in the Hanford Site groundwater. Consideration is not given to the potential impact of contaminants that may be in soils or facilities away from the Columbia River but that are not presently in the groundwater.

1.4 Preview of Report

The references used as data sources are annotated in Section 2.0 of this report. A composite list of radionuclides and chemicals identified as being present in environmental samples is presented in Section 3.0. The numerical approach to screening the several hundred analytes into a short list of contaminants of concern is presented in Section 4.0. The results of the screening process are listed in Section 4.3. A discussion of discrete radioactive particles in the sediment of the Columbia River shoreline and islands is given in Section 5.0. Section 6.0 addresses direct gamma irradiation from Hanford facilities located adjacent to the river. Section 7.0 addresses existing and potential future contaminants from groundwater sources away from the river. Contaminants of possible continued public interest are acknowledged in Section 8.0. The overall conclusions, listed as the contaminants of concern, are given in Section 9.0. Supporting material is made available in the appendices at the end of the report.

2.0 Data Sources

An annotated bibliography of the sources used to identify the analytes sampled in environmental media are provided in this section. No single document or electronic database was available that covered the entire scope of contaminants for this research. Baseline efforts similar to the scope of our task were done in a project by Fowler et al. (1993). However, because that project covered all exposure pathways and numerous U.S. Department of Energy (DOE) sites, and identified only the presence of contaminants and not their concentrations, it is not directly applicable or as comprehensive as required for this task.

The CRCIA Project developed a compendium of existing data on Columbia River contamination (Eslinger et al. 1994). The compendium is a large bibliography of Hanford and non-Hanford sources that potentially contain relevant environmental monitoring information. This compendium was used as a starting point for data information.

This study is primarily concerned with the current risks from contaminants of Hanford origin. Therefore, the most recent sampling data provide the source term for the risk calculations. A secondary concern of this study is the potential for future contamination of the river from Hanford facilities away from the river. Summary information related to existing groundwater plumes that are farther than 150 meters (500 feet) from the Columbia River on the Hanford Site was also reviewed.

To understand some of the key terms in the bibliography, it is necessary to know that the radioactive, hazardous chemical, and mixed wastes are found in various individual waste sites, referred to as waste management units, located throughout the Hanford Site. These individual waste management units include past practice sites; surplus facilities; and treatment, storage, and disposal (TSD) facilities. Past practice sites and TSD facilities may take the form of spills, cribs, ditches, ponds, tanks, trenches, landfills, burial grounds, pits, French drains, and other means of intentional or unintentional disposal. Surplus facilities include contaminated buildings, exhaust stacks, and underground transfer lines. The individual waste management units are organized into "operable units" based on geographic proximity or similarity of waste disposal history.

The following annotated bibliography summarizes the sampling data sources and primary references used in the compilation of the monitoring data. The complete reference, sampling purpose, sampling time frame, media sampled, as well as supplementary comments, are provided. Documents of specific types are listed together, in alphabetical order. Appendix A presents a complete list of radionuclides and chemicals evaluated at Hanford.

2.1 General References

Dirkes, R. L. 1993. *Columbia River Monitoring: Distribution of Tritium in Columbia River Water at the Richland Pumphouse*. PNL-8531, Pacific Northwest Laboratory, Richland, Washington.

This document reports the results of a special investigation conducted by the PNL Surface Environmental Surveillance Project. Supplemental monitoring of tritium (hydrogen-3) in the Columbia River

was conducted in the summers of 1987 and 1988. The purpose of the monitoring was to provide information related to the dispersion and distribution of Hanford-originating contaminants entering the river through the seepage of groundwater along the Hanford Site.

Dirkes, R. L. 1994. *Summary of Radiological Monitoring of Columbia River Water along the Hanford Reach, 1980 through 1989*. PNL-9223, Pacific Northwest Laboratory, Richland, Washington.

A portion of PNL's Surface Environmental Surveillance Project is involved with monitoring the Columbia River. This document summarizes the river water monitoring activities of the Columbia River monitoring program during the 1980s. Routine and special monitoring projects and radiological and chemical constituents are reviewed. This report summarizes the information presented in the annual environmental reports.

Dirkes, R. L., G. W. Patton, and B. L. Tiller. 1993. *Columbia River Monitoring: Summary of Chemical Monitoring Along Cross Sections at Vernita Bridge and Richland*. PNL-8654, Pacific Northwest Laboratory, Richland, Washington.

Chemical monitoring was performed by PNL's Surface Environmental Surveillance Project at the Vernita Bridge and the Richland Pumphouse. Potential Hanford-originating chemicals of interest were selected for sampling; these included volatile organic compounds (VOCs), metals, and anions. Monthly samples were taken from August 1991 to December 1991. The sample frequency was reduced to quarterly during calendar year 1992. The monitoring results were benchmarked with those of the United States Geological Survey monitoring program, and no variants were found.

DOE - U.S. Department of Energy. 1992a. *Sampling and Analysis of 100 Area Springs*. DOE/RL-92-12, Rev. 1, U.S. Department of Energy, Richland, Washington.

This document provides validated monitoring data from the sampling of the Columbia River, seeps, springs, and sediment adjacent to the Hanford 100 Areas National Priorities List Site. The data were published as part of a Tri-Party Agreement milestone to evaluate how the contaminated seeps and springs impact the Columbia River. An assessment of the data is included. Samples were collected in September and October 1991 during the normal low-flow period of the Columbia River. Twenty-six locations were sampled along a 37-kilometer (22-mile) stretch of the river, ranging from just upstream of the 100-B/C Area water intake to the old Hanford townsite.

DOE - U.S. Department of Energy. 1992b. *Hanford Site Groundwater Background*. DOE/RL-92-23, U.S. Department of Energy, Richland, Washington.

This report is a preliminary evaluation of data and information related to the natural composition of groundwater in the unconfined aquifer system beneath the Hanford Site. This information is to be used as a baseline for distinguishing the presence and significance of contamination in the groundwater. The relevant part of the aquifer evaluated extended from the surface waters that potentially recharge the aquifer to the uppermost portion of the underlying confined aquifer. Surface waters were found, in

general, to have lower concentrations of constituents than the springs, unconfined groundwater, and confined groundwater. The provisional background threshold levels of background constituent concentrations in groundwater that are indicated in this report are likely to be conservatively low.

DOE - U.S. Department of Energy. 1994a. *Hanford Site Background: Part 1, Soil Background for Nonradioactive Analytes*. DOE/RL-92-24, Rev. 2, Vol. 1 of 2, U.S. Department of Energy, Richland, Washington.

This document was written to support environmental restoration, waste management, and facilities operations activities at Hanford. The background composition of Hanford Site soil is characterized for the purposes of identifying soil contamination and as a baseline in risk assessment processes used to determine soil cleanup and treatment levels. The compositions of naturally occurring soil in the zone above the groundwater level have been determined for nonradioactive inorganic and organic analytes and related physical properties. The range of inorganic and organic analytes that can be expected in Hanford Site background soil is evaluated. The highest measured background concentrations occur in three volumetrically minor soil types, the most important of which is topsoil adjacent to the Columbia River, which are rich in organic carbon. The chemical composition of more than 170 soil samples from 22 places on the Hanford Site and 3 places adjoining the Hanford Site was determined for inorganic analytes in accordance with EPA protocols. Twelve of the samples were analyzed for volatile and semivolatile organic chemicals, as well as for pesticides and polychlorinated biphenyls (PCB). Samples were collected from September through November 1991.

DOE - U.S. Department of Energy. 1994b. *Annual Report for RCRA Groundwater Monitoring Projects at Hanford Site Facilities*. DOE/RL-93-88, Rev. 0, U.S. Department of Energy, Richland, Washington.

This report is an annual hydrologic evaluation of 20 RCRA groundwater monitoring projects and one nonhazardous waste facility at the Hanford Site. The interpretation of groundwater data collected at 30 waste management units between October 1992 and September 1993 is included. Also, recent groundwater quality evaluations for the 100 and 300 Areas and the entire Hanford Site are described. Widespread contaminants include nitrate, chromium, carbon tetrachloride, tritium (hydrogen-3), and other radionuclides.

Eslinger, P. W., L. R. Huesties, A. D. Maughan, T. B. Miley, and W. H. Walters. 1994. *Data Compendium for the Columbia River Impact Assessment*. PNL-9785, Pacific Northwest Laboratory, Richland, Washington.

This document provides a bibliography of sources of existing data on Columbia River contamination. Approximately 4,500 documents and 13 major databases are listed that potentially contain information about contaminants in the Columbia River due to Hanford activities. The bibliography was further refined to highlight 60 key documents that contain data or describe analyses important in evaluating the health of the Columbia River. The work was performed to meet the Tri-Party Agreement milestone number M-13-80.

Ford, B. H. 1993. *Groundwater Field Characterization Report for the 200 Aggregate Area Management Study*. WHC-SD-EN-TI-020, Westinghouse Hanford Company, Richland, Washington.

This report provides contaminant plume maps for the unconfined aquifer of the 200 East and 200 West groundwater aggregate areas. Data deficiencies are identified with recommendations for additional sampling and well drilling. Individual plumes are identified for arsenic, chromium, cyanide, fluoride, nitrate, carbon tetrachloride, chloroform, trichloroethylene, tritium (hydrogen-3), gross beta, cobalt-60, strontium-90, technetium-99, iodine-129, cesium-137, gross alpha, uranium, and plutonium.

Fowler, K. M., K. B. Miller, M. O. Hogan, and J. F. Donaghue. 1993. *Risk-Based Standards Chemicals of Interest Database Documentation*. DRAFT. Prepared for the U.S. Department of Energy by the Pacific Northwest Laboratory, Richland, Washington.

A comprehensive set of risk-based standards are needed by the U.S. DOE to conduct its waste management, environmental restoration, and decontamination and decommissioning activities. The first step in developing the standards was to gather information on hazardous and radioactive substances that are found as contaminants or that are stored at DOE facilities. Twenty-six DOE sites were surveyed for substances that are generated, used, or present. Sources of information included Superfund Amendments and Reauthorization Act (SARA) Title III reports, remedial investigation/feasibility study reports, and other miscellaneous sources. The radionuclide and chemical names and media type in which they were found (i.e., air, groundwater, sediment, soil, surface water, tank wastes, and not specified/available) are indicated, but no quantitative sampling results are provided in this document. A total of 326 radionuclides and chemicals were identified for the Hanford Site.

Hartman, M. J., and K. A. Lindsey. 1993. *Hydrogeology of the 100-N Area, Hanford Site, Washington*. WHC-SD-EN-EV-027, Westinghouse Hanford Company, Richland, Washington.

The report primarily describes the hydrologic units beneath the 100-N Area. It includes descriptions of primary contaminants of interest, including strontium-90 and tritium (hydrogen-3) associated with the liquid waste disposal sites, sulfate and sodium, and petroleum products associated with leaks and spills. A total of eight petroleum (diesel oil) spills are documented between 1966 and 1988. Following the 1966 leak, an interceptor trench was built to collect migrating diesel oil, where it was periodically burned. A significant amount of free petroleum apparently remains in the zone above groundwater level; as much as 45 centimeters (1.5 feet) of petroleum product has been observed floating on top of the water in some of the monitoring wells. The petroleum seems to appear on the water table following periods of recharge to the aquifer.

Law, A. G. 1990. *Status of Groundwater in the 1100 Area*. Correspondence No. 8900604B R4, Westinghouse Hanford Company, Richland, Washington.

This document provides the quarterly results from the Westinghouse Hanford Company operational groundwater monitoring program for five wells installed in the vicinity of the 1100 Area. Results for approximately 380 analytes are presented; all are essentially undetected or at background levels.

Peterson, R. E., and V. G. Johnson. 1992. *Riverbank Seepage of Groundwater Along the 100 Areas Shoreline, Hanford Site*. WHC-EP-0609, Westinghouse Hanford Company, Richland, Washington.

Data were obtained during environmental surveillance activities and remedial investigations to characterize the influence of contaminated groundwater on the Columbia River. Radionuclides and metals in the seepage, sediment associated with the seepage, and near-shore Columbia River water were sampled. Samples collected in September and October of 1991 are compared with data collected in 1984 and 1988, as well as nearby groundwater data.

Rowley, C. A. 1993. *100-N Area Underground Storage Tank Closures*. WHC-SD-EN-TI-136, Westinghouse Hanford Company, Richland, Washington.

This report describes removal/characterization actions concerning underground petroleum storage tanks in the 100-N Area undertaken from 1990 through 1992. Instances of leaks from underground connections are noted. No groundwater contamination was found resulting from these tanks.

Weiss, S. G. 1993. *100 Area Columbia River Sediment Sampling*. WHC-SD-EN-TI-198, Rev. 0, Westinghouse Hanford Company, Richland, Washington.

To determine whether radiological and chemical contaminants are present in the Columbia River, 44 sediment samples were collected from 28 locations in the Hanford Reach in the fall of 1992. The sand-sized and smaller sediment samples were analyzed for metals and radionuclides from the near-shore and shoreline. Three of the sample locations were upriver from Hanford. Sediment was collected at depths of 0-15 centimeters (0-6 inches) and 30-60 centimeters (12-24 inches) below the surface. Contamination from arsenic, chromium, copper, lead, and zinc was found. The arsenic, lead, and zinc contamination may not be of Hanford origin. Cesium-137 and europium-152 were the most frequently detected radionuclides.

Wells, D. 1994. *Radioactivity in Columbia River Sediments and their Health Effects*. Special Report, Washington State Department of Health, Olympia, Washington.

This document addresses the current human health effects of artificial radioactivity in the Columbia River sediment. The Columbia River sediment data from the early 1960s to the present were provided by state agencies, federal agencies, and academic researchers. The sediment samples were collected from the Hanford area to the estuaries and coastlines of Oregon and Washington. Samples include surface sediment and deeper sediment behind the dams of the lower Columbia River. Ecological risks were not evaluated; nor were the human health risks from sediment contaminated with radioactive materials entering the Columbia River at riverbank seeps and springs.

2.2 Hanford Environmental Information System

DOE - U.S. Department of Energy. 1994c. *HEIS - Hanford Environmental Information System*. For documentation supporting the HEIS database, see DOE/RL-93-24, 9 volumes, U.S. Department of Energy, Richland, Washington. Queried: August 24, 1994.

The Hanford Environmental Information System (HEIS) is an electronic database that consolidates the data gathered during environmental monitoring and restoration of the Hanford Site. Data stored in HEIS are collected under several regulatory programs. The basis of HEIS is individual sample data for air, biota, groundwater, soil, sediment, surface water, and miscellaneous materials. The HEIS system was queried for information about maximum contaminant concentrations in groundwater within 150 meters (500 feet) of the Columbia River.

2.3 Remedial Investigation/Feasibility Studies

The EPA is the lead regulatory agency for the *Comprehensive Environmental Response, Compensation, and Liability Act of 1980* (CERCLA). Under CERCLA, a specific process has been established to identify potentially hazardous sites, characterize site contamination, assess treatment technologies, and then design and construct the appropriate treatment facilities. The remedial investigation/feasibility study (RI/FS) portion of the process defined in CERCLA requires determining the nature and extent of the threat posed by a release of hazardous substances to the environment and evaluating proposed remedies. The RI/FS studies which contributed information to the CRCIA Project are:

DOE - U.S. Department of Energy. 1990a. *Remedial Investigation/Feasibility Study Work Plan for the 300-FF-5 Operable Unit, Hanford Site, Richland, Washington*. DOE/RL 89-14, U.S. Department of Energy, Richland, Washington.

The 300-FF-5 operable unit consists of the groundwater aquifer beneath the 300-FF-1, 300-FF-2, and 300-FF-3 source operable units and adjacent areas defined by the extent of the groundwater contamination. The scope of the 300-FF-5 operable unit RI/FS focuses on groundwater, soil, surface water/sediment and aquatic biota and considers all contaminant sources in the 300 Area that contribute to the existing groundwater contamination beneath the 300 Area and the surrounding environment. The sample data upon which the RI/FS is based appear to have been taken in the mid-1980s. Groundwater monitoring for metals began in 1985.

DOE - U.S. Department of Energy. 1990b. *Remedial Investigation/Feasibility Study Work Plan for the 300-FF-1 Operable Unit, Hanford Site, Richland, Washington*. DOE/RL 89-31, U.S. Department of Energy, Richland, Washington.

The purpose of the 300-FF-1 operable unit remedial investigation was to provide sufficient information to conduct the feasibility study by determining the nature and extent of the threat to public health and the environment posed by releases of hazardous substances from 300-FF-1, a process liquid operable unit that contains all the liquid waste disposal facilities within the 300 Area. Hazardous and radioactive

materials from this operable unit contribute to groundwater contamination. Soil sampling data are provided for radionuclides, inorganics, and an extensive list of organics. Monitoring of groundwater analytes was more limited.

2.4 Hanford Site Environmental Reports

Every year, beginning in 1957, a report is prepared that summarizes environmental data, which characterize the Hanford Site environmental management performance and demonstrate compliance status. These reports summarize the activities and results of monitoring by the Surface Environmental Surveillance Project. In recent years, data have been provided in separate volumes. Annual reports used in the development of this project include the following:

Bisping, L. E. 1994. *Hanford Site Environmental Data for Calendar Year 1993 - Surface and Columbia River*. PNL-9824, Pacific Northwest Laboratory, Richland, Washington.

Bisping, L. E., and R. K. Woodruff. 1993. *Hanford Site Environmental Data for Calendar Year 1992 - Surface and Columbia River*. PNL-8683, Pacific Northwest Laboratory, Richland, Washington.

Bisping, L. E. 1992. *Hanford Site Environmental Data 1991 - Surface and Columbia River*. PNL-8149, Pacific Northwest Laboratory, Richland, Washington.

Dirkes, R. L., R. W. Hanf, R. K. Woodruff, and R. E. Lundgren. 1994. *Hanford Site Environmental Report for Calendar Year 1993*. PNL-9823, Pacific Northwest Laboratory, Richland, Washington.

Woodruff, R. K., R. W. Hanf, and R. E. Lundgren. 1993. *Hanford Site Environmental Report for Calendar Year 1992*. PNL-8682, Pacific Northwest Laboratory, Richland, Washington.

Woodruff, R. K., R. W. Hanf, and R. E. Lundgren. 1992. *Hanford Site Environmental Report for Calendar Year 1991*. PNL-8148, Pacific Northwest Laboratory, Richland, Washington.

2.5 Limited Field Investigations

Limited Field Investigations (LFIs) are conducted as part of Tri-Party Agreement activities to identify those Hanford waste sites that are recommended to remain as candidates for interim remedial measures. The assessments include consideration of whether contaminant concentrations pose an unacceptable risk that warrants action through interim remedial measures.

Each LFI is conducted on a single Hanford operable unit (e.g., operable unit 100-HR-3). Operable unit is the term used to identify specific areas designated for cleanup. The number and first letter in the operable unit name indicate the location of the operable unit; operable unit 100-HR-3 is in the 100-H Area. Many of the column headings in Appendix A correspond to the operable unit name.

The LFI reports annotated in this section are available to the public. The following list of LFI reports are those identified by Westinghouse Hanford Company's Environmental Data Management Control as undergoing final review and so not yet available to the public:

<u>Operable Unit</u>	<u>Document Number</u>
100-FR-3	DOE\RL-93-83
100-FR-1	DOE\RL-93-02
100-NR-2	DOE\RL-93-81
100-BC-2	DOE\RL-94-42
100-HR-2	DOE\RL-94-53

DOE - U.S. Department of Energy. 1994d. *Limited Field Investigation Report for the 100-BC-1 Operable Unit*. DOE/RL-93-06, U.S. Department of Energy, Richland, Washington.

This study was initiated to characterize the liquid and sludge at disposal sites associated with the B Reactor in the 100-BC Area. Groundwater sampling data are contained in the LFI, 100-BC-5 (see below). Surface water and sediment sampling are not applicable to the 100-BC-1 area. Media were sampled for VOCs, semivolatiles, inorganics, metals, PCBs, pesticides, radionuclides, and physical properties. Sampling data were collected from April 1992 through July 1992.

DOE - U.S. Department of Energy. 1993a. *Limited Field Investigation Report for the 100-BC-5 Operable Unit*. DOE/RL-93-37, Draft A, U.S. Department of Energy, Richland, Washington.

This study was initiated to further characterize the groundwater contamination in the 100-BC Area. Groundwater, surface water, sediment, and soil sampling data are provided. Volatile constituent concentrations were of primary interest, but the media were also sampled for radionuclides, organics, inorganics, and physical properties. The LFI groundwater sampling data are reported for July 1992, October 1992, and January 1993.

DOE - U.S. Department of Energy. 1993b. *Limited Field Investigation Report for the 100-DR-1 Operable Unit*. DOE/RL-93-29, Draft A, U.S. Department of Energy, Richland, Washington.

The purpose of this study was to characterize the waste facility sites associated with the D Reactor and the water retention basin systems for both the D and DR Reactors and in the 100-DR Area. Soil sampling results are reported. Groundwater sampling data for this same region are contained in the LFI, 100-HR-3 (see below). Media were sampled for VOCs, semivolatiles, inorganics, metals, PCBs, pesticides, radionuclides, specific anions, hexavalent chromium, and physical properties. Samples were collected in March 1993.

DOE - U.S. Department of Energy. 1993c. *Limited Field Investigation Report for the 100-HR-1 Operable Unit*. DOE/RL-93-51, Draft A, U.S. Department of Energy, Richland, Washington.

This study was initiated to characterize the waste units associated with facility sites supporting the H Reactor in the 100-H Area. This document provides sludge, sediment, and soil sampling data. Groundwater sampling data are contained in the LFI, 100-HR-3 (see below). Media were sampled for VOCs, semivolatiles, inorganics, metals, PCBs, pesticides, radionuclides, and physical properties. The media were sampled from December 1991 through August 1992.

DOE - U.S. Department of Energy. 1993d. *Limited Field Investigation Report for the 100-HR-3 Operable Unit*. DOE/RL-93-43, Draft A, U.S. Department of Energy, Richland, Washington.

This study was initiated to further characterize the groundwater contamination in the 100-HR-3 operable unit, which is inclusive of three sub-areas: 100-D, 100-H, and the 600 Area between the D and H Reactor areas. This document provides groundwater, sediment and soil sampling data for radionuclides, volatile and semivolatile organic compounds, inorganics, and pesticides. Media were sampled from May 1992 through March 1993.

DOE - U.S. Department of Energy. 1994e. *Limited Field Investigation Report for the 100-KR-1 Operable Unit*. DOE/RL-93-78, Draft A, U.S. Department of Energy, Richland, Washington.

This document provides soil sampling data. Groundwater sampling data are contained in the LFI, 100-KR-4 (see below). Surface water and sediment sampling are not applicable to the 100-KR-1 operable unit. Media were sampled for VOCs, inorganics, metals, radionuclides, hexavalent chromium, and physical properties. Samples were taken from October 1992 through March 1993.

DOE - U.S. Department of Energy. 1994f. *Limited Field Investigation Report for the 100-KR-4 Operable Unit*. DOE/RL-93-79, U.S. Department of Energy, Richland, Washington.

This LFI was initiated to further characterize the groundwater contamination in the 100-KR area operable units: 100-KR-1, 100-KR-2, and 100-KR-3. In addition to the groundwater samples, other sampling data include surface water, sediment, soil, and aquatic biotic impacted by the KE and KW reactors. The media were sampled for VOCs, semivolatiles, inorganics, metals, pesticides, and radionuclides. Samples were collected in October 1991, September 1992, December 1992, March 1993, and June 1993.

2.6 Discrete Radioactive Particles and Other Direct Exposure Sources

In addition to the routine environmental monitoring documented in the Hanford Site annual reports, occasional special studies are performed to evaluate particular conditions. Key studies are described here.

Cooper, A. T., and R. K. Woodruff. 1993. *Investigation of Exposure Rates and Radionuclide and Trace Metal Distributions Along the Hanford Reach of the Columbia River*. PNL-8789, Pacific Northwest Laboratory, Richland, Washington.

This report documents the first major field study to investigate exposure rates along the Columbia River shoreline since the Sula (1980) investigation of 1979. Radionuclides and trace metals were surveyed between Priest Rapids Dam and north Richland. A smaller number of discrete radioactive particles were also noted.

EG&G Energy Measurements. 1990. *An Aerial Radiological Survey of the Hanford Site and Surrounding Area, Richland, Washington*. EGG-10617-1062, EG&G Energy Measurements, The Remote Sensing Laboratory, Las Vegas, Nevada.

EG&G used a radiation detection system in a helicopter to conduct a radiological survey of the Hanford area. The detection system was calibrated to suppress natural background radiation and therefore only detected sources of anthropomorphic gamma-emitting radioactivity. The aerial data are presented as isopleths overlaid onto maps of the Hanford Site. The aerial survey is an aid in locating areas with elevated exposure rates but does not stringently define contaminated areas.

Sula, M. J. 1980. *Radiological Survey of Exposed Shorelines and Islands of the Columbia River Between Vernita and the Snake River Confluence*. PNL-3127, Pacific Northwest Laboratory, Richland, Washington.

This report describes a radiological survey performed to evaluate the magnitude and distribution of radioactive contamination on the exposed shorelines of the Columbia River. External exposure rate measurements were made at nearly 30,000 locations. In addition, discrete particles of radioactive material were discovered. Discrete metallic flakes containing cobalt-60 were found. The highest areal density of particles was found on an island near D-reactor, although the presence of particles was indicated as far downriver as the survey extended.

Wade, C. D., and M. A. Wendling. 1994. *100-D Island USRADS Radiological Surveys Preliminary Report Phase II*. BHI-00-134, Bechtel Hanford, Inc., Richland, Washington.

This report describes the results of radiological surveys made in April 1994, over the upstream third of the island adjacent to the 100-D reactor area. The survey used the Ultrasonic Ranging and Data System. A significant note is that, "with a few exceptions, every area which was determined to be gamma elevated was sampled and the sampling removed the entire contamination present. In these locations, extremely small 'hot particles' were removed from the silt layer beneath the river rock." Analyses of these particles showed them to contain almost entirely cobalt-60 activity, between 0.4 and 22 microcuries each. A total of 103 particles were recovered from an area of about 5 hectares (12.5 acres).

2.7 National Environmental Policy Act (NEPA) Documents

Quantifying the potential for future releases of contaminants to the Columbia River from surplus facilities or waste sites requires a significant investigation, one which is beyond the scope of this report. However, several major environmental impact statements (EIS) concerning Hanford facilities and waste management practices have been written. Each of these reports contains evaluations of potential future conditions based on current or projected Hanford Site status.

DOE - U.S. Department of Energy. 1987. *Final Environmental Impact Statement, Disposal of Hanford Defense High-Level, Transuranic, and Tank Wastes, Hanford Site, Richland, Washington*. DOE/EIS-0113, U.S. Department of Energy, Washington, D.C.

This EIS addressed the selection and implementation of final disposal actions for high-level, transuranic, and tank wastes at Hanford. Although a decision on the existing single-shell tanks was ultimately deferred, this EIS provides descriptions of the potential releases of radionuclides to the groundwater, and ultimately the Columbia River, for each of the major waste categories at Hanford.

DOE - U.S. Department of Energy. 1989. *Decommissioning of Eight Surplus Production Reactors at the Hanford Site, Richland, Washington, Draft Environmental Impact Statement*. DOE/EIS-0119D, U.S. Department of Energy, Washington, D.C.

and

DOE - U.S. Department of Energy. 1992c. *Decommissioning of Eight Surplus Production Reactors at the Hanford Site, Richland, Washington, (Final Environmental Impact Statement)*. DOE/EIS-0119F, U.S. Department of Energy, Washington, D.C.

This EIS, together with its addendum which constitutes the final EIS, describes the potential future releases of radionuclides to groundwater, and ultimately the Columbia River, from decommissioning the eight original Hanford reactors (excluding N Reactor) and the associated fuel storage basins. The preferred alternative for disposal was selected to be one-piece removal of the reactors from the riverside and burial in the 200 Areas.

DOE - U.S. Department of Energy. 1990c. *Low-Level Burial Grounds Dangerous Waste Permit Application: Request for Exemption from Lined Trench Requirements for Submarine Reactor Compartments*. DOE/RL-88-20, Supplement 1, U.S. Department of Energy, Richland, Washington.

and

DOE - U.S. Department of Energy. 1992d. *Low-Level Burial Grounds Dangerous Waste Permit Application: Request for Exemption from Lined Trench Requirements and from Land Disposal Restrictions for Residual Liquid at 218-E-12B Burial Ground Trench 94*. DOE/RL-88-20, Supplement 1, Revision 1, U.S. Department of Energy, Richland, Washington.

These two reports discuss decommissioned, defueled naval submarine reactor compartments containing radioactivity caused by exposure of structural components to neutrons during normal operation of the submarines. After all the alternatives were evaluated in the U.S. Department of the Navy 1984 environmental impact statement (Navy 1984), land burial of the submarine reactor compartments was selected as the preferred disposal option. The reactor compartments currently are sent to Trench 94 of the Hanford 218-E-12B Burial Ground. In addition to radioactivity, the reactor compartments disposed contain lead and PCBs as hazardous constituents. Modeling results indicate that release of contaminants to the groundwater or surface water will not occur until after long periods of time and that even after reaching the groundwater, contaminants will not be in excess of current regulatory limits, such as drinking water standards.

DOE - U.S. Department of Energy. 1994g. *Hanford Remedial Action Draft Environmental Impact Statement*. DOE/DEIS-0222. U.S. Department of Energy, Washington, D.C.

This EIS provides estimates of long-term risk resulting from the current groundwater plumes existing beneath the Site, as well as projections of future risks from non-tank, non-operating-facility waste management units.

Navy - U.S. Department of the Navy. 1984. *Final Environmental Impact Statement on the Disposal of Decommissioned, Defueled Naval Submarine Reactor Plants*. U.S. Department of the Navy, Washington, D.C.

This EIS discusses various alternatives for disposal of the radioactive portions of decommissioned nuclear submarines, leading to the selection of the Hanford Site as the location for permanent disposal. Estimates are presented for potential future radiation doses resulting from these activities.

Rhoads, K., B. N. Bjornstad, R. E. Lewis, S. S. Teel, K. J. Cantrell, R. J. Serne, J. L. Smoot, C. T. Kincaid, and S. K. Wurstner. 1992. *Estimation of the Release and Migration of Lead Through Soils and Groundwater at the Hanford Site 218-E-12B Burial Ground*. PNL-8356 Vol. 1, Pacific Northwest Laboratory, Richland, Washington.

This report evaluates the potential for radioactive and nonradioactive lead to migrate from buried submarine reactor compartments to the Columbia River. The estimated time of arrival of the contaminant plume ranges from 60,000 years to 4 million years.

Rhoads, K., B. N. Bjornstad, R. E. Lewis, S. S. Teel, K. J. Cantrell, R. J. Serne, L. H. Sawyer, J. L. Smoot, J. E. Szecsody, M. S. Wigmosta, and S. K. Wurstner. 1994. *Estimation of the Release and Migration of Nickel Through Soils and Groundwater at the Hanford Site 218-E-12B Burial Ground*. PNL-9791, Pacific Northwest Laboratory, Richland, Washington.

This report evaluates the potential for radioactive and nonradioactive nickel to migrate from buried submarine reactor compartments to the Columbia River. The estimated time of arrival of the contaminant plume ranges from 60,000 years to 4 million years.

3.0 Composite List of Identified Radionuclides and Chemicals

A data matrix (see Appendix A) was developed using the information found in the documents listed in Section 2.0. All radionuclides and chemicals analyzed in surface water (the Columbia River, springs, and seeps), sediment, groundwater, and soil samples in the 100, 300, and 1100 Areas are included. The data matrix is a composite list of all detected and not detected (i.e., analyzed for but not detected), radionuclides and chemicals from the reviewed literature. Sampling data from 1980 through 1994 were considered.

3.1 Risk-Based Standards Database

The development of the data matrix began with all chemicals identified in the Risk-Based Standards Database (Fowler et al. 1993). The Risk-Based Standards Database is a list of hazardous and radioactive substances reportedly found as contaminants or that are stored at DOE facilities nationwide. There are a total of 326 radionuclide and chemical entries for the Hanford Site. The radionuclides and chemicals in the database are sorted by their presence in the following media: Columbia River water, groundwater, soil, air, tank waste, and sediment. A total of 120 organic compounds, 133 inorganics, and 73 radionuclides were identified. These data formed the early basis for the data matrix.

Duplicate entries were removed from the database. Three mixtures (diesel fuel, hydrocarbons, and kerosene) are included. The primary database references were consulted for the concentration detected for each media. However, it was not possible to confirm the presence of the organics from the primary references cited in the database. Additional sources were reviewed to obtain information on the organic constituents.

3.2 Environmental Sampling Data Reports

The chemical analytical and radioanalytical data collected and presented in published environmental sampling reports were compiled and are presented in the data matrix in Appendix A. These reports include LFI reports, qualitative risk assessments, RI/FS reports, RCRA groundwater monitoring, and special studies reports. The titles and summaries of these documents are contained in Section 2.0. The scope was limited to the 100, 300, and 1100 Areas because they are most likely to have current impact.

The names of all radionuclides and chemicals examined (including those reported as nondetected) were added to the data matrix (Appendix A). The reported maximum concentration or activity, by media, is noted along with the background value, its reference, and the operable unit or geographical area where the sampling occurred. A total of 568 and 560 analytes were reported to be tested for in groundwater/Columbia River and soil/sediment, respectively, in the reviewed literature.

Of the analytes tested, 73 were detected in groundwater or Columbia River water, and 92 were detected in soil and sediment. Many of the analytes found are naturally occurring in groundwater and soil or are present as a result of global radioactive fallout.

A separate data matrix in Appendix A was prepared for incorporation of data related to existing groundwater plumes in areas outside the area of primary interest (i.e., the 200 Areas and 600 Area groundwater plumes).

3.3 Detected Analytes

Table 3.1 lists the 73 radionuclides and chemicals detected and their maximum concentration or activity in groundwater and Columbia River water. These maximum values are used in the screening process described in Section 4.0. Table 3.2 lists the 92 radionuclides and chemicals detected and their maximum concentration or activity in sediment and soil. Table 3.3 lists the maximum concentration or activity reported in existing Hanford groundwater plumes away from the river.

The data on radionuclide activity in sediment were compared with values reported by the Washington State Department of Health (Wells 1994). All contaminants included in Wells (1994) were included in the tables.

Tables 3.1, 3.2, and 3.3 are used in the screening criteria described in Section 4.0.

Table 3.1. Maximum Detected Concentrations in the Columbia River and Groundwater in the Hanford Site 100, 300, and 1100 Areas Near the Columbia River, 1980-1994

	Name of Analyte	Concentration in	
		Surface Water	Groundwater
1	ACETONE	11 µg/L (a)	30 µg/L
2	ALUMINUM		4,810 µg/L
3	AMERICIUM 241		0.021 pCi/L (b)
4	AMMONIA		70 µg/L
5	AMMONIUM		1,630 µg/L
6	ANTIMONY		60 µg/L
7	ANTIMONY 125		20 pCi/L
8	ARSENIC	3.4 µg/L	17 µg/L
9	BARIUM	48.2 µg/L	719 µg/L
10	BERYLLIUM		6 µg/L
11	BERYLLIUM 7 (c)		
12	BIS(2-ETHYLHEXYL) PHTHALATE		50 µg/L
13	BISMUTH (c)		
14	BORON (c)		
15	CADMIUM		31 µg/L
16	CALCIUM	35,900 µg/L	302,000 µg/L
17	CARBON 14		23,000 pCi/L
18	CESIUM 134	0.012 pCi/L	
19	CESIUM 137	0.13 pCi/L	0.5 pCi/L
20	CHLORIDE	870 µg/L	122,000 µg/L
21	CHLOROFORM		42 µg/L
22	CHROMIUM	22 µg/L	1,950 µg/L
23	COBALT		8 µg/L
24	COBALT 60	0.011 pCi/L	140 pCi/L
25	COPPER	22 µg/L	516 µg/L
26	CYANIDE		21.1 µg/L
27	DICHLOROETHYLENE, 1,2-		200 µg/L
28	DICHLOROETHYLENE, 1,2-trans-		130 µg/L
29	EUROPIUM 154		2 pCi/L
30	FLUORIDE	150 µg/L	2,080 µg/L
31	HYDRAZINE		7 µg/L
32	IODINE 129	0.16 pCi/L	
33	IRON	463 pCi/L	37,300 µg/L
34	LEAD		173 µg/L
35	LITHIUM (c)		
36	MAGNESIUM	9,860 µg/L	55,000 µg/L
37	MANGANESE	22.8 µg/L	400 µg/L
38	MERCURY		8.9 µg/L
39	METHYL ETHYL KETONE		18 µg/L
40	METHYLENE CHLORIDE		3,040 µg/L

Table 3.1. (contd)

	Name of Analyte	Concentration in	
		Surface Water	Groundwater
41	NICKEL	31 µg/L	479 µg/L
42	NITRATE	480 µg/L	90,000 µg/L
43	NITRITE		60,000 µg/L
44	PHOSPHATE		3,240 µg/L
45	PLUTONIUM 238		0.01 pCi/L
46	PLUTONIUM 239		0.03 pCi/L
47	POTASSIUM	2,430 µg/L	11,300 µg/L
48	RADIUM 226		0.3 pCi/L
49	RUTHENIUM 106 + D		34.4 pCi/L
50	SELENIUM		17.2 µg/L
51	SILICON (c)		
52	SILVER		19 µg/L
53	SODIUM	13,800 µg/L	200,000 µg/L
54	STRONTIUM		310 µg/L
55	STRONTIUM 90	28 pCi/L	80,000 pCi/L
56	SULFATE	8,600 µg/L	600,000 µg/L
57	SULFIDE		3,000 µg/L
58	TECHNETIUM 99		2,270 pCi/L
59	TETRACHLOROETHYLENE		39 µg/L
60	THALLIUM		4 µg/L
61	THORIUM 228		3 pCi/L
62	THORIUM 232		44.5 pCi/L
63	TITANIUM (c)		
64	TOLUENE	4.7 µg/L	2.9 µg/L
65	TRICHLOROETHYLENE		24.1 µg/L
66	TRITIUM (HYDROGEN 3)	4,430 pCi/L	1,900,000 pCi/L
67	URANIUM 233		3.3 pCi/L
68	URANIUM 234	18 pCi/L	120 pCi/L
69	URANIUM 235	0.01 pCi/L	17 pCi/L
70	URANIUM 238	19 pCi/L	93 pCi/L
71	VANADIUM		40 µg/L
72	XYLENE	4 µg/L	
73	ZINC	11 µg/L	8,800 µg/L
	(a) µg/L = micrograms per liter.		
	(b) pCi/L = picocuries per liter.		
	(c) Concentrations of these chemicals fall within their respectively occurring background levels.		

Table 3.2. Maximum Detected Concentrations in Soil and Sediment in the Hanford Site 100, 300, and 1100 Areas, 1980-1994

	Name of Analyte	Concentration in	
		Soil	Sediment
	Radionuclides		
1	AMERICIUM 241	34 pCi/g (a)	
2	ANTIMONY 124		1.2 pCi/g
3	CARBON 14	34 pCi/g	
4	CESIUM 134	0.04 pCi/g	0.29 pCi/g
5	CESIUM 137	2,900 pCi/g	6 pCi/g
6	COBALT 60	18,000 pCi/g	4.9 pCi/g
7	EUROPIUM 152	59,000 pCi/g	2.41 pCi/g
8	EUROPIUM 154	20,000 pCi/g	0.24 pCi/g
9	EUROPIUM 155	6,200 pCi/g	0.32 pCi/g
10	NEPTUNIUM 237		0.606 pCi/g
11	NICKEL 63	20,000 pCi/g	
12	PLUTONIUM 238	11 pCi/g	0.00115 pCi/g
13	PLUTONIUM 239	230 pCi/g	0.071 pCi/g
14	PLUTONIUM 240	(w/Pu239) (b)	
15	POTASSIUM 40	16 pCi/g	23 pCi/g
16	RADIUM 226	3.09 pCi/g	1.7 pCi/g
17	STRONTIUM 90	950 pCi/g	207 pCi/g
18	TECHNETIUM 99	0.67 pCi/g	0.5 pCi/g
19	THORIUM 228	1.61 pCi/g	3 pCi/g
20	THORIUM 232	1.1 pCi/g	3.2 pCi/g
21	THORIUM 234	ND (c)	0.812 pCi/g
22	TRITIUM (HYDROGEN 3)	1,600 pCi/g	
23	URANIUM 233	3.9 pCi/g	2.3 pCi/g
24	URANIUM 234		3.9 pCi/g
25	URANIUM 235	1.23 pCi/g	0.1 pCi/g
26	URANIUM 238	4.7 pCi/g	3.2 pCi/g
27	ZINC 65	ND	0.24 pCi/g
28	ZIRCONIUM 95	0.56 pCi/g	
	Chemicals		
29	ACENAPHTHENE	210 µg/kg (d)	
30	ALUMINUM	26,700,000 µg/kg	9,350,000 µg/kg
31	AMMONIA	12,800 µg/kg	12,000 µg/kg
32	ANTHRACENE	430 µg/kg	
33	AROCLOR 1248 (PCB)	9,900 µg/kg	
34	ARSENIC	47,000 µg/kg	7,500 µg/kg
35	BARIUM	672,000 µg/kg	120,000 µg/kg
36	BENZENE	4,500 µg/kg	
37	BENZO(G,H,I)PERYLENE	410 µg/kg	
38	BENZO(a)ANTHRACENE	940 µg/kg	
39	BENZO(a)PYRENE	810 µg/kg	
40	BENZO(b)FLUORANTHENE	890 µg/kg	
41	BENZO(k)FLUORANTHENE	760 µg/kg	
42	BENZOIC ACID	1,700 µg/kg	
43	BERYLLIUM	8,000 µg/kg	1,100 µg/kg
44	BIS(2-ETHYLHEXYL) PHTHALATE	68,000 µg/kg	
45	CADMIUM	1,800 µg/kg	2,700 µg/kg
46	CALCIUM	40,800,000 µg/kg	4,460,000 µg/kg
47	CHLORDANE	4,500 µg/kg	
48	CHLORIDE	1,100 µg/kg	

Table 3.2. (contd)

	Name of Analyte	Concentration in	
		Soil	Sediment
49	CHLORINE (e)		
50	CHROMIUM	259,000 µg/kg	12,200 µg/kg
51	CHRYSENE	920 µg/kg	
52	COBALT	34,100 µg/kg	11,500 µg/kg
53	COPPER	140,000,000 µg/kg	40,000 µg/kg
54	CYANIDE	1,050 µg/kg	
55	DIBENZOFURAN	130 µg/kg	
56	DIESEL FUEL	2,800,000 µg/kg	
57	ENDRIN ALDEHYDE	3.3 µg/kg	
58	ETHYL BENZENE	32,000 µg/kg	
59	FLUORANTHENE	1,800 µg/kg	
60	FLUORENE	190 µg/kg	
61	FLUORIDE	4,700 µg/kg	
62	FLUORINE (e)		
63	INDENO(1,2,3-CD)PYRENE	520 µg/kg	
64	IRON	33,500,000 µg/kg	71,000,000 µg/kg
65	KEROSENE	3,085,000 µg/kg	
66	LEAD	540,000 µg/kg	73,000 µg/kg
67	LITHIUM (e)		
68	MAGNESIUM	11,600,000 µg/kg	7,600,000 µg/kg
69	MANGANESE	839,000 µg/kg	578,000 µg/kg
70	MERCURY	4,300 µg/kg	
71	METHYL-2-PENTANONE, 4-	22,000 µg/kg	
72	METHYLENE CHLORIDE	120 µg/kg	
73	METHYLNAPHTHALENE, 2-	42 µg/kg	
74	NICKEL	221,000 µg/kg	19,700 µg/kg
75	NITRATE	30,400 µg/kg	
76	PHENANTHRENE	1,500 µg/kg	
77	POTASSIUM	4,980,000 µg/kg	1,900,000 µg/kg
78	PYRENE	1,200 µg/kg	
79	SELENIUM	4,200 µg/kg	
80	SILVER	1,900 µg/kg	2,500 µg/kg
81	SILVER CHLORIDE	17,300,000 µg/kg	
82	SODIUM	1,770,000 µg/kg	920,000 µg/kg
83	STRONTIUM	67,000 µg/kg	
84	STRONTIUM CHLORIDE	1 µg/kg	
85	SULFATE (SULFUR)	131,000 µg/kg	
86	TITANIUM (e)		
87	TOLUENE	350,000 µg/kg	
88	TOTAL PETROLEUM HYDROCARBON	1.26E + 08	
89	VANADIUM	389,000 µg/kg	82,200 µg/kg
90	XYLENE	1,800,000 µg/kg	
91	ZINC	309,000 µg/kg	397,000 µg/kg
92	ZIRCONIUM (e)		
	(a) pCi/g = picocuries per gram.		
	(b) w/Pu239 = concentration included in that reported for plutonium-239.		
	(c) ND = not detected.		
	(d) µg/kg = micrograms per kilogram.		
	(e) Concentrations of these chemicals fall within their respectively occurring background levels.		

Table 3.3. Maximum Detected Concentrations in Groundwater in the Hanford Site 100, 200, and 600 Areas Away from the Columbia River, 1980-1994

Name of Analyte	Number of Plumes	Concentration
100 Areas		
Chromium (+ 6)	3	1,570 ppb
Nitrate	10	130,000 ppb
Strontium-90	8	1,800 pCi/L
Tritium (Hydrogen-3)	4	80,000 pCi/L
200 West Area		
Arsenic	4	24 ppb
Carbon Tetrachloride	1	6,559 ppb
Chloroform	2	1,595 ppb
Chromium	5	323 ppb
Fluoride	3	10,067 ppb
Iodine-129	2	30 pCi/L
Nitrate	5	1,322,000 ppb
Technetium-99	5	26,602 pCi/L
Trichloroethylene	3	32 ppb
Tritium (Hydrogen-3)	3	6,193,000 pCi/L
Uranium	4	1,616 pCi/L
200 East Area		
Arsenic	4	24ppb
Cesium-137	1	1,326 pCi/L
Chloroform	1	7 ppb
Chromium	4	288 ppb
Cobalt-60	2	440 pCi/L
Cyanide	2	893 ppb
Iodine-129	3	20 pCi/L
Nitrate	7	397,000 ppb
Plutonium-239/240	1	69 pCi/L
Strontium-90	5	5,149 pCi/L
Technetium-99	2	22,163 pCi/L
Tritium (Hydrogen-3)	5	4,126,000 pCi/L
Uranium	1	27 pCi/L
600 Area (Solid Waste Landfill Site)		
Chloroform	1	0.5 ppb
Dichloroethane, 1, 1-	1	7 ppb
Tetrachloroethene	1	12 ppb
Trichloroethane, 1, 1, 1-	1	50 ppb
Trichloroethene	1	7 ppb
(a) pCi/L = picocuries per liter.		
(b) ppb = parts per billion.		

4.0 Screening Approach

The review of the available data indicated that concentrations of various radionuclides, carcinogenic chemicals, and hazardous chemicals had been measured in Columbia River water (Columbia River, springs, and seeps), groundwater, river sediment, and near-river soil. A multi-stage screening process to prioritize these various contaminants in terms of human health risk and ecosystem risk was developed. Each stage of the process identifies contaminants of interest. The combined results of the entire screening process then compose the total list of contaminants of concern.

The conceptual model for human health risk is associated with a scenario of a dedicated river user. The reference screening exposure scenario involves a person who frequents the shores of the river, drinks 2 liters/day of untreated river water, consumes about 0.25 kilograms/day (100 kilograms/year) (CRITFC 1994) of freshwater fish, and has an incidental sediment ingestion rate of 10 milligrams/day (almost 4 grams/year). This conceptual model is an adaptation and expansion of the Hanford Site risk assessment methodology (DOE 1992e).

The conceptual models for ecosystem risk are simpler, relying on the EPA Ambient Water Quality Criteria (EPA 1992) and on a fraction of the concentrations that result in mortality for fish.

All analytes found in the reviewed literature, which related to the 100, 300, and 1100 Areas, regions along the banks of the Columbia River, or inland contaminant plumes, were compiled (see Appendix A). Initial screening eliminated the contaminants on the list that showed no detectable levels of activity or concentration. In addition, analytes which were present only in tank wastes and not in environmental media were eliminated from the study.

4.1 Screening Equations

The screening process operates on one portion of the available data at a time. Separate screenings are used for measurements in Columbia River water, groundwater, river sediment, and near-river soil. Within each of these divisions, further subdivisions address radionuclides, carcinogens, human toxins, and fish toxins. All of the screenings rely on river water concentration or a surrogate as a starting point. Procedures for estimating the surrogates are described below.

4.1.1 Radionuclide Screening

The screening is based on a scenario of exposure to a dedicated river user (see definition above). Internal risks are estimated using the EPA slope factor for ingestion (EPA 1994a). The EPA slope factor represents the lifetime excess total cancer risk per unit of intake. External exposure to contaminated sediment is addressed by assuming the parameters associated with the EPA slope factor for external exposure are appropriate (EPA 1994a).

A relationship between the concentration of the contaminant in the water and the concentration in the sediment is required. For the screening, this relationship is assumed to be described by a ratio of 1:100,000 (i.e., the concentration of the contaminant in the sediment is assumed to be 100,000 times

higher than in the Columbia River waters). This assumption is based on review of the very limited number of samples for which both river water and sediment values were available, as well as on an empirical equation developed for radionuclides in the Columbia River incorporated in the GENII computer code (Napier et al. 1988, p. 4.82).

The screening equation for radionuclides is:

$$\text{SCREEN} = C_w \left[\frac{100,000 * SS}{1000} + (730 + 100 * \text{BCF} + 100,000 * 0.0036) * \text{IS} \right] \quad (1)$$

where C_w = measured or surrogate water concentration, pCi/L
 100,000 = sediment/water ratio, L/kg
 SS = radionuclide slope factor for external exposure, risk/year per pCi/g
 1000 = unit conversion, g/kg
 730 = water consumption of 2 L/day for 1 year
 100 = fish consumption of 100 kg/year
 BCF = bioconcentration factor for fish, L/kg
 0.0036 = sediment consumption of 10 mg/day, giving 3.6 g/year
 IS = radionuclide slope factor for ingestion, risk/pCi.

Values resulting from this screening which approach or are greater than 10^{-6} imply radionuclides of potential concern.

4.1.2 Carcinogenic Chemical Screening

The conceptual exposure patterns for carcinogens in river water are the same as those for radionuclides; however, there is no factor for external exposure. Because the chemical cancer potency factors for oral exposure are in units of inverse milligram per kilogram per day, the consumption terms are put in daily, rather than annual, units (EPA 1994a).

$$\text{SCREEN} = C_w [2 + 0.27 * \text{BCF} + 100,000 * 1 \times 10^{-5}] (0.001) \frac{\text{CPF}}{70} \quad (2)$$

where C_w = measured or surrogate water concentration, $\mu\text{g/L}$
 2 = water consumption of 2 L/day
 0.27 = consumption of 100 kg/year of fish, on a daily basis 0.27 kg
 BCF = bioconcentration factor for fish, L/kg
 100,000 = sediment/water ratio, L/kg
 1×10^{-5} = consumption of 10 mg/day of sediment, kg
 0.001 = conversion factor, micrograms to milligrams
 CPF = cancer potency factor, $(\text{mg/kg/day})^{-1}$
 70 = assumed weight of an adult, 70 kg.

Values resulting from this screening which approach or are greater than 10^{-6} imply chemicals of potential concern.

4.1.3 Toxic Chemical Screening

For hazardous, but noncarcinogenic, chemicals, the ranking is based on a ratio of the estimated daily intake to the EPA chronic oral reference dose (EPA 1994a). The conceptual scenario is the same as for the radionuclides or carcinogens.

$$\text{SCREEN} = C_w [2 + 0.27 * \text{BCF} + 100,000 * 1 \times 10^{-5}] \frac{(0.001)}{70 * \text{RfD}} \quad (3)$$

where C_w = measured or surrogate water concentration, $\mu\text{g/L}$
2 = water consumption of 2 L/day
0.27 = consumption of 100 kg/year of fish, on a daily basis 0.27 kg
BCF = bioconcentration factor for fish, L/kg
100,000 = sediment/water ratio, L/kg
 1×10^{-5} = consumption of 10 mg/day of sediment, kg
0.001 = conversion factor, micrograms to milligrams
70 = assumed weight of an adult, 70 kg
RfD = EPA chronic oral reference dose, mg/kg/day.

Values resulting from this screening which approach or are greater than unity imply chemicals of potential concern.

4.1.4 Ambient Water Quality Criteria Screening

For aquatic biota, the measured or surrogate concentration of the contaminant in water is compared with the applicable EPA water quality criterion (EPA 1992). The ambient water quality criteria are values of the concentrations of chemicals in water that are considered by the EPA to be protective of aquatic life. The screening equation is

$$\text{SCREEN} = \frac{C_w}{\text{AWQC}} \quad (4)$$

where C_w = measured or surrogate water concentration, pCi/L
AWQC = ambient water quality criterion, $\mu\text{g/L}$.

Values resulting from this screening which approach or are greater than unity imply chemicals of potential concern.

4.1.5 Aquatic Biota Toxicity Screening

Limited data were available that identify the concentrations of certain chemicals that result in toxic effects to aquatic life. Where possible, the threshold concentration for fresh water at which any effect was noted was used. Although it would have been preferable to use information that related directly to the initiation of distress in aquatic life, rather than mortality, such information (e.g., the threshold limit value for the medium) was available for only a few chemicals. Therefore, the lowest concentration

lethal to 50 percent of small, freshwater fish (e.g., guppies, mosquito fish, rainbow trout) tested was also used (EPA 1985). To relate these lethal effects to less significant effects, the screening used a value of 1 percent of the LC50 in the determination. For a few analytes for which fish data were not available, test results for crayfish or insects were used as a surrogate. The equation is

$$\text{SCREEN} = \frac{C_w}{(\text{LD50} / 100)} \quad \text{else} \quad \frac{C_w}{\text{TLM}} \quad (5)$$

where C_w = measured or surrogate water concentration, pCi/L
LD50 = concentration of contaminant lethal to 50 percent of the tested fish population in time periods ranging from 48 to 96 hours (LC_{50}), $\mu\text{g/L}$
TLM = threshold limit for fresh water (TLM), $\mu\text{g/L}$.

Values using this screening approach or values greater than unity imply chemicals of potential concern.

A concern has been raised that groundwater, filtering through gravel beds into the waters of the Columbia River, could directly impact fish eggs laid in the gravels without prior dilution by Columbia River water. Sources of data related to the impact of the listed contaminants on fish eggs were sought. Very few positive connections between research on fish egg survival and contaminant concentrations were found, making it impossible to screen directly on this concept.

4.2 Estimation of Contaminant Concentrations in River Water

All of the screening equations presented in the preceding section require an estimate of the contaminant's maximum measured concentration in river water. Only the direct river measurements provide this information. For the other media, an estimated, surrogate water concentration must be developed. Radionuclide concentrations compiled were generally given in units of picocuries/liter or picocuries/gram. Chemical concentrations were standardized to units of micrograms/liter or micrograms/kilogram. Therefore, separate conversions were developed for radionuclides and chemicals.

4.2.1 Radionuclides

Separate sets of assumptions were needed to prepare screening surrogates for concentrations in river water for measurements in groundwater, river sediment, and near-river soil.

4.2.1.1 Groundwater

Groundwater adjacent to the Columbia River can flow into the river, and Columbia River water can flow into the groundwater, depending on river flow. Therefore, concentrations of contaminants in groundwater near the river are difficult to predict, and concentrations measured near the shore differ from those measured further inland. Flow rates from groundwater to the Columbia vary from location to location; individual springs may have very low flow rates. An average groundwater discharge to the Columbia River of 3 cubic feet per second (cfs) was modeled by Kipp et al. (1976) for a 8.3-kilometer

(5-mile) length of the river near the Hanford townsite. Raymond et al. (1976) and Cline et al. (1985) report an estimated discharge of 100 cfs over the entire Hanford Reach. More recent research (Wuestner and Devary 1993) indicates that 100 cfs is an upper bound. For conservatism (i.e., to provide an overestimate of the resulting concentration in the river), this upper value of 100 cfs was adopted for the screening. In effect, this implies that the entire volume of groundwater that flows from beneath Hanford to the Columbia River is contaminated to the maximum level reported. Thus, the conversion used is

$$C_w^o = C_{gw} * \frac{100}{100,000} \quad (6)$$

where C_w^o = surrogate river water concentration used in the screening, pCi/L
 C_{gw} = measured groundwater concentration, pCi/L
 100 = groundwater discharge rate, cfs
 100,000 = approximate annual average flow rate of the Columbia River at Hanford, cfs.

4.2.1.2 River Sediment

Sediment within the river is both a reservoir of contaminants and a source of contamination of the river water, as the material desorbs or resuspends into the water column. Accurate representation of this process requires detailed knowledge of the chemical interactions of the contaminant and the water. Information at this level of detail is not available for most of the contaminants considered. For consistency with the dose estimation assumptions, this relationship is assumed to be described by an equilibrium ratio of 1:100,000 (i.e., the concentration of the contaminant in the sediment is assumed to be 100,000 times higher than in the Columbia River water). The conversion used is then

$$C_w^o = \frac{C_{sed} * 1000}{100,000} \quad (7)$$

where C_w^o = surrogate river water concentration used in the screening, pCi/L
 C_{sed} = sediment concentration, pCi/g
 1000 = unit conversion, g/kg
 100,000 = assumed concentration ratio, L/kg.

4.2.1.3 Near-River Soil

Contaminants in waste sites or other sites adjacent to the Columbia River may not pose a current hazard to down-river users of the river, but they may pose a threat of future contamination of the river. The possibility also exists that such sources may be contributing as-yet undetected contamination to the river. One of the goals of the Columbia River Comprehensive Impact Assessment is to tie Hanford cleanup activities to the potential for river contamination. In this spirit, contaminated soil near the river is included as a possible source of contaminants. Adequate consideration of these contaminants must include site-specific details about how they could be transported from their current locations into the groundwater and hence into the Columbia River. For the purpose of screening, all contaminants are assumed to be environmentally mobile and potentially soluble in groundwater (contrast this assumption to that used for contaminants in sediment, where they are assumed to be tightly bound).

Based on this assumption, the surrogate groundwater contamination is assumed to have the same concentration of contaminants as the soil. The total area of industrial activity comprises approximately 6 percent of the Hanford Site (Dirkes et al. 1994, p. 5). Because it is unreasonable to assume that all of Hanford soil is contaminated to the maximum concentration reported, an effective area of 1 percent is assumed. The set of assumptions used to convert groundwater to river water concentrations is then also applied. The resulting equation for surrogate river water concentration resulting from soil is

$$C_w^o = C_{\text{soil}} * \frac{(1000 * 1 * 100 * 0.01)}{100,000} \quad (8)$$

where C_w^o = surrogate river water concentration used in the screening, pCi/L
 C_{soil} = concentration in soil, pCi/g
 1000 = unit conversion, g/kg
 1 = assumption of soil/groundwater concentration equivalency, kg/L
 100 = groundwater discharge rate, cfs
 0.01 = fraction of total area contaminated, dimensionless
 100,000 = approximate annual average flow rate of the Columbia River at Hanford, cfs.

4.2.2 Chemicals

Conversions from measured values to surrogate river water concentrations are also required for carcinogenic and hazardous chemical contaminants. The assumptions are the same as for radionuclides; however, the measured units are generally in micrograms/kg, rather than pCi/g, and some conversions differ by factors of 1000.

4.2.2.1 Groundwater

The conversion is numerically identical to that for radionuclides:

$$C_w^o = C_{\text{gw}} * \frac{100}{100,000} \quad (9)$$

where C_w^o = surrogate river water concentration used in the screening, $\mu\text{g/L}$
 C_{gw} = measured groundwater concentration, $\mu\text{g/L}$
 100 = groundwater discharge rate, cfs
 100,000 = approximate annual average flow rate of the Columbia River at Hanford, cfs.

4.2.2.2 River Sediment

The conversion is similar to that for radionuclides with the g/kg conversion removed:

$$C_w^o = \frac{C_{\text{sed}}}{100,000} \quad (10)$$

where C_w^o = surrogate river water concentration used in the screening, $\mu\text{g/L}$
 C_{sed} = sediment concentration, $\mu\text{g/kg}$
 100,000 = assumed concentration ratio, L/kg.

4.2.2.3 Near-River Soil

The conversion is similar to that for radionuclides with the g/kg conversion removed:

$$C_w^o = C_{\text{soil}} * \frac{(1 * 100 * 0.01)}{100,000} \quad (11)$$

where C_w^o = surrogate river water concentration used in the screening, $\mu\text{g/L}$
 C_{soil} = concentration in soil, pCi/g
 1 = assumption of soil/groundwater concentration equivalency, kg/L
 100 = groundwater discharge rate, cfs
 0.01 = fraction of total area contaminated, dimensionless
 100,000 = approximate annual average flow rate of the Columbia River at Hanford, cfs.

4.3 Screening Results

Application of the equations and assumptions defined above results in a series of complementary, but not necessarily intercomparable, screening values for each contaminant. The varying numbers of assumptions and associated varying degrees of conservatism require that each of the screenings be evaluated separately. The results of the combined screenings, however, then define the overall list of contaminants of interest. The complete list of radionuclides and chemicals entered into the project database is presented in Appendix A. The parameters used in the calculation are presented in Appendix B. The complete numerical results are presented in Appendix C. The overall results and interpretation of the screening are given here.

During the screening process, a few radionuclides and chemicals were identified as of potential interest, but not carried forward. Some items were measurements determined to be within the naturally occurring background levels of these materials. These materials included the radionuclides beryllium-7 and potassium-40 and the chemicals barium, bismuth, boron, chlorine, fluorine, lithium, silicon, silver, sulfide, titanium, vanadium, and zirconium. In addition, several materials were identified by the screening process that the EPA (EPA 1991; EPA 1989) considers nonhazardous under environmental conditions. These materials removed from further consideration included aluminum, calcium, iron, magnesium, potassium, and sodium.

4.3.1 River Water Sample Screening

Of the thousands of available environmental samples, relatively few show positive identification of contaminants directly in the waters of the Columbia River. A screening level was used to account for over 1) 95 percent of the carcinogenic risk for each result, above a cutoff of 10^{-6} , or 2) a non-carcinogenic hazard ranking of greater than 0.1. The individual screenings and the contaminants identified via each are listed in Table 4.1.

Table 4.1. Contaminants of Potential Interest Identified via Screening of Columbia River Samples

Radionuclide Screening	Carcinogenic Chemical Screening	Hazard Index Screening	Ambient Water Quality Criteria Screening	Aquatic Toxicant Screening
Cesium-134	Arsenic	Arsenic	Copper ^(a)	Arsenic
Cesium-137		Copper ^(a)	Nickel ^(a)	Copper ^(a)
Cobalt-60		Manganese	Zinc	Nickel ^(a)
		Nickel ^(a)		Nitrate
	Nitrate		Xylene ^(b)	
	Toluene ^(b)		Zinc	
	Xylene ^(b)			
	Zinc			

(a) See discussion in Section 4.4 on samples near limit of detection.
(b) See discussion in Section 4.4 on suspect samples.

The two isotopes of radiocesium, cesium-134 and cesium-137, are present in worldwide fallout. It is likely that these two contaminants are largely derived from non-Hanford sources. The Hanford Environmental Dose Reconstruction Project did not identify these two radionuclides as resulting from significant Hanford releases (Napier 1993).

Several contaminants are highlighted in Table 4.1 with footnotes. These indicate a potential problem with the screening result on the basis of source information. These difficulties are described in Section 4.4.

4.3.2 Groundwater Sample Screening

A very large fraction of available Hanford-related environmental samples are of groundwater. Only those taken within about a kilometer of the river were used in compiling the database used for the screening. Even so, many positive samples were noted. Most of the samples were derived from investigations of the Hanford operating areas (100, 300), but many were from wells located near the river but far from the reactor, fuel fabrication, and research sites. Contaminants identified for investigation include several metals. The individual screenings and the contaminants identified via each are listed in Table 4.2.

4.3.3 River Sediment Sample Screening

Because the Hanford Reach is a relatively fast-flowing portion of the river, there is actually little accumulation of sediment at Hanford. Accordingly, sediment samples represent a very small portion of the historical Hanford data. This is a clear area for future sampling work. Nevertheless, the sediment

samples did provide sufficient information to apply the screening technique. The individual screenings and the contaminants identified via each are listed in Table 4.3. Like the river water screening, this process identified two isotopes of cesium, both of which are most likely associated with global fallout.

Table 4.2. Contaminants of Potential Interest Identified via Screening of Groundwater Near the Columbia River

Radionuclide Screening	Carcinogenic Chemical Screening	Hazard Index Screening	Ambient Water Quality Criteria Screening	Aquatic Toxicant Screening
Cobalt-60 Strontium-90	Chromium	Antimony Copper Mercury Nitrate/Nitrite Phosphate	Chromium Mercury	Chromium Copper Nitrate/Nitrite Zinc

Table 4.3. Contaminants of Potential Interest Identified via Screening of Columbia River Sediment Samples

Radionuclide Screening	Carcinogenic Chemical Screening	Hazard Index Screening	Ambient Water Quality Criteria Screening	Aquatic Toxicant Screening
Cesium-134 Cesium-137 Cobalt-60 Europium-152	Chromium	Arsenic Copper Lead Zinc	Chromium Lead	Chromium Zinc

4.3.4 Near-River Soil Sample Screening

Contaminants measured in soil near the Columbia River are generally not an immediate hazard because they are currently in the soil and not subject to mass transport to the river, and subsequent human and biotic exposure. However, their existence is the primary reason for continuing cleanup of the Hanford operating areas, and it is useful to have a screening prioritization. It is also useful to direct future sampling efforts to determine if any of the contaminants most likely to cause problems are beginning to reach the river. Because of the nature of the contamination (generally solids in or associated with soil) and the nature of the activities carried out at Hanford over its history, these contaminants differ somewhat from those actually found in more mobile media (river water, groundwater, and

sediment). Even so, it is informative to note the similarities in the list generated via the soil screening with those lists generated for the other media. The individual screenings and the contaminants identified via each are listed in Table 4.4.

Table 4.4. Contaminants of Potential Interest Identified via Screening of Soil Near the Columbia River

Radionuclide Screening	Carcinogenic Chemical Screening	Hazard Index Screening	Ambient Water Quality Criteria Screening	Aquatic Toxicant Screening
Cesium-137	Arochlor 1248 (PCB)	Arsenic	Arochlor 1248 (PCB)	Chlordane
Cobalt-60	Benzo(a)pyrene ^(a)	Chlordane	Chlordane	Mercury
Europium-152	Chromium	Copper	Chromium	Zinc
Europium-154	Indeno(1,2,3-CD) pyrene ^(a)	Lead	Copper	Diesel Fuel
		Mercury	Lead	
		Nitrate	Mercury	
		Silver Chloride		
		Zinc		
		Diesel Fuel		

(a) See discussion in Section 4.4.

4.4 Use of Suspect Measurements

The majority of the measurements taken over the past 15 years were collected in accordance with modern quality assurance procedures (Dirkes et al. 1994). The data from the references used in this report are traceable and of high quality. All data recorded in the referenced studies were used in the development of the screening approach reported here.

During the evaluation of tens of thousands of media samples for hundreds of analytes over a period of many years, it is statistically expected that an occasional analysis will result in incorrect identification of an analyte or its quantity. The quality assurance procedures in place on the major Hanford Site databases generally serve to identify these abnormal values. For scientific completeness, the reported values are generally included in the databases with an indicator that they are potentially spurious. In the course of the evaluations for this report, six potential constituents of concern with single, questionable, measured results were encountered with the potential to influence the selection criteria, two in soil and four in Columbia River water.

Two of the chemicals labeled with a footnote in Table 4.1 are toluene and xylene. These two chemicals were identified as coming from a single sample which may have been contaminated during sampling or analysis because these and other chemicals identified in that one sample are common laboratory and industrial solvents (Dirkes et al. 1993, p. 4.1). Since the suspect sample was paired with another suspect sample from upstream of Hanford, which also indicated high concentrations of organic contaminants, it is unlikely that these compounds are elevated in river water as a result of releases from Hanford.

Two other chemicals labeled with a footnote in Table 4.1 are copper and nickel. These two chemicals and several more identified in Table C.1 (see SW-LD notations) were very near the lower limits of detection in a series of samples at the Richland pumphouse (Dirkes et al. 1993). This reference compared concentrations of 20 volatile organic chemicals, 19 metals, and 7 anions upstream from Hanford (Vernita Bridge) and downstream (Richland). No volatile organic chemicals were routinely detected at either location. The concentrations of most metals were also very low. However, copper and nickel were each reported one time (out of nine sampling periods) as being slightly above the limit of detection. The limit of detection for copper for this study was 20 micrograms/liter. The single reported positive sample was 22 micrograms/liter. The limit of detection for nickel was 30 micrograms/liter. The single reported positive sample was 31 micrograms/liter. These values probably do not represent the actual level of river contamination.

Two chemicals labeled with a footnote in Table 4.4 are benzo(a)pyrene and indeno(1,2,3-CD)pyrene. Both of these chemicals appear only once in the database of samples, and both are analytes from the same physical sample. This one sample is noted in the historical record as being "suspect" because the analysis results for all contaminants evaluated were very high and not repeated in other nearby samples. It is likely that these two chemicals do not need to be on the master list for further evaluation.

5.0 Discrete Radioactive Particles

The presence of small, discrete particles of radioactive material was discovered by Sula during a shoreline survey in 1978-1979 (Sula 1980). In the 1978-1979 survey, Sula reported finding 188 discrete particles of contaminated material. The majority of the discrete particles were found buried in rocky, flat areas with little or no vegetation. Sula recovered 14 particles for special study. Laboratory analysis identified the gamma radiations emitted from the particles to be entirely due to cobalt-60, with activities ranging from 1.7 to 24 microcuries. Sula (1980, p. 36) describes the particles as

When isolated, the particles were barely visible to the naked eye, appearing as small, dark colored chips or flakes of roughly equal size. Microscopic examination of three particles showed them to be metallic appearing flakes with diameters of approximately 0.1 mm. The particles were found to vary in elemental composition, but all contained significant proportions of chromium, iron, and cobalt characteristic of the alloy stellite, used in valve and pump components in all of the production reactors.

Sula declined to predict how many particles exist in the Columbia River but did note that "the number of particles found per square meter of ground surveyed decreases as one travels downstream from the reactor areas" (Sula 1980, p. 36).

The next attempt to measure these particles came in 1993 (Cooper and Woodruff 1993). Although the area surveyed was somewhat less than that surveyed by Sula, the 1993 survey also found 11 particles: 10 on one island near the reactors and one further downstream. Two particles were recovered for further analysis. The activities of these two particles were 1.7 and 16 microcuries of cobalt-60.

Most recently, cleanup efforts have been initiated on the island closest to and downstream of the 100-D Area, the island noted in both the Sula and Cooper and Woodruff surveys as having the highest concentration of particles. To date, 103 particles have been recovered, with activities ranging from 0.13 to 22 microcuries of cobalt-60, and minor amounts of other Hanford radionuclides (Wade and Wendling 1994).

Cooper and Woodruff (1993) included an evaluation of the potential for radiation dose from inhalation or ingestion of a discrete particle and from external exposure. It is concluded that, although the possibility of inhalation is remote, the dose-limiting exposure pathway is the inhalation of a particle at the upper end of the range of activity that would remain lodged in the nasal passages for up to 48 hours, resulting in a dose about 10 times the limit for occupational exposure (NCRP 1989).

6.0 Direct Irradiation from Hanford Facilities

For the last several years, the highest direct radiation exposure rates from Hanford operations observed at locations where the public currently has access have been on the Columbia River along the shoreline at the 100-N Area (e.g., Dirkes et al. 1994). Thermoluminescent dosimeter measurements have been reported annually in the Hanford Site annual environmental reports for this location since 1990. The source of the elevated exposure rates is radiation from facilities located above the river in the 100-N Area. The shoreline is not currently accessible to the public, but the adjacent river is open to the public for recreational uses.

Elevated dose rates at the shoreline are reported in Dirkes et al. (1994, pp. 76, 168). The highest values were measured adjacent to the N Reactor itself and also near the 1301-N Liquid Waste Disposal Facility. The highest readings along the shoreline in 1994 ranged up to about 100 microroentgen/hour in an area where background exposure rates are in the range of 7-10 microroentgen/hour. Dirkes et al. (1994, p. 75) qualify this number to be a probable overestimate. The dose rates have fallen significantly since the closure of the N Reactor in 1988. Dose rates are also elevated near the 100-K Area because of radiologically contaminated materials such as internally contaminated ion-exchange modules used in maintaining water quality in the nearby 105-KE fuel storage basin. A third area of elevated exposure rates is adjacent to the 300 Area.

In 1993, measurements were also made by boat on the Columbia River adjacent to the N Reactor facilities, about 75 meters (250 feet) from the Hanford shoreline (Cooper and Woodruff 1993, p. 4.12-4.13). At this distance, the exposure rates along a 1500-meter (5000-foot) track parallel to the facility ranged from essentially background levels (5 microroentgen/hour) to about 20 microroentgen/hour. Exposure rates on the north shore of the river, across from N Reactor, were all essentially background.

In 1988, EG&G performed an aerial survey of direct exposure rates on the Hanford Site, including the Columbia River and adjacent facilities (EG&G 1990). A low-level, generalized increase in exposure rates is indicated for the shorelines of most of the river. The individual facilities are distinctly noticeable. The 100-N Area evidences the highest exposure rates of river locations.

7.0 Potential Future Groundwater Sources


Certain contaminants now in soil or groundwater distant from the Columbia River at Hanford may some time in the future pose a source of contamination to the river. Some distant contaminants are essentially certain to reach the river, and others are, at this time, only potential, in part because planned remedial actions will either immobilize or remove them. The contaminants that are already in groundwater are quite likely to reach the Columbia River in the future. Those contaminants contained in Hanford tank farms or burial grounds may not pose a future hazard. For the Columbia River Comprehensive Impact Assessment, only those currently in the groundwater as defined in Section 7.1 are considered. Brief reference is given in Section 7.2 to documentation of the other categories of materials.

7.1 Existing Groundwater Plumes

More than 105 plumes, containing 20 contaminants, are readily observable in groundwater beneath the Hanford Site (Ford 1993; DOE 1994b). A summary of the nature of the existing groundwater contaminant plumes, their general locations, and maximum measured concentrations is given in Table 3.3. Maps of these plumes are provided in Ford (1993), DOE (1994b), and Dirkes et al. (1994). (Note that each of the authors of these reports draws the outlines of the plumes somewhat differently, depending on the purpose of the reports.) An example of one of the most widely dispersed contaminants, nitrate, is shown in Figure 7.1 (Dirkes et al. 1994).

Because those existing contaminant plumes addressed in this section of the report are not in direct contact with the Columbia River, they do not yet constitute a source of contaminants in the river. The window for future concern varies depending both on the location of the plumes and the material in them. Groundwater travel times from the current location to discharge in the river vary by location. Travel times in the 100 Areas generally are less than 1 year. Travel times for groundwater carrying the plumes in the 200 East Area are generally in the range of 20 to 200 years. Travel times for the contaminants in the 600 Area evolving from the Central Landfill Site (see Figure 7.1) are probably about 10 years. Travel times for plumes in the 200-West Area may be as long as 80 to 300 years (Freshley and Graham 1988). All of these estimated times depend on future groundwater conditions and influences such as quantity of water discharged from Hanford operating facilities.

Most of the contaminants listed in Table 3.1 are relatively mobile in groundwater. However, cobalt-60, strontium-90, and cesium-137 have significant chemical interactions with the soil and move much more slowly than the groundwater. (They exist in the groundwater in the 200 Areas because they were essentially injected there directly during waste disposal rather than arriving via percolation from a surface source.) The chemical interactions add to the delay that these materials will experience, particularly those in the distant 200 Areas, before the plumes begin to discharge to the Columbia River. Because the half-lives of cobalt-60 (5.3 years), strontium-90 (28.8 years), and cesium-137 (30.2 years) are relatively short compared to the travel time from the 200 Areas to the Columbia River, they will decay before ever reaching the river. The strontium-90 in the 100 Areas will likely reach the river or continue to enter the river as is the case at the 100-N Area.



S9501004.3

Figure 7.1. Nitrate Plume

Application of the equations and assumptions defined in Section 4.2 to the groundwater plumes results in a series of complementary, but not necessarily intercomparable, screening values for each contaminant. The varying numbers of assumptions and associated varying degrees of conservatism require that each of the screenings be evaluated separately. The combined results of the screenings, however, then define the overall list of contaminants of interest. The complete list of radionuclides and chemicals of concern entered into the project database is presented in Table 3.3. The parameters used in the calculations are presented in Appendix B. The complete numerical results are presented in Appendix C.

The overall screening results for existing groundwater plumes away from the river are given in Table 7.1.

Table 7.1. Contaminants of Potential Interest Identified via Screening of Groundwater Away from the Columbia River

Radionuclide Screening	Carcinogenic Chemical Screening	Hazard Index Screening	Ambient Water Quality Criteria Screening	Aquatic Toxicant Screening
-	Chromium (100 Areas)	Nitrate (100 Areas)	-	Chromium (100 Areas)
-	Chromium (200-West Area)	Nitrate (200-West Area)	-	Nitrate (100 Areas)
-	Chromium (200-East Area)	Nitrate (200-East Area)	-	Fluoride (200-West Area)
-	-	Carbon Tetrachloride (200-West Area)	-	Nitrate (200-West Area)
-	-	-	-	Nitrate (200-East Area)

7.2 Potential Future Groundwater Sources

A very large number of radionuclides and chemicals are contained in Hanford facilities, waste management sites, or other contaminated areas. Remedial actions are planned or under way by the DOE under the provisions of the *Hanford Federal Facility Agreement and Consent Order* (Tri-Party Agreement) (Ecology et al. 1994) to bring the Hanford Site into compliance with the applicable requirements of CERCLA, RCRA, and the Washington State Hazardous Waste Management Act. The DOE program responsible for conducting remedial actions at the Hanford Site is referred to as the Richland Environmental Restoration Project. The scope of the Richland Environmental Restoration Project (DOE 1994h) encompasses the following groups of actions:

- radiation area remedial actions/underground storage tanks (UST)
- RCRA closures

- single-shell tank (SST) closures
- past-practice site operable unit (source and groundwater) remedial actions
- surplus facilities decontamination and decommissioning
- storage and disposal facilities.

Radiation area remedial actions address the management and control of inactive waste sites to minimize the spread of surface soil contamination. The UST program addresses the management of state-regulated, nonradioactive USTs in accordance with Washington State regulations. RCRA closures address actions at certain waste management units classified under RCRA as treatment, storage, and disposal units (TSD). (At Hanford there are over 50 groups of TSD units.) Units subject to regulation as TSDs must either receive a RCRA operating permit or be closed in accordance with the RCRA closure process.

Single-shell tank closures address the development and implementation of final disposal of the 149 SSTs at Hanford. The *Tank Waste Remediation System (TWRS) Environmental Impact Statement (EIS)* is addressing the management, treatment, storage, and disposal of waste in the SSTs. The Notice of Intent for the TWRS-EIS was published in the *Federal Register* on January 28, 1994 (59 FR 4052).

Past-practice operable unit remedial actions address the investigation and remediation of units where waste or other substances have been disposed (intentionally or unintentionally) and are not subject to regulation as TSDs. Over 1000 past-practice units have been identified at the Hanford Site (Ecology et al. 1994).

The Surplus Facilities Decontamination and Decommissioning Program addresses the safe management and final disposition of facilities, such as surplus production reactors and chemical processing buildings, that have been retired and declared surplus. Decontamination and decommissioning of the reactors along the Columbia River are addressed in the *Decommissioning of Eight Surplus Production Reactors at the Hanford Site, Richland, Washington (Final Environmental Impact Statement)* (DOE 1992c). Storage and disposal facilities address the planning, construction, and operation of facilities required for the success of the Richland Environmental Restoration Project (DOE 1994h). These facilities are being addressed individually through CERCLA, RCRA, and NEPA requirements.

Descriptions of the various potential impacts and releases to the Columbia River from the Richland Environmental Restoration Project (DOE 1994h) are provided in the *Hanford Remedial Action Environmental Impact Statement* (DOE 1994g). In addition to the Richland Environmental Restoration Project efforts (DOE 1994h), additional documentation on high-level waste and transuranic waste facilities is covered in the *Final Environmental Impact Statement, Disposal of Hanford Defense High-Level, Transuranic, and Tank Wastes, Hanford Site, Richland, Washington* (DOE 1987).

The future of the many existing waste sites is undergoing review. Very few will remain in their current condition. It is nearly impossible to predict the future impact of these sites until additional planning and activities occur. The reader is directed to the various references for further information on the potential contaminants and their potential future impact on the Columbia River.

8.0 Materials of Additional Public Interest

As information has been released describing past operations and current conditions, public interest in the Hanford Site has increased. Some of the first questions raised during the public review of the *Columbia River Impact Evaluation Plan* (DOE 1993e) were about radiological contamination upriver from the Hanford Site. Questions were asked about the inclusion of chromium, nitrate, and sulfate ions, and the radionuclides cobalt-60 (dispersed as well as discrete particles), rubidium-86, molybdenum-96, ruthenium-106, cesium-137, europium-154, uranium and its decay progeny (specifically radium-226), and plutonium (from fuel failures as well as from decay of neptunium-239).

The majority of these topics have been addressed in this report. Background radiation is attributable to fallout from nuclear weapons testing or naturally occurring radionuclides: potassium-40, radium, tritium (hydrogen-3), thorium, and uranium. In fact, at background levels, it is possible to calculate that nearly 90,000 kilograms (100 tons) of uranium from natural sources alone pass the Hanford Site in the Columbia River every year. The isotope rubidium-86 has an 18-day half-life, and any released from historical Hanford operations would have long ago decayed. Molybdenum-96 is a stable isotope and, therefore, is not radioactive. The half-life of ruthenium-106 (367-day half-life) is similarly short. The half-lives of uranium isotopes are all in excess of 100,000 years (uranium-238, the progenitor of radium-226, has a half-life of 4.5 billion years), and no appreciable decay or progeny accumulation is expected to have occurred. During Hanford operations, about 6.3 million curies of neptunium-239 were released to the Columbia River (Heeb 1994, p. vii). All of that has now decayed into plutonium-239. Because each atom of neptunium becomes one atom of plutonium following the decay, there are no more atoms of plutonium in the river than there were neptunium atoms released. By ratio of the decay constants, that is shown to be no more than 1.7 curies of plutonium-239. Extremely low levels of plutonium have been measured in the sediment behind McNary Dam, enriched by about 30 percent in plutonium-239 over what would be expected from background radiation derived from global fallout.

Public meetings were held in December 1993 and summer 1994 regarding the CRCIA efforts. At these meetings, questions were asked about tritium (hydrogen-3), iodine-129, and uranium. Each of these contaminants has been addressed in this report.

A report produced by a public interest group provides details on Hanford contamination by arsenic, carbon tetrachloride, chloroform, chromium, cyanide, iodine-129, nitrate, plutonium, strontium-90, technetium-99, trichloroethylene, tritium (hydrogen-3), and uranium (Columbia River United circa 1994). All of these contaminants have been addressed by the CRCIA Project and the results presented in this report (see Appendix A).

Iodine-129, plutonium, technetium-99, tritium (hydrogen-3), uranium, and volatile organic compounds (e.g., chloroform and trichloroethylene) are routinely analyzed in Columbia River water samples by the Surface Environmental Surveillance Project (SESP) and the concentrations and resulting exposures reported annually (e.g., Dirkes et al. 1994). Currently, radiation doses to maximally exposed off-site individuals via the river pathway are estimated to be 0.01 mrem/year (Dirkes et al. 1994, p. 220), corresponding to a maximum individual risk of approximately 10^{-8} per year (a probability of an additional fatal cancer of 1 in 100,000,000). The concentrations of volatile organics are near or below detection levels.

Of the contaminants of potential concern raised by the public, some are of concern, but several would have been eliminated by the screening process because they are shown to be of minimal potential hazard. However, those of continued public interest will continue to be evaluated in the CRCIA Project.

These contaminants of probable continued public interest are

- chloroform
- cyanide
- iodine-129
- plutonium-239/240
- technetium-99
- trichloroethylene
- tritium (hydrogen-3)
- uranium.

9.0 Conclusions

More than 600 different radionuclides or chemicals have been sought in Hanford-related environmental samples. A large number of potential contaminants have never been detected in the Hanford/Columbia River environments. For the roughly 100 compounds that have been detected at some level, screening on the basis of potential impact on human health or the health of Columbia River ecosystems has been performed. Several different types of screenings were employed. The results were consistent in that the same compounds were identified numerous times by the various screenings. Application of the screenings for contaminants within 150 meters (500 feet) of the Columbia River yields a list of 20 contaminants of concern, plus direct irradiation. These contaminants are given in the first column of Table 9.1.

Existing Hanford groundwater contamination farther than 150 meters (500 feet) away from the Columbia River has also been addressed. The contaminants identified by the screening process (second column of Table 9.1) are not yet entering the Columbia River but have the potential to do so within 10 to 200 years (Freshley and Graham 1988). Two contaminants (chromium and nitrate) are common with those identified as being already in or near the river, and two (carbon tetrachloride and fluoride) are unique. Continued evaluation of the contaminants of concern (first column of Table 9.1) should cover most of the potential risk from the distant plumes.

Although the screenings did not indicate a potential risk, several potential or existing contaminants are of high interest to the public (third column in Table 9.1). Essentially all of these are the object of ongoing evaluation by SESP conducted by PNL at Hanford. The CRCIA Project should remain current on SESP activities and include SESP results in all project reports.

Each of the identified contaminants can be considered to have resulted from the past plutonium-production operations at Hanford. The radionuclides on the list generally represent those identified with river water or Hanford Reach sediment. The radionuclides resulted from activation of materials in the old production reactors. Although it is likely that the cesium isotopes are related to global fallout (Dirkes et al. 1994). Most of the metals identified in Hanford groundwater or sediment can be related to various Hanford operations in the 100 Areas. The PCB, Arochlor 1248, is used in equipment and the insecticide, Chlordane, has been used in Hanford facilities, but both are still essentially associated with soil near the river. The nitrate groundwater plumes result from past Hanford operations in the 100 and 200 Areas.

The reduction from more than 600 potential chemicals of concern to the final list of 20, plus direct irradiation, was based on several complementary screening techniques and illustrates that future sampling and environmental analyses are both possible and tractable for the CRCIA Project.

Table 9.1. List of Identified Contaminants of Concern^(a)

In Columbia River, Groundwater, ^(b) Sediment, and Soil	Groundwater Plumes Away from the Columbia River ^(c)	Continued Public Interest
Antimony Arochlor 1248 (PCB) Arsenic Cesium-134 Cesium-137 Chlordane Chromium ^(d) Cobalt-60/particles Copper Diesel Fuel Europium-152 Europium-154 Lead Manganese Mercury Nitrate/nitrite ^(d) Phosphate Silver Chloride Strontium-90 Zinc	Carbon Tetrachloride Fluoride	Chloroform Cyanide Iodine-129 Plutonium-239/240 Technetium-99 Trichloroethylene Tritium (Hydrogen-3) Uranium
<p>(a) Direct irradiation is also identified as being of concern. (b) Hanford groundwater within 150 meters (500 feet) of the Columbia River. (c) Hanford groundwater farther than 150 meters (500 feet) from the Columbia River. (d) These contaminants are also of concern in groundwater plumes away from the Columbia River but are not repeated in that list to avoid duplication.</p>		

10.0 Perspective

The identification of the radionuclides and chemicals of concern to the CRCIA Project should not imply that each or all of these compounds is necessarily a contamination or exposure problem for those who live downstream or the ecosystem of the Columbia River. The screening and selection process described in this report is a conservative (cautious) process designed to focus the resources of the project on those contaminants with potential risk.

Recent sampling has been performed in sediment of the Snake and Columbia Rivers as part of the studies underway concerning reservoir drawdowns for enhancement of salmon stocks. A study by Pinza et al. (1992) included grain size, total organic carbon, total volatile solids, ammonia, phosphorus, sulfides, oil and grease, total petroleum hydrocarbons, metals, polynuclear aromatic hydrocarbons, pesticides, PCBs, and 21 types of polychlorinated dibenzodioxins and dibenzofurans. Samples were taken from the Columbia River at the Port of Kennewick, the Boise Cascade facility below the confluence of the Snake and Columbia Rivers, and at Wallula Gap, as well as from 24 stations on the Snake River.

The study by Pinza et al. (1992) found most measured concentrations of all contaminants to be quite low in Columbia River sediment downstream of Hanford. The concentrations in this CRCIA Project report show most metals in Columbia River sediment to be within the ranges found by Pinza et al. (1992) in Snake River sediment. The few exceptions never differed from the extremes of the range found in the Snake River by more than a factor of 2. One of the pesticides identified by the CRCIA Project as of potential concern, chlordane, was undetected by Pinza et al. (1992) in Columbia River sediment. The PCB, Arochlor 1248, identified by the CRCIA Project as of potential concern was also undetected by Pinza et al. (1992) in Columbia River sediment. The two polynuclear aromatic hydrocarbons discussed in Section 4.4 of this CRCIA report, benzo(a)pyrene and indeno(1,2,3-cd)pyrene, were undetected by Pinza et al. (1992) at Kennewick or Wallula Gap. The frequent inability to detect contaminants at the Boise Cascade facility make it impossible to make a comparison at that location. Petroleum products measured at Kennewick were the lowest found by Pinza et al. (1992) at any location.

Contaminants in the Columbia River, groundwater, sediment, and soil may have potential for impacts on human or ecological health in areas immediately adjacent to the Hanford shorelines, or throughout the Hanford Reach. However, it is evident from the results presented by Pinza et al. (1992) that Columbia River concentrations are similar to those in other rivers not associated with Hanford releases. Whereas Pinza et al. (1992) sampled for non-radionuclides, Wells (1994) examined data for radionuclides and concluded that the potential risk is lower than that allowed by the federal drinking water standards.

11.0 References

59 FR 4052. January 28, 1994. "Intent to Prepare Hanford Tank Waste Remediation System Environmental Impact Statements, Richland, WA." *Federal Register*.

Ayres, J. M. 1993. *Qualitative Risk Assessment for 100-HR-1 Source Operable Unit*. WHC-SD-EN-RA-004, Rev. 0, Westinghouse Hanford Company, Richland, Washington.

Beasley, T. M., L. A. Ball, J. E. Andrews III, and J. E. Halverson. 1981. "Hanford-Derived Plutonium in Columbia River Sediments." *Science* 214(4523):913-915.

Bisping, L. E. 1992. *Hanford Site Environmental Data 1991-Surface and Columbia River*. PNL-8149, Pacific Northwest Laboratory, Richland, Washington.

Bisping, L. E., and R. K. Woodruff. 1993. *Hanford Site Environmental Data for Calendar Year 1992 - Surface and Columbia River*. PNL-8683, Pacific Northwest Laboratory, Richland, Washington.

Bisping, L. E. 1994. *Hanford Site Environmental Data for Calendar Year 1993 - Surface and Columbia River*. PNL-9824, Pacific Northwest Laboratory, Richland, Washington.

Cline, C. S., J. T. Rieger, and J. R. Raymond. 1985. *Ground-Water Monitoring at the Hanford Site, January - December 1984*. PNL-5408, Pacific Northwest Laboratory, Richland, Washington.

Columbia River United. Circa 1994. *Hanford and the River*. Hood River, Oregon.

Cooper, A. T., and R. K. Woodruff. 1993. *Investigation of Exposure Rates and Radionuclide and Trace Metal Distributions Along the Hanford Reach of the Columbia River*. PNL-8789, Pacific Northwest Laboratory, Richland, Washington.

CRITFC - Columbia River Inter-Tribal Fish Commission. 1994. *A Fish Consumption Survey of the Umatilla, Nez Perce, Yakama, and Warm Springs Tribes of the Columbia River Basin*. CRITFC Technical Report No. 94-3, Portland, Oregon.

Dirkes, R. L. 1993. *Columbia River Monitoring: Distribution of Tritium in Columbia River Water at the Richland Pumphouse*. PNL-8531, Pacific Northwest Laboratory, Richland, Washington.

Dirkes, R. L. 1994. *Summary of Radiological Monitoring of Columbia River Water along the Hanford Reach, 1980 through 1989*. PNL-9223, Pacific Northwest Laboratory, Richland, Washington.

Dirkes, R. L., G. W. Patton, and B. L. Tiller. 1993. *Columbia River Monitoring: Summary of Chemical Monitoring Along Cross Sections at Vernita Bridge and Richland*. PNL-8654, Pacific Northwest Laboratory, Richland, Washington.

Dirkes, R. L., R. W. Hanf, R. K. Woodruff, and R. E. Lundgren. 1994. *Hanford Site Environmental Report for Calendar Year 1993*. PNL-9823, Pacific Northwest Laboratory, Richland, Washington.

DOE - U.S. Department of Energy. 1987. *Final Environmental Impact Statement, Disposal of Hanford Defense High-Level, Transuranic, and Tank Wastes, Hanford Site, Richland, Washington*. DOE/EIS-0113, U.S. Department of Energy, Washington, D.C.

DOE - U.S. Department of Energy. 1989. *Decommissioning of Eight Surplus Production Reactors at the Hanford Site, Richland, Washington, Draft Environmental Impact Statement*. DOE/EIS-0119D, U.S. Department of Energy, Washington, D.C.

DOE - U.S. Department of Energy. 1990a. *Remedial Investigation/Feasibility Study Work Plan for the 300-FF-5 Operable Unit, Hanford Site, Richland, Washington*. DOE/RL 89-14, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1990b. *Remedial Investigation/Feasibility Study Work Plan for the 300-FF-1 Operable Unit, Hanford Site, Richland, Washington*. DOE/RL 88-31, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1990c. *Low-level Burial Grounds Dangerous Waste Permit Application: Request for Exemption from Lined Trench Requirements for Submarine Reactor Compartments*. DOE/RL-88-20, Supplement 1, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1992a. *Sampling and Analysis of 100 Area Springs*. DOE/RL-92-12, Rev. 1, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1992b. *Hanford Site Groundwater Background*. DOE/RL-92-23, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1992c. *Decommissioning of Eight Surplus Production Reactors at the Hanford Site, Richland, Washington (Final Environmental Impact Statement)*. DOE/EIS-0119F, U.S. Department of Energy, Washington, D.C.

DOE - U.S. Department of Energy. 1992d. *Low-Level Burial Grounds Dangerous Waste Permit Application: Request for Exemption from Lined Trench Requirements and from Land Disposal Restrictions for Residual Liquid at 218-E-12B Burial Ground Trench 94*. DOE/RL-88-20, Supplement 1, Revision 1, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1992e. *Hanford Site Baseline Risk Assessment Methodology*, DOE/RL-91-45, Rev. 2., U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1993a. *Limited Field Investigation Report for the 100-BC-5 Operable Unit*. DOE/RL-93-37, Draft A, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1993b. *Limited Field Investigation Report for the 100-DR-1 Operable Unit*. DOE/RL-93-29, Draft A, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1993c. *Limited Field Investigation Report for the 100-HR-1 Operable Unit*. DOE/RL-93-51, Draft A, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1993d. *Limited Field Investigation Report for the 100-HR-3 Operable Unit*. DOE/RL-93-43, Draft A, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1993e. *Columbia River Impact Evaluation Plan*. DOE/RL-92-28, Rev. 0, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1994a. *Hanford Site Background: Part 1, Soil Background for Nonradioactive Analytes*. DOE/RL-92-24, Rev. 2, Vol. 1 of 2. U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1994b. *Annual Report for RCRA Groundwater Monitoring Projects at Hanford Site Facilities*. DOE/RL-93-88, Rev. 0, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1994c. *HEIS - Hanford Environmental Information System*. For documentation supporting the HEIS database, see DOE/RL-93-24, 9 volumes, U.S. Department of Energy, Richland, Washington. Queried: August 24, 1994.

DOE - U.S. Department of Energy. 1994d. *Limited Field Investigation Report for the 100-BC-1 Operable Unit*. DOE/RL-93-06, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1994e. *Limited Field Investigation Report for the 100-KR-1 Operable Unit*. DOE/RL-93-78, Draft A, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1994f. *Limited Field Investigation Report for the 100-KR-4 Operable Unit*. DOE/RL-93-79, U.S. Department of Energy, Richland, Washington.

DOE - U.S. Department of Energy. 1994g. *Hanford Remedial Action Draft Environmental Impact Statement*. DOE/DEIS-0222, U.S. Department of Energy, Washington, D.C.

DOE - U.S. Department of Energy. 1994h. *Richland Environmental Restoration Project Plan*. DOE/RL-92-08, U.S. Department of Energy, Richland, Washington.

Droppo, J. G., Jr., D. L. Streng, J. W. Buck, B. L. Hoopes, R. D. Brockhaus, M. B. Walter, and G. Whelen. 1991. *MEPAS™ Multimedia Environmental Pollutant Assessment System, Application Guidance*. PNWD-1857, Vol. 1, Battelle, Pacific Northwest Laboratories, Richland, Washington.

Ecology - Washington State Department of Ecology, U.S. Environmental Protection Agency, and U.S. Department of Energy. 1994. *Hanford Federal Facility Agreement and Consent Order*. Document No. 89-10, Rev. 3 (The Tri-Party Agreement), Washington State Department of Ecology, Olympia, Washington.

EG&G Energy Measurements. 1990. *An Aerial Radiological Survey of the Hanford Site and Surrounding Area, Richland, Washington*. EGG-10617-1062, EG&G Energy Measurements, The Remote Sensing Laboratory, Las Vegas, Nevada.

EPA - U.S. Environmental Protection Agency. 1985. *OHM/TADS - Oil and Hazardous Materials/Technical Assistance Data System*, in TOMES--Toxicology, Occupational Medicine, and Environmental Series (CD-ROM). Database used: Micromedex.

EPA - U.S. Environmental Protection Agency. 1989. *Risk Assessment Guidance for Superfund; Human Health Evaluation Manual Part A*. Interim Final, EPA/540/1-89/002, U.S. Environmental Protection Agency, Washington D.C.

EPA - U.S. Environmental Protection Agency. 1991. *EPA Region 10 Supplemental Risk Assessment Guidance for Superfund*. Published Draft. U.S. Environmental Protection Agency, Washington, D.C.

EPA - U.S. Environmental Protection Agency. 1992. *Water Quality Standards; Establishment of Numeric Criteria for Priority Toxic Pollutants; States' Compliance; Final Rule (Federal Register)*. Vol. 57, No. 246. 12-22-92. U.S. Environmental Protection Agency, Washington, D.C.

EPA - U.S. Environmental Protection Agency. 1994a. *Health Effects Assessment Summary Tables*. FY-1994 Annual. EPA 540/R/94/020, U.S. Environmental Protection Agency, Washington, D.C.

EPA - U.S. Environmental Protection Agency. 1994b. *IRIS - Integrated Risk Information System*, in TOMES-Toxicology, Occupational Medicine, and Environmental Series (CD-ROM). Database used: Micromedex.

Eslinger, P. W., L. R. Huesties, A. D. Maughan, T. B. Miley, and W. H. Walters. 1994. *Data Compendium for the Columbia River Impact Assessment*. PNL-9785, Pacific Northwest Laboratory, Richland, Washington.

Ford, B. H. 1993. *Groundwater Field Characterization Report for the 200 Aggregate Area Management Study*. WHC-SD-EN-TI-020, Westinghouse Hanford Company, Richland, Washington.

Fowler, K. M., K. B. Miller, M. O. Hogan, and J. F. Donaghue. 1993. *Risk-Based Standards Chemicals of Interest Database Documentation*. Draft, prepared for the U.S. Department of Energy by Pacific Northwest Laboratory, Richland, Washington.

Freshley, M. D., and M. J. Graham. 1988. *Estimation of Ground-Water Travel Time at the Hanford Site: Description, Past Work, and Future Needs*. PNL-6328, Pacific Northwest Laboratory, Richland, Washington.

Hartman, M. J., and K. A. Lindsey. 1993. *Hydrogeology of the 100-N Area, Hanford Site, Washington*. WHC-SD-EN-EV-027, Westinghouse Hanford Company, Richland, Washington.

Heeb, C. M., and D. J. Bates. 1994. *Radionuclide Releases to the Columbia River from Hanford Operations, 1944-1971*. PNWD-2223 HEDR, Battelle, Pacific Northwest Laboratories, Richland, Washington.

Jaquish, R. E. 1989. "Uranium-236 as an Indicator of Full-Cycle Uranium in Ground Water." Presented at the 35th Annual Conference on Bioassay, Analytical and Environmental Radiochemistry. PNL-SA-17410, Pacific Northwest Laboratory, Richland, Washington.

Kipp, K. L., A. E. Reisenauer, C. R. Cole, and C. A. Bryan. 1976. *Variable Thickness Transient Groundwater Flow Model Theory and Numerical Implementation*. BNWL-1703, Pacific Northwest Laboratory, Richland, Washington.

Law, A. G. 1990. *Status of Groundwater in the 1100 Area*. Correspondence Number 8900604B R4, Westinghouse Hanford Company, Richland, Washington.

Napier, B. A., R. A. Peloquin, D. L. Streng, and J. V. Ramsdell. 1988. *GENII - The Hanford Environmental Radiation Dosimetry Software System; Volume 1: Conceptual Representation*. PNL-6584 Vol. 1, Pacific Northwest Laboratory, Richland, Washington.

Napier, B. A. 1993. *Determination of Key Radionuclides and Parameters Related to Dose from the Columbia River Pathway*. BN-SA-3768 HEDR, Battelle, Pacific Northwest Laboratories, Richland, Washington.

National Council on Radiation Protection and Measurements (NCRP). 1989. *Limit for Exposure to "Hot Particles" on the Skin*. NCRP Report No. 106, National Council on Radiation Protection and Measurements, Bethesda, Maryland.

Navy - U.S. Department of the Navy. 1984. *Final Environmental Impact Statement on the Disposal of Decommissioned, Defueled Naval Submarine Reactor Plants*. United States Department of the Navy, Washington, D.C.

NPS - National Parks Service. 1992. *Hanford Reach of the Columbia River, Draft Comprehensive River Conservation Study and Environmental Impact Statement*. U.S. Department of Interior, Seattle, Washington.

Peterson, R. E., and V. G. Johnson. 1992. *Riverbank Seepage of Groundwater Along the 100 Areas Shoreline, Hanford Site*. WHC-EP-0609, Westinghouse Hanford Company, Richland, Washington.

Pinza, M. R., J. Q. Word, E. S. Barrows, H. L. Mayhew, and D. R. Clark. 1992. *Snake and Columbia Rivers Sediment Sampling Project*. PNL-8479, Pacific Northwest Laboratory, Sequim, Washington.

Raymond, J. R., D. A. Myers, J. J. Fix, V. L. McGhan, and P. M. Schrotke. 1976. *Environmental Monitoring Report on Radiological Status of the Ground Water Beneath the Hanford Site, January - December 1974*. BNWL-1970, Pacific Northwest Laboratory, Richland, Washington.

Rhoads, K., B. N. Bjornstad, R. E. Lewis, S. S. Teel, K. J. Cantrell, R. J. Serne, J. L. Smoot, C. T. Kincaid, and S. K. Wurstner. 1992. *Estimation of the Release and Migration of Lead Through Soils and Groundwater at the Hanford Site 218-E-12B Burial Ground*. PNL-8356 Vol. 1., Pacific Northwest Laboratory, Richland, Washington.

Rhoads, K., B. N. Bjornstad, R. E. Lewis, S. S. Teel, K. J. Cantrell, R. J. Serne, L. H. Sawyer, J. L. Smoot, J. E. Szecsody, M. S. Wigmosta, and S. K. Wurstner. 1994. *Estimation of the Release and Migration of Nickel Through Soils and Groundwater at the Hanford Site 218-E-12B Burial Ground*. PNL-9791, Pacific Northwest Laboratory, Richland, Washington.

Rowley, C. A. 1993. *100-N Area Underground Storage Tank Closures*. WHC-SD-EN-TI-136, Westinghouse Hanford Company, Richland, Washington.

Sula, M. J. 1980. *Radiological Survey of Exposed Shorelines and Islands of the Columbia River Between Vernita and the Snake River Confluence*. PNL-3127, Pacific Northwest Laboratory, Richland, Washington.

Wade, C. D., and M. A. Wendling. 1994. *100-D Island USRADS Radiological Surveys Preliminary Report Phase II*. BHI-00-134, Bechtel Hanford, Inc., Richland, Washington.

Weiss, S. G. 1993. *100 Area Columbia River Sediment Sampling*. WHC-SD-EN-TI-198, Rev. 0, Westinghouse Hanford Company, Richland, Washington.

Wells, D. 1994. *Radioactivity in Columbia River Sediments and their Health Effects*. Special Report, Washington State Department of Health, Olympia, Washington.

Woodruff, R. K., R. W. Hanf, and R. E. Lundgren. 1992. *Hanford Site Environmental Report for Calendar Year 1991*. PNL-8148, Pacific Northwest Laboratory, Richland, Washington.

Woodruff, R. K., R. W. Hanf, and R. E. Lundgren. 1993. *Hanford Site Environmental Report for Calendar Year 1992*. PNL-8682, Pacific Northwest Laboratory, Richland, Washington.

Wurstner, S. K., and J. L. Devary. 1993. *Hanford Site Ground-Water Model: Geographical Information System Linkages and Model Enhancements, FY 1993*. PNL-8991, Pacific Northwest Laboratory, Richland, Washington.

Appendix A

Complete List of Analytes Evaluated at Hanford

Appendix A

Complete List of Analytes Evaluated at Hanford

Table A.1 provides a complete listing of all radionuclides and chemicals for which monitoring has been reported in the reviewed literature of samples from the Columbia River and groundwater in the Hanford Site 100, 300, and 1100 Areas within 150 meters (500 feet) of the Columbia River. For those contaminants which had a detected level, the highest concentration reported is listed. A total of 568 analytes are listed. The 73 analytes for which detected levels were reported are listed in Table 3.1.

Table A.2 provides a complete listing of all radionuclides and chemicals for which monitoring has been reported in the reviewed literature of samples from soil and sediment in the Hanford Site 100, 300, and 1100 Areas. For those contaminants which had a detected level, the highest concentration reported is listed. A total of 560 analytes are listed. The 92 analytes for which detected levels were reported are listed in Table 3.2.

Table A.3 provides a listing of the major radionuclides and chemicals for which monitoring has been reported in the reviewed literature of samples from groundwater in the Hanford Site 100, 200, and 600 Areas farther than 150 meters (500 feet) away from the Columbia River. The listing is not comprehensive for all analytes, as described in Section 7.0.

The following abbreviations are used in the tables. All units are as reported in the reviewed literature. The column headings, such as 100-KR-4, refer to sampling locations at operable units, described in Section 2.0.

- aCi/L = attocuries per liter (one one-millionth of a pCi/L).
- CAS# = Chemical Abstract Service number, a unique numerical identifier for chemicals.
- HEIS = Hanford Environmental Information System database.
- $\mu\text{g}/\text{kg}$ = micrograms per kilogram.
- $\mu\text{g}/\text{L}$ = micrograms per liter.
- mg/kg = milligrams per kilogram.
- ND = not detected in sample; not all data compilers used this convention; some analytes show no entry where an ND is appropriate.
- pCi/kg = picocuries per kilogram.
- pCi/L = picocuries per liter.
- ppb = parts per billion.
- SD = sediment.
- SW = surface water.
- w/Pu239 = concentration included in the value reported for plutonium-239.
- w/U233 = concentration included in the value reported for uranium-233.
- * = laboratory results marked as suspect data (see Section 4.4).

Table A.1. (contd)

Name of Analyte	CAS #	Background (a)	Background Reference	GROUNDWATER							COLUMBIA RIVER			
				HEIS (DOE 1994c)	100-KR-4 (DOE 1994f)	100-HR-3 (DOE 1993d)	100-BC-5 (DOE 1993a)	100-N (Hartman & Lindsey 1993)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	Richland PumpHouse (c)	
66 BENZICACRIDINE										ND				
67 BENZENE	71-43-2						ND	ND (d)	ND		ND	ND	ND	ND
68 BENZENETHIOL									ND					
69 BENZO(G,H,I)PERYLENE	191-24-2						ND		ND		ND			
70 BENZO(J)FLUORANTHENE									ND					
71 BENZO(a)ANTHRACENE	56-55-3						ND		ND		ND			
72 BENZIDINE									ND					
73 BENZO(a)PYRENE	50-32-8						ND		ND		ND			
74 BENZO(b)FLUORANTHENE	205-99-2						ND		ND		ND			
75 BENZO(k)FLUORANTHENE	207-08-9						ND		ND		ND			
76 BENZOIC ACID	65-85-0										ND			
77 BENZYL ALCOHOL	100-51-6										ND			
78 BENZYL CHLORIDE									ND					
79 BERYLLIUM	7440-41-7	ND (SW)	Dirkes et al. 1993	1 µg/L				6 ppb	5ppb		1.4 µg/L	ND	ND	
80 BERYLLIUM 7	7440-41-7	< 5 ppb	DOE 1992b											
81 BETA-BHC	319-85-7								ND		ND			
82 BIS(2-CHLORO-1-METHYLETHYL)ETH									ND					
83 BIS(2-CHLOROETHOXY)METHANE	111-91-1								ND		ND			
84 BIS(2-CHLOROETHYL)ETHER	111-44-4								ND		ND			
85 BIS(2-CHLOROISOPROPYL)ETHER	39835-32-9								ND		ND			
86 BIS(2-ETHYLHEXYL) PHTHALATE	117-81-7			50 µg/L			11 µg/L			50 µg/L	ND			
87 BISMUTH	7440-68-9	< 5 ppb	DOE 1992b											
88 BISMUTH 212	14913-49-6													
89 BISMUTH 214	14733-03-0													
90 BORON	7440-42-8	< 100 ppb	DOE 1992b						64ppb					
91 BROMIDE		ND (SW)	Dirkes et al. 1993					ND	ND					ND
92 BROMOACETONE									ND					
93 BROMODICHLOROMETHANE	75-27-4						ND		ND		ND	ND		
94 BROMOFORM	75-25-2						ND		ND		ND	ND		
95 BROMOMETHANE	74-83-9						ND		ND		ND	ND		
96 BROMOPHENYL-PHENYLETHER, 4-	101-55-3								ND		ND	ND		
97 BUTANOL, 1-	71-36-3	ND (SW)	Dirkes et al. 1993						ND		ND	ND	ND	ND
98 BUTANONE	78-93-3						ND		ND		ND			
99 BUTANONE, 2-	78-93-3											ND		
100 BUTYL BENZYL PHTHALATE	85-68-7						ND		ND					
101 BUTYNOL, 1-									ND					
102 CADMIUM	7440-43-9	< 10 ppb	DOE 1992b	10 µg/L				31 ppb	ND	6.6 µg/L	3 µg/L	ND	ND	ND
103 CADMIUM 109	14109-32-1													
104 CADMIUM NITRATE	10325-97-7													
105 CALCIUM	7440-70-2	63600 ppb	DOE 1992b	100000 µg/L	94600 µg/L	130000 µg/L		302000ppb	81400 ppb	21200 µg/L	76600 µg/L	35900 µg/L	19000 µg/L	
106 CALCIUM 41	14092-95-6													
107 CALCIUM BICARBONATE	1317-65-3													
108 CARBAZOLE	86-74-8										ND			
109 CARBAZOLE, 9H-	86-74-8						ND							
110 CARBON 14	14782-75-5			200 pCi/L	23000 pCi/L		110 pCi/L							
111 CARBON DISULFIDE	75-15-0						ND		ND		ND	ND	ND	
112 CARBON TETRACHLORIDE	56-23-5	ND (SW)	Dirkes et al. 1993				ND		ND		ND	ND	ND	ND
113 CARBOPHENOTHION									ND					
114 CERIUM	7440-45-1													
115 CERIUM 141	13967-74-3													
116 CERIUM 144	14782-78-8	ND (SW)	Dirkes 1994											ND
117 CESIUM 134	13967-70-9	ND (SW)	Dirkes et al. 1994				ND				ND	ND	0.012 pCi/L	
118 CESIUM 135	15726-30-4													
119 CESIUM 137	10045-97-3	ND (SW)	Dirkes et al. 1994	0.5 pCi/L			ND				ND	ND	0.13 pCi/L	
120 CHLOR-2, 3-EPOXYPROPANE, 1-									ND					
121 CHLORDANE	57-74-9								ND		ND			
122 CHLORIDE	16887-00-6	8690 ppb (b)	DOE 1992b	30000 µg/L				18000 ppb	43400 ppb	122000 µg/L	ND	ND	870 µg/L	
123 CHLORINE	7782-50-5								ND					
124 CHLORINAPHAZINE									ND					
125 CHLOROALKYL ETHERS									ND					
126 CHLOROANILINE, 4-	106-47-8										ND			
127 CHLOROBENZENE	108-90-7						ND		ND		ND	ND		
128 CHLOROBENZILATE									ND					
129 CHLORODIBROMOMETHANE	124-48-1													
130 CHLOROETHANE	75-00-3						ND		ND		ND	ND		

A.3

Table A.1. (contd)

Name of Analyte	CAS #	Background (a)	Reference	HEIS	100-KR-4 (DOE 1994f)	100-HR-3 (DOE 1993d)	100-BC-5 (DOE 1993a)	100-N (Lindsay 1993)	1100 Area (Law 1990)	300-F-1 (DOE 1990b)	300-F-5 (DOE 1990a)	Richard Pumphouse (c)
131 CHLOROETHOXY ETHENE, 2-	110-75-8	ND (SW)	Dikes et al. 1993	10 µg/L	17 µg/L	ND	6 ppb	42 µg/L	18 µg/L	ND	ND	ND
132 CHLOROTHYLVINYL ETHER, 2-	110-75-8	ND (SW)	Dikes et al. 1993	10 µg/L	17 µg/L	ND	6 ppb	42 µg/L	18 µg/L	ND	ND	ND
133 CHLOROFORM	67-66-3	ND (SW)	Dikes et al. 1993	10 µg/L	17 µg/L	ND	6 ppb	42 µg/L	18 µg/L	ND	ND	ND
134 CHLOROMETHANE	74-87-3	ND (SW)	Dikes et al. 1993	10 µg/L	17 µg/L	ND	6 ppb	42 µg/L	18 µg/L	ND	ND	ND
135 CHLOROMETHYL METHYL ETHER	35421-08-0	ND (SW)	Dikes et al. 1993	8 µg/L	ND	ND	ND	5.4 µg/L	ND	ND	ND	ND
136 CHLOROMETHYLPHENOL, 4-3-	13981-38-9	ND (SW)	Dikes et al. 1993	8 µg/L	ND	ND	ND	5.4 µg/L	ND	ND	ND	ND
137 CHLORONAPHTHALENE, 2-	13981-38-9	ND (SW)	Dikes et al. 1993	8 µg/L	ND	ND	ND	5.4 µg/L	ND	ND	ND	ND
138 CHLOROPHENOL, 2-	95-57-8	ND (SW)	Dikes et al. 1993	8 µg/L	ND	ND	ND	5.4 µg/L	ND	ND	ND	ND
139 CHLOROPHENYL PHENYL ETHER, 4-	7008-72-3	ND (SW)	Dikes et al. 1993	8 µg/L	ND	ND	ND	5.4 µg/L	ND	ND	ND	ND
140 CHLOROPROPIONITRILE, 3-	7738-94-6	ND (SW)	Dikes et al. 1993	8 µg/L	ND	ND	ND	5.4 µg/L	ND	ND	ND	ND
141 CHROMIC ACID	7738-94-6	ND (SW)	Dikes et al. 1993	8 µg/L	ND	ND	ND	5.4 µg/L	ND	ND	ND	ND
142 CHROMIUM	7440-47-3	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
143 CHROMIUM (VI)	17523-28-1	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
144 CHROMIUM (VI)	18640-29-9	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
145 CHROMIUM (VI)	14392-02-0	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
146 CHROMIUM NITRATE	13548-38-4	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
147 CHROMIUM SULFATE	10101-53-8	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
148 CHRYSENE	218-01-9	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
149 CITRUS RED	7440-48-4	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
150 COBALT	7440-48-4	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
151 COBALT 58	13981-38-9	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
152 COBALT 60	10198-40-0	< 30 ppb	DOE 1992b	500 µg/L	1950 µg/L	490 µg/L	180 ppb	257 µg/L	363 µg/L	ND	22 µg/L	ND
153 COPPER	7440-50-8	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
154 COPPER NITRATE	3251-23-8	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
155 COPPER SULFATE	7558-98-7	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
156 CRESOLS	7558-98-7	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
157 CROTONALDEHYDE	15510-73-3	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
158 CURIUM 242	15510-73-3	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
159 CURIUM 244	13981-15-2	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
160 CURIUM 245	15621-76-8	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
161 CYANIDE	57-12-6	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
162 CYCLOHEXYL-4,6-DINITROPHENOL, 2-	57-12-6	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
163 D, 2,4-	72-54-8	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
164 DDD, 4,4'-	72-54-8	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
165 DDE, 4,4'-	72-54-8	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
166 DDT, 4,4'-	50-29-3	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
167 DELTA-BHC	319-88-8	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
168 DIALLATE	319-88-8	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
169 DI-N-BUTYLPHTHALATE	84-74-2	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
170 DI-N-PROPYLPHTHALAMINE	84-74-2	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
171 DI-N-OCTYLPHTHALATE	117-84-0	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
172 DIBENZ(a,h)ACRIDINE	117-84-0	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
173 DIBENZ(a,h)ACRIDINE	117-84-0	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
174 DIBENZ(a,h)ANTHRACENE	63-70-3	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
175 DIBENZ(a,h)ANTHRACENE	63-70-3	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
176 DIBENZ(a,h)CARBAZOLE, 7H-	63-70-3	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
177 DIBENZ(a,h)PYRENE	63-70-3	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
178 DIBENZ(a,h)PYRENE	63-70-3	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
179 DIBENZ(a,h)PYRENE	63-70-3	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
180 DIBENZOFURAN	132-64-9	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
181 DIBROMO-3-CHLOROPROPANE, 1,2-	132-64-9	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
182 DIBROMOCHLOROPROPANE, 1,2-	124-48-1	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
183 DIBROMOETHANE, 1,2-	124-48-1	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
184 DIBROMOMETHANE	124-48-1	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
185 DIBUTYLPHOSPHATE	124-48-1	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
186 DICHLORO-2-BUTENE, 1,4-	124-48-1	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
187 DICHLOROBENZENE, p-	95-50-1	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
188 DICHLOROBENZENE, 1,2-	95-50-1	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
189 DICHLOROBENZENE, 1,3-	541-73-1	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
190 DICHLOROBENZENE, 1,4-	106-46-7	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
191 DICHLOROBENZENE, 3,3'	91-94-1	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
192 DICHLORODIFLUOROMETHANE	75-71-8	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
193 DICHLOROETHANE, 1,1-	75-34-3	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
194 DICHLOROETHANE, 1,2-	107-66-2	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L
195 DICHLOROETHYLENE, 1,1-	75-35-4	< 30 ppb	DOE 1992b	80 µg/L	ND	40 µg/L	91 ppb	64 µg/L	516 µg/L	14.7 µg/L	ND	0.011 µg/L

Name of Analyte	CAS #	Background (a)	Reference	HEIS	100-KR-4	100-HR-3	100-BC-5	100-N	1100 Area	300-FE-1	300-FE-5	300-FE-5	Richard	Pumphouse (c)
196 DICHOROETHYLENE, 1,2-	540-69-0	ND (SW)	Dikes et al. 1993	200 µg/L	ND	ND	ND	ND	150 µg/L	ND	ND	ND	ND	ND
199 DICHOROETHYLENE, 1,2-trans-	156-60-5													
199 DICHOROETHYLENE, 1,2-cis-	156-60-5													
199 DICHOROETHYLENE, 1,2-cis-	170-83-2													
200 DICHOROPHENOL, 2,6-	87-65-0													
200 DICHOROPHENOL, 2,4-	94-75-7													
201 DICHLOROPHENOXACETIC ACID, 2,4-	120-83-2													
202 DICHLOROPROPANE, 1,2-	78-87-5													
203 DICHLOROPROPANE, 1,3-	10081-02-8													
204 DICHLOROPROPENE, 1,3-cis-	10081-01-5													
205 DICHLOROPROPENE, 1,3-trans-	10081-01-5													
206 DIELDIN	60-57-1													
207 DIESEL FUEL														
208 DIETHYL-O,2-PYRAZINYL PHOSPHATE, 0,0-														
209 DIETHYLPHthalate	84-66-2													
210 DIETHYLSTIBESTEROL														
211 DIHYDROSAFROLE														
212 DIMETHOATE														
213 DIMETHOXYBENZENZINE, 3,3'														
214 DIMETHYLBENZEN(AN)THRACENE, 7,12-														
215 DIMETHYLBENZIDINE, 3,3'														
216 DIMETHYLHYDRAZINE, 1,1-														
217 DIMETHYLHYDRAZINE, 1,2-														
218 DIMETHYLPHENOL, 2,4-	105-67-9													
219 DIMETHYLPHthalate	131-11-3													
220 DINITRO-2-METHYLPHENOL, 4,6-	534-42-1													
221 DINITROBENZENE														
222 DINITRO-O-CRESOL, 4,6- and salts														
223 DINITROPHENOL, 2,4-	51-28-5													
224 DINITROPHENOL, p-														
225 DINITROTOLUENE, 2,4-	121-14-2													
226 DINITROTOLUENE, 2,6-	606-20-2													
227 DIMOSEB														
228 DIOXANE														
229 DIPHENYLAMINE														
230 DIPHENYLHYDRAZINE, 1,2-														
231 DISULFOTON														
232 ENDOSULFAN I	959-98-8													
233 ENDOSULFAN II	33213-65-9													
234 ENDOSULFAN SULFATE	1031-07-8													
235 ENDRIN	72-20-8													
236 ENDRIN ALDEHYDE	7421-93-4													
237 ENDRIN KETONE	53494-70-5													
238 ETHANOL														
239 ETHYLBENZENE	100-41-4													
240 ETHYLCARBAMATE														
241 ETHYL CYANIDE														
242 ETHYLENE GLYCOL														
243 ETHYLENE OXIDE														
244 ETHYLENIMINE														
245 ETHYLENETHIOUREA														
246 ETHYLMETHACRYLATE														
247 ETHYL METHANESULFONATE														
248 EURONIUM 152	14683-23-9													
249 EURONIUM 154	15585-10-1													
250 EURONIUM 155	14391-16-3													
251 FERRIC NITRATE	10421-48-4													
252 FERRIC SULFATE	10028-22-5													
253 FERROCYANIDE	13408-63-4													
254 FERROUS AMMONIUM SULFATE	7783-85-9													
255 FERROUS SULFATE	7720-78-7													
256 FLUORANTHENE	206-44-0													
257 FLUORENE	86-73-7													
258 FLUORIDE	7782-41-4	160 µg/L (SW)	Dikes et al. 1993	1000 µg/L	ND	ND	ND	ND	2080 µg/L	ND	ND	ND	150 µg/L	ND
259 FLUORINE	7782-41-4	175 ppb	DOE 1992b											
260 FLUOROTRICHLOROMETHANE	75-69-4													

Table A.1. (cont)

Table A.1. (contd)

Name of Analyte	CAS #	Background (a)	Background Reference	GROUNDWATER								COLUMBIA RIVER		
				HEIS (DOE 1994c)	100-KR-4 (DOE 1994f)	100-HR-3 (DOE 1993d)	100-BC-5 (DOE 1993a)	100-N (Hartman & Lindsey 1993)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	Richland Pumphouse (c)	
261 FORMALIN														
262 FUEL OIL #2	68476-34-6									ND				
263 GAMMA-BHC(LINDANE)	58-89-9													
264 GAMMA-CHLORDANE	5103-74-2									ND	ND			
265 HEPTACHLOR	76-44-8										ND			
266 HEPTACHLOR EPOXIDE	1024-67-3									ND	ND			
267 HEPTACHLOR EPOXIDE (ENDO)										ND	ND			
268 HEPTACHLOR EPOXIDE (EXO)														
269 HEXACHLOROBENZENE	118-74-1						ND			ND		ND		
270 HEXACHLOROBUTADIENE	87-68-3						ND			ND		ND		
271 HEXACHLOROCYCLOPENTADIENE	77-47-4						ND			ND		ND		
272 HEXACHLOROETHANE	67-72-1						ND			ND		ND		
273 HEXACHLOROPHENE										ND		ND		
274 HEXACHLOROPROPENE										ND		ND		
275 HEXANONE, 2-	591-78-6									ND				
276 HEXYL METHANOATE	629-33-4						ND			ND		ND	ND	
277 HYDRAZINE	302-01-2				7 µg/L					ND				
278 HYDROCARBONS														
279 HYDROCHLORIC ACID	7647-01-0													
280 HYDROCYANIC ACID	74-90-8													
281 HYDROFLUORIC ACID	7664-39-3													
282 HYDROGEN SULFIDE	7783-06-4													
283 INDENO(1,2,3-CD)PYRENE	193-39-5							ND						
284 IODINE 129	15046-84-1	5 µCi/L (SW)	Dirkes et al. 1994							ND		ND		
285 IODINE 131	10043-66-0	ND (SW)	Dirkes et al. 1994									ND	ND	160 pCi/L
286 IODOMETHANE														ND
287 IRON	7439-89-6	86 ppb (b)	DOE 1992b							ND				
288 IRON 59						5400 µg/L			37300 ppb	394 ppb	8300 µg/L	9570 µg/L	463 µg/L	92 µg/L
289 ISOBUTYL ALCOHOL							ND					ND	ND	
290 ISODRIN										ND				
291 ISOPHORONE	78-59-1									ND				
292 ISOSAFROLE							ND					ND		
293 KEPONE										ND				
294 KEROSENE	8008-20-6									ND				
295 KRYPTON 85										ND				
296 LANTHANUM	7439-91-0													
297 LEAD	7439-92-1	< 5 ppb	DOE 1992b	40 µg/L		5.1 µg/L			16 ppb	ND	173 µg/L	5.6 µg/L	ND	
298 LEAD 210	14255-04-0													
299 LEAD 212	15092-94-1													
300 LEAD NITRATE	10099-74-8													
301 LITHIUM	7439-93-2													
302 LITHIUM CHLORIDE	7447-41-8									ND				
303 MALEIC HYDRIZIDE														
304 MALONONITRILE										ND				
305 MAGNESIUM	7439-95-4	16480 ppb	DOE 1992b		30000 µg/L									
306 MANGANESE	7439-96-5	24.5 ppb (b)	DOE 1992b		400 µg/L	69.6 µg/L	180 µg/L		55000 ppb	15200 ppb	11800 µg/L	14600 µg/L	9860 µg/L	4200 µg/L
307 MANGANESE 54	13966-31-9								212 ppb	21 ppb	191 µg/L	332 µg/L	22.6 µg/L	11 µg/L
308 MELPHALAN										ND				
309 MERCURIC NITRATE	10045-95-0													
310 MERCURIC THIOCYANATE	592-85-8													
311 MERCURY	7439-97-6	< 0.1 ppb	DOE 1992b	0.2 µg/L					ND	ND	8.9 µg/L	ND	ND	ND
312 METHACRYLONITRILE										ND				
313 METHANAL	50-00-0									ND				
314 METHANETHIOL														
315 METHANOL	67-58-1									ND				
316 METHAPYRILENE														
317 METHOLONYL										ND				
318 METHOXYCHLOR	72-43-6									ND				
319 METHYLAZIRIDINE, 2-										ND				
320 METHYL BROMIDE										ND				
321 METHYL CHLORIDE										ND				
322 METHYL ETHYL KETONE		ND (SW)	Dirkes et al. 1993							ND				
323 METHYL 2-(METHYLTHIO)PROPIO, 2-										ND	18 µg/L			ND
324 METHYL 2-PENTANONE, 4-	108-10-1	ND (SW)	Dirkes et al. 1993					ND						ND
325 METHYLCHOLANTHRENE, 3-										ND				ND

Table A.1. (contd)

Name of Analyte	CAS #	Background (a)	Background Reference	GROUNDWATER								COLUMBIA RIVER		
				HEIS (DOE 1994c)	100-KR-4 (DOE 1994f)	100-HR-3 (DOE 1993d)	100-BC-5 (DOE 1993a)	100-N (Hartman & Lindsey 1993)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	Richland Pumphouse (c)	
391 OXYBIS(1-CHLOROPROPANE), 2,2'-														
392 P-BENZOQUINONE											ND			
393 P-CHLORO-M-CRESOL											ND			
394 P-CHLOROANILINE											ND			
395 P-DIMETHYLAMINOAZOBENZENE											ND			
396 P-NITROANILINE											ND			
397 PALLADIUM	7440-05-3													
398 PARALDEHYDE											ND			
399 PARATHION											ND			
400 PCDDs											ND			
401 PCDFs											ND			
402 PENTACHLOROBENZENE											ND			
403 PENTACHLOROETHANE											ND			
404 PENTACHLORONITROBENZENE											ND			
405 PENTACHLOROPHENOL	87-86-5										ND			
406 PHENANTHRENE	85-01-8										ND			
407 PHENACETIN											ND			
408 PHENOL	108-95-2										ND			
409 PHENYLENEDIAMINE											ND			
410 PHORATE											ND			
411 PHOSPHATE	7601-54-9	< 1000 ppb(c)	DOE 1992b	200 µg/L							ND		ND	3240 µg/L
412 PHOSPHORIC ACID	7664-38-2													
413 PHOSPHORUS	7723-14-0													
414 PHOSPHORUS 32											ND			
415 PHTHALIC ACID ESTERS											ND			
416 PICOLINE, 2-											ND			
417 PLUTONIUM 238	13981-16-3	ND (SW)	Dirkes et al. 1994									.01 pCi/L	ND	ND
418 PLUTONIUM 239	15117-48-3			0.03 pCi/L								ND	ND	
419 PLUTONIUM 240	14119-32-5											ND	ND	
420 PLUTONIUM 241	14119-32-6													
421 PLUTONIUM 242	13982-10-0													
422 POLONIUM 210	13981-52-8													
423 POLONIUM 212	15389-34-1													
424 POLONIUM 216	16766-58-8													
425 POLYCHLORINATED BIPHENYLS	1336-36-3													
426 POTASSIUM	7447-40-7	7975 ppb	DOE 1992b	9000 µg/L		9300 µg/L			10100 ppb	8020ppb	11300 µg/L	10200 µg/L	2430 µg/L	690 µg/L
427 POTASSIUM 40												240 pCi/L	ND	
428 POTASSIUM CHLORATE	3811-04-9													
429 POTASSIUM CYANIDE	151-50-8													
430 POTASSIUM DICHROMATE	7778-50-9													
431 POTASSIUM FLUORIDE	7789-23-3													
432 POTASSIUM HYDROXIDE	1310-58-3													
433 POTASSIUM NITRATE	7767-79-1													
434 POTASSIUM PERMANGANATE	7722-64-7													
435 PROMETHIUM 147	7440-12-2													
436 PRONAMIDE											ND			
437 PROPANOL, 1-											ND			
438 PROPIONITRILE											ND			
439 PROPYN-1-OL, 2-											ND			
440 PROTACTINIUM 231	14331-85-2													
441 PROTACTINIUM 233	13981-14-1													
442 PYRENE	129-00-0										ND			
443 PYRIDINE	110-86-1										ND			
444 RADIUM	7440-14-4	0.23 pCi/L	DOE 1992b	0.3 pCi/L							ND			
445 RADIUM 226	13982-63-3										ND			
446 RADIUM 228												.0825 pCi/L	ND	
447 RADON 220	22481-48-7											95 pCi/L	ND	
448 RESERPINE											ND			
449 RESORCINOL											ND			
450 RUTHENIUM 103	13968-53-1													
451 RUTHENIUM 106	13967-48-1	ND (SW)	Dirkes et al. 1994	20 pCi/L									34.4 pCi/L	ND
452 SAFROL											ND			
453 SAMARIUM 151	15705-94-3										ND			
454 SCANDIUM 46														
455 SEC-BUTYL-4,6-DINITROPHENOL											ND			

A.8

Table A.1. (contd)

Name of Analyte	CAS #	Background (a)	Reference	GROUNDWATER		COLUMBIA RIVER	
				100-N	100-Area	100-F-1	100-F-5
521 TITANIUM	7440-32-8			ND	ND	ND	ND
522 TITANIUM CHLORIDE	10048-06-6			ND	ND	ND	ND
523 TOLUENE	108-88-3			ND	ND	ND	ND
524 TOLUENEDIAMINE	8001-35-2			ND	ND	2.9 µg/L	4.7 µg/L
525 TOXAPHENE				ND	ND	ND	ND
526 TP(2,4)SILVEX	93-72-1			ND	ND	ND	ND
527 TRIBROMOMETHANE	75-25-2			ND	ND	ND	ND
528 TRIBUTYL PHOSPHATE	126-73-8			ND	ND	ND	ND
529 TRIBUTYLPHOSPHONIC ACID				ND	ND	ND	ND
530 TRICHLOROETHYLENE, 1,2,3-				ND	ND	ND	ND
531 TRICHLOROETHYLENE, 1,2,4-				ND	ND	ND	ND
532 TRICHLOROETHYLENE, 1,3,5-				ND	ND	ND	ND
533 TRICHLOROETHYLENE, 1,1,1-				ND	ND	ND	ND
534 TRICHLOROETHYLENE, 1,1,2-				ND	ND	ND	ND
535 TRICHLOROETHYLENE				ND	ND	24.1 µg/L	18 µg/L
536 TRICHLOROMETHANETHIOL				ND	ND	ND	ND
537 TRICHLOROMONOFLUOROMETHANE				ND	ND	ND	ND
538 TRICHLOROPHENOL, 2,4,5-				ND	ND	ND	ND
539 TRICHLOROPHENOL, 2,4,6-				ND	ND	ND	ND
540 TRICHLOROPROPANE, 1,2,3-				ND	ND	ND	ND
541 TRIS(2,3-DIBROMOPROPYL) PHOSPH				ND	ND	ND	ND
542 TRITIUM (HYDROGEN 3)				ND	ND	ND	ND
543 TUNGSTEN				ND	ND	ND	ND
544 URANIUM				ND	ND	ND	ND
545 URANIUM (TOTAL ACTIVITY)				ND	ND	ND	ND
546 URANIUM 233				ND	ND	ND	ND
547 URANIUM 234				ND	ND	ND	ND
548 URANIUM 235				ND	ND	ND	ND
549 URANIUM 238				ND	ND	ND	ND
550 URANIUM 238				ND	ND	ND	ND
551 VANADIUM				ND	ND	ND	ND
552 VANADIUM PENTOXIDE				ND	ND	ND	ND
553 VINYL ACETATE				ND	ND	ND	ND
554 VINYL CHLORIDE				ND	ND	ND	ND
555 WAFRARIIN				ND	ND	ND	ND
556 XYLENE				ND	ND	ND	ND
557 XYLENE, m-				ND	ND	ND	ND
558 XYLENE, o,p-				ND	ND	ND	ND
559 YTRIUM 90				ND	ND	ND	ND
560 ZINC				ND	ND	ND	ND
561 ZINC 66				ND	ND	ND	ND
562 ZINC AMALGAM				ND	ND	ND	ND
563 ZINC CHLORIDE				ND	ND	ND	ND
564 ZINC COMPOUNDS				ND	ND	ND	ND
565 ZINC NITRATE				ND	ND	ND	ND
566 ZINC CONIUM				ND	ND	ND	ND
567 ZINC CONIUM 93				ND	ND	ND	ND
568 ZINC CONIUM 95				ND	ND	ND	ND
569 ZINC				ND	ND	ND	ND
570 ZINC				ND	ND	ND	ND
571 ZINC				ND	ND	ND	ND
572 ZINC				ND	ND	ND	ND
573 ZINC				ND	ND	ND	ND
574 ZINC				ND	ND	ND	ND
575 ZINC				ND	ND	ND	ND
576 ZINC				ND	ND	ND	ND
577 ZINC				ND	ND	ND	ND
578 ZINC				ND	ND	ND	ND
579 ZINC				ND	ND	ND	ND
580 ZINC				ND	ND	ND	ND
581 ZINC				ND	ND	ND	ND
582 ZINC				ND	ND	ND	ND
583 ZINC				ND	ND	ND	ND
584 ZINC				ND	ND	ND	ND
585 ZINC				ND	ND	ND	ND
586 ZINC				ND	ND	ND	ND
587 ZINC				ND	ND	ND	ND
588 ZINC				ND	ND	ND	ND
589 ZINC				ND	ND	ND	ND
590 ZINC				ND	ND	ND	ND
591 ZINC				ND	ND	ND	ND
592 ZINC				ND	ND	ND	ND
593 ZINC				ND	ND	ND	ND
594 ZINC				ND	ND	ND	ND
595 ZINC				ND	ND	ND	ND
596 ZINC				ND	ND	ND	ND
597 ZINC				ND	ND	ND	ND
598 ZINC				ND	ND	ND	ND
599 ZINC				ND	ND	ND	ND
600 ZINC				ND	ND	ND	ND
601 ZINC				ND	ND	ND	ND
602 ZINC				ND	ND	ND	ND
603 ZINC				ND	ND	ND	ND
604 ZINC				ND	ND	ND	ND
605 ZINC				ND	ND	ND	ND
606 ZINC				ND	ND	ND	ND
607 ZINC				ND	ND	ND	ND
608 ZINC				ND	ND	ND	ND
609 ZINC				ND	ND	ND	ND
610 ZINC				ND	ND	ND	ND
611 ZINC				ND	ND	ND	ND
612 ZINC				ND	ND	ND	ND
613 ZINC				ND	ND	ND	ND
614 ZINC				ND	ND	ND	ND
615 ZINC				ND	ND	ND	ND
616 ZINC				ND	ND	ND	ND
617 ZINC				ND	ND	ND	ND
618 ZINC				ND	ND	ND	ND
619 ZINC				ND	ND	ND	ND
620 ZINC				ND	ND	ND	ND
621 ZINC				ND	ND	ND	ND
622 ZINC				ND	ND	ND	ND
623 ZINC				ND	ND	ND	ND
624 ZINC				ND	ND	ND	ND
625 ZINC				ND	ND	ND	ND
626 ZINC				ND	ND	ND	ND
627 ZINC				ND	ND	ND	ND
628 ZINC				ND	ND	ND	ND
629 ZINC				ND	ND	ND	ND
630 ZINC				ND	ND	ND	ND
631 ZINC				ND	ND	ND	ND
632 ZINC				ND	ND	ND	ND
633 ZINC				ND	ND	ND	ND
634 ZINC				ND	ND	ND	ND
635 ZINC				ND	ND	ND	ND
636 ZINC				ND	ND	ND	ND
637 ZINC				ND	ND	ND	ND
638 ZINC				ND	ND	ND	ND
639 ZINC				ND	ND	ND	ND
640 ZINC				ND	ND	ND	ND
641 ZINC				ND	ND	ND	ND
642 ZINC				ND	ND	ND	ND
643 ZINC				ND	ND	ND	ND
644 ZINC				ND	ND	ND	ND
645 ZINC				ND	ND	ND	ND
646 ZINC				ND	ND	ND	ND
647 ZINC				ND	ND	ND	ND
648 ZINC				ND	ND	ND	ND
649 ZINC				ND	ND	ND	ND
650 ZINC				ND	ND	ND	ND
651 ZINC				ND	ND	ND	ND
652 ZINC				ND	ND	ND	ND
653 ZINC				ND	ND	ND	ND
654 ZINC				ND	ND	ND	ND
655 ZINC				ND	ND	ND	ND
656 ZINC				ND	ND	ND	ND
657 ZINC				ND	ND	ND	ND
658 ZINC				ND	ND	ND	ND
659 ZINC				ND	ND	ND	ND
660 ZINC				ND	ND	ND	ND
661 ZINC				ND	ND	ND	ND
662 ZINC				ND	ND	ND	ND
663 ZINC				ND	ND	ND	ND
664 ZINC				ND	ND	ND	ND
665 ZINC				ND	ND	ND	ND
666 ZINC				ND	ND	ND	ND
667 ZINC				ND	ND	ND	ND
668 ZINC				ND	ND	ND	ND
669 ZINC				ND	ND	ND	ND
670 ZINC				ND	ND	ND	ND
671 ZINC				ND	ND	ND	ND
672 ZINC				ND	ND	ND	ND
673 ZINC				ND	ND	ND	ND
674 ZINC				ND	ND	ND	ND
675 ZINC				ND	ND	ND	ND
676 ZINC				ND	ND	ND	ND
677 ZINC				ND	ND	ND	ND
678 ZINC				ND	ND	ND	ND
679 ZINC				ND	ND	ND	ND
680 ZINC				ND	ND	ND	ND
681 ZINC				ND	ND	ND	ND
682 ZINC				ND	ND	ND	ND
683 ZINC				ND	ND	ND	ND
684 ZINC				ND	ND	ND	ND
685 ZINC				ND	ND	ND	ND
686 ZINC				ND	ND	ND	ND
687 ZINC				ND	ND	ND	ND
688 ZINC				ND	ND	ND	ND
689 ZINC				ND	ND	ND	ND
690 ZINC				ND	ND	ND	ND
691 ZINC				ND	ND	ND	ND
692 ZINC				ND	ND	ND	ND
693 ZINC				ND	ND	ND	ND
694 ZINC				ND	ND	ND	ND
695 ZINC				ND	ND	ND	ND
696 ZINC				ND	ND	ND	ND
697 ZINC				ND	ND	ND	ND
698 ZINC				ND	ND	ND	ND
699 ZINC				ND	ND	ND	ND
700 ZINC				ND	ND	ND	ND
701 ZINC				ND	ND	ND	ND
702 ZINC				ND	ND	ND	ND
703 ZINC				ND	ND	ND	ND
704 ZINC				ND	ND	ND	ND
705 ZINC				ND	ND	ND	ND
706 ZINC				ND	ND	ND	ND
707 ZINC				ND	ND	ND	ND
708 ZINC				ND	ND	ND	ND
709 ZINC				ND	ND	ND	ND
710 ZINC				ND	ND	ND	ND
711 ZINC				ND	ND	ND	ND
712 ZINC				ND	ND	ND	ND
713 ZINC				ND	ND	ND	ND
714 ZINC							

Table A.2. Radionuclide and Chemical Activity/Concentrations in Soil and Sediment

Name of Analyte	CAS #	Background(a)	Background Reference	SOIL						SEDIMENT				
				100-KR-4 (DOE 1994f)	100-HR-1 (DOE 1993c)	100-BC-1 (DOE 1994d)	100-BC-5 (DOE 1993a)	100-N (b)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	100 Areas (Weiss 1993)	
1 ACENAPHTHENE	83-32-9	ND	DOE 1994a		210 µg/kg			ND						
2 ACENAPHTHYLENE	208-96-8	ND	DOE 1994a					ND						
3 ACETONE	67-64-1	ND	DOE 1994a					ND						
4 ACETOPHENONE	98-86-2	ND	DOE 1990b								ND	ND		
5 ACETYLAMINOFLUORENE, 2-	53-96-3	ND	DOE 1990b								ND			
6 ACRYLAMIDE	79-06-1	ND	DOE 1990b								ND			
7 ACROLEIN	107-02-8	ND	DOE 1990b								ND			
8 ACRYLONITRILE	107-13-1	ND	DOE 1990b								ND			
9 ACTINIUM 227	14952-40-0													
10 ALDRIN	309-00-2	ND	DOE 1994a											
11 ALLYL ALCOHOL	107-18-6	ND	DOE 1990b											
12 ALPHA, ALPHA-DIMETHYLPHENETHYLAMINE	99-88-9	ND	DOE 1990b								ND			
13 ALPHA-BHC	319-84-6	ND	DOE 1994a											
14 ALPHA-CHLORDANE	5103-71-9	ND	DOE 1994a											
15 ALUMINIUM	7429-90-5	13621 mg/kg	DOE 1994a	7700 mg/kg	9070 mg/kg	12500 mg/kg					26,700 mg/kg	8760 mg/kg	6750 mg/kg	9,350 mg/kg
16 ALUMINIUM NITRATE	13473-90-0													
17 ALUMINIUM SULFATE	10043-01-3													
18 AMERICIUM 241	7440-35-9				0.72 pCi/g	34 pCi/g								ND
19 AMERICIUM 242M	13981-54-9													
20 AMERICIUM 243	14993-75-0													
21 AMINOBYPHENYL, 4-	92-67-1	ND	DOE 1990b								ND			
22 AMINOMETHYL-3 ISOAZOLOL, 5-	2763-96-4	ND	DOE 1990b								ND			
23 AMITROLE	61-82-5	ND	DOE 1990b								ND			
24 AMMONIA	7664-41-7	16.0 mg/kg	DOE 1994a									12.8 mg/kg	12 mg/kg	
25 AMMONIUM	14798-03-9	ND	DOE 1990b											
26 AMMONIUM ACETATE	631-61-8													
27 AMMONIUM CARBONATE	506-87-6													
28 AMMONIUM CHLORIDE	12125-02-9													
29 AMMONIUM FLUORIDE	12125-10-8													
30 AMMONIUM NITRATE	6484-52-2													
31 AMMONIUM OXALATE	1113-38-8													
32 AMMONIUM SILICOFLUORIDE	1309-32-6													
33 AMMONIUM SULFATE	7783-20-2													
34 AMMONIUM SULFITE	10186-04-0													
35 AMMONIUM THIOSULFATE	7783-18-8													
36 ANILINE	62-53-3	ND	DOE 1990b								ND			
37 ANTHRACENE	120-12-7	ND	DOE 1994a		430 µg/kg			ND						
38 ANTIMONY	7440-36-0	ND	DOE 1994a			ND					ND	ND	ND	ND
39 ANTIMONY (III) NITRATE	20328-86-5													
40 ANTIMONY 124	7440-36-0													1.2 pCi/kg
41 ANTIMONY 125	14234-35-6													
42 ANTIMONY CHLORIDE	10025-9-19													
43 ARAMITE	140-57-8	ND	DOE 1990b								ND			
44 AROCHLOR-1221	11104-28-2	ND	DOE 1994a								ND		ND	
45 AROCHLOR-1232	11141-16-5	ND	DOE 1994a								ND		ND	
46 AROCHLOR-1260	11096-82-5	ND	DOE 1994a								ND		ND	
47 AROCLOR 1016 (PCB)	12674-11-2	ND	DOE 1994a								ND		ND	
48 AROCLOR 1242 (PCB)	53469-21-9	ND	DOE 1994a								ND		ND	
49 AROCLOR 1248 (PCB)	12672-29-6	ND	DOE 1994a								9.9 mg/kg		ND	
50 AROCLOR 1254 (PCB)	11091-69-1	ND	DOE 1994a								ND		ND	
51 ARSENIC	7440-38-2	7.6 mg/kg	DOE 1994a		47 mg/kg	2.2 mg/kg					ND	9.3 mg/kg	7.5 mg/kg	
52 ARSENIC TRIOXIDE	1327-53-3													
53 ASBESTOS	332-21-4													
54 AURAMINE	492-80-8	ND	DOE 1990b								ND			
55 BARIUM	7440-39-3	155.9 mg/kg	DOE 1994a	85 mg/kg	672 mg/kg *	484 mg/kg					133 mg/kg	260 mg/kg	67.3 mg/kg	120 mg/kg
56 BARIUM 133	13981-41-4													
57 BARIUM 140	7440-39-3					ND						ND		
58 BARIUM NITRATE	10022-31-8													

Table A.2. (contd)

Name of Analyte	CAS #	Background(a)	Background Reference	SOIL							SEDIMENT			
				100-KR-4 (DOE 1994f)	100-HR-1 (DOE 1993c)	100-BC-1 (DOE 1994d)	100-BC-5 (DOE 1993a)	100-N (b)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	100 Areas (Weiss 1993)	
59 BENZENE	71-43-2	ND	DOE 1994a					4.5 mg/kg		ND	ND	ND		
60 BENZENETHIOL	108-98-5	ND	DOE 1990b							ND				
61 BENZIDINE	92-87-5	ND	DOE 1990b							ND				
62 BENZO[a]ANTHRACENE	56-55-3	ND	DOE 1994a							ND				
63 BENZO[a]PYRENE	50-32-8	ND	DOE 1994a		940 µg/kg				ND	ND				
64 BENZO[b]FLUORANTHENE	205-99-2	ND	DOE 1994a		810 µg/kg *				ND	ND				
65 BENZO[G,H,I]PERYLENE	191-24-2	ND	DOE 1994a		890 µg/kg				ND	ND				
66 BENZO[j]FLUORANTHENE	94-58-6	ND	DOE 1990b		410 µg/kg				ND					
67 BENZO[k]FLUORANTHENE	207-08-9	ND	DOE 1994a		760 µg/kg				ND					
68 BENZOIC ACID	65-85-0	ND	DOE 1994a			1700 µg/kg								
69 BENZOQUINONE, P-	106-51-4	ND	DOE 1990b							ND				
70 BENZYL ALCOHOL	100-51-6	ND	DOE 1994a					ND						
71 BENZYL CHLORIDE	100-44-7	ND	DOE 1990b							ND				
72 BERYLLIUM	7440-41-7	1.6 mg/kg	DOE 1994a		4.7 mg/kg	0.49 mg/kg				6 mg/kg	.93 mg/kg	ND	1.1 mg/kg	
73 BERYLLIUM 7	7440-41-7					ND					ND			
74 BETA-BHC	319-85-7	ND	DOE 1994a											
75 BIS(2-CHLOROETHOXY)METHANE	111-91-1	ND	DOE 1994a					ND		ND				
76 BIS(2-CHLOROETHYL)ETHER	111-44-4	ND	DOE 1994a					ND		ND				
77 BIS(2-CHLOROISOPROPYL)ETHER	39635-32-9	ND	DOE 1994a					ND		ND				
78 BIS(2-ETHYLHEXYL) PHTHALATE	117-81-7	ND	DOE 1994a		68 mg/kg			ND		ND				
79 BIS(CHLOROMETHYL) ETHER	542-88-1	ND	DOE 1990b							ND				
80 BISMUTH	7440-69-9													
81 BISMUTH 212	14913-49-6													
82 BISMUTH 214	14733-03-0													
83 BORON	7440-42-8													
84 BROMOACETONE	598-31-2	ND	DOE 1990b							ND				
85 BROMODICHLOROMETHANE	75-27-4	ND	DOE 1994a					ND			ND	ND		
86 BROMOFORM	75-25-2	ND	DOE 1994a					ND		ND	ND	ND		
87 BROMOMETHANE	74-83-9	ND	DOE 1994a					ND		ND	ND	ND		
88 BROMOPHENYL-PHENYLETHER, 4-	101-56-3	ND	DOE 1994a					ND		ND				
89 BUTANOL, 1-	71-36-3													
90 BUTANONE	78-93-3							ND						
91 BUTANONE, 2-	78-93-3	ND	DOE 1994a								ND	ND		
92 BUTYL BENZYL PHTHALATE	85-68-7	ND	DOE 1994a					ND		ND				
93 BUTYL PHTHALATE, DI-N-	84-74-2	ND	DOE 1990b											
94 CADMIUM	7440-43-9	ND	DOE 1994a			1.8 mg/kg				1 mg/kg	ND	ND	2.70 mg/kg	
95 CADMIUM 109	14109-32-1													
96 CADMIUM NITRATE	10325-97-7													
97 CALCIUM	7440-70-2	21012 mg/kg	DOE 1994a	7730 mg/kg	8620 mg/kg	14500 mg/kg				33,200 mg/kg	40800 mg/kg	4460 mg/kg	9000 mg/kg	
98 CALCIUM 41	14092-95-6													
99 CALCIUM BICARBONATE	1317-65-3													
100 CARBAZOLE	86-74-8													
101 CARBAZOLE, 9H-	86-74-8													
102 CARBON 14	14762-75-5				34 pCi/g	2.48 pCi/g								
103 CARBON DISULFIDE	75-15-0	ND	DOE 1994a					ND		ND	ND	ND		
104 CARBON TETRACHLORIDE	56-23-5	ND	DOE 1994a					ND		ND	ND	ND		
105 CARBOPHENOTHION		ND	DOE 1990b											
106 CERIUM	7440-45-1													
107 CERIUM 141	13987-74-3					ND					ND		ND	
108 CERIUM 144	14762-78-6					ND					ND			
109 CESIUM 134	13987-70-9				0.04 pCi/g	ND					ND		0.29 pCi/g	
110 CESIUM 135	15726-30-4													
111 CESIUM 137	10045-97-3				2900 pCi/g	800 pCi/g					0.23 pCi/g	.23 pCi/g	6.0 pCi/g	
112 CHLORAL	75-87-6	ND	DOE 1990b											
113 CHLORDANE	57-74-9	ND	DOE 1990b								4.5 mg/kg	ND		
114 CHLORIDE	16887-00-6									1.1 mg/kg				
115 CHLORINE	7782-50-5	331.3 mg/kg	DOE 1994a											
116 CHLORNAPHAZINE	494-03-1	ND	DOE 1990b											

A.12

Table A.2. (contd)

Name of Analyte	CAS #	Background(a)	Background Reference	SOIL						SEDIMENT			
				100-KR-4 (DOE 1994f)	100-HR-1 (DOE 1993c)	100-BC-1 (DOE 1994d)	100-BC-5 (DOE 1993a)	100-N (b)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	100 Areas (Weiss 1993)
117 CHLORO-2,3-EPOXYPROPANE, 1-		ND	DOE 1990b							ND			
118 CHLORO-M-CRESOL, P-	59-50-7	ND	DOE 1990b							ND			
119 CHLOROACETALDEHYDE	107-20-0	ND	DOE 1990b										
120 CHLOROALKYL ETHERS		ND	DOE 1990b							ND			
121 CHLOROANILINE, 4-	106-47-8	ND	DOE 1994a					ND					
122 CHLOROBENZENE	108-90-7	ND	DOE 1994a					ND		ND	ND		
123 CHLOROBENZILATE	510-18-6	ND	DOE 1990b										
124 CHLORODIBROMOMETHANE	124-48-1	ND	DOE 1994a										
125 CHLOROETHANE	75-00-3	ND	DOE 1994a					ND		ND	ND		
126 CHLOROETHOXY ETHENE, 2-	110-75-8									ND	ND	ND	
127 CHLOROETHYL VINYL ETHER, 2-	110-75-8	ND	DOE 1990b							ND			
128 CHLOROFORM	67-66-3	ND	DOE 1994a			ND		ND		ND	ND	ND	
129 CHLOROMETHANE	74-87-3	ND	DOE 1994a					ND		ND	ND	ND	
130 CHLOROMETHYLMETHYL ETHER	107-30-2	ND	DOE 1990b							ND			
131 CHLOROMETHYLPHENOL, 4-3-	35421-08-0							ND					
132 CHLORONAPHTHALENE, 2-	91-58-7	ND	DOE 1994a					ND		ND			
133 CHLOROPHENOL, 2-	96-57-8	ND	DOE 1994a					ND		ND			
134 CHLOROPHENYL-PHENYL ETHER, 4-	7005-72-3	ND	DOE 1994a					ND					
135 CHLOROPROPIONITRILE, 3-	542-76-7	ND	DOE 1990b										
136 CHROMIC ACID	7738-94-5												
137 CHROMIUM	7440-47-3	24.1 mg/kg	DOE 1994a		114 mg/kg	20.2 mg/kg				259 mg/kg	28.9 mg/kg	13.8 mg/kg	122 mg/kg
138 CHROMIUM (IV)	15723-28-1												
139 CHROMIUM (VI)	18540-29-9												
140 CHROMIUM 51	14392-02-0					ND					ND	ND	ND
141 CHROMIUM NITRATE	13548-38-4												
142 CHROMIUM SULFATE	10101-53-8												
143 CHRYSENE	218-01-9	ND	DOE 1994a		920 µg/kg			ND		ND			
144 CITRIS RED #2	6358-53-8	ND	DOE 1990b										
145 COBALT	7440-48-4	17.6 mg/kg	DOE 1994a	14.2 mg/kg	9.9 mg/kg	16.4 mg/kg					34.1 mg/kg	ND	11.5 mg/kg
146 COBALT 58	13981-38-9					ND					ND		
147 COBALT 60	10198-40-0	ND	DOE 1990b		18000 pCi/g	310 pCi/g					0.78 pCi/g	0.78 pCi/g	4.9 pCi/g
148 COPPER	7440-50-8	25.9 mg/kg	DOE 1994a	9 mg/kg	140000 mg/kg	27.8 mg/kg				2850 mg/kg	ND	16.1 mg/kg	40 mg/kg
149 COPPER NITRATE	3251-23-8												
150 COPPER SULFATE	7558-98-7												
151 CRESOLS	1319-77-3	ND	DOE 1990b							ND			
152 CROTONALDEHYDE	123-73-9	ND	DOE 1990b							ND			
153 CURIUM 242	15510-73-3												
154 CURIUM 244	13981-15-2												
155 CURIUM 245	15621-76-8												
156 CYANIDE	57-12-5	ND	DOE 1990b			1.05 mg/kg					ND	ND	
157 CYANOGEN	460-19-5	ND	DOE 1990b										
158 CYANOGEN CHLORIDE	506-77-4	ND	DOE 1990b										
159 CYANOGEN BROMIDE	508-68-3												
160 CYCLOHEXYL-4,6-DINITROPHENOL, 2-	131-89-5	ND	DOE 1990b							ND			
161 D(2,4)	94-75-7	ND	DOE 1990b										
162 DDD, 4,4'	72-54-8	ND	DOE 1994a										
163 DDE, 4,4'	72-55-9	ND	DOE 1994a										
164 DDT, 4,4'	50-29-3	ND	DOE 1994a										
165 DELTA-BHC	319-86-8	ND	DOE 1994a										
166 DI-N-BUTYLPHTHALATE	84-74-2	ND	DOE 1994a					ND		ND			
167 DI-N-OCTYLPHTHALATE	117-84-0	ND	DOE 1994a					ND		ND			
168 DI-N-PROPYLNITROSAMINE	621-64-7	ND	DOE 1990b							ND			
169 DIBENZ(A,G)ANTHRACENE	53-70-3												
170 DIBENZ(A,H)ACRIDINE	226-36-8	ND	DOE 1990b							ND			
171 DIBENZ(A,H)ANTHRACENE	53-70-3	ND	DOE 1994a					ND		ND			
172 DIBENZ(A,I)ACRIDINE	224-42-0	ND	DOE 1990b							ND			
173 DIBENZO(A,E)PYRENE	192-65-4	ND	DOE 1990b							ND			
174 DIBENZO(A,H)PYRENE	189-64-0	ND	DOE 1990b							ND			

A.13

Table A.2. (contd)

Name of Analyte	CAS #	Background(a)	Background Reference	SOIL						SEDIMENT		
				100-KR-4 (DOE 1994f)	100-HR-1 (DOE 1993c)	100-BC-1 (DOE 1994d)	100-BC-5 (DOE 1993a)	100-N (b)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)
175 DIBENZO(A,H)PYRENE	189-55-9	ND	DOE 1990b							ND		
176 DIBENZO(C,G)CARBAZOLE, 7H-		ND	DOE 1990b							ND		
177 DIBENZOFURAN	132-64-9	ND	DOE 1994a		130 µg/kg					ND		
178 DIBROMOMETHANE	74-95-3	ND	DOE 1990b							ND		
179 DIBROMO-3-CHLOROPROPANE, 1,2-	98-12-8	ND	DOE 1990b							ND		
180 DIBROMOCHLOROMETHANE	124-48-1	ND	DOE 1994a					ND		ND		
181 DIBROMOETHANE, 1,2-	106-93-4	ND	DOE 1990b							ND		
182 DICHLORO-2-BUTANE, 1,4-	616-21-7	ND	DOE 1990b							ND		
183 DICHLOROBENZENE, 1,2-	95-50-1	ND	DOE 1994a					ND		ND		
184 DICHLOROBENZENE, 1,3-	541-73-1	ND	DOE 1994a					ND		ND		
185 DICHLOROBENZENE, 1,4-	106-46-7	ND	DOE 1994a					ND		ND		
186 DICHLOROBENZIDINE, 3,3'	91-94-1	ND	DOE 1994a					ND		ND		
187 DICHLORODIFLUOROMETHANE	75-71-8	ND	DOE 1990b					ND		ND		
188 DICHLOROETHANE, 1,1-	75-34-3	ND	DOE 1994a					ND		ND		
189 DICHLOROETHANE, 1,2-	107-08-2	ND	DOE 1994a					ND		ND		
190 DICHLOROETHENE	25323-30-2	ND	DOE 1990b					ND		ND		
191 DICHLOROETHYLENE, 1,1-	75-35-4	ND	DOE 1994a					ND		ND		
192 DICHLOROETHYLENE, 1,2-	540-59-0	ND	DOE 1994a					ND		ND		
193 DICHLOROETHYLENE, 1,2-cis-	156-59-2									ND		ND
194 DICHLOROETHYLENE, 1,2-trans-	156-60-5											ND
195 DICHLOROMETHYLBENZENE		ND	DOE 1990b							ND		
196 DICHLOROPHENOL, 2,4-	120-83-2	ND	DOE 1994a					ND		ND		
197 DICHLOROPHENOL, 2,6-	87-65-0							ND		ND		
198 DICHLOROPHENOXYACETIC ACID, 2,4-	94-75-7									ND		
199 DICHLOROPROPANE, 1,2-	78-87-5	ND	DOE 1994a					ND		ND		
200 DICHLOROPROPANE, 1,3-	142-28-9	ND	DOE 1990b							ND		
201 DICHLOROPROPANOL	26545-73-3	ND	DOE 1990b							ND		
202 DICHLOROPROPENE, 1,3-cis-	10061-02-6	ND	DOE 1994a					ND		ND		
203 DICHLOROPROPENE, 1,3-trans-	10061-01-5	ND	DOE 1994a					ND		ND		
204 DIELDRIN	60-57-1	ND	DOE 1994a							ND		ND
205 DIESEL FUEL								2800 mg/kg				
206 DIETHYLARSINE	692-42-2	ND	DOE 1990b							ND		
207 DIETHYLHYDRAZINE, N,N	1615-80-1	ND	DOE 1990b							ND		
208 DIETHYLPHthalate	84-86-2	ND	DOE 1990b					ND		ND		
209 DIMETHOXYBENZIDINE, 3,3'	119-90-4	ND	DOE 1994a							ND		
210 DIETHYLSTILBESTEROL	56-53-1	ND	DOE 1990b							ND		
211 DIHYDROSAFROLE	94-58-6	ND	DOE 1990b							ND		
212 DIMETHOATE	60-51-5	ND	DOE 1990b							ND		
213 DIMETHYLBENZIDINE, 3,3'	119-93-7	ND	DOE 1990b							ND		
214 DIMETHYLHYDRAZINE, 1,1-	57-14-7									ND		
215 DIMETHYLHYDRAZINE, 1,2-		ND	DOE 1990b									
216 DIMETHYLPHENOL, 2,4-	105-67-9	ND	DOE 1994a					ND		ND		
217 DIMETHYLPHthalate	131-11-3	ND	DOE 1994a					ND		ND		
218 DIMENTYLAMINOAZOBENZENE, P-	60-11-7	ND	DOE 1990b							ND		
219 DIMENTYLBENZ(A)ANTHRACENE, 7,12-		ND	DOE 1990b							ND		
220 DINITRO-2-METHYLPHENOL, 4,6-	534-42-1	ND	DOE 1994a					ND		ND		
221 DINITRO-O-CRESOL, 4,6- and salts	534-52-1	ND	DOE 1990b							ND		
222 DINITROBENZENE	25154-54-5	ND	DOE 1990b							ND		
223 DINITROPHENOL, 2,4-	51-28-5	ND	DOE 1994a					ND		ND		
224 DINITROPHENOL, 2-SEC-BUTYL-4,6-		ND	DOE 1990b							ND		
225 DINITROTOLUENE, 2,4-	121-14-2	ND	DOE 1994a					ND		ND		
226 DINITROTOLUENE, 2,6-	606-20-2	ND	DOE 1994a					ND		ND		
227 DIOXANE		ND	DOE 1990b							ND		
228 DIOXIN		ND	DOE 1990b							ND		
229 DIPHENYLAMINE	122-39-4	ND	DOE 1990b							ND		
230 DIPHENYLHYDRAZINE, 1,2-	122-66-7	ND	DOE 1990b							ND		
231 DISULFOTON	298-04-4	ND	DOE 1990b							ND		
232 ENDOSULFAN I	959-98-8	ND	DOE 1994a									

Table A.2. (contd)

Name of Analyte	CAS #	Background(a)	Background Reference	SOIL						SEDIMENT				
				100-KR-4 (DOE 1994f)	100-HR-1 (DOE 1993c)	100-BC-1 (DOE 1994d)	100-BC-5 (DOE 1993a)	100-N (b)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	100 Areas (Weiss 1993)	
233 ENDOSULFAN II	33213-65-9	ND	DOE 1994a											
234 ENDOSULFAN SULFATE	1031-07-8	ND	DOE 1994a											
235 ENDRIN	72-20-8	ND	DOE 1994a											
236 ENDRIN ALDEHYDE	7421-93-4	ND	DOE 1994a			3.3 µg/kg								
237 ENDRIN KETONE	53494-70-5	ND	DOE 1994a											
238 ETHYL CARBAMATE	51-79-6	ND	DOE 1990b											
239 ETHYL CYANIDE	107-12-0	ND	DOE 1990b											
240 ETHYL METHANESULFONATE	62-50-0	ND	DOE 1990b							ND				
241 ETHYLBENZENE	100-41-4	ND	DOE 1990b					32 mg/kg						
242 ETHYLENE GLYCOL	107-21-1	ND	DOE 1990b											
243 ETHYLENE OXIDE	100-41-4	ND	DOE 1994a											
244 ETHYLENEIMINE	151-56-4	ND	DOE 1990b							ND	ND	ND		
245 ETHYLENE THIOUREA	96-45-7	ND	DOE 1990b											
246 ETHYL METHACRYLATE	97-63-2	ND	DOE 1990b							ND				
247 EUROPIUM 152	14683-23-9				59000 pCi/g	1400 pCi/g					0.17 pCi/g	.17 pCi/g	2.41 pCi/g	
248 EUROPIUM 154	15585-10-1				20000 pCi/g	410 pCi/g							ND	0.24 pCi/g
249 EUROPIUM 155	14391-16-3				6200 pCi/g	41 pCi/g								0.32 pCi/g
250 FERRIC NITRATE	10421-48-4													
251 FERRIC SULFATE	10028-22-5													
252 FERROCYANIDE	13408-63-4													
253 FERROUS AMMONIUM SULFATE	7783-85-9													
254 FERROUS SULFATE	7720-78-7													
255 FLUORANTHENE	206-44-0	ND	DOE 1994a		1800 pCi/g			ND		ND				
256 FLUORENE	86-73-7	ND	DOE 1994a		190 pCi/g			ND						
257 FLUORIDE	7782-41-4										2.0 mg/kg	4.7 mg/kg	ND	
258 FLUORINE	7782-41-4	5.3 mg/kg	DOE 1994a											
259 FLUOROACETIC ACID	144-49-0	ND	DOE 1990b											
260 FLUOROTRICHLOROMETHANE	75-69-4													
261 FORMALIN		ND	DOE 1990b							ND				
262 FUEL OIL #2	68476-34-6													
263 GAMMA-BHC(LINDANE)	58-89-9	ND	DOE 1994a											
264 GAMMA-CHLORDANE	5103-74-2	ND	DOE 1994a											
265 GASOLINE								ND						
266 GLYCIDYLALDEHYDE	765-34-4	ND	DOE 1990b											
267 HAPHTHYLAMINE, 2-		ND	DOE 1990b											
268 HEPTACHLOR	76-44-8	ND	DOE 1994a											
269 HEPTACHLOR EPOXIDE	1024-57-3													
270 HEPTACHLOR EPOXIDE (ENDO)	1024-57-3	ND	DOE 1994a											
271 HEPTACHLOR EPOXIDE (EXO)	1024-57-3	ND	DOE 1994a											
272 HEXACHLOROBENZENE	118-74-1	ND	DOE 1994a						ND	ND				
273 HEXACHLOROBUTADIENE	87-68-3	ND	DOE 1994a						ND	ND				
274 HEXACHLOROCYCLOPENTADIENE	77-47-4	ND	DOE 1994a						ND	ND				
275 HEXACHLOROETHANE	67-72-1	ND	DOE 1994a						ND	ND				
276 HEXACHLOROPHENE	70-30-4	ND	DOE 1990b							ND	ND			
277 HEXACHLOROPROPENE	1888-71-7	ND	DOE 1990b							ND	ND			
278 HEXANONE, 2-	591-78-6	ND	DOE 1994a					ND			ND	ND		
279 HEXONE	108-10-1									ND				
280 HEXYL METHANOATE	629-33-4													
281 HYDRAZINE	302-01-2	ND	DOE 1990b											
282 HYDROCARBONS														
283 HYDROCHLORIC ACID	7647-01-0													
284 HYDROCYANIC ACID	74-90-8													
285 HYDROFLUORIC ACID	7664-39-3													
286 HYDROGEN SULFIDE	7783-06-4	ND	DOE 1990b							ND				
287 INDENO(1,2,3-CD)PYRENE	193-39-5	ND	DOE 1994a		520 µg/kg *			ND		ND				
288 IODINE 129	15046-84-1													
289 IODINE 131	10043-66-0							ND				ND		
290 IODOMETHANE		ND	DOE 1990b							ND				

Table A.2. (contd)

Name of Analyte	CAS #	Background(a)	Background Reference	SOIL						SEDIMENT				
				100-KR-4 (DOE 1994f)	100-HR-1 (DOE 1993c)	100-BC-1 (DOE 1994d)	100-BC-5 (DOE 1993a)	100-N (b)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	100 Areas (Weiss 1993)	
291 IRON	7439-89-6	35746 mg/kg	DOE 1994a	25500 mg/kg	19000 mg/kg	44600 mg/kg					33,500 mg/kg	19500 mg/kg	17000 mg/kg	171000 mg/kg
292 IRON 59						ND						ND	ND	ND
293 ISOBUTYL ALCOHOL	78-83-1	ND	DOE 1990b											
294 ISOPHORONE	78-59-1	ND	DOE 1994a					ND						
295 ISOSAFROLE	120-58-1	ND	DOE 1990b								ND			
296 KEROSENE	8008-20-6	ND	DOE 1990b					3085 mg/kg			ND			
297 KRYPTON 85														
298 LANTHANUM	7439-91-0													
299 LEAD	7439-92-1	12.6 mg/kg	DOE 1994a	7.6 µg/L	540 mg/kg	4.8 mg/kg					ND	15.6 mg/kg	17.4 mg/kg	73 mg/kg
300 LEAD 210	14255-04-0													
301 LEAD 212	15092-94-1													
302 LEAD NITRATE	10099-74-8													
303 LITHIUM	7439-93-2	35 mg/kg	DOE 1994a											
304 LITHIUM CHLORIDE	7447-41-8													
305 MAGNESIUM	7439-95-4	8169 mg/kg	DOE 1994a	5030 mg/kg	4720 mg/kg	6390 mg/kg					11,600 mg/kg	8540 mg/kg	4020 mg/kg	7600 mg/kg
306 MALEIC HYDRAZIDE	123-33-1	ND	DOE 1990b								ND			
307 MALONONITRILE	109-77-3	ND	DOE 1990b								ND			
308 MANGANESE	7439-96-5	548 mg/kg	DOE 1994a	330 mg/kg	3050 mg/kg	839 mg/kg					396 mg/kg	403.2 mg/kg	327 mg/kg	578 mg/kg
309 MANGANESE 54	13966-31-9					ND						ND		0.057 pCi/g
310 MELPHALAN	148-82-3	ND	DOE 1990b								ND			
311 MERCURIC NITRATE	10045-95-0													
312 MERCURIC THIOCYANATE	592-85-8													
313 MERCURY	7439-97-6	0.61 mg/kg	DOE 1994a	1.4 mg/kg	1.1 mg/kg	4.3 mg/kg					2.77 mg/kg	.54 mg/kg	ND	ND
314 METHACRYLONITRILE	126-98-7	ND	DOE 1990b								ND			
315 METHANAL	50-00-0													
316 METHANETHIOL	74-93-1	ND	DOE 1990b								ND			
317 METHANOL	67-56-1													
318 METHAPYRILENE	91-80-5	ND	DOE 1990b								ND			
319 METHOLONYL			DOE 1990b								ND			
320 METHOXYCHLOR	72-43-5	ND	DOE 1994a											
321 METHYL BROMIDE	74-83-9	ND	DOE 1990b								ND			
322 METHYL CHLORIDE	74-87-3	ND	DOE 1990b								ND			
323 METHYL ETHYL KETONE	78-93-3	ND	DOE 1990b								ND			
324 METHYL METHACRYLATE	80-62-6	ND	DOE 1990b								ND			
325 METHYL METHANESULFONATE	66-27-3	ND	DOE 1990b								ND			
326 METHYL PARATHION	298-00-0	ND	DOE 1990b								ND			
327 METHYL-2-(METHYLO)PROPIONALDEHYDE, 2-		ND	DOE 1990b								ND			
328 METHYL-2-PENTANONE, 4-	108-10-1	ND	DOE 1994a					ND				22 mg/kg	ND	
329 METHYLAZIRIDINE, 2-	75-55-8	ND	DOE 1990b								ND			
330 METHYLCHOLANTHRENE, 3-	56-49-5	ND	DOE 1990b								ND			
331 METHYLENE Bis(3,4,6-TRICHLOROPHENOL)	70-30-4													
332 METHYLENE CHLORIDE	75-09-2	ND	DOE 1994a	120 µg/kg		ND		ND			ND	ND	ND	
333 METHYLENE BIS(2-CHLOROANILINE), 4-4'	101-14-4	ND	DOE 1990b								ND			
334 METHYLHYDRAZINE		ND	DOE 1990b											
335 METHYLLACTONITRILE, 2-	75-86-5	ND	DOE 1990b								ND			
336 METHYLNAPHTHALENE, 2-	91-57-6	ND	DOE 1994a		42 µg/kg			ND						
337 METHYLPHENOL, 2-	95-48-7	ND	DOE 1994a					ND						
338 METHYLPHENOL, 4-	106-44-5	ND	DOE 1994a					ND						
339 METHYLPHENOL, 4-CHLORO-3-	59-50-7	ND	DOE 1994a											
340 METHYLTHIOURACIL	56-04-2	ND	DOE 1990b								ND			
341 METHOXYCHLOR	72-43-5	ND	DOE 1990b											
342 MOLYBDENUM	7439-98-7	ND	DOE 1994a											
343 PROPYLAMINE, N-	107-10-8	ND	DOE 1990b											
344 NAPHTHALENE	91-20-7	ND	DOE 1994a					ND			ND			
345 NAPHTHOQUINONE, 1,4-	130-15-4	ND	DOE 1990b								ND			
346 NAPHTHYLAMINE, 1-	91-59-8	ND	DOE 1990b								ND			
347 NAPHTHYLAMINE, 2-		ND	DOE 1990b											
348 NEPTUNIUM 237	13994-20-2													0.608 pCi/g

Table A.2. (contd)

Name of Analyte	CAS #	Background(a)	Background Reference	SOIL						SEDIMENT			
				100-KR-4 (DOE 1994f)	100-HR-1 (DOE 1993c)	100-BC-1 (DOE 1994d)	100-BC-5 (DOE 1993a)	100-N (b)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	100 Areas (Weiss 1993)
349 NEPTUNIUM 239	13968-59-7												
350 NICKEL	7440-02-0	22.2 mg/kg	DOE 1994a	18 mg/kg	132 mg/kg	24.3 mg/kg				221 mg/kg	17.2 mg/kg	13.3 mg/kg	19.7 mg/kg
351 NICKEL 59	14336-70-0												
352 NICKEL 63	13981-37-8				20000 pCi/g								
353 NICKEL FERROCYANIDE	14874-78-3												
354 NICKEL NITRATE	13138-45-9												
355 NICKEL SULFATE	7786-81-4												
356 NICOTINIC ACID		ND	DOE 1990b							ND			
357 NIOBIUM 95	13967-76-5												
358 NITRATE	14797-55-8			4.3 mg/kg		5.9 mg/kg				30.4 mg/kg	12.7 mg/kg	ND	
359 NITRIC ACID	7697-37-2												
360 NITRITE	14797-65-0											ND	ND
361 NITRO-O-TOLUIDINE, 5-	99-55-8	ND	DOE 1990b										
362 NITROANILINE, 2-	88-74-4	ND	DOE 1994a					ND					
363 NITROANILINE, 3-	99-09-2	ND	DOE 1994a					ND					
364 NITROANILINE, 4-	100-01-6	ND	DOE 1994a					ND					
365 NITROBENZENE	98-95-3	ND	DOE 1994a					ND					
366 NITROGEN OXIDE	10024-97-2									ND			
367 NITROPHENOL, 2-	88-75-5	ND	DOE 1994a					ND					
368 NITROPHENOL, 4-	100-02-7	ND	DOE 1994a					ND					
369 NITROSO-DI-N-PROPYLAMINE, N-	621-64-7	ND	DOE 1994a					ND					
370 NITROSO-N-METHYLURETHANE, N-	815-53-2	ND	DOE 1990b							ND			
371 NITROSODI-N-BUTYLAMINE, N-	924-16-3	ND	DOE 1990b							ND			
372 NITROSODIETHANOLAMINE, N-	1116-54-7	ND	DOE 1990b							ND			
373 NITROSODIETHYLAMINE, N-	55-18-5	ND	DOE 1990b							ND			
374 NITROSODIMETHYLAMINE, N-	62-75-9	ND	DOE 1990b							ND			
375 NITROSODIPHENYLAMINE, N-	86-30-6	ND	DOE 1994a					ND					
376 NITROSOMETHYLETHYLAMINE, N-	10595-95-6	ND	DOE 1990b							ND			
377 NITROSOMETHYLVINYLAMINE, N-	4549-40-0	ND	DOE 1990b							ND			
378 NITROSOMORPHOLINE, N-	59-89-2	ND	DOE 1990b							ND			
379 NITROSONORNICOTINE, N-	16543-55-8	ND	DOE 1990b							ND			
380 NITROSOPERIDINE, N-	100-75-4	ND	DOE 1990b							ND			
381 NITROSOPYRROLIDINE	930-55-2	ND	DOE 1990b							ND			
382 ORTHO-PHOSPHATE													
383 OSMIUM		ND	DOE 1990b							ND			
384 OXYBIS(1-CHLOROPROPANE), 2,2'-													
385 PALLADIUM	7440-05-3												
386 PARALDEHYDE	123-63-7	ND	DOE 1990b										
387 PARATHION	56-38-2	ND	DOE 1990b										
388 PENTACHLORO BENZENE	608-93-5	ND	DOE 1990b							ND			
389 PENTACHLOROETHANE	76-01-7	ND	DOE 1990b							ND			
390 PENTACHLORONITROBENZENE	82-68-8	ND	DOE 1990b							ND			
391 PENTACHLOROPHENOL	87-86-5	ND	DOE 1994a					ND		ND			
392 PERCHLORATE		ND	DOE 1990b										
393 PERCHLOROETHYLENE	127-18-4	ND	DOE 1990b							ND			
394 PHENACETIN	62-44-2	ND	DOE 1990b							ND			
395 PHENANTHRENE	85-01-8	ND	DOE 1994a		1500 µg/kg			ND					
396 PHENOL	108-95-2	ND	DOE 1994a					ND		ND			
397 PHENYLENEDIAMINE	25265-76-3	ND	DOE 1990b							ND			
398 PHENYLTHIOUREA	103-85-5	ND	DOE 1990b							ND			
399 PHOSPHATE	7601-54-9	ND	DOE 1990b							ND	ND	ND	
400 PHOSPHORIC ACID	7664-38-2												
401 PHOSPHORUS	7723-14-0												
402 PHOSPHORUS 32													
403 PHTHALIC ACID ESTERS		ND	DOE 1990b							ND			
404 PICOLINE, 2-	109-06-8	ND	DOE 1990b							ND			
405 PLUTONIUM 238	13981-16-3				11 pCi/g	0.047 pCi/g						ND	0.00115 pCi/g
406 PLUTONIUM 239	15117-48-3			0.16 pCi/g	230 pCi/g	ND						ND	0.071 pCi/g

A.17

Table A.2. (contd)

Name of Analyte	CAS #	Background(a)	Background Reference	SOIL						SEDIMENT			
				100-KR-4 (DOE 1994f)	100-HR-1 (DOE 1993c)	100-BC-1 (DOE 1994d)	100-BC-5 (DOE 1993a)	100-N (b)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	100 Areas (Weiss 1993)
407 PLUTONIUM 240	14119-32-5				(w/Pu239)	ND							
408 PLUTONIUM 241	14119-32-6											ND	
409 PLUTONIUM 242	13982-10-0												
410 POLONIUM 210	13981-52-8												
411 POLONIUM 212	15389-34-1												
412 POLONIUM 216	15756-58-8												
413 POLYCHLORINATED BIPHENYLS	1336-36-3												
414 POTASSIUM	7447-40-7	2676 mg/kg	DOE 1994a	1360 mg/kg	13000 pCi/g	2130 mg/kg				1830 mg/kg	4980 mg/kg	ND	1900 mg/kg
415 POTASSIUM 40				16 pCi/g	15 pCi/g	13.86 pCi/g					ND	15 pCi/g	23 pCi/g
416 POTASSIUM CHLORATE	3811-04-9												
417 POTASSIUM CYANIDE	151-50-8												
418 POTASSIUM DICHROMATE	7778-50-9												
419 POTASSIUM FLUORIDE	7789-23-3												
420 POTASSIUM HYDROXIDE	1310-58-3												
421 POTASSIUM NITRATE	7757-79-1												
422 POTASSIUM PERMANGANATE	7722-64-7												
423 PROMETHIUM 147	7440-12-2												
424 PRONAMIDE	23950-58-5	ND	DOE 1990b							ND			
425 PROPYN-1-O1, 2-	107-19-7	ND	DOE 1990b										
426 PROTACTINIUM 231	14331-85-2												
427 PROTACTINIUM 233	13981-14-1												
428 PYRENE	129-00-0	ND	DOE 1994a		1200 µg/kg			ND					
429 PYRIDINE	110-86-1	ND	DOE 1990b							ND			
430 RADIUM	7440-14-4											ND	
431 RADIUM 223													
432 RADIUM 226	13982-63-3	ND	DOE 1990b	0.53 pCi/g	0.85 pCi/g	0.84 pCi/g					3.09 pCi/g	.71 pCi/g	1.7 pCi/g
433 RADIUM 228													ND
434 RADON 220	22481-48-7												
435 RESERPINE	50-56-5	ND	DOE 1990b							ND			
436 RESORCINOL	108-46-3	ND	DOE 1990b							ND			
437 RUTHENIUM 103	13968-53-1					ND					ND		
438 RUTHENIUM 106	13967-48-1					ND					ND	ND	ND
439 SAFROL	94-59-7	ND	DOE 1990b							ND			
440 SAMARIUM 151	15705-94-3												
441 SCANDIUM 46													
442 SECBUTYL-4,6-DINITROPHENOL													
443 SELENIUM	7782-49-2	ND	DOE 1994a			4.2 mg/kg				ND	ND	ND	
444 SELENIUM 79	15758-45-9												
445 SELENIUM CHLORIDE	10025-68-0												
446 SELENIUM NITRATE													
447 SILVER	7440-22-4	1.48 mg/kg	DOE 1994a			1.9 mg/kg				18 mg/kg	ND	ND	2.5 mg/kg
448 SILVER CHLORIDE	7783-90-6										17300 mg/kg		
449 SILVER NITRATE	7781-88-8												
450 SILVER OXIDE	20667-12-3												
451 SODIUM	7440-23-5	969 mg/kg	DOE 1994a	1770 mg/kg		779 mg/kg				401 mg/kg		ND	920 mg/kg
452 SODIUM 22	7440-23-5				ND								0.13 pCi/g
453 SODIUM ALUMINATE													
454 SODIUM CHLORIDE	7647-14-5												
455 SODIUM DICHROMATE	10588-01-9												
456 SODIUM FLUORIDE	7681-49-4												
457 SODIUM HYDROXIDE	1310-73-2												
458 SODIUM HYPOCHLORITE	7681-52-9												
459 SODIUM NITRATE	7631-99-4												
460 SODIUM PHOSPHATE, TRIBASIC	7601-54-9												
461 SODIUM SILICATE	1344-09-8												
462 SODIUM SULFATE	7757-82-6												
463 SODIUM SULFIDE	1313-82-2												
464 SODIUM THIOCYANATE	540-72-7												

Table A.2. (contd)

Name of Analyte	CAS #	Background(a)	Background Reference	SOIL						SEDIMENT				
				100-KR-4 (DOE 1994f)	100-HR-1 (DOE 1993c)	100-BC-1 (DOE 1994d)	100-BC-5 (DOE 1993a)	100-N (b)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	100 Areas (Weiss 1993)	
465 STRONTIUM	10476-85-4													
466 STRONTIUM 89	14158-27-1										67 mg/kg			
467 STRONTIUM 90	10098-97-2													
468 STRONTIUM CHLORIDE	10476-85-4			1 µg/kg	950 pCi/g	770 pCi/g						ND	ND	207pCi/g
469 STRYCHNINE	67-24-9	ND	DOE 1990b								ND			
470 STYRENE	100-42-6	ND	DOE 1994a					ND						
471 SULFATE	12808-79-8											ND	ND	ND
472 SULFIDE	18496-25-8	ND	DOE 1990b			32 mg/kg					52 mg/kg	131 mg/kg		ND
473 SULFUR OXIDES	20901-21-7													
474 SULFURIC ACID	7664-93-9	ND	DOE 1990b											
475 T (2,4,4)	7664-93-9													
476 SYM-TRINITROBENZENE		ND	DOE 1990b								ND			
477 T(2,4,5)	93-76-5													
478 TECHNETIUM 99	14133-76-7				0.67 pCi/g	ND							ND	0.5 pCi/g
479 TETRACHLOROBENZENE, 1,2,3,4-		ND	DOE 1990b								ND			
480 TETRACHLOROBENZENE, 1,2,3,5-		ND	DOE 1990b								ND			
481 TETRACHLOROBENZENE, 1,2,4,5-	95-94-3	ND	DOE 1990b								ND			
482 TETRACHLORODIBENZO-p-DIOXIN, 2,3,7,8-	1748-01-6													
483 TETRACHLOROETHANE, 1,1,1,2-	630-20-6	ND	DOE 1990b								ND			
484 TETRACHLOROETHANE, 1,1,2,2-	79-34-5	ND	DOE 1994a								ND			
485 TETRAETHYL PYROPHOSPHATE	107-49-3	ND	DOE 1994a					ND			ND		ND	
486 TETRACHLOROETHYLENE	127-18-4	ND	DOE 1994a					ND				ND	ND	
487 TETRACHLOROMETHANE	56-23-6													
488 TETRAHYDROFURAN	109-99-9													
489 THALLIUM	7440-28-0	ND	DOE 1994a			ND					ND	ND	ND	
490 THALLIUM 208	14913-50-9													
491 THILFANOX		ND	DOE 1990b								ND			
492 THIOUREA	62-56-6	ND	DOE 1990b											
493 THIOUREA, 1-(O-CHLOROPHENYL)-		ND	DOE 1990b											
494 THIOUREA, 1-ACETYL-2-		ND	DOE 1990b											
495 THIOUREA, 1-NAPHTHY-2-		ND	DOE 1990b											
496 THIRAM	137-26-8	ND	DOE 1990b								ND			
497 THORIUM 228					0.95 pCi/g	1.1 pCi/g						1.61 pCi/g	1.4 pCi/g	3 pCi/g
498 THORIUM 229	15595-54-4													
499 THORIUM 230	14268-63-7													
500 THORIUM 231														0.454 pCi/g
501 THORIUM 232				1.1 µg/kg	0.89 pCi/g	0.8 pCi/g								3.2 pCi/g
502 THORIUM 234								ND				1.1 pCi/g	1.1 pCi/g	0.812 pCi/g
503 TIN	7440-31-5												ND	
504 TIN 113	13966-06-8													
505 TIN 126	15832-50-6													
506 TITANIUM	7440-32-6	2925 mg/kg	DOE 1994a											
507 TITANIUM CHLORIDE	10049-06-6													
508 TOLUENE	108-88-3	ND	DOE 1994a		49 µg/kg			350 mg/kg			ND	ND	ND	
509 TOLUENEDIAMINE	496-72-0	ND	DOE 1990b								ND			
510 TOLUIDINE HYDROCHLORIDE, O-	636-21-5	ND	DOE 1990b								ND			
511 TOTAL ORGANIC CARBON		ND	DOE 1990b											
512 TOTAL ORGANIC HALIDE		ND	DOE 1990b											
513 TOTAL PETROLEUM HYDROCARBONS									125920 mg/kg					
514 TOXAPHENE	8001-35-2	ND	DOE 1994a											
515 TPI(2,4,5)SILVEX	93-72-1	ND	DOE 1990b											
516 TRIBROMOMETHANE	75-25-2													
517 TRIBUTYL PHOSPHATE	126-73-8													
518 TRIBUTYLPHOSPHORIC ACID		ND	DOE 1990b								ND			
519 TRICHLOROBENZENE		ND	DOE 1990b											
520 TRICHLOROBENZENE, 1,2,3-	87-61-6	ND	DOE 1990b								ND			
521 TRICHLOROBENZENE, 1,2,4-	120-82-1	ND	DOE 1994a					ND			ND			
522 TRICHLOROBENZENE, 1,3,5-	108-70-3	ND	DOE 1990b											

A19

Table A.2. (contd)

Name of Analyte	CAS #	Background(a)	Background Reference	SOIL						SEDIMENT			
				100-KR-4 (DOE 1994f)	100-HR-1 (DOE 1993c)	100-BC-1 (DOE 1994d)	100-BC-5 (DOE 1993a)	100-N (b)	1100 Area (Law 1990)	300-FF-1 (DOE 1990b)	300-FF-5 (DOE 1990a)	300-FF-5 (DOE 1990a)	100 Areas (Weiss 1993)
523 TRICHLOROETHANE, 1,1,1-	71-55-6	ND	DOE 1994a					ND		ND	ND	ND	
524 TRICHLOROETHANE, 1,1,2-	79-00-5	ND	DOE 1994a					ND		ND	ND	ND	
525 TRICHLOROETHYLENE	79-01-6	ND	DOE 1994a			ND		ND		ND	ND	ND	
526 TRICHLOROMETHANETHIOL	75-70-7	ND	DOE 1990b							ND			
527 TRICHLOROMONOFUOROMETHANE	75-96-4	ND	DOE 1990b							ND			
528 TRICHLOROPHENOL, 2,4,5-	95-95-4	ND	DOE 1994a					ND		ND			
529 TRICHLOROPHENOL, 2,4,6-	88-06-2	ND	DOE 1994a					ND		ND			
530 TRICHLOROPROPANE	25735-29-9	ND	DOE 1990b							ND			
531 TRICHLOROPROPANE, 1,2,3-	96-18-4	ND	DOE 1990b							ND			
532 TRIETHYLPHOSPHOROTHIOATE, O,O,O-	126-68-1	ND	DOE 1990b							ND			
533 TRIS (2,3 DIBROMOPROPYL) PHOSPHATE	126-72-7	ND	DOE 1990b							ND			
534 TRITIUM (HYDROGEN 3)	10028-17-8				1600 pCi/g	ND							
535 TUNGSTEN	7440-33-7												
536 URANIUM	7440-61-1												
537 URANIUM (TOTAL ACTIVITY)										ND	ND		
538 URANIUM 233	13968-55-3				0.59 pCi/g	0.53 pCi/g	0.6 pCi/g			3.9 pCi/g		2.3 pCi/g	
539 URANIUM 234	13966-29-5				(w/U233)		(w/U233)			w/U233	3.9 pCi/g	w/U233	
540 URANIUM 235	16117-96-1						0.0016 pCi/g	0.02 pCi/g		0.23 pCi/g	ND	0.1 pCi/g	
541 URANIUM 236	13982-70-2												
542 URANIUM 238	24678-82-8				0.59 pCi/g	4.7 pCi/g	0.62 pCi/g			3.2 pCi/g	3.2 pCi/g	2.3 pCi/g	
543 VANADIUM	7440-62-2	96.7 mg/kg	DOE 1994a	55.8 mg/kg	389 mg/kg	76.9 mg/kg				73 mg/kg	ND	44.4 mg/kg	82.2 mg/kg
544 VANADIUM PENTOXIDE	13140-62-1												
545 VINYL ACETATE	108-05-4	ND	DOE 1994a					ND			ND	ND	
546 VINYL CHLORIDE	75-01-4	ND	DOE 1994a					ND		ND	ND	ND	
547 WARFARIN	81-81-2	ND	DOE 1990b							ND			
548 XYLENE	1330-20-7	ND	DOE 1994a						1800 mg/kg		ND	ND	
549 XYLENE, m-	108-38-3	ND	DOE 1990b							ND			
550 XYLENE, o,p-		ND	DOE 1990b							ND			
551 YTTRIUM 90	10098-91-6												
552 ZINC	7440-66-6	74.7 mg/kg	DOE 1994a	24.3 mg/kg	520 mg/kg	309 mg/kg				97 mg/kg	70.7 mg/kg	118 mg/kg	397 mg/kg
553 ZINC 65	13982-39-3					ND					ND	ND	0.24 pCi/g
554 ZINC AMALGAM													
555 ZINC CHLORIDE	1646-85-7												
556 ZINC COMPOUNDS	7646-85-7												
557 ZINC NITRATE	7779-88-6												
558 ZIRCONIUM	7440-67-7	45.4 mg/kg	DOE 1994a										
559 ZIRCONIUM 93	15751-77-6												
560 ZIRCONIUM 95	13967-71-0					0.56 pCi/g	ND				ND		

(a) Provisional values estimated to be the background concentrations.

(b) Hartman and Lindsey (1993); Rowley 1993.

Table A.3. Maximum Detected Concentrations in Groundwater in the Hanford Site 100, 200, and 600 Areas Away from the Columbia River, 1980-1994

Name of Analyte	Number of Plumes	Concentration
100 Areas		
Chromium (+6)	3	1,570 ppb
Nitrate	10	130,000 ppb
Strontium-90	8	1,800 pCi/L
Tritium (Hydrogen-3)	4	80,000 pCi/L
200 West Area		
Arsenic	4	24 ppb
Carbon Tetrachloride	1	6,559 ppb
Chloroform	2	1,595 ppb
Chromium	5	323 ppb
Fluoride	3	10,067 ppb
Iodine-129	2	30 pCi/L
Nitrate	5	1,322,000 ppb
Technetium-99	5	26,602 pCi/L
Trichloroethylene	3	32 ppb
Tritium (Hydrogen-3)	3	6,193,000 pCi/L
Uranium	4	1,616 pCi/L
200 East Area		
Arsenic	4	24ppb
Cesium-137	1	1,326 pCi/L
Chloroform	1	7 ppb
Chromium	4	288 ppb
Cobalt-60	2	440 pCi/L
Cyanide	2	893 ppb
Iodine-129	3	20 pCi/L
Nitrate	7	397,000 ppb
Plutonium-239/240	1	69 pCi/L
Strontium-90	5	5,149 pCi/L
Technetium-99	2	22,163 pCi/L
Tritium (Hydrogen-3)	5	4,126,000 pCi/L
Uranium	1	27 pCi/L
600 Area (Solid Waste Landfill Site)		
Chloroform	1	0.5 ppb
Dichloroethane, 1, 1-	1	7 ppb
Tetrachloroethene	1	12 ppb
Trichloroethane, 1, 1, 1-	1	50 ppb
Trichloroethene	1	7 ppb

Appendix B

Parameter Values Used in Screening Analyses

Appendix B

Parameter Values Used in Screening Analyses

The equations detailed in Section 4.0 require parameters for each radionuclide and chemical evaluated. The parameters used to screen samples from the Columbia River and groundwater within 150 meters (500 feet) of the Columbia River are provided in Table B.1. The parameters used to screen samples of soil and sediment are provided in Table B.2. The parameters used to screen samples of groundwater farther than 150 meters (500) feet from the Columbia River are provided in Table B.3.

The following abbreviations are used in the tables:

- LC50 = lowest concentration reported to be lethal to aquatic life, as reported in EPA 1985.
- RfD = EPA chronic oral reference dose value.
- TLM = lowest concentration below which no effects on aquatic life are observed, as reported in EPA 1985.

Table B.1. Parameters Used to Screen Columbia River and Groundwater Near the Columbia River

Name of Analyte	Maximum Concentration in		Ingestion	External	RfD (mg/kg/day)	Cancer	Fish	LC50 (µg/L)	TLM (µg/L)	Notes on	Water Quality
	Surface Water	Groundwater	Slope Factor (Risk/pCi)	Slope Factor (Risk/pCi)		Potency Factor (1)/(mg/kg/day)	Bioaccumulation (L/kg)			Fish Toxicity	Criteria (µg/L)
1 ACETONE	11 µg/L	30 µg/L			0.1		0.2		4,000,000		
2 ALUMINUM		4,810 µg/L			0.0004		10	5,000		8	
3 AMERICIUM 241		0.021 pCi/L	2.40E-10	4.90E-09			250				
4 AMMONIA		70 µg/L			0.029		0.2	1,800		as ammonium	
5 AMMONIUM		1,830 µg/L			0.09		0.2		1,800	9	
6 ANTIMONY		60 µg/L			0.0004		200				
7 ANTIMONY 125		20 pCi/L	8.40E-13	1.20E-06			200				
8 ARSENIC	3.4 µg/L	17 µg/L			0.0003	1.75	100	1,100			190
9 BARIUM	48.2 µg/L	719 µg/L			0.07		200	400,000			
10 BERYLLIUM		6 µg/L			0.005	4.3	19	200			
11 BERYLLIUM 7										10	
12 BIS(2-ETHYLHEXYL) PHTHALATE		50 µg/L			0.02	0.014	70		32,000		
13 BISMUTH											
14 BORON		64 µg/L			0.09						
15 CADMIUM		31 µg/L			0.0005	6.3	200		30,000		1.1
16 CALCIUM	35,900 µg/L	302,000 µg/L									
17 CARBON 14		23,000 pCi/L	9.00E-13	0			4600				
18 CESIUM 134	0.012 pCi/L		4.10E-11	5.20E-06			2000				
19 CESIUM 137	0.13 pCi/L	0.5 pCi/L	2.80E-11	2.00E-06			2000				
20 CHLORIDE	870 µg/L	122,000 µg/L					50				
21 CHLOROFORM		42 µg/L			0.01	0.006	100	100,000			
22 CHROMIUM	22 µg/L	1,950 µg/L			1	41	200	1,000			11
23 COBALT		8 µg/L			0.0081		50		10,000,000		
24 COBALT 60	0.011 pCi/L	140 pCi/L	1.50E-11	8.60E-06			330				
25 COPPER	22 µg/L	516 µg/L			0.0003		50	500			12
26 CYANIDE		21.1 µg/L			0.02		0.2				5.2
27 DICHLOROETHYLENE, 1,2-		200 µg/L				0.009	2.9		5000		
28 DICHLOROETHYLENE, 1,2-trans-		130 µg/L			0.02	1.2	20				
29 EUROPIUM 154		2 pCi/L	3.00E-12	4.10E-06			25				
30 FLUORIDE	150 µg/L	2,080 µg/L			0.06		10		2,300	11	
31 HYDRAZINE		7 µg/L				3	0.5		2,000		
32 IODINE 129	0.16 pCi/L		1.90E-10	4.10E-09			15				
33 IRON	463 pCi/L	37,300 µg/L			1.3		2000				
34 LEAD		173 µg/L			0.0014		100		530		3.2
35 LITHIUM											
36 MAGNESIUM	9,860 µg/L	55,000 µg/L			50		50				
37 MANGANESE	22.8 µg/L	400 µg/L			0.07		400		500,000	12	
38 MERCURY		8.9 µg/L			0.0003		1000	10			0.012
39 METHYL ETHYL KETONE		18 µg/L			1		50	5,600,000			
40 METHYLENE CHLORIDE		3,040 µg/L			0.06	0.0075	2.5	550,000		13	
41 NICKEL	31 µg/L	479 µg/L			0.02		100		380		160

B.2

Table B.2. Parameters Used to Screen Soil and Sediment

	Maximum Concentration in		Ingestion	External	RfD (mg/kg/day)	Cancer	Fish	LC50 (µg/L)	TLM (µg/L)	Notes on	Water Quality
	Soil	Sediment	Slope Factor (Risk/pCi)	Slope Factor (Risk/pCi)		Potency Factor (1)/(mg/kg/day)	Bioaccumulation (L/kg)			Fish Toxicity	Criteria (µg/L)
Name of Analyte											
Radionuclides											
1 AMERICIUM 241	34 pCi/g		2.40E-10	4.90E-09			250				
2 ANTIMONY 124		1.2 pCi/g	2.90E-12	6.50E-06			200				
3 CARBON 14	34 pCi/g		9.00E-13	0			4,600				
4 CESIUM 134	0.04 pCi/g	0.29 pCi/g	4.10E-11	5.20E-08			2,000				
5 CESIUM 137	2,900 pCi/g	6 pCi/g	2.80E-11	2.00E-06			2,000				
6 COBALT 60	18,000 pCi/g	4.9 pCi/g	1.50E-11	8.60E-06			330				
7 EUROPIUM 152	59,000 pCi/g	2.41 pCi/g	2.10E-12	3.60E-06			25				
8 EUROPIUM 154	20,000 pCi/g	0.24 pCi/g	3.00E-12	4.10E-06			25				
9 EUROPIUM 155	6,200 pCi/g	0.32 pCi/g	4.50E-13	5.90E-08			25				
10 NEPTUNIUM 237		0.606 pCi/g	2.20E-10	7.80E-09			250				
11 NICKEL 63	20,000 pCi/g		2.40E-13	0			100				
12 PLUTONIUM 238	11 pCi/g	0.00115 pCi/g	2.20E-10	2.80E-11			250				
13 PLUTONIUM 239	230 pCi/g	0.071 pCi/g	2.30E-10	1.70E-11			250				
14 PLUTONIUM 240	(w/Pu239)		2.30E-10	2.70E-11			250				
15 POTASSIUM 40	16 pCi/g	23 pCi/g	1.10E-11	5.40E-07			1,000				
16 RADIUM 226	3.09 pCi/g	1.7 pCi/g	1.20E-10	1.20E-08			70				
17 STRONTIUM 90	950 pCi/g	207 pCi/g	3.30E-11	0			50				
18 TECHNETIUM 99	0.67 pCi/g	0.5 pCi/g	1.30E-12	6.00E-13			15				
19 THORIUM 228	1.61 pCi/g	3 pCi/g	1.10E-11	5.50E-10			100				
20 THORIUM 232	1.1 pCi/g	3.2 pCi/g	1.20E-11	2.60E-11			100				
21 THORIUM 234	ND	0.812 pCi/g	4.00E-12	3.50E-09			100				
22 TRITIUM (HYDROGEN 3)	1,600 pCi/g		5.40E-14	0			3,000				
23 URANIUM 233	3.9 pCi/g	2.3 pCi/g	1.60E-11	4.20E-11			50				
24 URANIUM 234		3.9 pCi/g	1.60E-11	3.00E-11			50				
25 URANIUM 235	1.23 pCi/g	0.1 pCi/g	1.60E-11	2.40E-07			50				
26 URANIUM 238	4.7 pCi/g	3.2 pCi/g	1.60E-11	2.10E-11			50				
27 ZINC 65	ND	0.24 pCi/g	8.50E-12	2.00E-06			2,500				
28 ZIRCONIUM 95	0.56 pCi/g		9.90E-13	2.50E-06			200				
Chemicals											
29 ACENAPHTHENE	210 µg/kg				0.06		300	4,000		1	
30 ALUMINIUM	26,700,000 µg/kg	9,350,000 µg/kg			0.004		10	5,000		7	
31 AMMONIA	12,800 µg/kg	12,000 µg/kg			0.029		0	1,800			
32 ANTHRACENE	430 µg/kg				0.3		3,000	4,000		1	
33 AROCLOR 1248 (PCB)	9,900 µg/kg					7.7	10,000		278		0.014
34 ARSENIC	47,000 µg/kg	7,500 µg/kg			0.0003	1.75	100	1,100			190

B.4

Table B.2. (contd)

	Name of Analyte	Maximum Concentration in		Ingestion	External	RfD	Cancer	Fish	LC50	TLM	Notes on	Water Quality
		Soil	Sediment	Slope Factor (Risk/pCI)	Slope Factor (Risk/pCI)	(mg/kg/day)	Potency Factor (1)/(mg/kg/day)	Bioaccumulation (L/kg)	(µg/L)	(µg/L)	Fish Toxicity	Criteria (µg/L)
35	BARIUM	672,000 µg/kg	120,000 µg/kg			0.07		200	400,000			
36	BENZENE	4,500 µg/kg					0.029	10		20		
37	BENZO(G,H,I)PERYLENE	410 µg/kg							4,000		1	
38	BENZO(a)ANTHRACENE	940 µg/kg					0.84	12,000	4,000		1	
39	BENZO(a)PYRENE	810 µg/kg					5.79	20,000	4,000		1	
40	BENZO(b)FLUORANTHENE	890 µg/kg					0.81	20,000	4,000		1	
41	BENZO(k)FLUORANTHENE	760 µg/kg					0.38	20,000	4,000		1	
42	BENZOIC ACID	1,700 µg/kg				4		6	180,000			
43	BERYLLIUM	8,000 µg/kg	1,100 µg/kg			0.005	4.3	19	200			
44	BIS(2-ETHYLHEXYL) PHTHALATE	68,000 µg/kg				0.02	0.014	70		32,000		
45	CADMIUM	1,800 µg/kg	2,700 µg/kg			0.0005	6.3	200		30,000		1.1
46	CALCIUM	40,800,000 µg/kg	4,460,000 µg/kg									
47	CHLORDANE	4,500 µg/kg				0.00006	1.3	322	8			0.0043
48	CHLORIDE	1,100 µg/kg				0.011		50				
49	CHLORINE (a)											
50	CHROMIUM	259,000 µg/kg	12,200 µg/kg			1	41	200	1,000			11
51	CHRYSENE	920 µg/kg					0.0255	20,000	4,000		1	
52	COBALT	34,100 µg/kg	11,500 µg/kg			0.0081		50		10,000,000		
53	COPPER	40,000,000 µg/kg	40,000 µg/kg			0.0003		50	500			12
54	CYANIDE	1,050 µg/kg				0.02		0				5.2
55	DIBENZOFURAN	130 µg/kg										
56	DIESEL FUEL	2,800,000 µg/kg				0.36		300	1,000			
57	ENDRIN ALDEHYDE	3.3 µg/kg				0.0003		1,480	0		2	
58	ETHYL BENZENE	32,000 µg/kg				0.1		100		30		
59	FLUORANTHENE	1,800 µg/kg				0.04		3,000	4,000		1	
60	FLUORENE	190 µg/kg				0.04		713	4,000		1	
61	FLUORIDE	4,700 µg/kg				0.04		10		2,300	3	
62	FLUORINE (a)											
63	INDENO(1,2,3-CD)PYRENE	520 µg/kg					1.34	40,000	4,000		1	
64	IRON	33,500,000 µg/kg	71,000,000 µg/kg			1.3		2,000				
65	KEROSENE	3,085,000 µg/kg				0.7		300		200		
66	LEAD	540,000 µg/kg	73,000 µg/kg			0.0014		100		530		3.2
67	LITHIUM (a)											
68	MAGNESIUM	11,600,000 µg/kg	7,600,000 µg/kg									
69	MANGANESE	839,000 µg/kg	578,000 µg/kg							500,000		
70	MERCURY	4,300 µg/kg				0.0003		1,000	10			0.012

B.5

Table B.2. (contd)

	Maximum Concentration In		Ingestion	External	RfD	Cancer	Fish	LC50	TLM	Notes on	Water Quality
	Soil	Sediment	Slope Factor	Slope Factor		Potency Factor	Bioaccumulation			Fish	Criteria
Name of Analyte			(Risk/pCI)	(Risk/pCI)	(mg/kg/day)	(1)/(mg/kg/day)	(L/kg)	(µg/L)	(µg/L)	Toxicity	(µg/L)
71 METHYL-2-PENTANONE, 4-	22,000 µg/kg										
72 METHYLENE CHLORIDE	120 µg/kg				0.06	0.0075	3	550,000		4	
73 METHYLNAPHTHALENE, 2-	42 µg/kg							4,000		1	
74 NICKEL	221,000 µg/kg	19,700 µg/kg			0.02		100		380		160
75 NITRATE	30,400 µg/kg				1.6		150,000	20,000		5	
76 PHENANTHRENE	1,500 µg/kg				0.04		1,000	4,000		1	
77 POTASSIUM	4,980,000 µg/kg	1,900,000 µg/kg						80,000			
78 PYRENE	1,200 µg/kg				0.03		2,800	4,000		1	
79 SELENIUM	4,200 µg/kg				0.005		170	2,500			5
80 SILVER	1,900 µg/kg	2,500 µg/kg						4			
81 SILVER CHLORIDE	17,300,000 µg/kg				0.005		2				
82 SODIUM	1,770,000 µg/kg	920,000 µg/kg							4,720,000		
83 STRONTIUM	67,000 µg/kg				0.6		50		200,000	6	
84 STRONTIUM CHLORIDE	1 µg/kg				0.6		50		200,000		
85 SULFATE (SULFUR)	131,000 µg/kg				71		750		80,000		
86 TITANIUM (a)											
87 TOLUENE	350,000 µg/kg				0.2		20	60,000			
88 TOTAL PETROLEUM HYDROCARBON	1.26E + 08										
89 VANADIUM	389,000 µg/kg	82,200 µg/kg							55,000		
90 XYLENE	1,800,000 µg/kg				0.2		150		4,000		
91 ZINC	309,000 µg/kg	397,000 µg/kg			0.3		2,500	430			110
92 ZIRCONIUM (a)											
(a) Concentrations of these chemicals fall within their respectively occurring background levels.											
Notes on Fish Toxicity											
1	assume naphthalene										
2	assume endrine										
3	assume fluorine										
4	assume chloromethane										
5	assume ferric nitrate										
6	assume strontium chloride										
7	assume aluminum hydroxide										

B.6

Table B.3. Parameters Used to Screen Groundwater Away from the Columbia River

Name of Analyte	Number of Plumes	Maximum Concentration	Reference	Ingestion Slope Factor (Risk/pCi)	External Slope Factor (Risk/pCi)	RfD (mg/kg/day)	Cancer Potency Factor (mg/kg/day)	Fish Bioaccumulation (L/kg)	LC50 (µg/L)	TLM (µg/L)	Water Quality Criteria (µg/L)
100 Areas											
Chromium (+ 6)	3	1,570 ppb	DOE 1994b			1	41	200	1,000		11
Nitrate	10	130,000 ppb	DOE 1994b			2		150,000	20,000		
Strontium-90	8	1,800 pCi/L	DOE 1994b	0	0			50			
Tritium (Hydrogen-3)	4	80,000 pCi/L	DOE 1994b	0	0			1			
200 West Area											
Arsenic	4	24 ppb	Ford 1993			0	2	100	1,100		190
Carbon Tetrachloride	1	6,559 ppb	Ford 1993			0	0	150	125,000		
Chloroform	2	1,595 ppb	Ford 1993			0	0	100	100,000		
Chromium	5	323 ppb	Ford 1993			1	41	200	1,000		11
Fluoride	3	10,067 ppb	Ford 1993			0		10		2,300	
Iodine-129	2	30 pCi/L	Ford 1993	0	0			15			
Nitrate	5	1,322,000 ppb	Ford 1993			2		150,000	20,000		
Technetium-99	5	26,602 pCi/L	Ford 1993	0	0			15			
Trichloroethylene	3	32 ppb	Ford 1993				0	11	55,000		
Tritium (Hydrogen-3)	3	6,193,000 pCi/L	Ford 1993	0	0			1			
Uranium	4	1,616 pCi/L	DOE 1994b	0	0			50			
200 East Area											
Arsenic	4	24 ppb	Ford 1993			0	2	100	1,100		190
Cesium-137	1	1,326 pCi/L	Ford 1993	0	0			2,000			
Chloroform	1	7 ppb	DOE 1994b			0	0	100	100,000		
Chromium	4	288 ppb	Ford 1993			1	41	200	1,000		11
Cobalt-60	2	440 pCi/L	Ford 1993	0	0			330			
Cyanide	2	893 ppb	Ford 1993			0		0			5
Iodine-129	3	20 pCi/L	Ford 1993	0	0			15			
Nitrate	7	397,000 ppb	Ford 1993			2		150,000	20,000		
Plutonium-239/240	1	69 pCi/L	Ford 1993								
Strontium-90	5	5,149 pCi/L	Ford 1993	0	0			50			
Technetium-99	2	22,163 pCi/L	Ford 1993	0	0			15			
Tritium (Hydrogen-3)	5	4,126,000 pCi/L	Ford 1993	0	0			1			
Uranium	1	27 pCi/L	Ford 1993	0	0			50			
600 Area (Solid Waste Landfill Site)											
Chloroform	1	0.5 ppb	DOE 1994b			0	0	100	100,000		
Dichloroethane, 1, 1-	1	7 ppb	DOE 1994b			0		7	220,000		
Tetrachloroethene	1	12 ppb	DOE 1994b			0	0	100	13,000		
Trichloroethane, 1, 1, 1-	1	50 ppb	DOE 1994b			0	0	39	50,000		
Trichloroethene	1	7 ppb	DOE 1994b			0	0	52	55,000		

B.7

Appendix C

Complete Numerical Results

Appendix C

Complete Numerical Results

This appendix provides the numerical results of applying the screening equations in Section 4.0 to the detected analytes described in Sections 3.0 and 7.0. Table C.1 presents the numerical results of screening samples at the Columbia River and groundwater within 150 meters (500 feet) of the Columbia River. Table C.2 presents the numerical results of screening soil and sediment samples. Table C.3 presents the numerical results of screening samples from groundwater farther than 150 meters (500 feet) from the Columbia River. Application of the equations and assumptions defined in Section 4.0 results in a series of complementary, but not necessarily intercomparable, screening values for each contaminant. The varying numbers of assumptions and associated varying degrees of conservatism require that each of the screenings be evaluated separately. The results of the combined screenings, however, then define the overall list of contaminants of concern.

Each table includes a "notes" column. The notes consist of abbreviated designations. The following are the full descriptions of each designation as well as explanations of the column headings.

Bkg	= background denotes that the highest concentration found was at background level so eliminated from consideration.
EPA-10	= eliminated based on the guidance in <i>EPA Region 10 Supplemental Risk Assessment Guidance for Superfund</i> (EPA 1991).
I	= parameters derived from the Integrated Risk Information System (IRIS) database (EPA 1994b).
Inadequate?	= insufficient information available to classify as toxic or having carcinogenic properties.
LC50/100	= lowest concentration reported to be lethal to aquatic life 100 days after exposure, as reported in EPA 1985.
LD	= near limit of detection.
M	= parameters derived from the Multimedia Environmental Pollutant Assessment System (MEPAS) database (Droppo et al. 1991).
ND	= not detected.
Non-Haz.?	= analyte not designated in database as containing hazardous properties.
Suspect	= noted in the source database as being unreliable (see Section 4.4).
SW	= surface water (Columbia River water).
SW-LD	= reported sample in surface water very near the limit of detection and, therefore, unreliable.
T 1/2	= half-life of analyte indicates that any concentration present at sampling should now be decayed to insignificance.
TLM	= lowest concentration below which no effects on aquatic life are observed, as reported in EPA 1985.
Unclass?	= not classified in MEPAS or IRIS as hazardous.
WQC	= water quality criteria.

Table C.1. Results for the Columbia River and Groundwater Near the Columbia River

	Name of Analyte	Notes	Carcinogenic Risk Ranking		Hazard Index Ranking		WQC Screen Ranking		LC50/100 Screen Ranking		TLM Screen Ranking	
			Surface	Ground-	Surface	Ground-	Surface	Ground-	Surface	Ground-	Surface	Ground-
1	ACETONE	M, SW-LD	M, EPA-10	4.80E-03	1.31E-05						2.75E-04	7.50E-09
2	ALUMINIUM											
3	AMERICIUM 241			1.42E-10								
4	AMMONIA											
5	AMMONIUM	M			1.05E-04						3.89E-03	
6	ANTIMONY											
7	ANTIMONY 125											
8	ARSENIC	I		2.58E-03	1.29E-05	4.92E+00	2.46E-02	1.79E-02	8.95E-05	3.09E-01	1.55E-03	
9	BARIUM	I, SW-Bkg			8.48E-03					1.21E-02	1.80E-04	
10	BERYLLIUM	I			3.02E-06						3.00E-03	
11	BERYLLIUM 7	Bkg										
12	BIS(2-ETHYLHEXYL) PHTHALATE	I			2.22E-07							
13	BISMUTH	Bkg, M										
14	BORON	Bkg, I										
15	CADMIUM	I			1.61E-04							1.03E-06
16	CALCIUM	Bkg, M, EPA-10										
17	CARBON 14											
18	CESIUM 134				6.34E-06							
19	CESIUM 137				2.67E-05							
20	CHLORIDE	M, SW-Bkg										
21	CHLOROFORM	I			1.09E-07							
22	CHROMIUM	I, SW-LD			6.60E-02						1.95E-01	
23	COBALT	M										
24	COBALT 60				9.47E-06							
25	COPPER	M, SW-LD			1.75E+01	4.10E-01	1.83E+00	4.30E-02	4.40E+00	1.03E-01		
26	CYANIDE	M										
27	DICHLOROETHYLENE, 1,2-	I			9.76E-08							4.00E-05
28	DICHLOROETHYLENE, 1,2-trans-	I			1.89E-05							
29	EUROPIUM 154				8.20E-07							
30	FLUORIDE	M, SW-Bkg										9.04E-04
31	HYDRAZINE											3.50E-06

Table C.1. (contd)

	Name of Analyte	Notes	Carcinogenic Risk Ranking		Hazard Index Ranking		WQC Screen Ranking		LC50/100 Screen Ranking		TLM Screen Ranking	
			Surface	Ground-	Surface	Ground-	Surface	Ground-	Surface	Ground-	Surface	Ground-
			Water	water	Water	water	Water	water	Water	water	Water	water
32	IODINE 129		1.44E-07									
33	IRON	M,EPA-10										
34	LEAD	M				5.37E-02		5.41E-02				3.26E-04
35	LITHIUM	Bkg,M										
36	MAGNESIUM	M,EPA-10										
37	MANGANESE	M			5.24E-01	9.19E-03					4.56E-03	8.00E-07
38	MERCURY	M				1.17E-01		7.42E-01		8.90E-02		
39	METHYL ETHYL KETONE	I				4.29E-06				3.21E-07		
40	METHYLENE CHLORIDE	I		1.20E-06		2.67E-03				5.53E-04		
41	NICKEL	M, SW-LD			6.73E-01	1.04E-02	1.94E-01	2.99E-03			8.16E+00	1.26E-03
42	NITRATE	M			1.76E+02	3.30E+01			2.40E+00	4.50E-01		
43	NITRITE					1.04E+03				3.00E-01		
44	PHOSPHATE	M				1.93E+00						5.49E-05
45	PLUTONIUM 238			5.74E-11								
46	PLUTONIUM 239			1.80E-10								
47	POTASSIUM	Bkg,M,EPA-10										
48	RADIUM 226			6.51E-10								
49	RUTHENIUM 106 + D			2.31E-06								
50	SELENIUM	M				2.44E-03		3.44E-03				6.88E-06
51	SILICON	Bkg,M										
52	SILVER	Bkg,I										
53	SODIUM	M,EPA-10										
54	STRONTIUM	M				1.23E-04						1.55E-06
55	STRONTIUM 90		5.63E-06	1.61E-05								
56	SULFATE	M, SW-Bkg				2.52E-02						7.50E-03
57	SULFIDE											3.75E-05
58	TECHNETIUM 99			7.79E-09								
59	TETRACHLOROETHYLENE	M		8.64E-07						2.17E-04		
60	THALLIUM									1.00E-02		
61	THORIUM 228			1.67E-06								
62	THORIUM 232			6.04E-09								

Table C.1. (contd)

	Name of Analyte	Notes	Carcinogenic Risk Ranking		Hazard Index Ranking		WQC Screen Ranking		LC50/100 Screen Ranking		TLM Screen Ranking	
			Surface	Ground-	Surface	Ground-	Surface	Ground-	Surface	Ground-	Surface	Ground-
			Water	water	Water	water	Water	water	Water	water	Water	water
63	TITANIUM	Bkg,M										
64	TOLUENE	SW sample suspect			5.61E-03	3.46E-06			7.83E-03	4.83E-06		
65	TRICHLOROETHYLENE	M		2.28E-08						4.38E-05		
66	TRITIUM (HYDROGEN 3)		2.86E-07	1.23E-07								
67	URANIUM 233			3.36E-10								
68	URANIUM 234		1.81E-06	1.21E-08								
69	URANIUM 235		2.41E-07	4.10E-07								
70	URANIUM 238		1.89E-06	9.26E-09								
71	VANADIUM	Bkg,M										7.27E-07
72	XYLENE	SW sample suspect			1.26E-03						1.00E-01	
73	ZINC	M, SW-LD			3.60E-01	2.88E-01	1.00E-01	8.00E-02	2.56E+00	2.05E+00		

C4

Table C.2. Results for Soil and Sediment

Name of Analyte	Notes	Carcinogenic Risk Ranking		Hazard Index Ranking		WOC Screen Ranking		LC50/100 Screen Ranking		TLM Screen Ranking	
		Soil	Sediment	Soil	Sediment	Soil	Sediment	Soil	Sediment	Soil	Sediment
1 AMERICIUM 241		2.30E-06									
2 ANTIMONY 124	T 1/2 = 60d										
3 CARBON 14		1.41E-07									
4 CESIUM 134		2.11E-07	1.53E-06								
5 CESIUM 137		5.96E-03	1.23E-05								
6 COBALT 60		1.55E-01	4.22E-05								
7 EUROPIUM 152		2.12E-01	8.68E-06								
8 EUROPIUM 154		8.20E-02	9.84E-07								
9 EUROPIUM 155		3.66E-04	1.89E-08								
10 NEPTUNIUM 237			3.95E-08								
11 NICKEL 63		5.33E-07									
12 PLUTONIUM 238		6.32E-07	6.61E-11								
13 PLUTONIUM 239		1.38E-05	4.26E-09								
14 PLUTONIUM 240											
15 POTASSIUM 40	Bkg										
16 RADIUM 226		6.71E-08	3.69E-08								
17 STRONTIUM 90		1.91E-06	4.16E-07								
18 TECHNETIUM 99		2.30E-11	1.72E-11								
19 THORIUM 228		2.85E-09	5.31E-09								
20 THORIUM 232		1.49E-09	4.34E-09								
21 THORIUM 234			3.20E-09								
22 TRITIUM (HYDROGEN 3)		2.60E-07									
23 URANIUM 233		3.97E-09	2.34E-09								
24 URANIUM 234			3.92E-09								
25 URANIUM 235		2.96E-07	2.41E-08								
26 URANIUM 238		4.68E-09	3.19E-09								
27 ZINC 65	Suspect		4.85E-07								
28 ZIRCONIUM 95		1.40E-06									
Chemicals											
29 ACENAPHTHENE	M		4.26E-05						5.25E-05		
30 ALUMINUM	Bkg, M, EPA-10										
31 AMMONIA	M										6.67E-03

Table C.2. (contd)

	Name of Analyte	Notes	Carcinogenic Risk Ranking		Hazard Index Ranking		WQC Screen Ranking		LC50/100 Screen Ranking		TLM Screen Ranking	
			Soil	Sediment	Soil	Sediment	Soil	Sediment	Soil	Sediment	Soil	Sediment
32	ANTHRACENE	M			1.69E-04				1.08E-04			
33	AROCOR 1248 (PCB)	M	2.99E-02				7.07E+00				3.56E-04	
34	ARSENIC	I	3.57E-04	5.70E-05	6.80E-01	1.09E-01	2.47E-03	3.95E-04		6.82E-03		
35	BARIUM	SD-Bkg,I			7.93E-02				1.68E-03			
36	BENZENE	M	1.07E-07								2.25E-03	
37	BENZO(G,H,I)PERYLENE	Non-Haz?,M							1.03E-04			
38	BENZO(a)ANTHRACENE	M	3.71E-04						2.35E-04			
39	BENZO(a)PYRENE	M,Suspect	3.67E-03						2.03E-04			
40	BENZO(b)FLUORANTHENE	M	5.65E-04						2.23E-04			
41	BENZO(k)FLUORANTHENE	M	2.26E-04						1.90E-04			
42	BENZOIC ACID	M			2.82E-07				9.44E-06			
43	BERYLLIUM	I	4.03E-05	5.54E-06	1.88E-03	2.58E-04			4.00E-02	5.50E-03		
44	BIS(2-ETHYLHEXYL) PHTHALATE	I	3.02E-06		1.08E-02							
45	CADMIUM	I	9.36E-05	1.40E-04	2.97E-02	4.46E-02	1.64E-02	2.45E-02			6.00E-07	9.00E-07
46	CALCIUM	Bkg,M,EPA-10										
47	CHLORDANE	I	7.62E-05		9.77E-01		1.05E+01		5.49E-01			
48	CHLORIDE	Bkg,M										
49	CHLORINE (a)	Bkg,I										
50	CHROMIUM	I	8.77E-02	4.13E-02	2.14E-03	1.01E-03	2.35E-01	1.11E-01		1.22E-01		
51	CHRYSENE	M	1.84E-05									
52	COBALT	M			1.00E-02	3.39E-03					3.41E-08	1.15E-08
53	COPPER	M			1.11E+03	3.18E-01	1.17E+02	3.33E-02		8.00E-02		
54	CYANIDE	M			2.29E-05		2.02E-03					
55	DIBENZOFURAN	Inadequate?,M										
56	DIESEL FUEL	M			9.47E-02				2.80E+00			
57	ENDRIN ALDEHYDE	M			6.42E-04				1.65E-02			
58	ETHYL BENZENE	M			1.39E-03						1.07E-02	
59	FLUORANTHENE	I			5.30E-03				4.50E-04			
60	FLUORENE	I			1.35E-04				4.75E-05			
61	FLUORIDE	M			9.63E-05						2.04E-05	
62	FLUORINE (a)	Bkg,I										
63	INDENO(1,2,3-CD)PYRENE	M, Suspect	1.09E-03						1.30E-04			
64	IRON	M,EPA-10										

Table C.2. (contd)

	Name of Analyte	Notes	Carcinogenic Risk Ranking		Hazard Index Ranking		WQC Screen Ranking		LC50/100 Screen Ranking		TLM Screen Ranking	
			Soil	Sediment	Soil	Sediment	Soil	Sediment	Soil	Sediment	Soil	Sediment
65	KEROSENE	M			5.36E-02						1.54E-01	
66	LEAD	M			1.67E+00	2.26E-01	1.69E+00	2.28E-01			1.02E-02	1.38E-03
67	LITHIUM (a)	Bkg,M										
68	MAGNESIUM	Bkg,M,EPA-10										
69	MANGANESE	Bkg,M										
70	MERCURY	M			5.67E-01		3.58E+00		4.30E-01			
71	METHYL-2-PENTANONE, 4-	Non-haz?,M										
72	METHYLENE CHLORIDE	I	4.74E-10		1.05E-06				2.18E-07			
73	METHYLNAPHTHALENE, 2-	Unclass?,M							1.05E-05			
74	NICKEL	M			4.80E-02	4.28E-03	1.38E-02	1.23E-03			5.82E-03	5.18E-04
75	NITRATE	M			1.12E-01				1.52E-03			
76	PHENANTHRENE	M			1.48E-03				3.75E-04			
77	POTASSIUM	Bkg,M,EPA-10										
78	PYRENE	M			4.40E-03				3.00E-04			
79	SELENIUM	M			5.95E-03		8.40E-03		1.68E-03			
80	SILVER	Bkg,M										
81	SILVER CHLORIDE	M			1.79E+00							
82	SODIUM	Bkg,M,EPA-10										
83	STRONTIUM	M			2.66E-04						3.35E-06	
84	STRONTIUM CHLORIDE	M			3.98E-09						5.00E-11	
85	SULFATE (SULFUR)	M			5.50E-05						1.64E-05	
86	TITANIUM (a)	Bkg,M										
87	TOLUENE	M			2.12E-03				5.83E-03			
88	TOTAL PETROLEUM HYDROCARBONS											
89	VANADIUM	Bkg,M										
90	XYLENE	M			5.67E-02						4.50E-03	
91	ZINC	M			1.01E-01	1.30E-01	2.81E-02	3.61E-02	7.19E-01	9.23E-01		
92	ZIRCONIUM (a)	Bkg,M										
(a)	Concentrations of these chemicals fall within their respectively occurring background levels.											

C.7

Table C.3. Results for Groundwater Away from the Columbia River

		Carcinogenic	Hazard	WQC	LC50/100	TLM
		Risk	Index	Screen	Screen	Screen
Name of Analyte	Notes	Ranking	Ranking	Ranking	Ranking	Ranking
100 Areas						
Chromium (+6)	I	5.31E-02	1.30E-03	1.43E-01	1.57E-01	
Nitrate	M		4.77E+01		6.50E-01	
Strontium-90		3.62E-07				
Tritium (Hydrogen-3)		5.16E-09				
200 West Area						
Arsenic	I	1.82E-05	3.47E-02	1.26E-04	2.18E-03	
Carbon Tetrachloride	M	5.37E-04	5.90E+00		5.25E-03	
Chloroform	I	4.16E-06	6.93E-02		1.60E-03	
Chromium	I	1.09E-02	2.67E-04	2.94E-02	3.23E-02	
Fluoride	M		1.38E-02			4.38E-03
Iodine-129		2.71E-08				
Nitrate	M		4.85E+02		6.61E+00	
Technetium-99		9.13E-08				
Trichloroethylene	M	3.02E-08			5.82E-05	
Tritium (Hydrogen-3)		4.00E-07				
Uranium		1.61E-07				
200 East Area						
Arsenic	I	1.82E-05	3.47E-02	1.26E-04	2.18E-03	
Cesium-137		2.73E-04				
Chloroform	I	1.82E-08	3.04E-04		7.00E-06	
Chromium	I	9.75E-03	2.38E-04	2.62E-02	2.88E-02	
Cobalt-60		3.79E-04				
Cyanide	M		1.95E-03	1.72E-01		
Iodine-129		1.81E-08				
Nitrate	M		1.46E+02		1.99E+00	
Plutonium-239/240						
Strontium-90		1.04E-06				
Technetium-99		7.61E-08				
Tritium (Hydrogen-3)		2.66E-07				
Uranium		2.69E-09				
600 Area (Solid Waste Landfill Site)						
Chloroform	I	1.30E-09	2.17E-05		5.00E-07	
Dichloroethane, 1, 1-	M		4.92E-06		3.18E-06	
Tetrachloroethene	M	2.66E-07	5.21E-04		9.23E-05	
Trichloroethane, 1, 1, 1-	M	5.60E-07	2.44E-03		1.00E-04	
Trichloroethene	M	1.90E-08			1.27E-05	

Distribution

<u>No. of Copies</u>		<u>No. of Copies</u>
	Offsite	F. R. Cook 2552 Harris Richland, WA 99352
12	DOE Office of Scientific and Technical Information	J. P. Corley 2213 Torbett Richland, WA 99352
	M. Bauer Confederated Tribes and Bands of the Yakama Indian Nations P.O. Box 151 Toppenish, WA 98948	2 DNR Hanford Projects Attn: J. R. Wilkinson Confederated Tribes of the Umatilla Indian Nation P.O. Box 638 Pendleton, OR 97801
	S. Benz 600 S. Kent #71 Kennewick, WA 99336	G. deBruler Columbia River United P.O. Box 667 Bingen, WA 98605
	L. Block U.S. Fish and Wildlife Service Moses Lake Field Office P.O. Box 1157 Moses Lake, WA 98837	C. Denniston Greenpeace 11815 - 20th SW Seattle, WA 98146
	D. Bradshaw Audobon Society of Portland 5151 Northwest Cornell Road Portland, OR 97210	B. Drost USGS Water Resources Division Washington District 1201 Pacific Avenue Suite 600 Tacoma, WA 98402
	N. Buske Search Technical Services Star Route Box 17 Davenport, WA 99112	D. Dunning Oregon Dept. of Energy 625 Marion Street NE Salem, OR 97310
	D. Cellarius Sierra Club Northwest Office 1516 Melrose Avenue Seattle, WA 98122	3 J. Erickson Division of Radiation Protection Washington State Dept. of Health P.O. Box 47827 Olympia, WA 98504-7827
	C. Cline Washington State Department of Ecology P.O. Box 47600 Olympia, WA 98504-7600	

No. of
Copies

No. of
Copies

M. Fox, President
Hanford Family
P.O. Box 1462
Richland, WA 99352

2 D. Powaukee
Nez Perce Tribe
P.O. Box 365
Lapwai, ID 83540

R. Gardiner
614 Sherman Drive
The Dalles, OR 97058

L. Stembridge
HEAL
1408 West Broadway Avenue
Spokane, WA 99201-1902

N. J. Germond
League of Women Voters
224 Iron Mountain Boulevard
Lake Oswego, OR 97034

C. Stephan
1520 Torthay Court
Richland, WA 99352

D. Maez
4214 W. John Day Place
Kennewick, WA 99336

3 D. Stewart-Smith
Oregon Department of Energy
625 Marion Street NE
Salem, OR 97310

C. Mebane
National Oceanic and Atmospheric
Administration
c/o EPA Seattle Office
1200 6th Avenue, HW-113
Seattle, WA 98101

J. P. Thomas
HHIN Resource Center
1719 Smith Tower
506 Second Avenue
Seattle, WA 98104

J. Monteith
Oregon Natural Resources Council
1161 Lincoln Street
Eugene, OR 97401

Onsite

11 DOE Richland Operations Office

R. F. Brich (5) H4-83
P. W. Willison A4-52
DOE Public Reading Room (5) H2-53

D. Nichols
WNP-1
Building 52
Richland, WA 99352

6 Washington State Department of Ecology

S. M. Alexander B5-18
D. P. Holland (4) B5-18
J. W. Yokel B5-18

R. Patt
Water Resources Department
158 - 12th Street N.E.
Salem, OR 97310-0210

4 Westinghouse Hanford Company

Administrative Record (3) H6-08
A. S. Carlson B3-35

G. Pollet
Heart of America Northwest
1305 4th Avenue, 208
Seattle, WA 98101

<u>No. of Copies</u>		<u>No. of Copies</u>	
6	U. S. Environmental Protection Agency	L. R. Huesties	K6-61
		A. D. Maughan	K6-52
	L. E. Gadbois (5)	T. B. Miley	K5-12
	D. R. Sherwood	L. F. Morasch	K6-62
		B. A. Napier (5)	K3-54
34	Pacific Northwest Laboratory	D. B. Shipler	K9-13
		S. F. Snyder	K3-54
	S. D. Cannon	W. L. Templeton	K9-13
	R. L. Dirkes	W. H. Walters	K9-33
	P. W. Eslinger	L. K. Wenrich (5)	K3-70
	S. L. Friant	B. K. Wise	K9-04
	M. S. Hanson	Records Center (2)	K3-70
	G. L. Harvey	Publishing Coordination	K1-06
	P. C. Hays	Technical Library (5)	P8-55