

LA-UR-97- 3049

Approved for public release;
distribution is unlimited.

TITLE: LOCALIZED EXCITATIONS IN UPdSn

CONF-970814--

RECEIVED

NOV 12 1997

OSTI

AUTHOR(S): H. Nakotte, LANSCE 12
R.A. Robinson, LANSCE 12
M. Bull, Dept. of Physics, Birkbeck College, Univ. of London, Van der
Waals-Zeeman Inst. Univ. of Amsterdam
K.A. McEwen, Dept. of Physics, Birkbeck College, Univ. of London
T. Swan, LANSCE 12
T. Kelley, LANSCE 12
R.A. Ecclestone, ISIS Division, Rutherford Appleton Laboratory
E. Bruck, Van der Waals-Zeeman Institute, Rutherford Appleton
Laboratory

SUBMITTED TO: ICNS '97, Toronto, Aug 18-22
Physica B

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED *ph*

Los Alamos
NATIONAL LABORATORY

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. The Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Localized excitations in UPdSn

H. Nakotte^(a), R.A. Robinson^(a), M. Bull^(b,d), K.A. McEwen^(b), T. Swan^(a), T. Kelley^(a)
R.A. Ecclestone^(c) and E. Brück^(d)

(a) *Manuel Lujan Jr. Neutron Scattering Center, Los Alamos National Laboratory, Los Alamos NM 87545, USA*

(b) *Department of Physics, Birkbeck College, University of London, Malet Street, London WC1E 7HX, UK*

(c) *ISIS Division, Rutherford Appleton Laboratory, Chilton OX11 0QX, UK*

(d) *Van der Waals-Zeeman Institute, University of Amsterdam, Valckenierstraat 65, 1018 XE Amsterdam, The Netherlands*

We have measured the inelastic neutron-scattering response of UPdSn at various temperatures using the HET and PHAROS spectrometers at the ISIS and MLNSC facilities, respectively. UPdSn shows some quasielastic scattering, which may be attributed to the hybridization of the 5f electrons with the conduction electrons. Furthermore, we find a clear excitation around 40meV above 40K in addition to the phonon contribution. While this excitation may be indicative of crystal fields in UPdSn, its strong temperature dependence seems to contradict a simple crystal-field picture. Below T_N , the unusual temperature dependence may be attributed to magnetically-driven distortions (and subsequent changes in the local surrounding of the U ions), but there is some evidence that other additional mechanism(s) may contribute above T_N . Some possible mechanisms will be discussed.

KEYWORDS: inelastic neutron scattering, crystal fields, UPdSn

Among the uranium intermetallics, UPdSn has attracted recent attention as it is believed to be one of the rare examples where the 5f electrons are believed to be fairly localized.¹ Therefore, UPdSn is a prime candidate for crystal-field effects, and its observation in uranium intermetallics is extremely rare. UPdSn has two antiferromagnetic transition at about 40 and 25K.²

Here, we report on inelastic-neutron-scattering data on UPdSn using the HET and PHAROS spectrometers at the ISIS and MLNSC facilities, respectively. The experiments were performed on 25g of polycrystalline UPdSn, and data were at various temperatures using an incident energy of 100meV.

The HET data are displayed in Fig. 1, where the scattering response of UPdSn at 20 and 100K is shown. The symbols represent the response as seen in HET's low-angle detector bank (sensitive to scattering with small momentum transfer, Q), while the solid line represents the response seen in the high-angle detectors (large Q). Scaled appropriately the high-angle response provides an estimate of the scattering contribution due to lattice vibrations (phonons), while any additional scattering seen in the low-angle detectors must be magnetic in origin. Clearly, there are additional 'magnetic' features. First, at 100K, there is some quasielastic scattering (below 10meV), which is barely present at 20K. Secondly, there is a pronounced excitation around 40-50meV.

The quasielastic term seen at 100K may originate in the hybridization of 5f electrons with the conduction electrons.³ In the magnetically-ordered state (at 20 K), on the other hand, some kind of a 'hybridization gap' may have formed. Recent band-structure calculations on UPdSn by Sandratskii and Kübler⁴ find indeed a local minimum in the density of states at E_F . Furthermore, as the 5f electrons carry a magnetic moment, one would expect subsequent gapping in the magnon-excitation spectrum, and bulk results provide evidence for its presence.⁵

The excitation around 40meV (at 100K) may signal crystal fields in UPdSn. The strong energy shift ($\sim 10\text{meV}$) on going to 20K could be a consequence of magnetically-driven distortions at the magnetic transitions,⁶ because of subsequent changes in the local symmetry of the U ions. For better understanding of the nature of this excitation, we performed additional neutron-scattering studies at various temperatures using PHAROS (Fig. 2). Although the statistics is relatively poor (because of limited beam time), the excitation is clearly visible at all temperatures. After subtraction of a phonon background (estimated from HET data), we fitted the excitation to a single Gaussian (solid lines in Fig. 2). There is only little temperature dependence in the location of the excitation above T_N , which is followed by a strong shift below T_N (where also the magnetically-driven distortion occur). While this can be expected within crystal-field picture, we also find an unusual temperature dependence of the integrated intensities in the paramagnetic range. Only for temperatures above 100K, we find a meaningful value (~ 3) for the ratio of the degeneracies of the first excited and the ground state. Any extension toward lower temperatures yield unphysical large values. On the basis of the present results, however, it is not unambiguous whether the observed feature consists out of a single or possibly more excitations. Nevertheless, the above observation may indicate that there is some other physics which contributes to this excitation.

It may be tempting to ascribe the excitation to an anisotropy gap in a similar manner as done recently for URhAl.⁷ However, the fact that the excitation almost vanishes at 300K contradicts bulk magnetic results, where huge magnetic anisotropy even at room temperature is found.¹

Another alternative explanation is that there is some interaction in terms of spin-waves or magnon-phonon hybridization.⁸ However, one expects such correlations to be largely absent above T_N .

Therefore, it seems likely that the observed excitation is of crystal-field origin, but some other physics interfere with the crystal fields. Some insight may be gained by

comparing the inelastic-neutron-scattering data with the bulk observations. Such analysis is presently underway.

In conclusion, we believe that the 5f electrons in UPdSn are somewhat itinerant (yielding a quasielastic response), but that the hybridization of 5f electrons with the conduction electrons is too weak to completely smear out localized crystal-field excitations.

figure captions:

Fig. 1: Inelastic-scattering response of UPdSn using 100meV incident energy at 20 and 100K, taken on the HET spectrometer. The solid line represents the 20K phonon estimate obtained from the high-angle detector bank (see text).

Fig. 2: Inelastic-scattering response of UPdSn using 100meV incident energy at various temperatures, as taken on the PHAROS spectrometer. The solid lines represent the Gaussian fits to the excitation after subtraction of an estimated phonon background.

references

- [1] F.R. de Boer, E. Brück, H. Nakotte, A.V. Andreev, V. Sechovsky, L. Havela, P. Nozar, C.J.M. Denissen, K.H.J. Buschow, B. Vaziri, M. Meissner, H. Maletta and P. Rogl, *Physica B* 176 (1992) 275.
- [2] R.A. Robinson, A.C. Lawson, K.H.J. Buschow, F.R. de Boer, V. Sechovsky and R.B. Von Dreele, *J. Magn. Magn. Mater.* 98 (1991) 147.
- [3] E. Holland-Moritz and G.H. Lander, in *Handbook of Rare Earths*, Vol 19, eds: K.A. Gschneidner, L. Eyring, G.H. Lander and G.R. Choppin (Elsevier, Amsterdam, 1994) p. 1.
- [4] L.M. Sandratskii and J. Kübler, preprint (1997).
- [5] L. Havela, V. Sechovsky, H. Nakotte, Brück and F.R. de Boer, *IEEE Transact. on magnetics* 30 (1994) 1130.
- [6] R.A. Robinson, A.C. Lawson, J.A. Goldstone and K.H.J. Buschow, *J. Magn. Magn. Mater.* 1226 (1993) 143.
- [7] A. Hiess, L. Havela, K. Prokes, R.S. Eccleston and G.H. Lander, *Physica B* 230-232 (1997) 89.
- [8] S. W. Lovesey, *Theory of neutron scattering from condensed matter*, Vol. 2, (Clarendon Press, Oxford, 1984) p. 97.



