

FIG. 9. Mean numbers of arthropods caught with sweep net from cowpea plots (1992)




FIG. 10. Mean numbers of arthropods caught with pitfall traps from cowpea plots (1993)

TABLE VI. MEAN NUMBERS OF ARTHROPODS (\pm SE) CAUGHT WITH SWEEP NET FROM COWPEA PLOTS, 1993

DAIT	0		1		10		15		22		26[4]		42[20]	
	C	En	C	En	C	En	C	En	C	En	C	En	C	En
Ants	0.3	0.5	1.0	0.3	1.8	0.0	0.5	1.0	1.0	0.3	0.3	0.8	1.8	0.5
	\pm	\pm	\pm	\pm	\pm		\pm	\pm	\pm	\pm	\pm	\pm	\pm	\pm
	0.5	0.6	1.2	0.5	3.5		0.6	1.4	1.1	0.5	0.5	0.9	0.9	0.6
Spiders	0.8	0.0	0.5	1.0	0.5	0.3	1.0	1.0	0.8 ^a	0.0 ^b	0.8	1.0	0.3	0.0
	\pm		\pm	\pm	\pm	\pm	\pm	\pm	\pm		\pm	\pm	\pm	
	0.9		0.5	0.8	0.6	0.5	0.8	1.4	0.9		0.9	1.1	0.5	
Coccinellids	0.0	0.0	0.3	0.0	0.0	0.0	0.8	0.3	0.5	0.0	0.3	0.3	0.0	0.0
			\pm				\pm	\pm	\pm		\pm	\pm		
			0.5				1.5	0.5	1.0		0.5	0.5		

DAIT; days after initial treatment; [] days after endosulfan treatment

Means within rows for the same day followed by different letters are statistically different ($p < 0.05$)

C = Control

En = cypermethrin + endosulfan

3.9.4. Arthropods caught with sweep net from cowpea plots, 1993

The endosulfan treatment seemed to have caused reductions in ants, and spiders caught 20 DAET but the differences were not statistically significant ($P > 0.05$) (Table VI).

3.9.5. Arthropods caught with pitfall traps from cowpea plots, 1994

Arthropods collected with pitfall traps from cowpea plots included ants, spiders and Collembola. All were significantly reduced by the endosulfan treatment 14 DAET ($P < 0.05$) (Table VII).

3.9.6. Arthropods caught with D-VAC machine from cowpea plants, 1994

Endosulfan treatment significantly reduced ants caught from cowpea plants 1 DAET ($P > 0.05$) (Table VIII) but it appeared recovery commenced 14 DAET. Spiders were, however, not significantly reduced.

3.9.7. Arthropods caught with D-VAC machine from the ground in cowpea plots, 1994

From Table IX, it appears endosulfan had caused a non-significant reduction in ant catches ($P > 0.05$). Prior to the application of endosulfan, comparatively higher numbers of ants had been collected from the plot. Spider catches 14 DAET were relatively lower in the endosulfan-treated plots than those from the other plots but there was no significant reduction ($P > 0.05$).

3.10. General discussion

3.10.1. Fluctuations in arthropod populations

In addition to arthropod natural enemies and other predators such as birds, lizards and frogs and microorganisms in regulating insect populations in ecosystems, the scarcity of resource and space can also cause population changes. Abiotic factors such as temperatures, light, rainfall, soil and other environmental conditions could also influence the extent of arthropod activities on day to day basis and hence the numbers trapped. These factors could partly account for the fluctuations of arthropod numbers recorded in both and unsprayed plots.

3.10.2. Apparent recoveries

Reports by other researchers [7, 8, 9] had shown that dissipation rates of the organochlorine pesticides were relatively faster under tropical conditions. The observed recoveries in some of the arthropods after the application of lindane and endosulfan could very likely be attributed to the degradation and or dissipation of the pesticides from the maize and cowpea agro-ecosystems. This is supported by the residue data reported by Yeboah et al. (this TecDoc).

TABLE VII. MEAN NUMBERS OF ARTHROPODS (\pm SE) CAUGHT WITH PITFALL TRAPS FROM COWPEA PLOTS (1994)

DAIT	0			11			21 [1]			35 [14]		
	C	Cy	En	C	Cy	En	C	Cy	En	C	Cy	En
Ants	15.5 ^a ± 2.5	17.6 ^b ± 3.7	9.4 ^b ± 1.0	7.1 ± 1.0	9.0 \pm 1.2	6.3 \pm 0.8	7.2 \pm 0.7	8.6 ± 1.2	6.5 ± 0.6	10.1 ^a ± 1.2	18.7 ^b ± 2.8	6.7 ^c ± 0.7
Spiders	0.6 \pm 0.2	0.6 \pm 0.6	0.7 \pm 0.2	0.5 \pm 0.1	0.5 \pm 0.1	0.6 \pm 0.1	0.6 \pm 0.1	0.8 \pm 0.3	0.6 \pm 0.1	1.0 ^a ± 0.1	0.8 ^{ab} ± 0.1	0.6 ^b ± 0.1
Collembola	6.5 ^a ± 1.7	15.8 ^b ± 2.2	7.9 ^a ± 1.0	5.0 \pm 0.5	6.0 \pm 0.5	5.5 \pm 0.5	8.0 \pm 0.7	8.2 \pm 0.7	8.3 ± 0.6	7.7 ^a ± 0.9	10.8 ^a ± 1.6	3.5 ^b ± 0.4

DAIT; days after initial treatment; [] days after endosulfan treatment

Means within rows for the same day followed by different letters are statistically different ($p < 0.05$)

C = control

Cy = cypermethrin

En = cypermethrin + endosulfan

TABLE VIII. MEAN NUMBERS OF ARTHROPODS (\pm SE) CAUGHT WITH D-VAC MACHINE FROM COWPEA PLANTS (1994)

DAIT	0			1			11			21[1]			35[14]		
	C	Cy	En	C	Cy	En	C	Cy	En	C	Cy	En	C	Cy	En
Ants	4.3 \pm	3.7 \pm	4.8 \pm	3.2 \pm	2.1 \pm	2.0 \pm	1.1 \pm	0.8 \pm	0.6 \pm	1.7 ^a	0.63 ^b	0.06 ^b	0.75	1.1 \pm	0.75
	1.3	1.1	1.7	0.4	0.7	0.6	0.5	0.2	0.3	\pm 0.3	\pm 0.4	\pm 0.1	\pm 0.2	0.5	\pm 0.3
Spiders	0.7 \pm	0.5	0.8 \pm	0.4 \pm	0.5 \pm	0.8 \pm	0.9 ^a	0.1 ^b	0.3 ^{ab}	0.5 \pm	0.7 \pm	0.3 \pm	1.1 ^{ab}	1.8 ^a \pm	0.6 ^b \pm
	2.1	\pm 0.2	0.2	0.1	0.1	0.4	\pm 0.5	\pm 0.1	\pm 0.1	0.3	0.2	0.1	\pm 0.4	0.1	0.2

DAIT; days after initial treatment; [] days after endosulfan treatment

Means within rows for the same day followed by different letters are statistically different ($p < 0.05$)

C = control

Cy = cypermethrin

En = cypermethrin + endosulfan

TABLE IX. MEAN NUMBERS OF ARTHROPODS (\pm SE) CAUGHT WITH D-VAC MACHINE FROM THE GROUND IN COWPEA PLOTS (1994)

DAIT	0			1			11			21[1]			35 [14]		
	C	Cy	En	C	Cy	En	C	Cy	En	C	Cy	En	C	Cy	En
Ants	1.4 ^a \pm	2.1 ^b \pm	2.3 ^b \pm	1.4 \pm	0.7 \pm	0.8 \pm	0.7 ^a	1.6 ^b \pm	1.3 ^b \pm	1.1 \pm	1.6	2.6 \pm	0.7	1.3 \pm	0.6 \pm
	0.4	0.4	0.6	0.4	0.2	0.2	\pm 0.2	0.3	0.3	0.4	\pm 0.5	0.6	\pm 0.2	0.3	0.2
Spiders	0.5 \pm	0.7 \pm	0.8 \pm	0.3 \pm	0.2 \pm	0.1 \pm	0.1 \pm	0.2 \pm	0.2 \pm	0.0	0.2 \pm	0.1 \pm	0.6 \pm	0.9 \pm	0.3 \pm
	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1		0.1	0.1	0.3	0.4	0.1

DAIT; days after initial treatment; [] days after endosulfan treatment

Means within rows for the same day followed by different letters are statistically different ($p < 0.05$)

C = control

Cy = cypermethrin

En = cypermethrin + endosulfan

4. CONCLUSIONS

Collembola seemed to be the most susceptible insect to both lindane and endosulfan ($P < 0.05$). Ants and spiders were also significantly reduced by these organochlorine pesticides but most often recovered from the adverse effects between 29 and 57 days in maize plots and after 21 days in cowpea plots. Lindane treatments did not consistently increase maize yields or reduce stemborer attacks and three consecutive endosulfan treatments did not significantly increase cowpea yields ($P > 0.05$). Endosulfan application preceded by two successive cypermethrin treatments significantly increased cowpea yields and reduced seed damage ($P < 0.05$). Neither lindane nor endosulfan seemed to have significantly affected soil microbial activity ($P > 0.05$) based on the rates of decomposition of organic matter buried in maize and cowpea plots and retrieved between 84 and 259 days later.

ACKNOWLEDGEMENTS

The author wishes to thank the IAEA, the FAO and the Swedish International Development Agency (SIDA) for financially sponsoring this project. Thanks also go to my employers, The Ghana Atomic Energy Commission, for permitting me to undertake the project and for the material and financial contributions. Special gratitude goes to my colleague Dr. P.O. Yeboah for partly supporting this project in the initial stages with part of his grant and for his encouragement. I wish also to thank Mr. Stephen Yeboah for helping me with the data processing. The contributions of Messrs. C.F.S. Edmundson, John Apatey, Titus Vuore, Anyetei Akrama, my daughter Efua, and all the other numerous colleagues and farm hands who helped me to execute this project are gratefully acknowledged.

REFERENCES

- [1] EDWARDS, C.A., Persistent pesticides in the Environment (2nd Ed.), CRC Press, Ohio (1973a) 170pp.
- [2] JAMAL, A., Pesticides, Hazards and Alternatives, African Development Foundation United States of America Draft Pesticides Factsheet (1991).
- [3] GRAINS AND LEGUMES DEVELOPMENT BOARD - MAIZE, Ministry of Agriculture Technical Bulletin No. GDB/M1.
- [4] PHILIPS, T.A., An Agricultural Note Book, Green and Co. Ltd, Longmans, Accra (1964) 480 pp.
- [5] TINDALL, H.D., Commercial Vegetable Growing. Oxford University Press. (1968) 300 pp.
- [6] GHANA/CIDA GRAINS DEVELOPMENT PROJECT, Maize and Cowpea Production Guide (1991).
- [7] WANDIGA, S.O., MGHENYI, J.M., "Persistence of gamma- 1,2,3,4,5,6-hexachlorocyclohexane (γ -HCH) in tropical soils of Kenya", Isotope Techniques for Studying the Fate of Persistent Chemicals in the Tropics, IAEA TECDOC-476, Vienna (1988).
- [8] SAMUEL, T., PILLAI, M.K.K., Persistence and fate of ^{14}C -gamma-HCH in an Indian sandy loam soil under field and laboratory conditions, Ibid. (1988) 93-102.
- [9] BOLAÑOS DE MORENO, M., ESPINOSA, L., PASTOR, Y., MOLINEROS, J., MERINO, R., "Persistence of lindane in the Ecuadorian environment", Ibid. (1988) 85-91.

- [10] SINGH, S.R., TAYLOR, T.A., Pests of grain legumes and their control in Nigeria: (Singh, S.R., Van Emden, H.F., Taylor, T.A., Eds), Pests of Grain Legumes: Ecology and Control. Academic Press. London.
- [12] PERFECT, T.J., COOK, A.G., CRITCHLEY, B.R., CRITCHLEY, U., DAVIES, A.L., SWIFT, M.J., RUSSELL-SMITH, YEADON, R., The effect of DDT contamination on the productivity of a cultivated forest soil in the subhumid tropics. J.Appl. Ecol. 16 (1979) 705-719.

**NEXT PAGE(S)
left BLANK**

FATE AND DISTRIBUTION OF LINDANE AND ENDOSULFAN IN MAIZE AND COWPEA ECOSYSTEMS RESPECTIVELY



XA9743719

P.O. YEBOAH
Department of Chemistry,
Ghana Atomic Energy Commission

K.G. MONTFORD
Biotechnology and Nuclear Agriculture Research Institute,
Ghana Atomic Energy Commission

R.K.KLAKE, C.K AKPABLI
Department of Chemistry,
University of Ghana

Legon, Ghana

Abstract

A quantitative study is presented on lindane and endosulfan residues in maize and cowpea ecosystems respectively. Both pesticides were found to dissipate very fast under the tropical Ghanaian conditions. The high rate of dissipation in leaves is attributed to the fact that the leaves were exposed to sunshine and wind leading to increased volatilisation. Endosulfan was found to dissipate faster from the cowpea ecosystem than lindane did in the maize ecosystem. The mean residue levels of lindane in maize grains were $0.02 \mu\text{g g}^{-1}$; whilst residue levels of endosulfan in cowpea seeds were $0.05 \mu\text{g g}^{-1}$. These levels are lower than the maximum residue limits recognised as acceptable by the Codex Alimentarius Commission.

1. INTRODUCTION

As a result of increasing world population, there has been a proportionate increase in the demand for food. Currently, 58 percent of the world's population is undernourished [1]. There is therefore the need to increase food production. This can be achieved through the application of such products of modern technology, as high yielding crop varieties, fertilisers, irrigation and pesticides. Pesticides help boost food production by reducing pest populations to tolerable levels. In 1986, it was found that crop loss due to pests was 40%. Cotton yields have also been found to increase three fold through effective use of chemical pesticides for pest control in the Sudan [2].

Despite the benefits derived from pesticides, some of them tend to persist in the environment, and may thus pose a hazard to living things. Organochlorine pesticides constitute one group of such pesticides that are persistent. Examples of these include DDT, lindane and endosulfan. These compounds have also been found to (i) be highly fat soluble and hence accumulate in fatty tissues of animals, (ii) accumulate in food chains, and (iii) degrade into other toxic residues which can have long term adverse effects on the environment [3,4].

In Ghana, the surviving organochlorines in use are lindane, endosulfan, aldrin and dieldrin. Lindane, formulated under the trade name Gammalin 20, has been restricted to be used only in the cocoa growing industry. However, due to mixed cropping, other crops end up by being sprayed with lindane during its application. These crops include maize, cocoyam and cassava. Under such conditions, lindane can serve to control insects that bore into stems of the maize plant.

Notwithstanding the hazards posed by organochlorine pesticides to the environment and being banned or restricted in temperate regions, developing countries including Ghana, continue to use them to boost food production. They are also used in public health to control disease vectors. Their continued use has been attributed to their high efficiency, low cost, and easy availability. Furthermore, owing to the intense heat and high humidity in the tropics, organochlorine pesticides dissipate at faster rates than in temperate regions. Consequently, it is suggested that, there should be less danger of possible accumulation of their residues in soils, plants and animal matter, that could lead to the precipitation of environmental hazards.

Whereas considerable data on pesticide residues are available in the developed countries to assist in decision making regarding the continued use of particular pesticides, knowledge on pesticide residues and their effects are rarely available in many developing countries. Hence the need to generate data on the effects of these pesticides in various agroecosystems in the tropics to allow correct decisions on their use. These should include extensive knowledge about their fate, persistence, distribution, effect on yield, and the way they affect target and non target organisms to enable an assessment of their potential for human and environmental hazards.

Maize is a staple Ghanaian food which is taken in many forms. It may be boiled, roasted, or ground into dough. Since lindane is applied to maize, a knowledge of lindane residues in maize will therefore be of immense importance to Ghanaians. Cow peas are also taken in Ghanaian homes in the boiled form. They are also ground and fried. In northern Ghana, the leaves are used to prepare soup [5].

This report therefore covers studies to identify and quantify lindane and endosulfan residues in maize and cowpea ecosystem respectively, under Ghanaian conditions.

2. EXPERIMENTAL

2.1. Apparatus

A Hewlett packard gas chromatograph series II, model 5890, equipped with a ^{63}Ni electron-capture detector and a SPB 5 capillary column (30 m \times 0.5 mm id) was used for the study. The carrier gas, nitrogen, had a flow rate of 15 mL min^{-1} . The operating temperatures were: injector, 200°C; detector, 300 °C; and column, 200° C isothermal. The attenuation was set on the HP 3396A recording integrator at 8, while chart speed was 10 mm min^{-1} . Each injection had a volume of 1 μl .

Solid Phase Extraction (SPE) Bond Elute C-18 columns (500 mg in 2.8 mL) were purchased from Analytical Instruments Varian GmbH, Vosendorf, Austria.

2.2. Reagents and materials

All solvents were Nanograde, and purchased from Mallinckrodt Specialty Chemical Co. Paris, Kentucky, USA. All other reagents were Analar grade. Commercial lindane formulated as an emulsifiable concentrate under the trade name Gammalin 20 was obtained from ICI Ghana Ltd, whilst endosulfan formulated under the trade name Thiodan 35 was obtained from Hoechst Aktiengesellschaft Frankfurt, Germany. Cypermethrin was obtained from Shell Chemicals, Abidjan, Cote d' Ivoire.

The maize planted was a local variety called "Abelehi" which matures in 107 days; and the cowpea a variety called "Asutem". Lindane and endosulfan standards were obtained from the International Atomic Energy Agency (IAEA) in Vienna.

2.3. Experimental farm

The experiment was conducted on a 1.6 ha. plot situated at the Ghana Atomic Energy Commission, Kwabanya, which is about 20 km from the centre of Accra. The plot had been treated with 1 kg AI of lindane the previous year. Four plots each measuring 50 m × 50 m were marked out. Two diagonally opposite plots were sown with maize whilst the others were sown with cowpeas.

2.4. Pesticide treatment

The maize plot that had been earmarked for pesticide treatment was sprayed twice with a water emulsion of the commercial lindane at a rate of 0.5 kg ha⁻¹ in accordance with local practices. The split application was done to avoid the resurgence of insect pests that may survive the first application.

Pesticide treatment of the cowpea plot was also in accordance with the recommendations of the Plant Protection and Regulatory Services of Ghana's Ministry of Food and Agriculture. Two earlier treatments were carried out with the synthetic pyrethroid cypermethrin at a dose of 50 g AI ha⁻¹. per each treatment. This application was to control insect pests that destroy cowpea flowers. These were followed by the application of endosulfan (Thiodan 35) at a dose of 750 g AI ha⁻¹. The endosulfan application was to control the pod borers that destroy cowpea seeds. The weeding, planting, spraying and harvesting schedules are as shown in Table 1.

Table 1. Schedule of activities

Activity	Crop	Date	
		Control	Treated
planting	maize	2/6	3/6
	cowpea	17-18/6	19-20/6
spraying, lindane	maize	-	17/6
	maize	-	1/7
spraying, cypermethrin	cowpea	-	21/7
	cowpea	-	31/7
spraying, endosulfan	cowpea	-	12/8
weeding, manual	maize	21/6	23/6
	cowpea	4-6/6	2-3/6
fertiliser application	maize	22/6	25/6
harvesting	cowpea	9/9	9/9
	maize	24/9	24/9

2.5. Soil characteristics

2.5.1. Sampling

The method followed the FAO/IAEA protocol [Appendix I]. Eight cores of soils were sampled at random from each sub-plot using a soil auger of 25 mm diameter and a depth of 150 mm. These were combined, ground and passed through a 2 mm sieve by mechanical shaking. Duplicate sub-samples were then taken for the measurement of pH, moisture content, organic carbon and particle size.

2.5.2. Determination of pH

The method followed that prescribed by Davies[6]. A sample (20 g) of the air dried sample was washed into of water (100 mL). The suspension was stirred with a mechanical stirrer for 5 minutes and allowed to stand for 30 minutes. The pH was determined using a meter which had been calibrated with pH4 and 7 standard buffers.

2.5.3. Determination of soil moisture

The soil sample were weighed into moisture cans and dried in an oven at 105°C to constant weight. The moisture content was determined as a percentage loss in weight.

2.5.4. Determination of organic carbon

The method followed that prescribed in Appendix I. A sample of soil (0.5 g) passed through a 0.5 mm mesh sieve was added to 1M potassium dichromate solution (10 mL). Concentrated sulphuric acid (20 mL) was added to the suspension, thoroughly swirled and allowed to stand on asbestos for 30 minutes. Distilled water (100 mL) was added, followed by orthophosphoric acid (10 mL) and finally barium diphenylamine sulphonate indicator (2 mL). The resulting solution was then titrated with 0.2M ferrous ammonium sulphate solution until the solution changed from blue to green. A blank titration was carried out to standardise the potassium dichromate.

2.5.5. Particle size analysis

Particle size analysis was determined by standard hydrometer method following dispersion with sodium hexametaphosphate, as prescribed by Charmetski [7] and Dickson [8].

2.6. Residue analysis

2.6.1. Sampling

2.6.1.1. Soils

Triplicate samples (50 g) were taken for residue analysis. For lindane, sampling was made at the following times (crop growth stages): pre-treatment (4th leaf stage), day of first treatment, 1 day (ankle stage) and 14 days after first treatment (knee high stage). Samples were also taken on the day of second treatment, followed by 14, 30 and 60 days (post tusseling stage) after the second treatment. For the cowpea plots (including the endosulfan experiment), sampling was done on the pre treatment day (flowering stage), 1, 7 (pod formation stage), 14 (pod drying stage), and 22 (harvesting stage) days post-treatment.

2.6.1.2. Leaves

For the maize ecosystem, the youngest and oldest leaves were taken off randomly chosen plants from each of the sub-plots as follows: pre-treatment day of first treatment 1, 14 days after first treatment. Samples were also taken on the day of second treatment, followed by 14 and 30 days after second treatment. With cowpeas, leaves were randomly sampled from the various sub-plots as follows: 1, 7, 14 and 22 days post-treatment.

2.6.2. *Extraction and clean up*

2.6.2.1. Soils

The triplicate samples were accurately weighed and transferred into a single thickness cellulose extraction thimble, (43 mm i.d. × 123 mm long) and extracted in a Soxhlet through 10 cycles i.e. for ~5 h with methanol (250 mL). An aliquot (5 mL) of the methanol extract was evaporated to dryness over nitrogen. The residue was taken up in methanol (1 mL) and to this was added water (2.5 mL). Prior to this, a C-18 Bond Elut solid phase extraction column had been reconditioned by flushing through two volumes of methanol followed by two volumes of distilled water. The diluted extract was poured through the reconditioned column at a flow rate of 2 mL min⁻¹ and then washed with methanol + water (3+7 by volume, 1 mL) followed by distilled water (1 mL). The washed column was vacuum dried for 15 min and then the pesticide was eluted from the column with hexane (2 mL) into a glass vial. The eluent was made up to 2 mL with more hexane and then analysed by gas chromatography.

2.6.2.2. Leaves and seeds

Leaf tissue (20 g) and/or seeds were homogenised separately for 2 minutes in a Waring blender and mixed thoroughly with methanol (200 mL). The suspension was filtered and a volume (20 mL) of the methanol extract was dried over nitrogen and the residue was dissolved in methanol (1 mL) and diluted with water (2.5 mL). The aqueous methanol extract was passed through a preconditioned SPE column at a flow rate of 2 mL min⁻¹. The column was then washed with distilled water (1 mL). The washed column was vacuum dried for 15 min and then eluted with hexane (1.5 mL) into glass vials. The eluent was made up to 2 mL with hexane and analysed by gas chromatography.

2.6.3. *Recovery experiments*

2.6.3.1. Soils

Samples of pesticide-free soils (50 g) were air dried and sieved as before. Volumes (1 mL) of lindane and endosulfan standards (10 µg mL⁻¹) were added to the soils. The soil was then extracted using a Soxhlet for 5 hours (10 cycles) with methanol. The extracts were then cleaned up for GLC measurements as in Section 2.6.1.

2.6.3.2. Seed and leaves

Samples (10 g) of maize grains, cowpea seeds, maize leaves and cowpea leaves were homogenised using Waring blender. A volume (1 mL) of standard solutions (10 µg mL⁻¹) was added to the homogenised samples (maize leaves and grains, lindane; cowpea seeds and leaves, endosulfan). Methanol (100 mL) was added to each mixture and shaken mechanically for thorough mixing. The methanol extract was filtered and a volume (20 mL) of the extract cleaned up for GLC analysis as in Section 2.6.2.

3. RESULTS AND DISCUSSION

3.1. Recoveries and soil characteristics

Though the plot had been exposed to pesticides only a year before the investigation, initial analysis before application of any pesticide showed the absence of pesticide residues within the limit of detection of the gas chromatograph ($0.01 \mu\text{g g}^{-1}$). All blank samples were found to be below the detection limit. Spiked soil samples gave recoveries of 80.5 and 92.7% for lindane and endosulfan residues, respectively. Spiked leaf samples gave recoveries of 95.2 and 80.5% for lindane and endosulfan residues, respectively.

The persistence of pesticides in soils is known to depend on soil characteristics. The major ones include organic matter, moisture, texture and acidity. The soil analysis showed the soils to be slightly acidic with an average pH of 6.3 and had a sandy clay loam texture (Tables 2 and 3). The average moisture content of the soils from the four subplots during the sampling period ranges from 4.78-13.3 g kg^{-1} and 3.40-7.74 g kg^{-1} in the maize and cowpea soils respectively.

3.2. The maize ecosystem

The dissipation of lindane in maize soils was very fast (Table 5) with 46% of the initial residue detected on the day of treatment lost one day after the first application (6th leaf

Table 2 Soil characteristics of the maize plots.

Property ^{a)}	Plot	
	Control (SD)	Treated (SD)
pH	6.21 (0.17)	6.3 (0.15)
% sand	64.2 (5.4)	63.8 (7.3)
% silt	13.4 (2.8)	11.1 (3.9)
% clay	24.5 (4.7)	25.0 (3.8)
% organic matter	2.4 (0.8)	1.9 (0.8)
% organic carbon	1.4 (0.5)	1.1 (0.5)

a) texture class : sandy clay loam

Table 3. Soil characteristics of the cowpea plots

Property ^{a)}	Plot	
	Control (SD)	Treated (SD)
pH	6.4 (0.3)	6.3 (0.2)
% sand	57.0 (7.6)	63.0 (3.2)
% silt	12.2 (2.4)	11.0 (1.6)
% clay	29.0 (5.0)	26.2 (4.0)
% organic matter	2.0 (0.4)	2.2 (0.3)
% organic carbon	1.8 (0.7)	1.3 (0.2)

a) texture class : sandy clay loam

stage), and less than 5% remained after 14 days. This may be attributable to volatilisation under the hot windy and rainy conditions of the area (Table 4). Increase in temperature is known to increase pesticide loss from soils through volatilisation, chemical degradation and bacterial decomposition, all of which are temperature dependent [9].

Rainfall might also have contributed to the pesticide loss by causing surface run-off. Furthermore, water competes with pesticides for adsorption to the soil surface [10] and lindane will therefore be more easily volatilised from surfaces of soil that have high moisture content. The moisture content of 5.8 to 13.8 observed during the period of investigation were fairly high and could have been a contributing factor to the fast dissipation rate.

About 5% of the residue detected on the day of first treatment remained in the soils 14 days (knee high stage) after application (Table 5). This is less than the 13% of the residue detected in soils 14 days after the second application (above knee high stage). This relatively slower rate of lindane dissipation after the second application might have been due to the fact that the maize plants, which had matured at the time of the second spraying provided cover over the top soil and lowered temperatures and minimised loss through volatilisation. Only 2.8% of the initial residue of lindane remained 30 days after the second application.

Thirty percent of the initial lindane residues detected in the maize leaves on the day of first treatment remained one day after application and fell to 3% 14 days after the first application (Table 5). Similarly, 14 days after the second application of lindane, the residues had fallen to 1.7% of the initial level.

Table 4. Monthly total rainfall and mean temperatures

	May	June	July	August	September
total rain, mm	49	81	41	17	75
mean temp. °C	28.5	26.9	25.5	25.3	26.4

Table 5 Lindane residues in soil samples and plant samples of the maize ecosystem

Sampling time	Mean lindane residue, $\mu\text{g g}^{-1}$ (SD)	
	Soil	Plant leaves
pretreatment	nd	nd
day of treatment	0.22 (0.04)	0.22 (0.014)
1 day after treatment	0.12 (0.05)	0.067 (0.01)
14 days after treatment	0.011 (0.008)	0.006 (0.002)
day of second treatment	0.025 (0.013)	0.24 (0.02)
14 days after second treatment	0.032 (0.006)	0.004 (0.001)
30 days after second treatment	0.008 (0.003)	nd
60 days after second treatment	nd	-

nd = none detected

Mean lindane residues levels in the maize grains were $0.02 \mu\text{g g}^{-1}$ (Table 7), which is well below the acceptable maximum residue limit for grains of $1 \mu\text{g g}^{-1}$ [10].

Table 6. Endosulfan residues in soil samples and leaf samples of the cowpea ecosystem

Sampling time	Mean endosulfan residue, $\mu\text{g g}^{-1}$ (SD)	
	Soil	Plant leaves
pretreatment	nd	nd
day of treatment	0.29 (0.17)	0.33 (0.022)
7 days after treatment	0.022 (0.011)	0.011 (0.006)
14 days after treatment	0.008 (0.005)	0.002 (0.001)
22 days after treatment	nd	nd

nd = none detected

Table 7. Lindane and endosulfan residues in maize and cowpea grains

Crop	Mean residues, $\mu\text{g g}^{-1}$ (SD)
maize	0.020 (0.002)
cowpea	0.045 (0.002)

3.3 The cowpea ecosystem

The amount of endosulfan in cowpea soils at 7 days after spraying (i.e. pod formation stage) was found to be only 7.6% of the residue detected on the day of treatment, similar to that found with lindane in the maize soils (Table 6). The low persistence of endosulfan as explained in the section for the maize ecosystem can also be attributed to the low organic matter content of the soil (ranging from 1.91 to 2.18%), the fairly high temperature, the sandy clay loam texture of the soil as well as the fairly heavy rainfall (Tables 3, 4). The mean residue level of endosulfan in cowpea seeds was $0.045 \mu\text{g g}^{-1}$ (Table 7). This is higher than the residue levels of lindane in maize grains and could be due to the fact that endosulfan was applied close to the time of seed formation, making it possible for endosulfan to be quickly incorporated into the seeds before it could dissipate.

CONCLUSION

Lindane and endosulfan have been shown to dissipate quickly under Ghanaian climatic conditions which are characterised by heavy rainfall and fairly high temperatures. The residue levels of lindane in maize and endosulfan in cowpea were well below the acceptable maximum residue levels.

ACKNOWLEDGEMENTS

This work was supported by the SIDA through the International Atomic Energy Agency under Research Contract No. 6754. The authors also acknowledge the contribution of Messrs B.Q. Modzinuh and S. Afful of the Chemistry Dept. of the NNRI for their assistance during sampling.

REFERENCES

- [1] WARE, G. W., Pesticide, Theory and Applications. W. H. Freeman & Co., San Francisco (1978) 14.
- [2] FAO International Code of Conduct on the Distribution and Use of Pesticides (1986) FAO, Rome.
- [3] IAEA, Isotope Techniques for studying the Fate of Persistent Pesticides in the Tropics, IAEA-TECDOC-476, Vienna (1988).
- [4] FAO/IAEA Technical Report Series No 329 (1991) Vienna.
- [5] Personal Communication with people from Northern Ghana.
- [6] DAVIES, Soil Sci. **22** (1943) 405.
- [7] CHARMETSKI, W.A., LICHTENSTEIN, E.P., J. Econ. Entomol. **66** (1973) 344-349.
- [8] DICKSON, K.B., BENNEH, G., A New Geography of Ghana (1988) Longman, London 22-26.
- [9] EDWARDS, C.A., Residue Reviews (1966) **13** 83-132.
- [11] Codex Alimentarius Commission. Guide to Codex recommendations concerning pesticide residues (1986) 3rd prelim. issue FAO/CAC/VOL 8 Ed 2.

RESIDUES OF LINDANE AND ENDOSULFAN IN WATER AND FISH SAMPLES FROM RIVERS, FARMS IN BESEASE, AGOGO AND AKOMADAN IN THE ASHANTI REGION OF GHANA



S. OSAFO-ACQUAAH
Department of Chemistry,
University of Science and Technology

E. FRIMPONG
Department of Biological Sciences,
University of Science and Technology

Kumasi, Ghana

Abstract

Pesticide residue analyses were performed on water and fish samples from River Oda in Besese, River Afranso in Nobewam near Kumasi, River Atwetwe in Akomadan, and River Kowire at Agogo. Residues of lindane and endosulfan were found in water and fish (*Oreochromis niloticus*, *Tilapia zillii*, *Barbus trispulis*, *Heterobranchus sp.*, *Tilapia busumana*, *Ophiocephalus obscura* and *Chana obscura*) samples. The residues of lindane varied between the years and months in the year but were in the range of 0.3 - 15 ng L⁻¹ (1993-94) and 8.7-32.0 ng L⁻¹ (1995) for the water samples and 0.2 - 24 ng g⁻¹ (1993-93) and 8.4-120.4 ng g⁻¹ (1995) for the fish samples. Residues of endosulfan in the water and the fish samples were zero in 1993-1994 but, in 1995, were in the range of 6.4-35.2 ng L⁻¹ for the water samples and 5.0-267.5 ng g⁻¹ for the fish samples. In all cases the lindane and endosulfan concentrations in the water were 10,000-20,000 times lower than known toxic concentration levels and therefore unlikely to cause fish toxicity problems.

1. INTRODUCTION

Organochlorine pesticides have found wider application in agriculture for the control of pests and in public health for vector control [1]. Furthermore, there has been an increase in pesticide use in Ghana over the past ten years as a result of the efforts by the government to boost food production.

In agriculture, organochlorine pesticides are widely used by farmers due to their cost effectiveness and their broad spectrum of activity, with a neglect of their possible impact on the environment and man. Research in the temperate regions and some parts of Africa have found the pesticides to be very persistent in the environment and remain in the soil [2] for longer periods of time resulting in serious damage to soil flora and fauna [3].

Detailed information on the effects of organochlorine pesticide residues and the amounts present in water, food, soil, flora and fauna in Africa is however, very limited. In some African countries, organochlorine pesticides have been banned based on data obtained from the developed countries in the temperate regions, without any assessment of the effect of these pesticides in the tropical environment [4].

Lindane (Gammalin 20) is widely used in Ghana in the cocoa plantations for the control of capsids on vegetables and for the control of stemborers in maize [5]. Endosulfan is used in cotton-growing areas, as well as on vegetable farms and for the control of pests in coffee plantations.

Most of the vegetable farms in the commercial vegetable producing areas in Ghana like Akomadan and Agogo are situated along rivers which act as the source water supply for farming and for drinking purposes. Pollution of rivers is known to occur through run-off of pesticides from farms during heavy rain as well as through aerial spraying of pesticides and the improper disposal and handling of the chemicals during and after spraying.

Endosulfan and lindane are known to be toxic to fish. For example, lindane has an LC₅₀ (48h) of 0.16-0.3 mg L⁻¹ for guppies [6] and an LC₅₀ (96h) values of 1.2 mg L⁻¹ and 0.75 mg L⁻¹ for *Fundus heteroclitus* and *Oreochromis niloticus* respectively [7, 8].

This report is based on our work on the effects of organochlorine pesticide residues in water and fish, in the forest zone of Ghana, as part of the coordinated research work of the Joint IAEA/FAO research programme on the 'Adverse effects on flora and fauna from the use of organochlorine pesticides on the African continent'.

2. MATERIALS AND METHODS

2.1. Experimental materials

Pesticide grade hexane was obtained from Mallinckrodt Special Chemicals, Kentucky, USA. HPLC grade methanol and acetonitrile were obtained from Aldrich Chemical Company, USA. Bonded phase C-18 SPE columns were obtained from Varian, Sunnyvale, USA.

2.2. Experimental site

This general study covered about 500,000 hectares of land in the closed forest region of Ghana which is about 34.5% of the entire land area of Ghana. Three regions of intense farming activities were selected for the residue analysis work after the general survey work had established the nature of the agricultural activities undertaken in the areas under consideration and the extent of pesticide usage.

The farming community at Akomadan is involved in large-scale vegetable farming in addition to maize farming with tomato farming constituting about 80% of the farming activity in the area. Besease and Nobewam are used mainly for rice production and are the location of the Anum Valley Irrigation Project. Close to the rivers are over 200 rice fields covering more than 400 hectares of land. Agogo is noted for its vegetable farming, rice and maize production and has other surrounding villages involved in cocoa farming.

2.3. Sampling

The pH values of the river waters were in the range 7.0-8.2 and the average temperature was 28°C. Water samples were collected at a depth of 250mm from River Oda at Besease, River Aframso at Nobewam near Kumasi, River Kowire at Agogo and River Aframso and River Atwetwe at Akomadan in the Ashanti Region in one litre sterilized bottles closed with a stopper. The duplicate samples were collected between January 1993 and July 1994 and from June 1995 to October 1995. The rivers pass through regions which have intense cocoa production and other farming activities. The initial stage of work, during 1993-94, involved random sampling in the various rivers. In 1995, a much more systematic sampling was undertaken from June to October and involved the farming communities at

Agogo and Besease. No samples were taken from Nobewam because neither lindane nor endosulfan were being used in the farming in this area. The river Oda at Besease flows through several areas of cocoa farming where lindane is actively used. Endosulfan is used in the farming at Agogo while other nearby cocoa farms use lindane.

After collection the water samples were taken to the laboratory for analysis. Fish samples for pesticide analyses were collected from River Oda near the rice fields and River Aframso in Agogo where there is large scale vegetable production. The fish samples (seven species) were obtained by a hook and line method and also with cages and nets. The fish samples obtained were immediately washed with distilled water and stored in a freezer.

The samples were later taken to the Ghana Atomic Energy Commission laboratory and the Institute of Aquatic Biology in Accra, Ghana for pesticide residue analysis.

2.4. Sampling of other species

Although several varieties of birds, rodents (rats, squirrels, grasscutter) and reptiles (lizards) were identified, the project focused on residues in fish. The rodent (grasscutter) samples, which were collected from the area, are yet to be analysed.

2.5. Laboratory analytical procedures

2.5.1. Water residue analyses

Volumes (2 x 10 mL) of methanol followed by volumes (2 x 10 mL) of distilled water were passed through an SPE C-18 column via a reservoir mounted on top of the column before conditioning. A small pressure was applied to the top of the reservoir to help force each volume of liquid to flow through the column at a rate of at least 2 mL per minute.

A volume (800 mL) of each water sample was passed through a pre-conditioned column at a flow rate of 2 ml per min. Each column was then washed with water + methanol (1 mL, 3.3 + 1 by volume) followed by distilled water (1mL) and dried under vacuum for 20 minutes. The sample which was trapped in the column was eluted with hexane (5 x 0.5 mL) to recover the pesticide residues.

2.5.2. Fish residue analyses

A sample (10 g) of each fish was homogenized using a pestle and mortar. The sample was mixed with sodium sulphate (10 g) and then transferred to a separatory funnel (250 mL). Hexane (20 mL) was added followed by acetonitrile (100 mL) which had been saturated with hexane, and the mixture was shaken for about one minute. The acetonitrile was drained into a 1 litre separatory funnel. The mixture was further extracted with portions of acetonitrile (3 x 50 mL) and the extracts combined with those in the separatory funnel. These were washed with water (500 mL) followed by saturated sodium chloride solution (40 mL) and hexane (50 mL). The mixture was allowed to stand and the aqueous layer was drained into another 1 litre separatory funnel and extracted with a further volume (50 mL) of hexane. The hexane extracts were combined and passed through a plug of sodium sulphate (10 g) into a flask. The solvents were evaporated almost to dryness and the residue dissolved in methanol (10 mL) and diluted with water (25 mL). A volume (3.5 mL) of this solution was taken for pesticide residue analysis (equivalent to 1 g of the fish sample) and passed through a

previously pre-conditioned column at a flow rate of 2 ml per minute. The column was washed, dried and eluted with hexane as described for the water samples above. Volumes (1 μL) of this hexane eluent were taken for analysis by the gas chromatographic analysis method detailed in Section 2.5.3.

2.5.3. Gas chromatographic analysis

Gas chromatograph : Hewlett-Packard Series II model 5890; column : 30 m x 0.5 mm i.d.; stationary phase : SPB-5; carrier gas: nitrogen; flow rate : 30 ml min^{-1} ; temperatures: column, 200°C, injector, 250°C, detector, 300°C; detector, ^{63}Ni electron capture. The sample injection volume was 1 μL . The limit of detection in this work was 1pg mL^{-1} . Recovery of lindane and endosulfan from the fish and water samples was 90%. Lindane and endosulfan in the residues were identified only by comparison of the retention times with those of their analytical standards and confirmation by an alternative method was not attempted.

3. RESULTS AND DISCUSSION

Seven species of fish caught in the various rivers were subjected to residue analysis. The species of fish caught depended on the time of the year when the trapping was done. It appeared that due to unknown adverse effects of some material, the fish populations had decreased considerably in these rivers. There were whole days when our team could not trap any fish in some of the rivers using a combination of a hook and line method, cages and nets.

Our initial survey work indicated that farms near River Oda at Besease were sprayed up to six times during the year. Farms near River Atwetwe and River Kowire which were along tomato-producing areas were subjected to pesticide applications of up to six times over either one or two seasons. Although lindane is not used widely in the spraying of tomatoes and rice, endosulfan is one of the commonly used insecticides in these regions. Despite this, the levels of endosulfan in the water samples were below the limits of detection (1 pg mL^{-1}). The concentrations of both lindane and endosulfan, when present, were found to be significantly higher in the fish than in the water (cf Tables 1, 2 with 3, 4) though the pesticide levels were well below their lethal levels, as discussed later.

Residues whose retention times corresponded with DDT and other organochlorines were observed in the chromatogram but were not quantified since the focus of this project was on lindane and endosulfan.

Table 1. Pesticide residues in river water samples during 1993-94

Sample		Residue, ng L^{-1}	
river identity	date	lindane	endosulfan
Oda	15.01.93	14.2	0
Oda	21.12.93	15.0	0
Atwetwe	15.01.94	1.3	0
Aframso	12.03.94	3.9	0
Kowire	07.07.94	3.0	0
Aframso	22.07.94	0.3	0

Table 2. Pesticide residues in fish caught during 1993-94

Source of fish		Species	Residue, ng g ⁻¹	
river identity	date		lindane	endosulfan
Oda	15.01.93	<i>Heterobranchus sp</i>	14.2	0
Oda	21.12.93	<i>Heterobranchus sp</i>	24.3	0
Atwetwe	15.01.94	<i>Barbus trispulis</i>	23.2	0
		<i>Chana obscura</i>	0.8	0
Aframso	12.03.94	<i>Tilapia sp</i>	0.2	0
		<i>Barbus tripulis</i>	0.2	0
Kowire	07.07.94	<i>Ophiocephalus obscura</i>	1.5	0

Table 3. Pesticide residues in river water samples during 1995

Sample		Residue, ng L ⁻¹	
river	month	lindane	endosulfan
Oda	June	25.0	35.2
	July	24.0	30.0
	August	22.0	15.0
	September	32.0	14.0
	October	8.7	6.4
Kowire	September	10.0	15.4
	October	14.8	25.6
Atwetwe	September	12.5	28.2
	October	20.2	14.6

Table 4. Pesticide residues in fish samples caught during 1995

Source of fish		Species	Residue, ng g ⁻¹	
river	month		lindane	endosulfan
Oda	June	<i>Oreochromis niloticus</i>	20.0	80.2
		<i>Barbus trispulis</i>	120.4	158.0
		<i>Heterobranchus sp</i>	14.0	160.0
	July	<i>Oreochromis niloticus</i>	22.5	267.5
		<i>Tilapia zilli</i>	34.0	142.5
		<i>Heterobranchus sp</i>	35.2	64.0
	August	<i>Oreochromis niloticus</i>	36.4	20.2
		<i>Barbus trispulis</i>	16.8	11.4
		<i>Heterobranchus sp</i>	24.8	6.5
	September	<i>Oreochromis niloticus</i>	100.0	186.0
		<i>Tilapia zilli</i>	72.0	77.0
		<i>Heterobranchus sp</i>	80.0	66.0
Kowire	October	<i>Oreochromis niloticus</i>	38.2	5.0
		<i>Barbus trispulis</i>	22.0	10.0
	September	<i>Oreochromis niloticus</i>	8.4	12.4
	October	<i>Oreochromis niloticus</i>	13.4	22.8

In none of the samples of river water collected during 1993 and 1994 was endosulfan found while lindane was found at low concentrations up to 15 ng mL^{-1} (Table 1). During these two years no endosulfan residues were found in the fish samples but lindane was found at concentrations up to 24.3 ng g^{-1} . This represents over 1000 fold concentration of lindane by the fish and is in general accord with, though exceeding, the known bioconcentration factor (153) for lindane and fish in water. Although the toxic levels of lindane to these particular species of fish are not known, the fact that they were surviving is not too surprising since the actual concentrations of lindane found in the water were around 10,000 to 20,000 times below the toxic concentrations to another species of fish (guppy, LC50 (48 h), $160,000 - 30,000 \text{ ng L}^{-1}$) [6].

In the more complete studies of 1995, both lindane and endosulfan were found in the river water samples. Concentrations varied with the location and the month of sampling but were in the ranges $8.7-25.0 \text{ ng L}^{-1}$ and $6.4-35.2 \text{ ng L}^{-1}$ for lindane and endosulfan respectively. As in the previous two years, lindane residues were found in the fish samples, ranging from $8.4-120.4 \text{ ng g}^{-1}$. These levels were generally higher than those of the previous two years but still orders of magnitude below likely toxicity levels.

In contrast to previous years, endosulfan was found in the fish samples and often at concentrations higher than lindane. In fact this is in general accord with its known bioconcentration factor for fish (583) which is approximately four times higher than that for lindane. In the case of endosulfan the aqueous concentration toxicity levels against local species of fish are known (LC50 (96 h), $1,200,000 \text{ ng L}^{-1}$, *Fundus heteroclitus*; $750,000 \text{ ng L}^{-1}$, *Oreochromis niloticus*). It can be seen that the concentrations in the river water samples are over 20,000 times below these toxicity levels and therefore present no toxic hazard to the fish.

So, in conclusion, although both lindane and endosulfan were found in river waters and it could be seen that the fish in these rivers absorbed and concentrated these insecticides, the levels that were actually found were so far below known toxicity levels that it is unlikely that they were the cause of the perceived low populations of fish in the rivers and that must be attributed to another cause.

ACKNOWLEDGEMENTS

We are grateful to the International Atomic Energy Agency for their grant support for this project and Dr. C.A. Edwards for his comments and assistance. We also acknowledge with gratitude the assistance of the Ghana Atomic Energy Agency and the Institute of Aquatic Biology, Accra in performing the residue analysis.

REFERENCES

- [1] EDWARDS, C.A., "Pesticide residues in soil and water", Environmental pollution by pesticides (Edwards, C.A., Ed), pp 409-458. Plenum Publ. New York (1973).
- [2] WHO technical report series, Research into Environmental pollution, No. 406, Geneva (1968).
- [3] COPE, O.B., Contamination of the fresh water ecosystem by pesticides, J Appl. Ecol., 3 (1966) 33-34.

- [4] TALEKAR, N.S., CHEN, J.S., KAO, H.T., Long term persistence of selected insecticides in subtropical soil: their adsorption by crop plants, *J. Econ. Entomol.*, **76** (1983) 207.
- [5] Ministry of Agriculture Technical Bulletin No.GDB/M1 Ghana.
- [6] Pesticide Manual, 10th Edition (C. Tomlin, Ed) The British Crop Protection Council, Surrey, UK (1994).
- [7] FELTZ, A.R., National monitoring program for the assessment of pesticide residues in water, *Pesticide Monitoring Journal* **5** (1971)54-62.
- [8] GODZI, A., Undergraduate thesis, University of Science and Technology, Kumasi,Ghana.
- [9] NEGRO, J.J, DONAZAR, J.A., HIRALDO, F., HERNANDEZ L.M., FERNANDEZ, M.A., Organochlorine and heavy metal contamination in non-viable eggs and its relation to breeding success in a Spanish population of lesser Kestrels (*Falco naumanni*), *Environmental Pollution* **82** (1993) 201-205.
- [10] SAMUEL,T. PILLAI, M.K.K., Impact of repeated application on the binding and persistence of [¹⁴C]-DDT and [¹⁴C]-HCH in a tropical soil, *Environmental Pollution* **74** (1991) 205-216.
- [11] WALISZEWSKI, S. M., Residues of lindane, HCH isomers and HCB in the soil after lindane application, *Environmental Pollution* **82** (1993) 289-293.

**NEXT PAGE(S)
left BLANK**



STUDIES ON THE DISTRIBUTION, SALE, HANDLING AND USAGE OF AGROCHEMICALS IN ASHANTI, BRONG AHAFO AND EASTERN REGIONS OF GHANA

S. OSAFO-ACQUAAH

Department of Chemistry,
University of Science and Technology

E. FRIMPONG

Department of Biological Sciences, University of Science and Technology

Kumasi, Ghana

Abstract

Studies were carried on the distribution, sale, handling and usage of agrochemicals in 19 major towns located in three regions noted for their intensive agricultural activities and covering 500,000 hectares of land area in Ghana. The most prevalent of the agrochemicals were insecticides (52%) followed by herbicides (28%) and fungicides (16%). Of these the organophosphates constituted 24% and the organochlorines 6% in terms of the total numbers of pesticides sold, though lindane and endosulfan were among the insecticides most frequently used by farmers. The fungicide, copper oxychloride as "Kocide", and the insecticide, λ cyhalothrin as "Karate", with 39% and 22% respectively of the sales by volume, were found to be the most widely used agrochemicals. Most of the merchants of agrochemicals and the farmers had only primary school education and had very little knowledge on the handling and safety aspects of the use of agrochemicals. The Extension Service to the farmers was inadequate and the farmers depended largely on the agrochemicals merchants for information on pesticide usage.

1. INTRODUCTION

This study was undertaken to provide additional information on the application, handling and use of pesticides in Ghana. With the launching of the Economic Recovery Programme there has been considerable emphasis on the development of agriculture in Ghana and pesticides usage has been increasing during the past ten years. Owing to costs associated with these chemicals and the lack of adequate information on the usage and application of pesticides, deleterious effects on the environment have been largely ignored. Furthermore, major law enforcement agencies like the Environmental Protection Council, Plant Protection and Quarantine Division and the Custom, Excise and Preventive Service have very little information at their disposal for effective legislation on the use of pesticides and the management of pesticides in general. Most of the data available are based on studies in developed countries. The objective of the study is to increase our knowledge of the distribution, usage and handling of pesticides in Ghana as part of the coordinated research programme on the adverse effect on the use of organochlorines (lindane and endosulfan) on the African Continent.

The study covered about 500,000 hectares of land area in the Ashanti, Brong Ahafo and Eastern Regions of Ghana located within the closed forest vegetation zone of Ghana; this constitutes about 34.5% of the total land area of Ghana. Sixteen major towns, noted for their intensive agricultural activities, within these three regions were selected for the study.

These centres were Agogo, Juaso, Konongo, Ejisu, Kumasi and Mampong in the Ashanti Region; Akomadan, Ejura, Kintampo, Sunyani, Techiman and Wenchi in the Brong Ahafo Region; and Nkawkaw, Anyinam, Koforidua, Nsawam, Suhum, Asamankase and Oda in the eastern Region.

The major crops cultivated in these areas include cocoa, palm fruits, rice, maize, cowpea and vegetables such as tomatoes, okro, eggplant and pepper.

2. MATERIALS AND METHODS

Questionnaires specifically designed for the study were used to collect data on the distribution, sale and usage of agrochemicals from the selected towns within the three Regions. Seventy-eight agrochemical sellers and one hundred and ninety-five farmers were interviewed. Besides the interviews, visits were made to selected farms to obtain additional information on the application of the agrochemicals.

3. RESULTS

3.1. Types and distribution of agrochemicals

The agrochemicals on sale and used by the farmers in the three regions were made up of insecticides (56%), herbicides (29%) and fungicides (16%) (Table 1). Insecticides always constituted the highest percentage of agrochemicals used in each of the three regions.

On the basis of their chemical composition the available agrochemicals fall into the following groups of compounds: organochlorines, organophosphates, carbamates, synthetic pyrethroids, inorganic compounds (mainly fungicides) and a miscellaneous category with a variety of chemical structures (Table 2). The organophosphates and organochlorines make up 30% of the total number of agrochemicals while the newer pyrethroids form a surprisingly large proportion (15%) of the market taking over from the older organochlorines and, to some extent, the organophosphates. DDT was one of the agrochemicals also found to be in use, though it is supposed to be banned in Ghana.

In terms of the volume of purchases the most commonly bought agrochemical was found to be Kocide, a copper oxychloride fungicide, and one of the least commonly bought was Sumithion, the organophosphate insecticide, fenitrothion (Table 3).

3.2. Socio-economic and other characteristics of merchants of agrochemicals (retailers and local distributors)

3.2.1. Educational background

The highest proportion of the merchants were middle school leavers with the remainder having secondary or other type of final education (Table 4). None of the merchants had a tertiary education.

3.2.2. Distribution of merchants by sex and age

Males (76%) were, by far, the greater proportion of the 78 merchants in the survey. The majority of them were between the ages of 20 and 40 years.

Table 1. Types of Agrochemicals with respect to their use, by farmers from three regions in Ghana.

Insecticides	Herbicides	Fungicides
Actellic (pirimiphos methyl)	Gesaprim (atrazine)	Baygon (propoxur)
Cymbush (cypermethrin)	Bellator	Caocobre
Decis(deltamethrin)	Garlon (triclopyr)	Champion
Diazinon	Gramoxone(paraquat)	Cobox(copper oxychloride)
Dimethoate 40EC	Ronstar (oxadiazon)	Dithane (mancozeb)
Dursban (chlorpyrifos)	Roundup(glyphosate)	Kocide (copper hydroxide)
Endosulfan	Samppi No3	Gastoxin
Fenitrothion	Stam F34T(propanil)	
Fenom C	Basta 20 SL (glufosinate)	
Primigram	Paracol	
Grofol	Saturnmate (thiobencarb)	
Karate (1 - cyhalothrin)	Bladex (cyanazine)	
Nogos (dichlorvos)	Garex	
Ofunack (pyridaphenthion)		
Perfekthion (dimethoate)		
Phostoxin (aluminium phosphide)		
Ripcord (cypermethrin)		
Secto		
Sumithion (fenitrothion)		
Thiodan (endosulfan)		
Trebon (etofenprox)		
Gammalin 20 (lindane)		
Callifan 50CE		
Furadan (carbofuran)		
Unden (propoxur)		
Totals 25	13	7

Common names of the active ingredient of commercial products are given in parenthesis where known.

3.2.3. Sales experience and training of merchants

Most of the merchants had been in the agrochemical business for less than 5 years (Table 6). Less than 3% had been associated with the sale of agrochemicals for more than 20 years. Most (95%) claimed to have had some basic training in handling of such chemicals. Almost all of them acquired their knowledge on the job through seminars and workshops organised by the Ministry of Agriculture, Extension Services, Plant Quarantine and Protection Services and Agrochemical Companies.

Table 2. Types of agrochemicals on the basis of their chemical composition

Organo-chlorine	Organo-phosphate	Carbamate	Synthetic Pyrethroid	Inorganic Compound	Miscellaneous
Thiodan 35EC (endosulfan) Gammalin 20 (lindane)	Actellic (pirimiphos- methyl) diazinon dimethoate 40EC Dursban (chlorpyrifos) Sumithion (fenitrothion) Nogos (dichlorvos) Ofunack (pyridathion) Perfecthion (dimethoate)	Baygon (propoxur) Furadan (carbofuran) Saturnmate (thiobencarb?) Sebon (?)	Cymbush (cypermethrin) Decis (deltamethrin) Karate (λ-cyhalothrin) Ripcord (cypermethrin) Fenom C (?)	Champion (?) Caocobre (?) Cobox (copper oxychloride) Kocide (copper hydroxide) Dithsne (mancozeb)	Gesaprim (atrazine) Garlon (triclopyr) Ronstar (oxadiazon) Secto (?) Stam F34T (propanil) Callifan 50CE (?) Gramoxone (paraquat) Roundup (glyphosate) Basta 20SL glufosinate
2 (6%)	8 (24%)	4(12%)	5 (15%)	5 (15%)	9 (27%)

Table 3. Commonly used agrochemicals by volume of purchase

Chemical	Percentage
Kocide	39.0
Karate	22.0
Perfekthion	11.0
Actellic	11.0
Karate	11.0
Sumithion	6.0

Table 4. Educational background of agro-chemical sellers in the three regions.

Level of education	Number	Percentage
Middle School Leaving Certificate	41	52.6
Secondary School	30	38.4
Others	7	9
University	-	-

Table 5. Length of work experience in retail selling

Duration	Number	Percentage
0 - 5 years	36	46.2
6 - 10 years	31	39.8
11 - 20 years	9	11.5
> 20 years	2	2.5

Table 6. Source of Supply of agrochemicals to retailers and farmers in Ashanti, Brong Ahafo and Eastern Regions of Ghana

Source of agrochemical	Percentage of the total market
Chemico (GH) Limited	23.1
Ministry of Agriculture	18.0
Reiss and Co.	6.4
Imperial Chemical Company	12.8
Wienco	5.1
Retailers	23.1

3.2.4. Handling of agrochemicals by merchants

(a) **Record keeping**

Less than 30% of the merchants kept records of their business including financial records, expiry dates of chemicals and stock controls.

(b) **Storage and display conditions**

Chemicals were mostly stored in the shops in their containers and some packed in original boxes. A few of the sellers had storage rooms. Sellers in kiosks and those operating on table tops displayed some of their items in the open sun.

(c) **Other activities undertaken by merchants in shops**

Apart from the sale of agrochemicals over 90% of the merchants stocked other items including seeds, fertilizers, etc., in their shops. In a few instances, food items such as bread, vegetables, gari and rice were also sold in agrochemical shops.

(d) **Accidents**

There were some reported cases of accidents which occurred as a result of the improper repackaging of agrochemicals into smaller packages for sale to farmers. Reports of occasional headaches and other health problems were considered by the sellers to be related to the chemicals.

3.2.5. Major sources of supply of agrochemicals to retailers and farmers

The supply of agrochemicals to farmers was from commercial sales outlets except for a few cases in the cocoa industry where some farmers obtained chemicals directly from the Ministry of Agriculture. Most of the supply to farmers was from retailers in open market places throughout the regions.

3.2.6. Educational background of farmers

Nearly half (47%) of the farmers had Middle School Leaving Certificate though a reasonable proportion (17%) had no formal education at all (Table 7) and most (92%) were male.

Table 7. Educational background of farmers using agrochemicals.

Level of education	Number	Percentage
Middle School leaving Certificate	92	47.2
G.C.E. O'Level	39	20.0
Others - Post Sec, Agric. Colleges	26	13.3
University	4	2.1
None	34	17.4

3.2.7. Tenure and size of farm

Over one third (38%) of farmers owned their own lands with 18% based on the extended family system. The largest proportion (42%) of the land being farmed was rented but only a very small fraction (0.5%) was under the cooperative system. In general, the size of most farms for individuals was between 0.8 and 2.5 hectares. However there were a few cases where farmers owned about 15-20 hectares of land.

3.2.8. Experience of use, application and storage of pesticides by farmers

Most (66%) of the farmers had used pesticides actively over the past 6 years, though only about 7% had used pesticides for a period greater than 15 years (Table 9).

Most of the individual farms did not receive any Extension Service except those involved in cocoa farming. Some of the rented farms and cooperatives benefitted from those extension services provided by the Ministry of Agriculture with farms in the cocoa producing areas benefitting most from them.

Table 8. Type of farm tenure

Type of tenure	Number	Percentage of total
Owner	75	39
Family	36	19
Cooperative	1	0.5
Rented	83	43

Table 9 Duration of use of pesticides on farms

Years of Use of Pesticide on farm	Number	Percentage
0 - 5 years	128	65.6
6 - 10 years	47	24.1
11 - 15 years	6	3.1
> 15 years	14	7.2

Half (51%) of the farmers did not have any training in the use of agrochemicals although the remainder have had at least some information on the use of particular pesticides on their farm. Such information was from friends, Agricultural Extension Officers and agrochemical merchants.

Nearly all (98%) of the farmers used knapsack sprayers while a proportion (12%) of these used Mistblowers in addition to the knapsack sprayer. Less than 5% used other applicators and sprinklers.

Agrochemicals were mostly stored in the house before use though excess agrochemicals were stored under a variety of conditions. A third (31%) of the farmers used the agrochemicals immediately on the farm. A larger proportion (54%) stored them on the farm or buried the containers in the ground on the farm for future use, while a significant fraction (12%) poured them away on the farm.

A very small proportion (<5%) of the farmers used adequate protective clothing. The majority of the farmers used a variety of protective measures which appeared inadequate, though a quarter (24%) used nose protectors or cloths to cover their nose. Only a few (18%) either washed their hands after using the chemicals or used wellington boots or gloves (15%). A significant fraction (22%) indicated they did not use any safety measures at all.

There were reports of fainting (12%) after the spraying exercise and all the farmers complained of at least one ailment (fever, headache, stomach pains, chest pains) which occurred very frequently.

4. DISCUSSION

Pesticides have been in use in the Ashanti, Brong Ahafo and Eastern regions of Ghana for a relatively long time, and at least 3% of the agrochemical shops have been in operation for over 20 years whilst some farmers have used pesticides on their farms for more than 15 years.

Insecticides form the most prevalent type of agrochemical used in all three regions of Ghana and constitute about 52% of the total number of agrochemicals in use. Lindane and endosulfan are among the insecticides being used most extensively. These organochlorines can be persistent and their accumulation could pose a threat to man and the environment. Persistent insecticides, such as DDT, which have been banned are still available in shops and are used extensively on farms. This emphasises the need for a more strict enforcement of laws regarding the importation and use of such chemicals in Ghana. The danger posed by the low educational background of a large number (52%) of the agrochemical merchants needs to be seriously addressed. These merchants not only mishandle pesticides through lack of knowledge but farmers depend on them as their main source of information. There is the need to intensify training programmes to include not only the handling of such chemicals but also the disposal of stocks pesticides that have exceeded their expiration dates. The tendency for farmers to purchase such agrochemicals and even banned pesticides at relatively reduced prices is closely related to a lack of education coupled with the fact that the merchants who advise them may themselves lack the requisite knowledge about the chemicals. The relatively large number of farmers (51%) who have not had any previous training in the use of agrochemicals may further compound the problem of mishandling of these agrochemicals.

Improper record keeping also posed a big problem in the collection of data on volume of usage of these agrochemicals. The lack of an extension service to family and self-owned farms needs to be remedied. It deprives such farmers of the opportunity of benefitting from training programmes in the correct application and handling of agrochemicals.

Safety measures taken by the farmer were found to be grossly inadequate with less than 5% using adequate protective clothing while 22% did not use any safety measures. A programme of workshops, seminars and other training programmes, aimed at enhancing the farmer's capacity to handle and use agrochemicals to their full potential, needs to be set up. Furthermore, since pesticides can have adverse effects on the environment, a better knowledge of the proper handling of these chemicals from the manufacturers through retail merchants to the farmers could ensure for safer working conditions for the distributor and the farmer and reduce the potential for environmental contamination.

This study presents some of the problems related to the management of pesticides in Ghana which we believe could be mitigated through the coordinated effort of policymakers and scientists on the African Continent with the support of interested International funding agencies and organizations involved in the protection of the environment.

ACKNOWLEDGEMENT

We are grateful to the International Atomic Energy Agency for their grant support for this project.

PERSISTENCE AND FATE OF SOIL APPLIED ^{14}C - LINDANE IN A MAIZE ECOSYSTEM



XA9743722

P.O. YEBOAH
Department of Chemistry,
Ghana Atomic Energy Commission

K.G. MONTFORD
Biotechnology and Nuclear Agricultural Research Institute,
Ghana Atomic Energy Commission

W. APOH, S. DODOO
Department of Chemistry,
University of Cape-Coast

Legon, Ghana

Abstract

^{14}C -lindane applied to soil surface in a maize ecosystem (one month after planting) was taken up by the plant. Within the first 25 days of treatment, ^{14}C -lindane or its metabolites were found within the entire plant with the greatest concentration in lower leaves (from the ground level); and a sharp build up of lindane concentration towards the tip of each leaf. Radioactivity and hence pesticide concentration was uniformly distributed in the plant with time; to the extent that measurable levels of ^{14}C -compounds were detected in the tassel cob and the grain. This indicated that soil applied lindane was available to the maize plant. The persistence of ^{14}C -lindane in soils of variable organic matter content was also studied. Evidence is presented to show that ^{14}C -lindane dissipated faster in soils of lower organic matter content. Levels of surface applied pesticides that became bound in the soil increased with time after application and also with increasing organic matter content. ^{14}C -activity was mainly associated with the top soil layer (0-30 mm).

1 INTRODUCTION

The following study was designed to complement observations made in the previous paper [1].

2 MATERIALS AND METHODS

2.1. Chemicals

Uniformly labelled ^{14}C -lindane (specific activity $647 \text{ MBq mmole}^{-1}$) was obtained from Sigma Chemical Co. St Louis, USA, through the International Atomic Energy Agency (IAEA) in Vienna. The ^{14}C -concentration of the lindane (supplied as toluene solution) was found to be 1 kBq ml^{-1} . The scintillators 2,5-diphenyloxazole (PPO) and 2,2-phenylenebis-(4-methyl-5-phenyloxazole) (DMPOPOP) were purchased from Eastman Kodak Co., Rochester, N.Y., USA. All other chemicals were Analar grade.

2.2. Studies on the persistence of ^{14}C -lindane

Fifty-one PVC tubes (length, 300 mm; internal diameter, 70 mm) were buried in the soil in the rows between plants (in another plot), with 30 mm projecting above ground level. The tubes were arranged into groups of fifteen and in accordance with the organic matter content of the soils they contained. The soil in each of these group tubes were treated with ^{14}C -lindane (185 kBq) at the same time that plants on the field were sprayed with

commercial lindane. Two tubes in each soil were not treated with ^{14}C -lindane to serve as the control. The tubes were dug out carefully at intervals, wrapped in polythene bags and stored in a refrigerator pending extraction and determination of pesticides. The soil in each of these tubes was sprayed with ^{14}C -lindane (185 kBq) in hexane (100 mL).

2.3. Studies on the uptake of ^{14}C -lindane into plants

Twenty plastic buckets (length, 220 mm; internal diameter, 240 mm and open at both ends) were filled with soil from one of three sources, as described previously [1]. They were buried in the crop row with 30 mm projecting above ground level. Fifteen tubes of each soil type were grouped together. Maize was planted in each bucket on the same day that planting was done in the field. The soils in the buckets were treated with ^{14}C -lindane (370 kBq) on the same day that the plants on the experimental farm were sprayed with commercial lindane. Three plants were taken periodically and analysed for uptake of ^{14}C -activity. Both anatomical and sectional distribution of ^{14}C -lindane residues in the maize plants were studied.

2.4. Extraction and analysis

2.4.1. Soils

The soil contained in the PVC tubes was scooped out from top (end exposed to the atmosphere) in successive 150 mm layers. Each soil layer was weighed, dried, ground separately (using a pestle and mortar) and sieved through a 2 mm mesh sieve. Samples (40 g) of ground soil were weighed into cellulose extraction thimbles (size 43 × 123 mm) and soxhlet extracted for 2.5 h using methanol. The extracts were concentrated at $50 \pm 5^\circ\text{C}$ under vacuum to a volume of 5 mL which was then transferred into scintillation vials and analysed for ^{14}C -activity. Recovery was 80.2%.

2.4.2. Plants

Anatomical sections of maize plants above ground level (leaves, stem, cob, etc.) were removed at intervals and stored in a refrigerator for analysis. Stem and leaves were carefully removed from points of attachments cut into 30 mm segments and analysed for ^{14}C -activity using the dry combustion method.

2.5. ^{14}C -activity measurements

Total and bound residues in soils as well as in plant parts were determined by combusting weighed samples in a model 600 Biological Material oxidiser (R.J. Harvey Instrument Corporation, New Jersey, USA). The $^{14}\text{CO}_2$ was absorbed in absorber + cocktail (1+1 by volume, 2 mL). The absorber was made up of ethanolamine + methanol (2.5 + 17.5 by volume) and the cocktail consisting of PPO (5 g) + DMPOPOP (50 g) dissolved in toluene (1 L). ^{14}C -activity was measured by liquid scintillation counting (LSC) in a Tri-Carb liquid scintillation Analyser model 100 using the sample channel ratio method for quench correction.

3. DISCUSSION

3.1. Uptake and translocation in maize

Within 25 days of pesticide application to the soil, it was observed that the plants had taken up a measurable amount of ^{14}C -activity with the greatest amount of 0.08% of applied found in the lowest leaf, i.e. the first leaf from the ground level (Fig. 1). There was then a gradual decrease in pesticide concentration in the higher leaves and the stem. The upwards movement of ^{14}C -activity in the maize plant was further shown by the gradual build up of ^{14}C -activity towards the tip of lower leaves (Figs 3, 4). With time, i.e. within 50 days after




Fig 1. Percentage of initial dose recovered from leaves and stem.




Fig 2. Anatomical distribution of ^{14}C lindane in maize plant.

pesticide treatment, the ^{14}C -activity tended to spread out more uniformly throughout the plant though the greatest concentrations of ^{14}C -activity remained within the bottom three leaves with smaller ^{14}C -activity in the uppermost leaves and the stem (Fig. 2). Some ^{14}C -activity was also found in the tussel, the cob and the grain 50 days after treatment (Fig. 2) indicating that soil applied lindane or its metabolites were available for uptake by the plant. The translocation of soil applied lindane and its metabolites have been investigated in detail [2, 4, 5, 6]. Neither lindane nor its metabolites were evenly distributed within the plants. Comparatively high residue levels were always detected in the leaves whereas small amounts were translocated into the stems, leaves, and fruits. Paasivirta et al. [5] showed that in water plants, lindane concentrations were similar in both the roots and the stem. Differences in residue levels have been shown to be dependent on plant species. Of a series of edible crops grown in soils containing lindane, carrots were shown to have higher levels than beans, tomatoes and potatoes in that order. In soils with low organic matter contents, as in this study, insecticide residues are more available and hence susceptible to uptake by plants [7].




Fig 3. Distribution of ^{14}C lindane in maize leaves (25 days).




Fig 4. Distribution of ^{14}C lindane in maize leaves (50 days).




Fig 5. Bound residues of ¹⁴C lindane in soils of different organic matter content.




Fig 6. Dissipation of ¹⁴C lindane in soils of different organic matter content.




Fig. 7. Reciprocal value of activity against time.

3.2. Movement in soils

Lindane dissipated faster in soils of lower organic matter contents (Fig.5). In the three soils tested, there was a rapid rate of dissipation of ^{14}C -lindane within the first 10 days of application followed by a gradual decline in this rate. This rate of dissipation was attributed to the low organic matter content of the soils in this study. Although there were copious amounts of rainfall during the period after pesticide application (104 mm in 2 months), residues of ^{14}C -lindane were found only within the first 0-30 mm layer of the soil surface in agreement with Cliath and Spencer [7] who observed no mobility of lindane in clay soils.

ACKNOWLEDGEMENTS

This work was supported by SIDA through the International AtomicEnergy Agency under Research Contract No. 6754. The authors acknowledge the contributions of Messrs B.Q. Modzinuh and S. Afful of the Chemistry Department of the NNRI for their assistance during sample collection.

REFERENCES

- [1] YEBOAH, P.O., MONTFORD, K.G., KLAKE, R.K., AKPABLI, C.K., this TECDOC.
- [2] ITOKAWA, H., SCHALA, A., WEISBERGER, I., KLEIN, W., KORTE, W., Tetrahedron **26** (1970) 763-773.
- [3] BALBA, M.H., SABA, J., Environ Lett. **7** (1974) 181-194.
- [4] BRADBURY, F., Ann. Appl. Biol. **52** (1963) 361-370.
- [5] PAASIVIRTA, J., PALM, H., PAUKKA, R., AKHABUHAYA, J., LODENIUS M., Chemosphere **17** (1988) 2055-2062.
- [6] FUHREMANN, T., LICHTENSTEIN, E., J. Agric. Food Chem. **28** (1980) 446-452.
- [7] CLIATH, M., SPENCER, W., Environ Sci. Technol. **6** (1972) 910-914.

**NEXT PAGE(S)
left BLANK**

PERSISTENCE OF LINDANE IN RICE AND MAIZE ECOSYSTEMS IN NIGERIA



XA9743723

G.U. OKEREKE, C.C. ONOCHIE
Biotechnology Research Centre, Nnamdi Azikiwe University,
Awka, Anambra State, Nigeria

E. ONYEAGBA
Anambra College of Agriculture, Igbariam, Anambra State, Nigeria

Y. DJE
Department of Food Technology and Science, University of Tennessee,
Knoxville, Tennessee, United States of America

Abstract

A three year field study was undertaken to examine the fate of residues of lindane in soil and crops after repeated seasonal applications of lindane to soil in which maize and rice crops were grown. In the 1993 rice study, the lindane residues in the treated soil were 0.6 mgkg⁻¹ (1day) and 0.04 mgkg⁻¹ (2 months) after application respectively. In the soil from the maize plots, lindane residues were 0.19 mg kg⁻¹ (1 day) and 0.16 mg kg⁻¹ (1month) after application and not detectable after 6 months. In the 1994 maize study, the lindane residues in the soil were 0.12 (1 day), 0.10 (1 month) and 0.033 mg kg⁻¹ (2 months) after application respectively while in the crop they were 0.19 (1 day), 0.105 (2 weeks) and 0.05 (4 weeks) mg kg⁻¹, respectively. In general, lindane residues were not detected in the soil one year after application indicating that there was no accumulation of lindane in the soil during the period 1992-1994 and therefore unlikely to be any long term effects on soil fauna. Lindane residues in rice and maize crops continued to decrease during the period up to harvest.

1. INTRODUCTION

Control of insect pests of rice and maize rely largely on the use of chemical pesticides. In addition to lowering the population levels of an insect, pesticides can cause changes within both the non-target organisms and co-inhabitants of its environment. Some pesticides can also accumulate in the soil and cause environmental pollution when they are leached to the underground water or runs off to streams and rivers.

Most pesticidal chemicals can be absorbed to varying degrees and in some cases can be translocated to the edible parts of plants growing in contaminated soils [1]. The presence of chlorinated hydrocarbon pesticide residues in human adipose tissue most often results from the consumption of contaminated food, either deliberately treated with chemicals for the control of pests or incidentally contaminated from environmental or industrial sources [2]. Despite the use of lindane in controlling insect pests, little is known about its fate in the soils and crops under conditions of Nigerian agriculture.

The publicity generated by the restrictions of certain pesticides, especially organochlorines, in developed countries has led to the ban of the use of these chemicals without sufficient information on their impact on other environments. Knowledge of the behaviour of pesticides and their residues in soils and crops in Nigeria is vital as it is not scientifically justifiable to make predictions of pesticidal residue accumulation using data generated elsewhere since pesticidal residue accumulation in will depend on factors such as the nature of the pesticide, soil type, moisture, soil pH, temperature, cultivation, mode of application and soil organisms [3]. The objective of the study was to determine lindane pesticide residues in soils and their absorption into crops in a field experiment after yearly applications of lindane.

2. MATERIALS AND METHODS

2.1. Materials

Lindane was obtained as the commercial formulation (200 g L⁻¹ emulsifiable concentrate, Gammalin 20, ICI Ltd, Nigeria). A solution containing standard lindane was obtained from Supelco, Baltenfonte, PA by the International Atomic Energy Agency, Vienna, Austria. The standard solution was stored in the refrigerator (4°C) until used. Analytical grade hexane, methanol and acetone were supplied by Aldrich Chemical Ltd, Gillingham, Dorset SP8 4JC. Sep Pak phase C-18 cartridges were obtained from Waters Chromatography Division, Millipore Cooperation, Milford, Massachusetts, 01757, USA.

2.2. Plot design and lindane application

The plot design and application procedures have been described elsewhere [see this Tecdoc, Umeh *et al.*].

2.3. Crop sampling

The aerial parts of 12 plants per subplot were collected 1 day and 2 months after lindane application from both the rice and maize fields in the 1992 and 1993 studies, while in 1994 the sampling intervals were 1 day, 2 weeks and 1 month after application. Amounts (~100 g) of plants were chopped into one mixture and stored in polyethylene bags in a freezer compartment until extracted.

2.4. Soil sampling

Soils were sampled at 1 day pre-treatment, 1 day, 2 and 6 months after lindane application. On each sampling occasion, 12 cores of soil were taken from a 5 m x 5 m area at the side of each subplot to a depth of 15 cm using a soil auger. The samples were ground to pass 0.25 mm mesh sieve and mixed to provide one bulk sample for each subplot. Duplicate soil samples were taken for measurement of moisture content and for residue analysis. All soil samples were stored in polyethylene bags in a freezer compartment until extracted.

2.5. Residue extraction and analysis

2.5.1. Extraction of plant tissue samples

The plant tissue samples were thawed at room temperature. Samples (20 g) were homogenised with of methanol (200 mL). These suspensions were shaken mechanically for 1 h in a screw-capped jar. The suspensions were filtered to remove plant debris. Volumes (10 mL, 1993; 50 mL, 1994) were dried by aspirating with nitrogen and redissolved in methanol + water (3.5 mL, 1+2.5 by volume).

2.5.2. Soil

Samples (20 g) of soil were placed in a screw-capped bottle, mixed with methanol (200 mL) and mechanically shaken for 1 h. The suspension was filtered and volumes (10 mL, 1993; 50 mL, 1994) were dried with nitrogen. Each residue was then redissolved in methanol + water (3.5 mL, 1 + 2.5 by volume).

2.5.3. Clean up procedure

Each aqueous methanol extract was passed through a preconditioned C-18 SPE column and then eluted with hexane. The hexane extract was dried with nitrogen and redissolved again in methanol and the clean up procedure repeated. The residue was dissolved in hexane (2 mL) and stored in a refrigerator until analyzed by the gas chromatographic method.

2.5.4. Gas chromatographic analysis

1992^{a)} : Instrument, Shimadzu 9; column, fused silica capillary, 30 m x 0.25 mm i.d.; temperatures, column, 240°C, injector, 250°C, detector, 350°C; carrier gas, nitrogen; detector, ⁶³Ni electron capture.

1993^{b)} : Instrument, Varian 3700; column, fused silica capillary; temperatures, column, 190°C; injector, 220°C; detector, 270°C; carrier gas, nitrogen; detector, ⁶³Ni electron capture.

1994^{a)} : Instrument, Hewlett-Packard 5890A; column, fused silica capillary, 30 m x 0.25 mm i.d.; temperatures, column, 240°C ; injector, 250°C; detector, 300°C; carrier gas, nitrogen; detector, ⁶³Ni electron capture.

^{a)} Dept. of Food Technology and Science, University of Tennessee, Knoxville, USA.

^{b)} Dept. of Chemistry, University of Ibadan, Nigeria.

3 RESULTS AND DISCUSSIONS

The limit of detection of lindane was 0.0044 mg kg⁻¹ for both crop and soil samples. The recovery from fortified soils after clean up was 93.5%. There were no detectable lindane residues before lindane application in 1992 for the rice plots and in 1992 and 1993 for the maize plots. Lindane residues were not detected in the untreated rice and maize soil because the field experimental design was oriented to avoid any contamination by erosion from treated to untreated plots. Absence of lindane residues at the beginning of the studies in each new site confirmed that lindane had not been used in the locations before the study was started.

Only data of the *extractable* lindane residues in the soil are reported. In the 1993 rice study, the lindane residues in the treated soil were 0.36 mg kg⁻¹ (1 day) and 0.04 mg kg⁻¹ (2 months) after application respectively. For the maize soil the residues were 0.19 mg kg⁻¹ (1 day) and 0.016 mg kg⁻¹ (1 month) after application and were either very low (< 0.09 mg kg⁻¹) or not detectable at 6 months after application. All these results suggest that there is no problem of long term residues of lindane building up in the soil from either rice or maize fields.

The 1994 maize study gave similar results to that of 1993 in that residues of lindane in the soil were not detectable after 6 months probably as a result of loss of lindane due to volatilization and microbial degradation. Some other studies have shown that organochlorine pesticides dissipate at higher rates under tropical and sub-tropical conditions than in temperate regions [4, 5]. The high lindane residues (0.25 mg kg⁻¹) found in maize foliage, 1 day after application, will have arisen from lindane sprayed onto the leaves, since it was applied as a surface treatment and not incorporated into the soil. The residues dissipated rapidly (0.094 mg kg⁻¹ at 2 months after application) but not as fast as they did in 1993 when

they were found to be as low as 0.0033 mg kg⁻¹ at the same interval after application. The cause of the differences between the years is unknown.

Table 1. Characteristics of the soils from the crop fields

Field Crop	% sand	%silt	%clay	% organic matter	pH	%N
rice ^{a)}	21	25	54	2.23	6.3	0.9
maize ^{a)}	25	20	55	1.85	6.1	0.08

a) classified as a clay

Table 2. Concentration of lindane samples taken from the rice field

Year	Crop	Sample	Lindane residues at period from spraying mgkg ⁻¹ (SD)				
			1 day before	1 day	1 month	2 months	6 months
1992	rice ^{a)}	soil ^{d)}	ND	NM	NM	0.17 (0.05)	0.09 (0.11)
		crop	-	0.53 (0.19)	NM	0.1 (0.04)	-
	maize ^{b)}	soil ^{d)}	ND	NM	NM	0.1 (0.02)	0.09 (0.01)
		crop	-	0.25 (0.19)	NM	0.094 (0.004)	-
1993	rice ^{a)}	soil ^{d)}	ND	0.36 (0.02)	NM	0.04 (0.05)	ND
		crop	-	NM	0.011 (0.01)	0.02 (0.002)	-
	maize ^{c)}	soil ^{d)}	ND	0.19 (0.013)	0.16 (0.013)	NM	ND
		crop	-	NM	0.0033 (0.003)	NM	-
1994	maize ^{c)}	soil ^{d)}	ND	0.12 (0.007)	0.103 (0.44)	0.033 (0.045)	NM
		crop	-	0.19 ^{c)} (0.06)	0.05 (0.03)	NM	-

ND = not detectable, NM = not measured

a) plot location, Igbariam; b) plot location, Ugwuoba; c) plot location, Enugu
d) there were no detectable residues of lindane in any of the samples of soils taken from the untreated control plots before and after application to treated plots on each occasion.

e) lindane residues at 2 weeks after application = 0.11 (SD,0.02) mgkg⁻¹.

4. CONCLUSIONS

In conclusion, our results indicate that lindane was not detected in the soil samples taken 6 months after application to rice and maize agro-ecosystems in Nigeria. The results also show that there was no accumulation of lindane or its metabolites in the soil during the study period thus giving little cause for concern as far as a build-up of lindane residues and long term effects on either soil fauna or transfer to the food chain is concerned. They also show that data from temperate countries should only be applied with caution to Nigerian conditions. The variability in the physical and biological properties of soil give rise to difficulties in the prediction of the fate of pesticides as their degradation is a complicated function of volatilisation loss by erosion, leaching, biotic and abiotic conversion [6].

ACKNOWLEDGEMENTS

The authors wish to thank the International Atomic Energy Agency for supporting the work under the funds provided by Swedish International Development Agency (SIDA). Thanks are also due to Professor O. Osibanjo of Department of Chemistry, University of Ibadan, Nigeria and Professor S.T. Melton of Department of Food Technology and Science, University of Tennessee, Knoxville, USA for use of their laboratory facilities and gas chromatograph to run the residue analyses.

REFERENCES

- [1] TALEKAR, N.S., SUM, L.T., LEE, E.M., CHAN, J.S.J., Persistence of some insecticides in the Tropics, Kenya. *J. Sci. Technol. (Ser A)* **5** (1977) 31-44.
- [2] SANBORN, J.R., CHILDOR, W.F., HANSEN, L.G., Uptake and elimination of ¹⁴C hexachlorobenzene (HBC) by the green sunfish (*Leponis cyanellus* Raf.) after feeding contaminated food, *J. Agric. Food Chem.* **25** (1977) 551-553.
- [3] LICHTENSTEIN, E.P., SCHULTZ, K.R., Persistence of insecticidal residues in soils, *J. Econ Ent.* **54** (1961) 617-622.
- [4] SLEICHER, C.A., HOPCRAFT, J., Persistence of pesticides in surface soil and relation to sublimation, *Environ. Sci. Technol.* **18** (1984) 514-518.
- [5] PERFECT, T.J., Environmental impact of DDT in a tropical agro-ecosystem, *Ambio*, **16**(1980) 16-21.
- [6] HORVATH, L., KLING, F., SIMON, L.P., Fate of pesticides in soilsof various organic matter, *Pesticides, Food and Environmental Implications. Proceedings of a symposium, Neuherberg 24-27 (Nov. 1987)* 177-183.

**NEXT PAGE(S)
left BLANK**

LINDANE RESIDUES IN FISH INHABITING NIGERIAN RIVERS

G.U. OKEREKE

Biotechnology Research Centre, Nnamdi Azikiwe University,
Awka, Anambra State, Nigeria



XA9743724

J.CHIDODBEM

Enugu State University of Science and Technology,
Enugu, Enugu State, Nigeria

Y. DJE

Department of Food and Technology and Science, University of Tennessee,
Knoxville, Tennessee United States of America

Abstract

Analysis for residues of lindane in fish collected from various rivers close to rice agroecosystems showed that the concentrations of lindane ranged from none detectable to 3.4 mg kg^{-1} . Fish from rivers where strict regulations prohibits its use had no detectable lindane residues while appreciable amounts of lindane were found in fish where such restriction was not enforced with the variation attributed to the extent of use of lindane in the area of contamination. The investigation confirms that the use of lindane in rice production in Nigeria can cause the contamination of fish in nearby rivers.

1. INTRODUCTION

Pesticides have benefitted man by reducing loss and increasing food production, but unfortunately have been a major source of pollution to fish [1, 2]. Lindane residues can be stored in tissues of birds, and mammals including man [3] and can cause death to organisms at the end of the food chain [4]. Pollution of the environment and its effects on the health of man, animals, fish, wildlife and the ecosystem in general have become a subject of concern[5]. Broad conclusions regarding lindane levels can be drawn from comparison of their distribution in biological materials from a given area. Because of the widespread distribution of fish, their opportunity for exposure to pesticides applied in rice agroecosystems, their role in Nigerian economy and their importance in the web of life, fish was chosen as a logical organism for monitoring lindane residues.

Reports show that the use of pesticides requires close monitoring of the effects on non-target organisms especially on the economically important fishery resources in rivers close to farm lands (6). In developed countries in the temperate regions there is a continuous effort to monitor pesticide residues in air, water, soil, man, wildlife and fish. On the other hand, the effects of pesticide residues on non-target organisms in Nigerian agriculture and its environment is still speculative because of a lack of quantitative data. The purpose of the present study was to monitor certain rivers in Anambra State to evaluate the level of lindane contamination in fish.

2. EXPERIMENTAL

2.1. Sample collection and storage

In 1991, fish samples of different species were collected from different rivers close to rice fields that were suspected to have received lindane applications (Table 1). Fish from locations thought to be untreated were also collected. The fish were collected from the river

with a cast net and transported to the laboratory in ice buckets where they were sorted out into species, wrapped in aluminum foil and frozen until analysed.

2.2. Sample preparation, extraction and clean-up

Each frozen fish sample was ground and homogenized in a meat grinder. Samples (20 g) of the homogenized fish were blended and mixed with anhydrous sodium sulphate (80 g). The pesticide was extracted for 1 hour in a screw-capped jar with acetone + hexane (1+1 by volume, 200 mL). After tumbling, the solids were allowed to settle and the solvent extract filtered into a separating funnel containing sodium sulphate solution (20 gL⁻¹, 200 mL) and shaken for about one minute. The lower aqueous layer was discarded and the hexane extract washed twice more with sodium sulphate solution (20 gL⁻¹, 50 mL). Quadruple hexane extracts were combined and evaporated to about 10mL and then

Table 1. Residues of lindane found in fish caught at the sampling stations in Anambra State, Nigeria

River	Location	Fish species	Lindane residues, mgkg ⁻¹
Ani	Ufuma ^{a)}	<i>Heterotis niloticus</i> ^{d)}	ND ⁱ⁾
		<i>Clarias zariepinus</i> ^{e)}	ND
		<i>Heterobranchus bidorsalis</i> ^{f)}	ND
		<i>Oreochromia niloticus</i> ^{g)}	ND
Ozzi	Enuguabo ^{a)}	<i>Heterotis niloticus</i>	ND
		<i>Clarias zariepinus</i>	ND
		<i>Heterobranchus bidorsalis</i>	ND
		<i>Oreochromia niloticus</i>	ND
Ozalla	Oyi ^{b)}	<i>Heterotis niloticus</i>	<1 ^{j)}
		<i>Clarias zariepinus</i>	<1
		<i>Heterobranchus bidorsalis</i>	<1
Omoh	Oyi ^{b)}	<i>Heterotis niloticus</i>	<1
		<i>Clarias zariepinus</i>	<1
		<i>Oreochromia niloticus</i>	<1
Anambra	Oyi ^{b)}	<i>Heterotis niloticus</i>	3.1
		<i>Clarias zariepinus</i>	2.1
		<i>Heterobranchus bidorsalis</i>	3.1
		<i>Oreochromia niloticus</i>	2.2
Nkissi	Onitsha ^{c)}	<i>Heterotis niloticus</i>	3.4
		<i>Cyprinus carpio</i> ^{h)}	1.9
		<i>Heterobranchus niloticus</i>	2.0

Local Government area :

a) Orumba North

b) Oyi

c) Onitsha

i) none detectable

j) detectable, but not quantifiable

Local name : d) African boy tongue (Okpo)

e) Mudfish

f) Catfish (Azu isi)

g) Tilapia (Ikpokpo)

h) Carp

transferred into vials and evaporated using a N₂ sparge. The residue was redissolved in hexane (2mL). This extract was passed through a column containing Florisil to remove non-pesticidal extracts by the procedure reported previously [7].

2.3. Gas chromatographic analysis

The hexane extracts were analysed by gas chromatography at the Department of Food Technology and Science, University of Tennessee, Knoxville, USA using a Shimadzu 9 AM; column: 30 m x 0.25 mm i.d. SPB-5 fused silica column (Supelco Inc. Belafonte, PA); temperatures : column, 70°C for 1.5min, increased to 200°C at 10°C min⁻¹ and then to 240°C for 20 min; injector, 250°C; detector, 250°C; carrier gas, helium; flowrate, 2 mL min⁻¹; detector, ⁶³Ni electron capture interfaced with a Shimadzu QP 1000 mass spectrometer. At the beginning of the analysis each day, several consecutive injections were done to prime the column [8].

3. RESULTS AND DISCUSSION

3.1. Questionnaire survey of lindane use in Anambra State

The preliminary questionnaire survey involved extensive visits and interviews with officials of the State Ministry of Agricultural Development Agencies, Pest Control Department, Office of Statistics as well as on-the-spot interviews of rice farmers in various locations in Anambra State, Nigeria. The survey revealed that there was no information on the occurrence, distribution, present usage, effects and the impact of lindane in the rice agroecosystem in Nigeria. It was observed that lindane, commonly called gammalin, is still sold in local markets even though the Federal Government of Nigeria has banned its use. Farmers still use it for pest control in various crops and also for killing fish. In some areas the Government ban is strictly observed while, in most other areas, farmers use it for pest control because it is effective and relatively cheap compared with other pesticides. The unavailability of authentic data on the usage and distribution of lindane in Nigeria as evidenced by our survey, justified the collection of fish from various rivers close to rice growing areas to begin the collection of baseline data of lindane residue levels in fish.

3.2. Lindane residues in fish

Comparison of lindane residues in fish from different rivers show that Anambra and Nkissi rivers were more exposed to pesticide contamination than the other rivers (Table 1) with the residues in fish ranging from none detectable to 3.4 mg kg⁻¹. Only small amounts of lindane were detected in fish from the Omoh and Ozalla rivers and no detectable residues in the fish from the Ani and Ozzi rivers. The absence of lindane residues in fish collected from the Ani and Ozzi rivers reflects the strong restrictions on the use of lindane for farming and fishing in these areas, while the low level of lindane found in fish from the Omoh and Ozalla rivers indicate that only small amounts of lindane, that is applied to the rice in this area, entered the rivers. These low levels of lindane are comparable to those found in areas subjected only to airborne contamination [9]. Such levels may not be harmless since it has been reported that DDT at low levels can interfere with the reproduction of certain fish species [2].

Fish obtained from Anambra and Nkissi river contained appreciable amounts of lindane probably because the river flowed through areas where lindane was frequently used

by farmers. Reports show that under certain conditions pesticide residues arising from normal agricultural use may contribute to diffuse (non-point) pollution of the aquatic environment [11]. In an experiment in Zimbabwe, all the fish sampled contained insecticide residues ranging from 0.055mg kg⁻¹ in the liver of living species up to 5.1mg kg⁻¹ in the liver of dead specimens, the latter presumably being the cause of death [12]. Individual fish vary in residue levels and this can be attributed to differences in fish movement, size, food habits and fat content [13]. Other field and laboratory studies showing vast differences in residue levels in fish have also been reported in Wisconsin [14]. Although a very limited number of fish samples were collected in this study, when present the levels of lindane were similar to those found in fish in the midwest of the USA, where total pesticide residue of 2.10 µgg⁻¹ were recorded [15] but higher than those in Utah where only 0.05-0.096 µgg⁻¹ of organochlorine residue was detected [16]. Because some fish continue to accumulate residues over a period of years, the levels observed in the fish from the Nkissi and Anambra rivers reflect the integrated history of exposure. Interviews with villagers indicate a direct contamination as the farmers in the Oyi Local Government areas, where Anambra and Nkissi rivers flow through, reported non-restricted use of the lindane in agriculture and in fish farming.

4. CONCLUSION

In the rivers studied, the extent of contamination of fish by lindane varied. Lindane which was found in fish collected from the Anambra, Nkissi and Ozalla rivers that may have originated from run-off from the rice fields or from direct contamination of the rivers by fishermen. The considerable variation in lindane residue levels in different fish samples, could be caused by variation of fish movement, food habits, species size, age and fat content. Because of these variations, there should be caution in interpretation and application of these data.

ACKNOWLEDGEMENTS

We wish to thank the International Atomic Energy Agency, Vienna and Swedish International Development Agency for providing the fund to support this study under the Coordinated Research Project on Adverse Effect on Flora and Fauna from the use of Organochlorine Pesticides in African Continent, IAEA Research Contract No. 6585 SD.

REFERENCES

- [1] INGLIS, A., HENDERSON, C., JOHNSON, W.L., Expanded program for pesticide monitoring of fish, *Monitoring Journal* **5** (1) (1971) 47-49.
- [2] JOHNSON, D.W., Pesticides in fish: a review of selected literature, *Trans. Amer. Fish. Soc.* **97** (1968) 398-424.
- [3] EDWARDS, C.A., *Environmental pollution by pesticides*. Plenum Press, London. (1973).
- [4] AN de WAERDT, J., *Drins and related insecticides: toxicology, use and alternatives with special reference to Latin America and South East Asia*, *Econoscript* **23** (1983) 1-176.
- [5] KLASSES, H.E., KADDOUM, A.M., Pesticide residues in natural fish population of the Smoky River of Western Kansas, 1967-69. *Pesticide Monit. J.* **7** (1973) 53-61.

- [6] ANTWI, L.A.E., "Studies on the effect of some insecticides on the brain acetylcholinesterase activity of *Tilapia zilli* in two treated tropical rivers", Pesticides, Food and Environmental Implications. IAEA, Vienna 247-256.
- [7] MELTON, S.L., "Pesticide report on Bush Brothers and Company, Vegetables, October 1, 1990 - October 30, 1991", Chemical fall out 2nd Edition (Muller, N.W., Berg, G.G., Eds) Thomas, C.C. Illinois USA (1991) 119.
- [8] WANDIGA, S.O., MUTERE, A., Determination of γ -BHC in breast milk of Kenyan women., Bull. Chem. Soc. Ethiop. 2 (1988) 39-44.
- [9] SODERGREN, A., Transport, distribution and degradation of organochlorine residue in a south Swedish lake ecosystem, Vatten (1973) 108-290.
- [10] SODERGREN, A., DSIRARAI, R., CHARIBZADEU N., MOIMPOUR, A., Organochlorine residues in aquatic environments in Iran, Pest Moni J. 12 (1978) 81- and 87.
- [11] KREUGER, J.K., BRINK, N., Losses of pesticides from agriculture, Pesticides: Food Environmental Implications IAEA, Vienna (1988) 101-112.
- [12] MULLER, P., Effects of pesticides on flora and fauna, Pesticides: Food and Environmental Implications. IAEA, Vienna (1988) 11-27.
- [13] HENDERSON, C., JOHNSON, W.L., INGLIS, A., Organochlorine insecticide residues in fish (National Pesticides Monitoring Program). Pest. Mon. J. 3 (1969) 145-171.
- [14] KLWINWER, A.J. DEGURSE, P.E., WIRTH, T.L., Occurrence and significance of DDT and dieldrin residues in Wisconsin fish, Tech Bull (1968) 41- 43. Dept of Natural Resources, Madison, Wisconsin.
- [15] HENDERSON, C., INGLIS, A., JOHNSON W.L., Organochlorine insecticide residue in fish. National Pesticide Monitoring Program, Pesticides Mon. J. 5 (1971).
- [16] HAW, J.S., LOVERRIDGE, E.L. RIPPEE, K.P., PETERSON, J.L., WHITE, D.A., BARTON, J.R., FUHRIMAN, D.K., Seasonal variations in residues of chlorinated hydrocarbon pesticides in the water of the Utah lake drainage system, 1970-1971. Pest Mon. J. 6 (1972) 167-170.

**NEXT PAGE(S)
left BLANK**



ADVERSE EFFECTS ON FLORA AND FAUNA FROM THE USE OF ORGANOCHLORINE PESTICIDES ON THE AFRICAN CONTINENT: THE NIGERIAN CASE

E.D.N. UMEH

Enugu State University of Science and Technology, Enugu

R.I. EGWUATU, E.N. NWAKEZE

Nnamdi Azikiwe University, Awka

P.E. AMULU

Institute of Management and Technology, Enugu

Nigeria

Abstract

Lindane was found to reduce stem borer damage significantly in both rice and maize plots in most years. However, a significant difference in yield was recorded only for maize in 1993 and 1995. There was a slight but steady decrease in the yield of maize over the years in the lindane treated plots although yields in these plots remained higher than in the control plots. Significant differences in the mean dry weight of retrieved leaf discs were recorded in the maize plots in 1994 and 1995. Neither insects nor spiders, were caught from plants in the treatment plots 1 day after lindane application, although almost equal numbers of insects and spiders were recovered from D-vac ground samples of treatment and control plots on the same day. Data from pitfall traps showed that lindane significantly affected the population of *Aranae*, *Formicidae*, *Collembola* and *Acarina* for up to 6 weeks after application. Its effect on *Carabidae* was inconsistent.

1. INTRODUCTION

Organochlorine pesticides are still widely used in Africa for agricultural and public health purposes. Their cost effectiveness and broad spectrum of activity have continued to encourage their demand and use, in spite of international pressures calling for their ban. Presently, there is no pesticide legislation in Nigeria and perhaps also, in some of the African countries resulting in unrestricted importation and use of various kinds of pesticides including organochlorines. Many organochlorine pesticides are persistent in the environment for a long time. They are also known to leave residues in the soil [1,2] resulting in the disruption of soil fauna and flora [3] in the temperate regions, and thus affecting organic matter breakdown and nutrient cycling [4,5]. Treatment of a cultivated forest soil in Nigeria with DDT resulted in a steady decline in yield of cowpea over a four year period [6] and affected activity of certain crickets, lycosid spiders and millipedes [7].

The objectives of this investigation are to determine the effects of lindane (γ -1,2,3,4,5,6 - hexachlorohexane) on:

- (i) the key pests of maize and upland rice, i.e stem borers,
- (ii) predatory and other non-target species in maize and upland rice agro-ecosystems,
- (iii) soil biology and biological activity.

2. MATERIALS AND METHODS

The experiments were carried out from 1992 to 1995 in maize (*Zea mays*) plots, and in 1992 and 1993 in rice (*Oryza sativa*) plots. In 1992, the maize and rice plots were located at Ugwuoba, Enugu State, and Igbariam, Anambra State, respectively, the two towns being about 20 km from each other. Both towns are located in the derived rain forest zone of

Nigeria. In 1993 to 1995, the maize plots were located in Enugu, Enugu State and is at the northern fringe of the derived rain forest. These towns fall within the zone whose annual rainfall does not fall below 1,500 mm and temperature ranges between 27 and 32°C during most of the year.

2.1. Experimental plots

The plots used in Enugu and Ugwuoba were under fallow for 1 and 2 years, respectively, before the experiments began in 1993 and 1992. The plots used in Igbariam were cropped to yam (*Dioscorea* sp.), maize, egusi melon (*Colocynthis vulgaris*), cassava (*Manihot esculenta*), and various leafy vegetables in a mixed farming system. These plots had no previous history of pesticide use.

At the beginning of each experimental season, all the vegetation was ploughed into the soil which was later harrowed twice before mapping out the plots. Soil samples were taken in the three locations and found to be clayey (48-57%), acidic (pH ranges from 6.2 to 6.8), with high organic matter content (Ugwuoba: 8.0%; Igbariam: 3.8% and Enugu: 2.1%).

In 1992 (for maize and rice) and 1993 (for rice), the plot size was 50 m x 50 m each, for treatment and controls, and separated by a 3-metre row of grass. From 1993, the plot sizes for maize were 32 m x 80 m for both treatment and controls, separated by a 1-metre row of grass. In these years (1992 and 1993) there were 4 sub-plots (5 m x 5 m) located within each of the two plots 10 m away from the edge of the plots, and used for sampling. In 1994 and 1995, there was randomised block design with 4 blocks and two treatments. Within each plot was a sub-plot (10 m x 10 m). Because of the possible interference of the various sampling activities within the sub-plots with yield, a yield area plot (2 m x 2 m) was located outside but close to each sub-plot.

2.2. Planting and fertiliser application

The maize variety used throughout these experiments was TZSR yellow while the rice varieties used were FARO 43 (1992), and ITA 306 (1993). Maize was planted in rows, 1 m apart and 50 cm between hills at 2 seeds/hill in 1992. In the subsequent years, the inter-row distance was reduced to 75 cm. Rice was planted in rows by dibbling seeds, 15 cm apart both between rows and hills at the rate of 3-4 seeds/hill. A compound fertiliser (NPK 15.15.15) was applied to rice and maize soils at the rate of 200 kg ha⁻¹, 21 days after planting. A second fertiliser application of urea was made at the tasselling stage (for maize) and the booting stage (for rice), at the rate of 50 kg ha⁻¹. Weeding was done twice manually during each experimental season.

2.3. Pesticide applications

Lindane (200 g L⁻¹ EC) was applied to the treatment plots in rice (1992 and 1993) and maize (1992) in a single dose of 1 kg AI ha⁻¹. From 1993 to 1995, 1 kg AI ha⁻¹ lindane was applied to maize in split doses of 0.5kg AI ha⁻¹. The first lindane application corresponded to the 3-leaf stage in the maize development (stage 13 in the decimal code [8]). The second lindane application was made two weeks later. High volume sprays (120-200 L of water ha⁻¹), using a knapsack sprayer(CP15) with a red nozzle were used in the lindane application. There was no lindane application to the control plots.

2.4. Sampling

Pre-treatment samples of stem borer damaged plants, insects, other arthropods, and soil fauna were taken from the sub-plots a day before lindane application. Thereafter, samples were taken at two weekly intervals for 2 months. In 1994 and 1995, insect and other arthropod samples were also taken a day, and up to week 10 after lindane application.

2.4.1. Plant damage by stem borers

Stem borer damage was assessed by counting all the affected plants in the maize sub-plots. Because of the high plant population in the rice sub-plots, only half the plants were randomly sampled for stem borer damage. At the early stages, window panes and dead hearts were used as indices for stem borer damage. At the later stages, holes in the stem, lodging, and white heads (dried up tassels and panicles) arising from stem borer damage were also used in assessing plant damage. In 1993, the populations of stem borer larvae was estimated 4 weeks after the last lindane application by dissecting 50 stalks, randomly selected outside the sub-plots, and counting all the larvae and tunnels found. Cob damage was also estimated by counting all the damaged cobs in the yield area plots.

2.4.2. Sweep netting, D-vac sampling and pitfall traps for arthropods

Insect, spider and mite populations were sampled using sweep net and pitfall traps in 1992 and 1993. In 1994 and 1995, sweep net sampling was replaced with use of a D-vac sampler which was used to sample both plants and ground. Sweep netting was carried out along two diagonal lines in each sub-plot at the rate of 5 sweeps/diagonal line. After 5 sweeps, the collections were removed and dropped into 70% alcohol until counted. Collections from the sub-plots were stored separately.

D-vac samples for plants and soil surface were taken at the rate of 4 samples per sub-plot. Each sample consisted of 5 sub-samples. For the ground samples, each sub-sample consisted of a 5 second suck with the sampling cone lightly pressed to the ground. For the plant samples, each hill containing 1 or 2 plants, chosen at random, represented one sub-sample. At the earlier stages, the D-vac cone was placed over the plants while sampling. Later, as the plants became taller, the cone was used to sweep the plants from all sides. D-vac sampling of plants was carried out up to 6 week after lindane application while ground sampling continued up to the 10th week.

Four pitfall traps were set up at random, in each sub-plot. The plastic containers used (10 cm in diameter and 10 cm deep) were buried so that their rims were flush with the soil surface. The traps were half filled with soapy water. In 1992, 1993 and 1994, and 1995, the traps were left in the field for 1 day, 2 days and 1 week, respectively, before collection. In 1995, the traps were emptied twice during the sampling week.

In 1992, the target taxa included numbers of adult stem borers, *Carabidae*, *Staphylinidae*, *Coccinellidae*, *Chrysopidae*, *Formicidae*, *Dermaptera* and *Aranae*. In 1993, stem borers, *Carabidae* and *Aranae* were sampled, while in 1994 and 1995, stem borers, *Carabidae*, *Coccinellidae*, *Formicidae*, *Aranae*, *Acarina* and *Collembola* were sampled.

2.4.3. Breakdown of organic matter

Fifty leaf discs (each measuring 25 mm in diameter), cut from mango leaves (*Mangifera indica*), were put in each 12 cm² litter bag with 10 mm apertures. The litter bag was attached to a marker peg and buried to a depth of 50 mm. Four litter bags were buried randomly in each sub-plot prior to lindane application. The litter bags were retrieved three months later and the contents of each litter bag were washed separately, oven dried at 60°C for 24 hrs, and weighed.

2.4.4. Soil fauna

Four soil cores, 150 mm deep and 50 mm in diameter were taken at random from each sub-plot before pesticide application in 1992 and 1993. Each soil core was cut into 6 pieces and the soil fauna present were extracted using Tullgren type extractors. A second set of soil core samples were taken 2 months after lindane application and treated similarly.

3. ANALYSIS OF DATA

The various data obtained were analysed using a two-way analysis of variance incorporating effects of treatment and time. The sub-plots in 1992 and 1993 were regarded as real plots and the data generated were analysed similarly.

4. RESULTS

4.1 Maize 1992

There was heavy rainfall (31.6 mm), which lasted several hours on June 9, 1 hr after lindane application and this seemed to have affected the experiment. The important stem borer species of maize encountered in this work were *Busseola fusca* Fuller, *Sesamia calamistis* Hamp, and *Eldana saccharina* Wlk. Although damage caused by these stem borers was low in 1992, there was an increase in their damage with time in both treatment and control plots (Fig. 1) but the differences were only significant ($P < 0.05$) at one interval (6 weeks after treatment).

A few insects and spiders were caught with the sweep net (Table I). These included specimens from *Coccinellidae*, *Formicidae* and *Aranae*. The *Coccinellids* caught were the phytophagous species, *Epilachna* sp., known to scrape leaf tissues from young maize plants. More insects and spiders were caught in the control than in the treatment plots but their differences were not statistically significant ($P < 0.05$).

Pitfall trap collections included specimens from *Carabidae*, *Staphylinidae*, *Formicidae* and *Aranae* (Table II and Fig. 2). For *Carabidae* and *Staphylinidae* (Table II), there was no significant difference in the mean catches made on any of sampling dates in both treatment and control plots at a 5% level. For *Formicidae* (Fig. 2), only the second post-treatment sample, with mean catches of 3.5 and 8.0 for treatment and control plots, respectively, was found to be statistically significant. Analysis of the mean spider catches showed that significantly more spiders were trapped in the control than in the treatment plots in the 4th, 6th and 10th week after treatment. These results (*Formicidae* and *Aranae*), do not show any clear trends in their response to lindane application. *Chrysopidae* was caught neither in sweep nets




FIG. 1. Stem borer damage in maize and rice plots.




FIG. 1. (cont.).

nor trapped in pitfalls in this experiment. Overall, it would appear that the heavy rain which occurred soon after lindane application may have affected some of these results.

The soil fauna in the samples included *Collembola* (springtails), *Acarina* (mites) and other soil mesofauna (Table IV). Although the pre-treatment samples showed no significant differences in the mean numbers of the various soil fauna extracted from both treatment and

control plots, post-treatment samples showed significant differences in the mean numbers of *Collembola* and *Acarina* extracted from the lindane-treated compared with the control plots. Both *Collembola* and *Acarina* are micro-arthropods inhabiting the soil and leaf litter. They would have received lindane sprays directly and may have been killed before commencement of the rain on the treatment day. Their population seemed unable to recover from the impact of lindane, even at two months after application.

Mean dry weight of leaf discs retrieved (Table V) were 4.7g and 3.8g for treatment and control plots, respectively, and this difference was not statistically significant. Yields obtained from the lindane treated plot (5.12 t ha⁻¹) was not significantly different (P <0.05) from the yields obtained from control plots (4.99 t ha⁻¹). This result may be due partly to low pest infestation on maize this year but also due to the effect of rainfall which may have leached or washed away most of the lindane deposits on the treatment day.

Table I. Sweep net collections in maize and rice

a) Maize

Weeks after treatment	Mean number of specimens caught ±SE							
	<i>Coccinellidae</i>		<i>Aranae</i>		<i>Formicidae</i>		Stem borers	
	T	C	T	C	T	C	T	C
Pretreatment	0.00	0.00	0.00	0.00	0.00	0.00	-	-
2	0.75 ±0.5	1.50 ±1.3	0.50 ±0.6	0.50 ±0.6	0.50 ±0.5	1.25 ±0.5		
4	0.00	0.25 ±0.5	0.00	0.00	0.00	0.00		
6	0.00	0.50 ±0.7	0.00	0.00	0.25 ±0.5	0.75 ±0.5		
8	0.25 ±0.5	0.75 ±0.5	0.00	0.00	0.00	0.00		
10	0.75 ±0.5	0.75 ±0.5	0.25 ±0.5	0.25 ±0.5	6.75 ±1.0	6.50 ±1.3		

T = treated plot, C = untreated control plot

b) Rice

Weeks after treatment	Mean number of specimens caught ±SE							
	<i>Coccinellidae</i>		<i>Aranae</i>		<i>Formicidae</i>		Stem borers	
	T	C	T	C	T	C	T	C
Pretreatment	0.25 ±0.5	1.25 ±0.5	0.00	0.25 ±0.5	1.25 ±0.5	1.25 ±0.5	1.50 ±0.6	1.25 ±0.5
2	0.25 ±0.5	1.00 ±0.0	0.00	0.00	0.25 ±0.5	4.25 ±1.3*	1.00 ±0.8	6.25 ±1.7*
4	0.00	0.50 ±0.5	0.00	0.75 ±0.5	2.25 ±1.3	0.00	0.75 ±1.0	3.00 ±1.4*
6	0.50 ±0.6	0.50 ±0.6	0.25 ±0.5	0.75 ±0.5	1.25 ±0.5	3.00 ±0.8*	1.00 ±0.8	0.75 ±0.5
8	0.25 ±0.5	0.50 ±0.5	0.50 ±0.6	1.00 ±0.8	1.00 ±0.0	1.00 ±0.0	0.25 ±0.5	0.25 ±0.5
10	0.25 ±0.5	0.25 ±0.5	0.50 ±0.6	1.50 ±0.6	0.50 ±0.6	0.25 ±0.5	0.25 ±0.5	0.50 ±0.6

T = treated plot, C = untreated control plot

* significantly different from the treated plot at P = 0.05

Table II. Pitfall trap collections from the maize trials

Year	Weeks after treatment	Mean number of specimens caught \pm SE			
		<i>Carabidae</i>		<i>Staphylinidae</i>	
		T	C	T	C
1992	pretreatment	2.50 \pm 0.6	2.25 \pm 1.0	1.75 \pm 0.5	2.25 \pm 0.9
	2	1.25 \pm 1.0	2.25 \pm 0.5	0.00	1.00 \pm 0.8
	4	1.25 \pm 0.5	2.75 \pm 0.5	0.00	0.75 \pm 0.8
	6	1.25 \pm 0.5	1.25 \pm 1.0	0.50 \pm 0.6	1.00 \pm 0.8
	8	0.25 \pm 0.5	0.50 \pm 0.6	0.25 \pm 0.5	0.25 \pm 0.5
	10	2.00 \pm 1.1	2.00 \pm 1.2	0.50 \pm 0.6	0.75 \pm 1.0
1993	pretreatment	0.75 \pm 0.5	0.75 \pm 1.0		
	2	0.25 \pm 0.5	1.75 \pm 1.0		
	4	0.25 \pm 0.5	2.00 \pm 1.0		
	6	0.50 \pm 1.0	2.25 \pm 0.5		
	8	1.00 \pm 0.5	2.50 \pm 0.6		
	10				
1994	pretreatment	1.25 \pm 1.3	1.50 \pm 0.6		
	1 day	1.75 \pm 0.5	1.75 \pm 1.0		
	2	2.25 \pm 1.3	3.25 \pm 1.3		
	4	1.50 \pm 0.6	2.25 \pm 0.5		
	6	1.25 \pm 0.5	1.25 \pm 0.5		
	8	1.75 \pm 0.5	1.50 \pm 0.6		
1995	pretreatment	7.25 \pm 5.3	7.5 \pm 5.1		
	1 day	0.00	8.25 \pm 1.5		
	2	1.00 \pm 0.8	12.75 \pm 3.8		
	4	2.75 \pm 1.0	15.00 \pm 2.5		
	6	7.00 \pm 2.7	17.00 \pm 1.7		
	8	11.00 \pm 2.6	12.25 \pm 2.2		
10	7.75 \pm 3.9	10.25 \pm 2.9			

T = treated plot, C = untreated control plot


Table III. Pitfall trap collections from the rice trials

Year	Weeks after treatment	Mean number of specimens caught \pm SE			
		<i>Carabidae</i>		<i>Staphylinidae</i>	
		T	C	T	C
1992	pretreatment	1.50 \pm 0.6	1.50 \pm 0.6	0.75 \pm 0.5	0.50 \pm 0.6
	2	1.00 \pm 0.0	1.75 \pm 0.5	0.00	0.25 \pm 0.5
	4	2.50 \pm 0.6	1.75 \pm 1.0	0.00	0.75 \pm 1.0
	6	2.00 \pm 0.8	2.75 \pm 1.0	0.50 \pm 0.5	1.50 \pm 0.6
	8	3.75 \pm 1.0	5.00 \pm 0.8	0.25 \pm 0.5	1.25 \pm 0.5
	10	4.50 \pm 1.3	6.75 \pm 3.1	2.25 \pm 0.5	2.00 \pm 0.8
1993	pretreatment	4.00 \pm 1.6	3.50 \pm 1.3		
	2	1.50 \pm 1.0	4.50 \pm 1.3*		
	4	2.50 \pm 1.0	4.75 \pm 1.5*		
	6	4.00 \pm 0.8	4.75 \pm 1.3		
	8	3.25 \pm 0.5	3.00 \pm 0.8		
	10				


T = treated plot, C = untreated control plot

* significantly different from the treated plot at P = 0.05

1992 spiders



1994 spiders



1995 spiders






FIG. 2. Pitfall trap collections in maize plots.

1992, ants



1994 ants



1995 ants




FIG. 2. (cont.)






FIG. 2. (cont.)

4.2. Rice 1992

The major stem borers of upland rice are *Diopsis macrophthalma* Dalm., *Chilo zacconius* Bleszynsky, *S. calamistis* and *Scirpophaga* sp. During the vegetative stage of rice, plant damage by stem borers showed up as dead hearts while damage during the reproductive phase resulted in white heads. There was no significant difference in the mean number of dead hearts obtained in the pre-treatment, and in the week 2 samples (Fig. 1). Week 2 sample results may have been because damage attributable to the larvae arising from eggs laid in control subplots between lindane application in the treatment plots and the week 2 sample may not have been manifested. It takes about 12-20 days from egg deposition to appearance of dead hearts. Subsequent samples showed significant differences in the mean number of damaged plants recorded in the treatment compared to the control plots at 5% level.

1992 ants



1992 spider




FIG. 2. (cont.)

Table IV. Fauna found in the soil cores taken from the maize and rice fields

Year	Crop	Species	Mean number of species found \pm SE			
			pretreatment		posttreatment	
			T	C	T	C
1992	maize	Collembola	150.50 \pm 16.7	151.00 \pm 8.7	69.00 \pm 6.5	155.00 \pm 19.6*
		Acarina	16.50 \pm 5.2	16.75 \pm 2.5	10.25 \pm 1.9	11.75 \pm 1.7*
		others	9.75 \pm 3.1	10.50 \pm 3.9	9.75 \pm 3.1	11.00 \pm 3.4
1992	rice	Collembola	74.25 \pm 14.1	60.00 \pm 15.2	27.00 \pm 4.2	80.75 \pm 19.9*
		Acarina	10.00 \pm 3.2	11.00 \pm 1.4	4.75 \pm 3.3	13.50 \pm 6.6
		others	7.75 \pm 4.0	8.75 \pm 1.7	3.00 \pm 1.6	12.00 \pm 2.2*
1993	maize	Collembola	10.00 \pm 3.7	9.25 \pm 1.7	8.00 \pm 2.4	7.25 \pm 1.7
		Acarina	5.50 \pm 1.7	3.75 \pm 1.0	5.00 \pm 1.8	4.50 \pm 1.0
1993	rice	Collembola	52.50 \pm 5.1	47.25 \pm 3.3*	43.00 \pm 7.1	51.25 \pm 5.5*
		Acarina	12.75 \pm 1.7	13.50 \pm 1.7	18.25 \pm 4.1	27.00 \pm 3.7*

T = treated plot, C = untreated control plot

* significantly different from the treated plot at P = 0.05

Table V. Retrieval after 3 months of litter buried in the control and treated plots of maize and rice

Crop	Year	Mean retrieved weight, g \pm SE	
		Treated	Control
maize	1992	4.70 \pm 0.5	3.76 \pm 0.6
	1993	5.42 \pm 2.4	4.64 \pm 0.8
	1994	1.33 \pm 0.3	0.67 \pm 0.2*
	1995	2.40 \pm 0.0	1.71 \pm 0.2*
rice	1992	5.49 \pm 1.7	4.46 \pm 1.0
	1993	5.93 \pm 2.1	3.81 \pm 1.4*

* significantly different from the treated plot at P = 0.05

The rice plots had a richer faunal composition than the maize plots. Adult stem borers, *Coccinellidae*, *Formicidae* and *Aranae* were the main groups caught with the sweep net during the vegetative phase. At this stage, most coccinellids caught were the phytophagous species. In the later stages of rice development, a few *Carabidae*, *Dermaptera* and predaceous *Coccinellidae* were also caught in the sweep net. Statistical analyses of sweep net catches of adult stem borers showed significant differences in the mean catches from the lindane treated and the control plots during week 2 and week 4 sampling (Table I). Lindane affected stem borers of rice for up to one month after application. For all the other groups of insects and spiders, except the *Formicidae*, lindane application had no effect. The effects of lindane on *Formicidae* appeared inconsistent.

Pitfall trap collections (Table III and Fig. 2) showed that the important families trapped were *Carabidae*, *Staphylinidae*, *Formicidae* and *Aranae*. Stem borers, particularly, *D. macrophthalma* were also trapped in the pitfalls. In terms of numbers of arthropods caught in both pitfall traps and sweep nets, *D. macrophthalma* was the dominant stem borer species. Carabid beetles were trapped throughout the sample periods but no significant difference was found in their mean numbers in the lindane treated and the control plots. Similarly *Staphylinidae*, though caught on fewer occasions and in lower numbers than *Carabidae*, was also found not to be statistically significant in treatment and control plots at the 5% level of probability.

For *Formicidae* and *Aranae* (Fig. 2) significant differences in their means were recorded in treatment and control plots in all post treatment samples. Thus lindane remained effective against ants and spiders in the rice plots for more than two months after its application.

Soil fauna data (Table IV) showed a mean of 81 *Collembola* in the control plots against 27 in the lindane treated plots. These data were found to be statistically significant at the 5% level of probability. Mite populations, as well as rates of leaf disc disintegration, were unaffected by lindane in the rice plots. The difference in rice yield from treatment (2.48 t ha⁻¹) and control (2.10 t ha⁻¹) plots was not statistically significant.

4.3. Maize 1993

Stem borer damage increased steadily with time and ranged from 1.90% in the pre-treatment to 14.90% in the post-treatment control plots (Fig. 1). In lindane treated plots, plant

damage ranged from 1.35% before treatment to 3.53% after treatment. These differences between treated and control were statistically significant at a 5% level. Thus, the split dose application of 1 kg AI ha⁻¹ of lindane provided protection to the plants against stem borer attack for up to 6 weeks. Larval populations determined by stem dissection showed that more larvae ($X = 17$) were recovered from the control plots than from the lindane treated plots ($X = 4$) and these were found to be statistically significant at the 5% level. Similarly stem borer damage (stem tunnelling), also determined by stem dissection, showed that a mean of 27 and 18 stems were damaged in the control and the lindane treated plots, respectively, and these were found to be statistically significant. On the other hand, more cobs were damaged in treatment plots ($X = 12$) than in control plots ($X = 6.3$) and these data were also statistically significant. None of the target families were caught in the sweep net in 1993.

Generally, pitfall trap catches were lower in 1993. There was no significant difference in the mean numbers of *Carabidae* and *Aranae* caught in the pre-treatment samples. Post-treatment samples of *Carabidae* showed inconsistent trends thus making interpretation difficult. Except the first post treatment sample, lindane application affected spider catches significantly in the treatment plots throughout the sampling period.

Mean dry weights of leaf discs (Table V) retrieved from the lindane treated plots (5.42 g) were not statistically significant from the dry weight of leaf discs retrieved from control plots (4.64 g) at a 5% level. *Collembola* and *Acarina* populations extracted from the treatment and control soil cores two months after lindane application were not significantly different.

Maize yield was found to be significantly higher (5.03 t ha⁻¹) in the treatment than in the control plot (3.61 t ha⁻¹).

4.4. Rice 1993

The major stem borer species monitored was *D. macrophthalma*. However, stem borer damage reported here resulted from the combined effects of *D. macrophthalma* and other stem borers. Plant damage was found to be significantly lower in the treatment than the control plots, in all but the last two post-treatment samples. Sweep net samples showed low populations of *D. macrophthalma* in the treatment plots at 2 and 4 week after lindane application. Differences in the mean numbers of *D. macrophthalma* found in the samples from both the treated and the control plots were not statistically significant throughout.

Pitfall trap collections of *Carabidae* showed that the lindane application reduced carabid populations significantly, for up to 4 weeks after application. Post-treatment soil fauna samples showed that significantly fewer *Collembola* and *Acarina* were extracted from the lindane treated plots, and the mean dry weight of leaf discs retrieved was also significant ($P < 0.05$) in treatment compared with control plots. The yield obtained from treatment plots (2.45 t ha⁻¹) was not statistically different from that obtained from control plots (2.08 t ha⁻¹).

4.5. Maize 1994

Stem borer damage increased steadily with time although damage was lower in treatment than control plots. Damage caused by stem borers was significantly less in the lindane-treated compared with the control plots during the first 6 weeks after lindane application (Fig. 1). Another type of plant damage which became important this year was that

caused by the coccinellid, *Epilachna* sp. which feeds on the leaf tissues of young maize plants, leaving large strips of papery epidermis. Although this damage was not quantified, visual observation showed far greater damage in control than treatment plots.

D-vac samples of plants showed that *Coccinellidae*, *Formicidae* and *Aranae* were the main families caught. Pre-treatment samples showed that in these 3 families, there were no significant differences in the mean numbers caught in the lindane-treated and the control plots. Samples taken 1 d after lindane application showed that no insects or spiders were caught on plants in the lindane treated plots. Subsequent samples also showed significant differences in the numbers of insects and spiders caught in treatment compared with control plots for between 1 and 2 months after treatment (Table VI).

More faunal specimens were collected in D-vac ground samples than from plants. Specimens collected included those of *Collembola*, *Acarina*, *Carabidae*, *Formicidae* and *Aranae*. The mean number of insects, spiders, and mites collected in the pre-treatment and 1d post-treatment samples were statistically similar in all plots. Similarity in these results occurred because specimens killed by lindane the previous day were also picked up by the D-vac. Subsequent soil surface samples showed that significantly fewer *Collembola* and *Acarina* were caught in the treatment than control plots (Table VII) at the 5% level, throughout the sampling period. Carabid catches in D-vac samples were low and not statistically significant between the treatment and the control plots. Lindane application caused significantly fewer spiders and ants in the lindane treated than the control plots during the first 6 weeks of the initial lindane application.

Pitfall trap collection showed that lindane affected *Collembola*, *Aranae* and *Formicidae* populations adversely (Fig. 2). Spider and ant populations recovered by the 8th week. Springtail populations built up steadily in the treatment plots throughout the experiment, but on a significantly lower level than in the control plots. Pitfall catches of *Carabidae* were statistically similar in both plots throughout the experiment (Table II).

Table VI. Insect species caught from the maize plants by the D-VAC sampler

Year	Weeks after treatment	Mean number of specimens caught \pm SE					
		<i>Aranae</i>		<i>Coccinellidae</i>		<i>Formicidae</i>	
		T	C	T	C	T	C
1994	pretreatment	2.50 \pm 0.6	2.25 \pm 0.5	4.00 \pm 0.8	4.00 \pm 0.8	1.50 \pm 1.0	1.25 \pm 0.5
	1day	0.00	2.50 \pm 0.6*	0.00	3.00 \pm 0.8*	0.25 \pm 0.5	3.50 \pm 1.3*
	2	0.00	1.25 \pm 0.5*	2.75 \pm 0.5	5.00 \pm 0.8*	0.25 \pm 0.5	2.25 \pm 1.0*
	4	1.00 \pm 0.8	3.00 \pm 0.8*	4.75 \pm 1.0	6.75 \pm 1.3*	3.00 \pm 1.2	5.25 \pm 2.2
	6	1.00 \pm 0.8	2.75 \pm 0.5*	2.50 \pm 0.6	3.00 \pm 0.8	7.00 \pm 1.8	11.75 \pm 2.4
1995	pretreatment	0.75 \pm 0.5	0.50 \pm 0.6	1.25 \pm 0.6	1.50 \pm 0.6	0.00	0.25 \pm 0.5
	1day	0.00	0.75 \pm 0.5	0.00	1.50 \pm 0.6*	0.00	0.00
	2	0.00	0.75 \pm 0.5	0.00	2.00 \pm 0.8*	0.00	2.00 \pm 0.8*
	4	0.00	1.00 \pm 0.5	0.75 \pm 0.5	3.25 \pm 0.5*	0.50 \pm 0.6	5.25 \pm 2.2*
	6	1.50 \pm 1.0	1.25 \pm 0.5	4.25 \pm 1.0	3.75 \pm 1.7	2.00 \pm 1.4	9.75 \pm 1.3*

T = treated plot, C = untreated control plot

* significantly different from the treated plot at P = 0.05

Table VII. Insect species caught from the soil in the maize plots by the D-VAC sampler

Year	Weeks after treatment	Mean number of specimens caught \pm SE							
		<i>Acarina</i>		<i>Aranae</i>		<i>Carabidae</i>		<i>Collembola</i>	
		T	C	T	C	T	C	T	C
1994	pretreatment	46.8 \pm 10.1	40.2 \pm 1.8	1.0 \pm 0.8	1.50 \pm 1.3	1.25 \pm 1.0	1.00 \pm 0.8	469 \pm 100.9	4
	1 day	43.5 \pm 5.8	41.5 \pm 4.8	0.5 \pm 0.6	1.00 \pm 0.8	0.5 \pm 0.6	2.00 \pm 0.8	424 \pm 83.3	4
	2	12.8 \pm 3.6	43.8 \pm 4.3*	0.5 \pm 0.6	1.25 \pm 0.5*	0.25 \pm 0.6	1.00 \pm 1.0	41.8 \pm 3.2	5
	4	18.8 \pm 6.7	49.5 \pm 5.2*	0.5 \pm 0.6	1.75 \pm 0.5*	0.25 \pm 0.5	0.5 \pm 0.6	127 \pm 23.2	4
	6	30.2 \pm 5.8	51.8 \pm 2.1*	1.0 \pm 0.6	3.00 \pm 1.4*	0.5 \pm 0.6	0.75 \pm 0.5	141 \pm 37.2	5
	8	40.2 \pm 7.1	73.0 \pm 4.1*	2.0 \pm 0.8	3.00 \pm 0.8	0.75 \pm 0.5	1.75 \pm 0.5	160 \pm 43.1	5
	10	58.2 \pm 6.5	41.2 \pm 2.2*	3.5 \pm 1.3	2.75 \pm 0.5	1.25 \pm 0.5	1.5 \pm 0.6	295 \pm 34.2	5
1995	pretreatment	38.5 \pm 2.5	39.8 \pm 3.1	1.25 \pm 0.5	0.75 \pm 0.5	3.00 \pm 5.4	0.50 \pm 0.6	276 \pm 20.4	2
	1 day	39.3 \pm 2.6	34.5 \pm 3.0*	1.00 \pm 0.8	0.75 \pm 0.5	0.00	0.50 \pm 0.6	276 \pm 19.3	2
	2	8.8 \pm 1.8	38.0 \pm 2.2*	0.00	2.25 \pm 1.3*	0.00	0.00	21.8 \pm 2.2	2
	4	8.5 \pm 1.3	40.5 \pm 1.3*	0.25 \pm 0.5	1.25 \pm 1.0	0.25 \pm 0.5	0.50 \pm 0.6	43.0 \pm 13.0	2
	6	20.5 \pm 4.1	44.0 \pm 5.7*	1.00 \pm 0.8	0.75 \pm 1.0	0.25 \pm 0.5	0.25 \pm 0.5	78.0 \pm 14.8	2
	8	35.5 \pm 4.1	41.8 \pm 3.9*	0.75 \pm 1.0	1.00 \pm 0.8	0.00	0.00	245.5 \pm 23.6	3
	10	40.3 \pm 2.4	41.0 \pm 4.1	1.25 \pm 0.5	1.25 \pm 0.5	0.50 \pm 0.6	0.00	280.5 \pm 27.1	3

T = treated plots, C = untreated control plots

* significantly different from the treated plot at P = 0.05

Rates of leaf disc degradation were significantly lower in the lindane treated than in the control plots. Yields, though higher in the treatment (4.53 t ha⁻¹) than in the control plots (3.51 t ha⁻¹) were not statistically different. There was a drop in yield in the lindane treated plots from 5.03 t ha⁻¹ in 1993 to 4.53 t ha⁻¹ in 1994.

4.6. Maize 1995

As in the previous years, stem borer damage increased steadily with time in both treatment and control plots, although at a significantly lower level in treatment than control plots (Fig. 1).

Arthropod samples (Table VI) had no significant difference in the mean numbers of specimens of the various families obtained in the pre-treatment samples from both control and treatment plots. Similar to the results of the previous year, no insects or spiders were caught on the plants in treatment plots 1 day after treatment. *Epilachna* sp. was found to be an important pest of maize seedlings this year, but was not found on the plants in the treated plots two weeks after lindane application. By the 4th week, coccinellids were recorded on plants in treated plots but their number ($X = 0.75$) was significantly lower than control plots ($X = 3.25$) at 5% level.

Like in the previous year, there were no significant differences in the mean numbers of insects and spiders obtained in D-vac ground samples of the treated and the control plots in the pre-treatment and 1 day post treatment samples (Table VI). Subsequent samples showed that the lindane application significantly reduced *Formicidae*, and *Collembola* and *Acarina* populations for about 4 and 8 weeks, respectively, after application. The effects of lindane on carabid populations was not significant. In general, few *Aranae* were caught in the D-vac this year and analysis showed no significant difference in numbers caught in treated and control plots on most sampling dates. This result may be misleading because of the few numbers involved.

Pitfall traps were left in the field for 1 week and this resulted in richer collections of *Collembola* and *Formicidae* than in the previous year. *Collembola* trapped in pitfalls in the pre-treatment samples were enormous ($X = 5629$ and 5555 for control and treatment plots, respectively) compared to the previous years. Samples 1 day after treatment showed that the *Collembola* population in the lindane treated plots had crashed ($X=173$) compared with the controls ($X = 5651$). The *Collembola* population remained significantly lower in the lindane-treated than in the control plots throughout the experiment. For *Formicidae*, the mean catches in the treatment plots remained significantly lower up to 4 weeks after lindane application.

The mean dry weight of leaf discs retrieved from treatment and control plots were statistically different. Higher yields (4.03 t ha^{-1}) were obtained in treatment than control plots (3.60 t ha^{-1}), and their differences were statistically significant. Decrease in yield in treatment plot was also recorded from 4.53 t ha^{-1} in 1994 to 4.03 t ha^{-1} in 1995.

5. DISCUSSION

Lindane is used extensively in Nigeria on cocoa and sometimes also in rice farms. However, it is not recommended for use on maize. Throughout, the maize crop showed phytotoxicity after lindane application and this tended to retard plant growth for a while.

Lindane protected maize and rice plants effectively from stem borer damage in most years. However, this protection was not always reflected in crop yields. Indeed, significant differences in maize yields in the lindane-treated compared with the control plots were recorded only in 1993 and 1995. Rice yield in 1992 and 1993 were not statistically significant. This would suggest that stem borers and perhaps other pests controlled by lindane were not the only determining factors in yield in these experiments. It is possible that lindane may have had an indirect negative effect on yields which declined in the treatment plots over the years. Perfect et al. [6] recorded similar decline in yields of cowpea grown on soils treated with DDT in Nigeria over a 4-year period. They postulated that DDT adversely affected the numbers and structure of soil fauna as well as the decomposition of organic matter which affected some soil properties. Their postulation was based on an earlier paper [9] which showed that elimination of soil fauna by fumigation adversely affected plant growth. The indirect effects of lindane on yield, speculated in this work, may be the result of its adverse effects on some aspects of soil biology which in turn affected yield.

The target soil fauna in these experiments were *Collembola* (springtails) and *Acarina* (mites). The mites included both predatory and detritivorous species. These soil invertebrates are micro-arthropods which, by virtue of their sizes should easily be affected by pesticides. The degree of susceptibility of these micro-arthropods to lindane in these experiments appeared to depend on the concentration of the spray solution, the quantity of the spray solution that reached the soil and perhaps also, their intrinsic ability to resist adverse effects of lindane. In the maize plots (1992) with plant distances of 1m and 500mm between rows and hills, respectively, large areas of soil surface were exposed and received lindane (1 kg AI ha^{-1}) sprays directly, and this decimated both *Collembola* and *Acarina* populations, resulting in significantly fewer numbers recovered in the lindane treated than the control plots in the post-treatment samples. In 1993, with reduced inter-row distances, a closer plant canopy was formed, which reduced the exposed soil surface thereby giving greater protection to the soil fauna. The split dose application of 1 kg AI lindane per hectare this year may also have helped these micro-fauna to resist lindane toxicity resulting in no significant differences in the mean numbers of both *Collembola* and *Acarina* recorded in treatment and control plots.

In rice (1992) with plant distances of 150 mm both between rows and hills, a close plant canopy was formed, leaving limited exposed soil surfaces which received lindane sprays (1 kg AI ha⁻¹ in a single dose) directly. In this situation, the *Acarina* which probably had greater intrinsic resistance to lindane than the *Collembola* were unaffected. However in 1993, significantly fewer *Acarina* was recorded in the lindane treated than in the control plots. It is possible that rainfall in the season had spread lindane residues from where the lindane was deposited to other areas in the plot in 1992. Igbariam had a total annual rainfall of 2181 mm in 1992. Ploughing and harrowing in 1993 may have also helped to redistribute residual lindane in the farm. Samuel and Pillai [10] reported that about 15% of the initial concentration of lindane applied to an Indian soil remained as extractible and bound residues one year after application. Yeadon and Perfect [11] found that after 1 year of soil treatment with DDT, 4.2% of the initial concentration was detected as residues (chiefly DDE) and after 4 years the percentage of the detectable DDE rose to 24%. With 1993 spray, the level of lindane in the soil may have become high enough to also affect *Acarina*.

Micro-arthropods, particularly *Collembola*, which feed on leaf litter, are regarded as important in the fragmentation and incorporation of leaf litter into the soil. Surprisingly, in these experiments (1992 and 1993), there was no significant difference in the mean dry weight of leaf discs retrieved from treatment and control plots in spite of the significant differences in collembola population in treated and control plots. On retrieval of litter bags, they were found to be infested with young earthworms, nematodes, other unidentified mesofauna, roots of weeds and in a few cases, ants. These agents, together with some soil microflora (bacteria and fungi) probably played more important role in litter fragmentation inside the soil than collembola.

The effects of lindane on the agents of litter fragmentation when the same plot was used over the years appeared to be cumulative. For example, in 1992 (on rice) and 1993 (on maize), the effects of lindane on litter disintegration was not significant. In subsequent years however, the effect became progressively more significant ($P < 0.024$ in 1994 and $P < 0.005$ in 1995). This may suggest a cumulative effect due probably to lindane residue which may have increasingly accumulated over the years to a level that drastically affected agents of litter fragmentation in sprayed plots in 1995.

Throughout these experiments, spiders and ants were consistently adversely affected by lindane. Spiders are generalist predators known to range freely over sprayed vegetation and soil surfaces in search of their preys. Critchley et al. [5] also recorded adverse effects of DDT on spiders and ants.

In general, a sweep net appeared inappropriate for sampling maize plants as very few specimens were caught in sweep net compared with the numbers caught with the D-vac sampler.

REFERENCES

- [1] EDWARDS, C.A., Insecticide residue in soils, *Residue Rev.* **13** (1966) 83-132.
- [2] EDWARDS, C.A., 'Pesticide residues in soil and water', in *Environmental Pollution by Pesticides* (EDWARDS, C.A., Ed.), Plenum, London (1973) 1-7.
- [3] EDWARDS, C.A., THOMPSON, A.R., Residues and the soil fauna, *Residue Rev.* **45** (1973) 1-79.

- [4] KELSEY, J.M., ARLIDGE, E. Z., Effects of isobenzan on soil fauna and soil structure, New Zealand Jour. Agric. Res. **11** (1968) 245-260.
- [5] PALISA, A., Soil animals and soil fertility, Urania **20** (1668) 3- 33.[6] PERFECT, T.J., et.al., The effects of crop protection with DDT on some elements of the subterranean and surface active arthropod fauna of a cultivated forest soil in the humid tropics, Pedobiologia **20** (1980) 31-38.
- [7] CRITCHLEY, B.R., et al., The effects of crop protection with DDT on some elements of the subterranean and surface active arthropod fauna of a cultivated forest soil in the humid tropics, Pedobiologia **20** (1980) 31-38.
- [8] ZADOKS, J.C., *et al.*, A Decimal code for the growth stages of cereals, Weed Research **14** (1994) 415-421.
- [9] EDWARDS, C.A., LOFTY, J.R., The influence of arthropods and earthworms upon root growth of direct drilled cereals, Jour. Applied Ecol. **15** (1978) 789-795.
- [10] SAMUEL, T., PILLAI, M.K.K., "Persistence and fate of ¹⁴C-gamma-HCH in an Indian sandy loam soil under field and laboratory conditions", Isotope Techniques for Studying the Fate of Persistent Pesticides in the Tropics, IAEA-TECDOC-476, IAEA,Vienna (1988) 93-102.
- [11] YEADON, R., PERFECT, T.J., DDT residues in crop and soil resuting from application of cowpea *Vigna unguiculata* (L.) Walp.in the sub-humid tropics, Environ. Pollution **B2** (1981) 275-294.

**NEXT PAGE(S)
left BLANK**

THE DEVELOPMENT OF LABORATORY AND SEMI-FIELD METHODS TO TEST THE EFFECTS OF PESTICIDES ON PREDATORY BEETLES

P. CHIVERTON, H. WALLIN*
Department of Entomology,
Swedish University of Agricultural Sciences,
Uppsala, Sweden



XA9743726

Abstract

Following the sequential testing procedure adopted by the IOBC/WPRS Working Group Pesticides and Beneficial Organisms, two simple, robust methods are presented which were designed for testing the effects of pesticides on predatory beetles. In a laboratory initial toxicity test both DDT and lindane were found harmful to the carabid *Pterostichus cupreus*, whereas α -endosulfan was 'harmless'. DDT was found harmless to *P. melanarius*. Sub-lethal doses of both DDT and lindane incorporated in prey caused *P. cupreus* females to produce smaller eggs. In a semi-field test it was demonstrated that Lindane reduced the beneficial capacity of *P. cupreus*. Climatic conditions at the time of the test however were such that the majority of test animals in control treatments escaped. Caution was therefore advised in the choice of test animal and test design for the semi-field test.

1. INTRODUCTION

Testing or screening for the side-effects of pesticides on beneficial arthropods is gaining increasing attention by research workers in different parts of the world [1]. Results from such tests enable agronomists to choose safe preparations for use in Integrated Pest Management schemes. They can also be of use to authorities concerned with pesticide registration.

Recognising the need for standardised testing methods and greater international co-operation, a Working Group Pesticides and Beneficial Organisms was formed in 1974 within the framework of the International Organisation of Biological Control (IOBC), Western Palaearctic Regional Section (WPRS) [2]. Aware of the fact that no single test method would provide sufficient information to show the side-effects of a pesticide on a beneficial organism, this group devised a sequential procedure that includes laboratory, semi-field and field methods. These methods are carried out in a particular sequence progressing from laboratory to semi-field and, ultimately, to a field test only if a preparation under examination is found to be harmful to the test organism. The reasoning behind this is that pesticides found harmless to a particular beneficial in a stringent laboratory initial toxicity test are most likely to be harmless to the same organism in the field.

In this paper we adapt a laboratory and a semi-field test method, originally designed for and used in the IOBC Working Group, to examine the direct effects of organochlorine pesticides on predatory arthropods. We also examine whether any sub-lethal effects occur when predatory beetles consume organochlorine-tainted prey. The aim has been to produce protocols for robust test methods that could be used by African colleagues participating in the FAO/IAEA/SIDA Co-ordinated Research Programme on Adverse Effects on Flora and Fauna from the Use of Organochlorine Pesticides on the African Continent.

* Present address: Anticimex AB, Box 47025, S 100 74, Stockholm, Sweden.

2. MATERIALS AND METHODS

2.1. Selection and treatment of test animals

The beetles chosen to illustrate our test methods were two medium-sized carabids, *Pterostichus cupreus* L. and *Pterostichus melanarius* (Illiger) (Carabidae, Coleoptera). Both are common in agricultural fields in Sweden and both have been shown to prey upon the most serious aphid pest in Swedish cereals, *Rhopalosiphum padi* L.[3]. Large numbers of beetles were caught in May and June 1991 in dry pitfall traps. The latter were placed on uncultivated land where no previous pesticide treatments had been made. Beetles were held in plastic containers lined with moistened tissue paper, in a refrigerator (5°C) and fed dog biscuits. All beetles were acclimatised to test conditions (15°C, 16 h light, 8 h dark, 70% r.h.) at least 72 h prior to testing.

2.2. Determination of insecticidal activity in laboratory tests

The basic test method was as follows: Individual beetles were transferred to small plastic beakers (height, 60 mm; diameter, 60 mm) containing 20 mm of even-grained sterile sand. Twenty such beakers were then placed in four rows of five on a plastic seeding tray (400 x 600 mm x 80 mm deep) in a fume cupboard. The beakers (+ beetles) were then sprayed with a hand-held sprayer connected to a compressor set at 35 kPa. The spray nozzle was passed over the beakers at a constant speed and height, calculated to deliver the recommended concentration at the IOBC standard amount (for initial toxicity tests in laboratories) of 6 mg fluid per 100 mm² sand. Two test runs of 20 replicates per run were performed for each pesticide/dose (see below). Controls were sprayed with water + alcohol (19 + 1 by volume), as this was used to dissolve the organochlorines tested. Between sprays, the sprayer was thoroughly washed with repeated dosings of acetone followed by water and then dried.

Treated groups were transferred to a large, well ventilated controlled climate room set at the above conditions. The small beakers were then inserted into larger ones of the same diameter (i.e. height, 60 mm; diameter, 60 mm). This ensured that the beetles a) could not escape and b) did not come into contact with excess pesticide on the beaker walls. Each beetle received one dog biscuit as food, and this was moistened daily. Beetles were examined at 1, 6 and 24 h post treatment, and thereafter daily until the sixth day whereupon the test was terminated. Numbers of beetles found dead or moribund were recorded.

Two toxicity tests were performed using this basic method. Different dilutions of DDT, starting with a 'field dose' equivalent to 1000g AI ha⁻¹ (as this was the most common dose found in the literature) were tested on *P. melanarius*, and several organochlorines (i.e. DDT at a rate equivalent to 1000g AI ha⁻¹, α - and β -endosulfan : both at 1100g AI ha⁻¹ and lindane : at 350g AI ha⁻¹) were screened for toxicity to *P. cupreus*. In both tests water containing alcohol (19 + 1 by volume) was used as a control treatment.

2.3. Determination of sub-lethal effects in laboratory tests

In June 1991, 50 female *P. cupreus* were selected from our field-collected stock. These were placed individually in petri dishes containing moistened pulverised peat, and divided into five groups of ten. Every third day one group of ten beetles were each fed with

one (initially two) blowfly larvae (maggots) which had previously been sprayed with a 1:32 dilution of the 'field dose' DDT used above. The other four groups of ten beetles were fed maggots that had been sprayed with dilutions (1:32) of α - and β -endosulfan, lindane, and water + alcohol (19 + 1 by volume) as control, respectively. After spraying, all but two of the maggots were frozen for later use. At subsequent feeding events, maggots were thawed and then almost cut in half prior to feeding in order to ensure that the beetles could penetrate the larval cuticle. Twice a week beetles were temporarily removed from the petri dishes and the peat substrate was replaced. The used substrate was washed through a fine-meshed sieve. Any eggs the beetles had laid in the substrate were subsequently caught in the sieve. The number of eggs found per female were recorded. Eggs were placed briefly on a filter paper to dry and then weighed.

2.4. Determination of insecticidal activity in semi-field tests

The species of beetle used in this test (*P. cupreus*) was collected and held under the same conditions as above. In June 1992, 16 plastic seeding trays (400 x 600 x 80 mm) were half-filled with potting soil and sown with three rows of spring barley at normal field seed rates (4-500 per square meter, in this case ~50 plants per 600 mm row) and at a row width of 125 mm. Trays were then placed in a greenhouse to hasten germination. The inner edges of each tray, from the soil surface to the rim, were painted with Fluon (polytetra-fluoroethylene) to discourage beetles from climbing the tray walls and escaping. When the plants had reached growth stage 32 4 four of the trays were taken outdoors and sprayed with Lindane at a 'field' dose equivalent to 300 g AI ha⁻¹. Four trays were sprayed with a dilution (1:4) of the field dose, and a further four trays were sprayed with a second dilution (1:16). The remaining four trays were sprayed with water + alcohol (19 + 1 by volume) containing several drops of a wetting agent (Triton X-100) as controls. The trays were allowed to dry for 24 h, after which four *P. cupreus* were released onto each tray. The trays were then placed in a large, outdoor bird-proof cage and the beetles allowed to acclimatise for 48 h.

At 72 h post-treatment, each tray was provided with 100 freshly-frozen *R. padi* arranged on four moistened filter papers with 25 aphids per paper. The filter papers were placed on the soil surface and covered with a plastic roof to protect the aphids from the rain and wind. The number of aphids removed after 24 h was recorded. This procedure was repeated at weekly intervals for 3 weeks. Four fresh beetles were released onto each tray 24 h before each feeding occasion. The mean total number of aphids eaten per treatment and per feeding occasion was noted and the results examined by analysis of variance (ANOVA).

Dead beetles found on the soil surface were removed during the course of the experiment and numbers recorded. After the last feeding period, plants were cut at the stem base and removed from the trays. The soil surface was then carefully searched and surviving beetles removed with a pooter. Thereafter, the soil was sieved onto a larger tray and the beetles that had buried into the soil were recovered. The mean number of survivors per treatment was noted and the results analysed by ANOVA.

3. RESULTS

3.1. Insecticidal activities in laboratory tests

The chosen 'field' dose of 1000g AI ha⁻¹ DDT caused 40% mortality in *P. melanarius* with dilutions of this field dose giving a typical dose response curve (Fig. 1). In contrast, the

same field dose of DDT caused 100% mortality in *P. cupreus*, as did the chosen field dose of lindane, whereas α -endosulfan only caused 30% mortality even at the relatively high dose of 1100 g AI ha⁻¹ (Fig. 2).




Fig 1. Initial toxicity of DDT to the carabid beetle *Pterostichus melanarius*




Fig 2. Selectivity of organochlorine insecticides to the carabid beetle *Pterostichus cupreus* in a laboratory initial toxicity test

3.2. Sub-lethal effects in laboratory studies

No significant differences were found between the number of eggs laid by female *P. cupreus* fed on organochlorine tainted prey or fed on water treated control prey ($F = 0.28$, d.f. = 50, n.s.). Eggs from female beetles fed sub-lethal doses of DDT and lindane, however, were significantly smaller (Table 1).

3.3. Insecticidal activities in a semi-field test.

The mean number of aphids eaten per treatment for each feeding occasion are given in Table 2. Mean numbers of aphids eaten in the field dose treatment were always significantly

lower than numbers in the diluted (1:4) field dose treatment. Surprisingly, mean numbers eaten in the control were not significantly different from those in either the field dose or from the different dilutions of the field dose. An explanation was found at the end of the experiment when beetles were recovered. Only one beetle of a treatment total of 48 was recovered from the control trays. There was no significant difference in the number of survivors in the treated plots (Table 3), although dead beetles were only recovered from the field dose (1:1) treatment.

Table 1. Sub-lethal effects of organochlorine tainted prey on egg weights of the carabid beetle *Pterostichus cupreus*.

Pesticide	Egg wt. (mg)	N	P
DDT	0.65	291	0.0001
endosulfan	0.70	288	n.s.
endosulfan	0.69	261	n.s.
lindane	0.63	315	0.0001
water control	0.71	254	

n.s. = not significant in a Student's T-test against control

Table 2. Consumption of aphids by the carabid *Pterostichus cupreus* in arenas sprayed either with different dilutions of lindane or with water treated controls.

Week after treatment	Control	lindane (300g AI ha ⁻¹)		
		1:1	1:4	1:16
1	12.9 ab	4.3 b	20.9 a	16.4 ab
2	14.3 ab	4.5 b	20.8 a	20.0 a
3	14.6 bc	11.1 c	24.4 a	24.4 ab

Numbers with the same letter are not significantly different at $P < 0.05$ (Duncan's multiple range test)

Table 3. Recovery of alive and dead *Pterostichus cupreus* from arenas sprayed either with lindane dilutions or water treated controls.

Insect status	Control	lindane (300 g AI ha ⁻¹)		
		1:1	1:4	1:16
live	0.3 b	1.8 ab	3.5 a	2.5 a
dead	0	41	0	0

Numbers with the same letter are not significantly different from the control at $P < 0.05$

4. DISCUSSION

The carabid species chosen for the semi-field test, *P. cupreus*, is macropterous and can therefore fly. However, extensive behavioural studies of this species (in which beetles confined on similar arenas were video filmed at different temperatures in a controlled climate chamber 5) had revealed that the beetle rarely flies unless the soil surface temperature exceeds ~ 30°C, whereupon they climb to the top of the plants and fly away. Unfortunately the summer of 1992 was one of the hottest and driest on record in Sweden, with ambient temperatures often exceeding 26-28°C throughout the duration of the experiment! Soil surface temperatures, usually 10°C higher than ambient temperature, would therefore have exceeded the 30°C upper threshold for this species and caused it to flee. Interestingly, fewer beetles fled, or were capable of fleeing, from the 1:4 and 1:16 dilution treatments of lindane as significantly more were recovered from the arenas compared with controls. We have previously used this method successfully to screen the effects of pesticides on a small, brachypterous carabid, *Bembidion lampros* Herbst., with good results and would therefore encourage the use of the method, though with a caution that the beneficial insect should be carefully selected.

This laboratory test has proved useful, and is robust enough to be carried out in most laboratories. The basic method can even be adapted and used as a persistence test, in which beetles are placed in the beakers at intervals post-treatment. This would be a useful exercise particularly regarding the organochlorines, and might reveal differences between, for example, DDT and lindane regarding the persistence of harmful effects. Should a macropterous species be chosen for the laboratory test, we suggest that a simple muslin cloth coated with Fluon and held in place by a rubber band over the top of the beaker will suffice to prevent escape. This was demonstrated at the CRP Workshop in Arusha where the method to screen the effects of organochlorines was used on a coccinellid species.

The increase in egg production following ingestion of a carbamate (pirimicarb) has previously been observed in *P. cupreus* [6] and is thought to be due to a hormoligosis effect [7]. Egg size has been found to affect the survival of carabid larvae [6], and first instars hatching from large eggs can withstand starvation better than those hatching from small eggs. Therefore ingestion of DDT or lindane tainted prey by female ground beetles could have significant effects on the survival of their progeny. These and other sub-lethal effects are not exclusive to the organochlorines and need to be considered when describing the overall side effects of pesticides.

ACKNOWLEDGEMENTS

We thank Carol Högfeldt for technical assistance. This work was carried out within the project "The effects of organochlorine pesticides on the non-target fauna" and was supported by grants from the Swedish International Development Authority (SIDA).

REFERENCES

- [1] HASSAN, S.A., Procedure for testing the side effects of pesticides on beneficial arthropods as being considered by the International Working Group Pesticides and

- Beneficial Arthropods, Mitteilungen der Deutschen Gesellschaft fuer Allgemeine und Angewandte Entomologie **4** (1983) 86-88.
- [2] FRANZ, J.M., Pesticides and Beneficial Arthropods, IOBC/WPRS Bulletin **1** (1975)147-152.
- [3] CHIVERTON, P.A., Predation on Rhopalosiphum padi by polyphagous predatory arthropods during the aphids pre-peak period in spring barley, Ann. Appl. Biol. **111** (1987) 257-269.
- [4] ZADOKS, J.C., CHANG, T.T., KONZAK, D.E., A decimal code for the growthstages of cereals, Weed Research **14** (1974) 415-421.
- [5] CHIVERTON, P.A., Searching behaviour and cereal aphid consumption by Bembidion lampros and Pterostichus cupreus, in relation to temperature and prey density, Entomol. Exp. Appl. **47** (1988) 173-182.
- [6] WALLIN, H., CHIVERTON, P.A., EKBOM, B.S., BORG, A., Diet, fecundity and egg size in some polyphagous predatory beetles, Entomol. Exp. Appl. **65** (1992) 129-140.
- [7] LUCKEY, T.D., Insecticide hormoligosis, J. Econ. Entomol. **61** (1968) 7 - 12.

**NEXT PAGE(S)
left BLANK**



EFFECTS OF LINDANE ON A MAIZE ECOSYSTEM AT TPRI, ARUSHA, TANZANIA

E.M. MINJA, R. MAKUSI, F. TESHA
Tropical Pesticides Research Institute,
Arusha, Tanzania

Abstract

The effects of the organochlorine insecticide, lindane, on target and non-target organisms was studied in field plots of maize at Arusha, Tanzania for 4 growing seasons. A single application of $1 \text{ kg} \cdot \text{ha}^{-1}$ caused leaf scorch to the crop but two applications of $0.5 \text{ kg} \cdot \text{ha}^{-1}$ at an interval of 2 weeks were tolerated. The insecticide reduced damage by the stem borers *Busseola fusca* and *Sesamia calamistis* in all seasons and maize yields were higher on the treated plots although the differences were not always significant ($P > 0.05$). Collembola and sometimes ant and spider numbers were lower in treated plots early in the season but the differences did not persist. Differences in rates of decomposition of buried leaf litter were not significant.

1. INTRODUCTION

The use of synthetic pesticides in crop and forestry production as well as in the control of both animal and human disease vectors has been a common practice worldwide. These pesticides are harmful to the environment particularly those which persist for long periods. Persistent pesticides include insecticides in the organochlorine group (e.g. DDT, benzene hexachloride-BHC, chlordane, heptachlor, toxaphene, methoxychlor, aldrin, dieldrin, endrin and endosulfan) which are lipophilic. While most of these have been banned from use in many countries on the basis of their persistence and accumulation in animal and human tissues, some of them are still registered for use in Tanzania [1].

Lindane (gamma benzene hexachloride) is among the organochlorine insecticides registered for use in crop protection in Tanzania. In most developed countries the environmental impact of organochlorine pesticides has been monitored and documented [2]. It has been calculated that as much as 50% of pesticide sprays applied to crops miss their target and fall on to the soil surface. Also the air can easily become contaminated with pesticides during spraying operations [3]. In most developing countries including Tanzania, very little information is available on pesticide effects on the environment. The maize stem borer *Busseola fusca* Fuller is one of the major insect pests limiting maize production in Tanzania. DDT dust and high volume sprays were recommended and commonly used by farmers but after the banning of DDT due to its persistence and hazardous effects on the environment, farmers have been using various other insecticides including lindane. The work being presented was therefore proposed to investigate the effects of lindane on field maize arthropod fauna.

2. MATERIALS AND METHODS

The studies were conducted on a 0.7 ha field at the Tropical Pesticides Research Institute, Arusha for four planting seasons starting in 1991/92 short rain season. Planting dates were based on the on-set of the rains which was November/December (this applied to 1991/92, 1992/93 and 1994/95) for the short rains and March/April for the long rains (1994 trial). Planting of the trial was done in the 1994 long rain season (which is cooler and wetter, and therefore with lower insect populations compared with the warm short rain season) due to failure of the 1993/94 short rain season crop from a severe drought in January/February 1994. The field was first ploughed and for the first two seasons animal manure (cow dung) at 3 tons

per ha was broadcasted and harrowed into the volcanic sand loamy soil which is known to be low in organic matter. The cow dung was applied to the soil a month before planting maize. The field was then divided into four equal blocks which were each further divided into two 25 m x 25 m plots. The plots in a block were each randomly assigned two treatments i.e. sprayed and unsprayed.

Maize hybrid seed CG 4141 was sown at 90 cm x 30 cm at two seeds per hole and the seedlings were thinned to one per hill at three weeks after emergence (3WAE). All field operations apart from tractor ploughing and harrowing were carried out by hand. Fertilizer top dressing using urea at 25 kg N/ha was done twice at 3WAE and 6WAE.

In each of the two plots per block, 5 m x 5 m subplots were marked out permanently for different investigations. Peripheral subplots acted as guard rows. In one subplot 4 standard pitfall traps were installed in the ground for monitoring ground fauna population activity. The traps were set for three days with a weak soap solution. They were then emptied and arthropod fauna sorted out and recorded. A second subplot was used for aerial fauna monitoring in which arthropods were swept from plants using a sweep net (three seasons) and a D-Vac in the fourth season. A third subplot was used for visual fresh plant damage assessment where all plants with fresh borer damage were counted and converted into a proportion of total subplot plant population. A fourth subplot was left undisturbed until harvest for grain yield assessment. A fifth subplot was used for organic matter decomposition studies in which four litter bags containing *Grevillea robusta* A. Cunn leaves were randomly buried for three months. Twenty plants were randomly uprooted from the remaining four subplots at each assessment day for two seasons to assess stem borer larval population and parasite incidence. The stems were each dissected for stem borer larval counts.

The crop was constantly inspected for stem borer eggs and first instar larvae to help in setting out the date for initial insecticide sprays. Spraying was carried out at 4WAE when either eggs and or first instar stem borer larvae were recorded on 0.5-1.0% of the plants. Lindane 20 EC. (Gammalin 20% from Imperial Chemical Industries) was sprayed once at 1.0 kg AI.ha⁻¹ in 1991/92 and 1992/93. Because this caused severe leaf scorching the application was split and sprayed twice at 0.5 kg.AI.ha⁻¹ at two week intervals starting at 4WAE in 1994 and 1994/95. CP15 and CP20 knapsack hand sprayers with polijet cone type of nozzles and operated at a pressure of 3 x 10⁵ Pa (3 bar) were used at high volume rate (300 L water ha⁻¹). Arthropod populations and plant damage were monitored just before the first spray, 24 hours later and then every two weeks until harvest. All data was subjected to analysis of variance.

3. RESULTS

Field observations showed that the maize stem borer, *B. fusca* was the dominant pest early in the four seasons starting towards the end of the 3WAE. Towards the middle of each growing season, however, minor infestations of *Sesamia calamistis* Hamp. occurred and the two borers cohabited on the plants until harvest. Damage assessments were based on *B. fusca* which feeds in the leaf whorl, in the stems and on maturing seed on the cobs. *S. calamistis* normally rings the lower part of the stems and migrate to other stems or to the maturing seeds on the cobs.

The total monthly rainfall and mean temperatures for the institute where the experiments were carried out are shown in Figures 1-4 corresponding to the four seasons. The results on beneficial arthropod population groups represented by ants, spiders, and collembola,

and plant damage for 1991/92, 1992/93, 1994 and 1994/95 are shown in Tables 1-4, respectively. Buried leaf litter weight losses are shown in Table 5 and maize grain yields in Table 6.

The results for 1991/2 show that arthropod populations were very low in that season. The one lindane application was sufficient to check borer infestations for 7 weeks by which time the plants were old enough to withstand a degree of pest attack.

In 1992/3 lindane significantly reduced Collembola numbers initially but the population recovered to a similar level to that in the control plots by the end of the season. Ant and spider numbers were essentially the same in treated and control plots but treated plots showed significantly less crop damage.

The 1993/4 short season crop failed because of drought so a crop was grown in the 1994 long rainy season. The long rainy season is normally cooler and wetter with relatively low pest insect activity but more abundant beneficial soil arthropods. Splitting the lindane treatment into two applications prolonged the depression of arthropod populations but, as in the previous seasons, they recovered by harvest time. Again, pest damage to the crop was significantly lower on the treated plots. These trends were repeated in 1994/5. In addition a gregarious *Apanteles* spp. which is a parasite of *B. fusca* larvae was observed in similar numbers in treated and untreated plots.

Buried leaf litter decomposition (Table 5) was slightly, but, not significantly, lower in treated than control plots in all three years that litter bags were buried. Grain yields (Table 6) were 31% higher on average in the treated plots over the 4 seasons although the variability was such that the differences were not significant ($p > 0.05$).

4. DISCUSSION

The results indicate that lindane was effective in reducing *B. fusca* damage on maize. On the other hand it had adverse effects on most of the early season beneficial arthropod fauna monitored in the four seasons. It was especially so with Collembola, and in some seasons it also affected ants and spiders. Similar reductions in population activity for ants and spiders were observed in the humid tropical zone of Nigeria in plots treated with DDT [4].

The mid- and late-season beneficial arthropods including spiders (Araneae) and ants (Formicidae) were not significantly affected by lindane. Collembola populations were the most abundant beneficial soil microfauna at the beginning of each season but their numbers would normally decline gradually even in the controls as soil moisture decreased towards crop maturity. These results indicate that lindane could have short-lived adverse effects on the maize environment at the rate in which it was used in the warm and semi-humid climatic conditions in which the experiments were conducted. Edwards [5] observed that in a combination of wet and warm weather, the pesticides are released and may break down and disperse into the atmosphere much more rapidly than from dry soils; and that there was evidence that pesticides which persist for more than 10 years in temperate climates may disappear almost entirely in a year in the tropics. The results in this programme from Ghana, Nigeria and Zambia confirm this.

Lindane sprays in the present study seem to have reduced buried leaf litter decomposition rate (though not significantly) in the three seasons. Although there was no

Text cont. on p. 179.




Figure 1: Total monthly rainfall (mm) at T.P.R.I. for July 1991 - June 1992




Figure 2: Total monthly rainfall (mm) and mean monthly temperature at T.P.R.I. during 1992/1993




Figure 3: Total monthly rainfall (mm) at T.P.R.I. for July 1993 - June 1994




Figure 4: Total monthly rainfall (mm) at T.P.R.I. for July 1994 - June 1995

Table 1. Mean number of ants (Formicidae) populations in sprayed (lindane) and unsprayed plots

Sampling occasion	1991/92		1992/93		1994		1994/95	
	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed
1	0	0	36	16.75	56.25	65.75	7.25	11.75
2*	0	0	27.25	18.50	18.75	65.00	10.50	5.25
3	0	1.5	21.25	16.00	18.50	55.75	64.75	92.50
4**	1.75	6.0	28.25	33.25	8.00	53.50	44.75	53.00
5	3.25	0.75	52.50	47.00	27.50	47.25	22.25	25.50
6	8.75	12.00	30.00	82.25	37.00	41.75	23.75	23.00
7	0.75	11.75	61.00	54.25	65.75	52.75	0	0
8	0.0	13.00	66.00	74.25	98.25	220.00	0	0
9					113.50	238.75		
10					144.25	236.50		
Mean±SE	1.8±0.30	5.6±1.20	40.3±5.62	43.5±6.89	58.8±7.54	107.7±10.7	28.9±5.22	35.2±6.80

- Sampling occasion 2 corresponds to 24 hours after the first spray for all four seasons

** Sampling occasion 4 corresponds to two weeks after the second spray in 1994 and 1994/95

Mean values for 1992/92 are significantly different ($p < 0.05$), values for other years are not

Table 2. Mean number of spider populations in sprayed and unsprayed plots

Sampling occasion	1991/92		1992/93		1994		1994/95	
	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed
1	0	0	3.00	3.75	4.00	8.25	10.5	12.00
2*	0	0	3.75	2.00	1.25	6.75	1.75	5.00
3	0	0.75	1.25	5.50	1.25	8.25	3.50	4.00
4**	0.75	2.50	2.5	3.75	1.25	12.25	8.00	11.75
5	0.50	1.50	4.25	3.00	2.25	12.00	0.50	3.00
6	2.75	1.50	7.00	6.75	5.25	11.00	3.25	3.00
7	0.75	1.50	3.25	5.50	5.25	10.25	0	0
8	2.75	2.00	3.75	6.00	7.75	11.00	0	0
9					9.25	10.25		
10					10.00	11.00		
Mean±SE	0.94±0.21	1.2±0.40	3.6±0.84	4.5±0.67	4.8±0.96	10.1±2.90	4.6±1.10	6.5±1.21

Mean values for 1994 are significantly different ($p < 0.05$), values for other years are not

Table 3. Mean number of Collembola populations in sprayed and unsprayed plots

Sampling occasion	1992/93		1994		1994/95	
	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed
1	268.75	275.75	594.75	528.25	787.50	790.00
2*	190.00	262.25	228.00	603.50	17.50	12.50
3	189.25	247.50	204.50	688.00	275.00	430.00
4**	145.00	249.00	175.25	366.00	282.50	787.50
5	143.75	187.00	82.75	101.50	166.50	294.00
6	162.00	175.50	61.25	74.75	91.25	108.50
7	180.00	193.75	85.00	70.75	45.00	73.75
8	148.50	173.00	87.25	69.00	0	0
9			87.50	89.00		
10			94.50	98.50		
Mean \pm SE	178.4 \pm 12.8	220.5 \pm 20.6	170.1 \pm 18.7	288.05 \pm 40.63	237.9 \pm 17.5	356.6 \pm 20.6

Mean values for treated and control plots are significantly different ($p < 0.05$) for all years

Table 4. Fresh borer damage (% mean) on maize plants in sprayed and unsprayed plots

Sampling occasion	1991/92		1992/93		1994		1994/95	
	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed
1	1.5	0.7	1.2	1.4	0.25	0.37	1.10	1.00
2*	0	0.7	2.9	6.8	0	0.75	0.40	1.42
3	0	5.0	2.7	5.1	0	1.12	0	2.30
4**	0	6.2	4.4	9.0	0.25	1.37	1.12	3.55
5	2.2	8.0	0.6	3.0	0.62	1.62	1.92	7.72
6	1.1	6.0	0	0.5	1.12	1.87	2.70	12.40
7	1.0	3.2	0	0	1.25	2.50	0	0
8	1.0	3.4	0	0	1.50	2.62	0	0
9					2.00	2.75		
10					2.00	3.05		
Mean \pm SE	0.85 \pm 0.12	4.2 \pm 0.58	2.0 \pm 0.04	4.3 \pm 0.61	0.90 \pm 0.03	1.8 \pm 0.06	0.9 \pm 0.17	3.5 \pm 0.71

Mean values for treated and control plots are significantly different ($p < 0.05$) for all years

TABLE 5. MEAN PERCENTAGE BURIED LEAF LITTER WEIGHT LOSS IN SPRAYED AND UNSPRAYED PLOTS

Season	Sprayed	Unsprayed	P value
1992/93	34.4 ± 2.6	37.6 ± 3.2	Non significant
1994	25.8 ± 2.5	27.7 ± 1.9	Non significant
1994/95	9.4 ± 2.4	10.1 ± 1.6	Non significant

TABLE 6. MAIZE GRAIN YIELD (T HA⁻¹) IN SPRAYED AND UNSPRAYED PLOTS (MEANS ± SE)

Season	Sprayed	Unsprayed	%yield increase	P value
1991/92	2.20 ± 0.06	1.90 ± 0.03	15.79	Non significant
1992/93	2.40 ± 0.05	1.50 ± 0.02	60.00	Non significant
1994	2.56 ± 0.04	1.80 ± 0.04	42.22	Non significant
1994/95	2.99 ± 0.03	2.79 ± 0.04	7.00	Non significant

microarthropod extraction from the buried leaf litter in this experiment, the reduced Collembola activity in pitfall traps reported above could help to explain this slow-down in decomposition. Cook *et al.* [6] showed that Collembola and Acari account for 90-100% of fauna responsible for buried leaf litter decomposition and that their numbers were significantly reduced by DDT sprays.

The results further show that lindane reduced early season stem borer infestations thereby protecting the young maize plants at their critical stage of vegetative growth (4-8WAE). This protection is revealed by low plant damage. Protection of young maize from borer damage is important in that early infestations lead to direct plant loss due to lack of compensatory growth. Timing of spray application before borer larvae tunnel the stems is equally crucial. Delayed applications will not reach the protected larvae in the stems. Furthermore, many larvae which will have hatched from an egg batch on a single plant will have migrated to other plants if sprays are delayed until the larvae reach their third instar stage. Plant damage can be related to pest populations and insecticide applications. Severely damaged plants would certainly give poor grain yields. The insecticide sprays used in this study contributed substantially to grain yield increase in maize for the four seasons. These yield increases were however, not statistically significant in any of the seasons. Grain yield increases have been reported on cowpea after DDT applications [7]. The above results therefore show that although Lindane 20 e.c. reduced stem borer population and damage on maize thereby improving grain yields slightly, it also showed short-lived adverse effects on most of the early season beneficial arthropods which highlights the importance of detailed studies to monitor residues in the soils and on some of the beneficial arthropod fauna groups.

ACKNOWLEDGEMENTS

The authors are grateful to the Swedish International Development Agency (SIDA) for financial support and to the Director General, International Atomic Energy Agency (IAEA) Vienna, Austria for availing the funds. Sincere thanks to the Director, Tropical Pesticides Research Institute (TPRI), Arusha, Tanzania for providing facilities for the field work at the Institute. The assistance extended to us by Dr. P. Chiverton and other staff of the Swedish University of Agricultural Sciences (SUAS), Uppsala, Sweden is highly appreciated. Lastly but not least we are thankful to Mr. G. Kayange and Ms. G. Shayo for their keen supervision of all the field work.

REFERENCES

- [1] TANZANIA GAZETTE, 18th January 1991 pp. 29-63.
- [2] CAREY, A.E., GARVEN, J.A., TAI, H., MITCHELL, W.G., WIERSMA, G.B., Pesticide residue levels in soils and crops from 37 states - National Soils Monitoring Program (iv). Pesticide Monitoring Journal, **12** (1979) 209-233.
- [3] TABOR, E.C. . Contamination of urban air through the use of insecticides. Transactions of the New York Academy of Sciences, Series 2. **28** (1966); 569.
- [4] CRITCHLEY, B.R., COOK, A.G., CRITCHLEY, U., PERFECT, T.J., RUSSELL-SMITH, A.. The effects of crop protection with DDT on some elements of the subterranean and surface active arthropod fauna of a cultivated forest soil in the humid tropics., Pedobiologia **20** (1980) 31-38.
- [5] EDWARDS, C.A.. Environmental aspects of the usage of pesticides in developing countries. Med. Fac. landbouw. Rijksuniv. Gent 42/2. (1977) pp. 853-68.

- [6] COOK, A.G., CRITCHLEY, B.R., CRITCHLEY, U., PERFECT, T.J., RUSSELL-SMITH, A., YEADON, R., The effects of soil treatment with DDT on the biology of a cultivated forest soil in the sub-humid tropics., *Pedobiologia* **19** (1979) 279-292.
- [7] PERFECT, T.J., COOK, A.G., CRITCHLEY, B.R., CRITCHLEY, U., DAVIES, A.L., SWIFT, M.J., RUSSELL-SMITH, A., YEADON, R.. The effect of DDT contamination on the productivity of a cultivated forest soil in the semi-humid tropics. *Journal of Applied Ecology* **16** (1979) 705-719.

- [10] SINGH, S.R., TAYLOR, T.A., Pests of grain legumes and their control in Nigeria: (Singh, S.R., Van Emden, H.F., Taylor, T.A., Eds), Pests of Grain Legumes: Ecology and Control. Academic Press. London.
- [12] PERFECT, T.J., COOK, A.G., CRITCHLEY, B.R., CRITCHLEY, U., DAVIES, A.L., SWIFT, M.J., RUSSELL-SMITH, YEADON, R., The effect of DDT contamination on the productivity of a cultivated forest soil in the subhumid tropics. J.Appl. Ecol. 16 (1979) 705-719.

**NEXT PAGE(S)
left BLANK**

FATE AND DISTRIBUTION OF LINDANE AND ENDOSULFAN IN MAIZE AND COWPEA ECOSYSTEMS RESPECTIVELY



XA9743719

P.O. YEBOAH
Department of Chemistry,
Ghana Atomic Energy Commission

K.G. MONTFORD
Biotechnology and Nuclear Agriculture Research Institute,
Ghana Atomic Energy Commission

R.K.KLAKE, C.K AKPABLI
Department of Chemistry,
University of Ghana

Legon, Ghana

Abstract

A quantitative study is presented on lindane and endosulfan residues in maize and cowpea ecosystems respectively. Both pesticides were found to dissipate very fast under the tropical Ghanaian conditions. The high rate of dissipation in leaves is attributed to the fact that the leaves were exposed to sunshine and wind leading to increased volatilisation. Endosulfan was found to dissipate faster from the cowpea ecosystem than lindane did in the maize ecosystem. The mean residue levels of lindane in maize grains were $0.02 \mu\text{g g}^{-1}$; whilst residue levels of endosulfan in cowpea seeds were $0.05 \mu\text{g g}^{-1}$. These levels are lower than the maximum residue limits recognised as acceptable by the Codex Alimentarius Commission.

1. INTRODUCTION

As a result of increasing world population, there has been a proportionate increase in the demand for food. Currently, 58 percent of the world's population is undernourished [1]. There is therefore the need to increase food production. This can be achieved through the application of such products of modern technology, as high yielding crop varieties, fertilisers, irrigation and pesticides. Pesticides help boost food production by reducing pest populations to tolerable levels. In 1986, it was found that crop loss due to pests was 40%. Cotton yields have also been found to increase three fold through effective use of chemical pesticides for pest control in the Sudan [2].

Despite the benefits derived from pesticides, some of them tend to persist in the environment, and may thus pose a hazard to living things. Organochlorine pesticides constitute one group of such pesticides that are persistent. Examples of these include DDT, lindane and endosulfan. These compounds have also been found to (i) be highly fat soluble and hence accumulate in fatty tissues of animals, (ii) accumulate in food chains, and (iii) degrade into other toxic residues which can have long term adverse effects on the environment [3,4].

In Ghana, the surviving organochlorines in use are lindane, endosulfan, aldrin and dieldrin. Lindane, formulated under the trade name Gammalin 20, has been restricted to be used only in the cocoa growing industry. However, due to mixed cropping, other crops end up by being sprayed with lindane during its application. These crops include maize, cocoyam and cassava. Under such conditions, lindane can serve to control insects that bore into stems of the maize plant.

Notwithstanding the hazards posed by organochlorine pesticides to the environment and being banned or restricted in temperate regions, developing countries including Ghana, continue to use them to boost food production. They are also used in public health to control disease vectors. Their continued use has been attributed to their high efficiency, low cost, and easy availability. Furthermore, owing to the intense heat and high humidity in the tropics, organochlorine pesticides dissipate at faster rates than in temperate regions. Consequently, it is suggested that, there should be less danger of possible accumulation of their residues in soils, plants and animal matter, that could lead to the precipitation of environmental hazards.

Whereas considerable data on pesticide residues are available in the developed countries to assist in decision making regarding the continued use of particular pesticides, knowledge on pesticide residues and their effects are rarely available in many developing countries. Hence the need to generate data on the effects of these pesticides in various agroecosystems in the tropics to allow correct decisions on their use. These should include extensive knowledge about their fate, persistence, distribution, effect on yield, and the way they affect target and non target organisms to enable an assessment of their potential for human and environmental hazards.

Maize is a staple Ghanaian food which is taken in many forms. It may be boiled, roasted, or ground into dough. Since lindane is applied to maize, a knowledge of lindane residues in maize will therefore be of immense importance to Ghanaians. Cow peas are also taken in Ghanaian homes in the boiled form. They are also ground and fried. In northern Ghana, the leaves are used to prepare soup [5].

This report therefore covers studies to identify and quantify lindane and endosulfan residues in maize and cowpea ecosystem respectively, under Ghanaian conditions.

2. EXPERIMENTAL

2.1. Apparatus

A Hewlett packard gas chromatograph series II, model 5890, equipped with a ⁶³Ni electron-capture detector and a SPB 5 capillary column (30 m × 0.5 mm id) was used for the study. The carrier gas, nitrogen, had a flow rate of 15 mL min⁻¹. The operating temperatures were: injector, 200°C; detector, 300 °C; and column, 200° C isothermal. The attenuation was set on the HP 3396A recording integrator at 8, while chart speed was 10 mm min⁻¹. Each injection had a volume of 1 µl.

Solid Phase Extraction (SPE) Bond Elute C-18 columns (500 mg in 2.8 mL) were purchased from Analytical Instruments Varian GmbH, Vosendorf, Austria.

2.2. Reagents and materials

All solvents were Nanograde, and purchased from Mallinckrodt Specialty Chemical Co. Paris, Kentucky, USA. All other reagents were Analar grade. Commercial lindane formulated as an emulsifiable concentrate under the trade name Gammalin 20 was obtained from ICI Ghana Ltd, whilst endosulfan formulated under the trade name Thiodan 35 was obtained from Hoechst Aktiengesellschaft Frankfurt, Germany. Cypermethrin was obtained from Shell Chemicals, Abidjan, Cote d' Ivoire.

The maize planted was a local variety called "Abelehi" which matures in 107 days; and the cowpea a variety called "Asutem". Lindane and endosulfan standards were obtained from the International Atomic Energy Agency (IAEA) in Vienna.

2.3. Experimental farm

The experiment was conducted on a 1.6 ha. plot situated at the Ghana Atomic Energy Commission, Kwabanya, which is about 20 km from the centre of Accra. The plot had been treated with 1 kg AI of lindane the previous year. Four plots each measuring 50 m × 50 m were marked out. Two diagonally opposite plots were sown with maize whilst the others were sown with cowpeas.

2.4. Pesticide treatment

The maize plot that had been earmarked for pesticide treatment was sprayed twice with a water emulsion of the commercial lindane at a rate of 0.5 kg ha⁻¹ in accordance with local practices. The split application was done to avoid the resurgence of insect pests that may survive the first application.

Pesticide treatment of the cowpea plot was also in accordance with the recommendations of the Plant Protection and Regulatory Services of Ghana's Ministry of Food and Agriculture. Two earlier treatments were carried out with the synthetic pyrethroid cypermethrin at a dose of 50 g AI ha⁻¹. per each treatment. This application was to control insect pests that destroy cowpea flowers. These were followed by the application of endosulfan (Thiodan 35) at a dose of 750 g AI ha⁻¹. The endosulfan application was to control the pod borers that destroy cowpea seeds. The weeding, planting, spraying and harvesting schedules are as shown in Table 1.

Table 1. Schedule of activities

Activity	Crop	Date	
		Control	Treated
planting	maize	2/6	3/6
	cowpea	17-18/6	19-20/6
spraying, lindane	maize	-	17/6
	maize	-	1/7
spraying, cypermethrin	cowpea	-	21/7
	cowpea	-	31/7
spraying, endosulfan	cowpea	-	12/8
weeding, manual	maize	21/6	23/6
	cowpea	4-6/6	2-3/6
fertiliser application	maize	22/6	25/6
harvesting	cowpea	9/9	9/9
	maize	24/9	24/9

2.5. Soil characteristics

2.5.1. Sampling

The method followed the FAO/IAEA protocol [Appendix I]. Eight cores of soils were sampled at random from each sub-plot using a soil auger of 25 mm diameter and a depth of 150 mm. These were combined, ground and passed through a 2 mm sieve by mechanical shaking. Duplicate sub-samples were then taken for the measurement of pH, moisture content, organic carbon and particle size.

2.5.2. Determination of pH

The method followed that prescribed by Davies[6]. A sample (20 g) of the air dried sample was washed into of water (100 mL). The suspension was stirred with a mechanical stirrer for 5 minutes and allowed to stand for 30 minutes. The pH was determined using a meter which had been calibrated with pH4 and 7 standard buffers.

2.5.3. Determination of soil moisture

The soil sample were weighed into moisture cans and dried in an oven at 105°C to constant weight. The moisture content was determined as a percentage loss in weight.

2.5.4. Determination of organic carbon

The method followed that prescribed in Appendix I. A sample of soil (0.5 g) passed through a 0.5 mm mesh sieve was added to 1M potassium dichromate solution (10 mL). Concentrated sulphuric acid (20 mL) was added to the suspension, thoroughly swirled and allowed to stand on asbestos for 30 minutes. Distilled water (100 mL) was added, followed by orthophosphoric acid (10 mL) and finally barium diphenylamine sulphonate indicator (2 mL). The resulting solution was then titrated with 0.2M ferrous ammonium sulphate solution until the solution changed from blue to green. A blank titration was carried out to standardise the potassium dichromate.

2.5.5. Particle size analysis

Particle size analysis was determined by standard hydrometer method following dispersion with sodium hexametaphosphate, as prescribed by Charmetski [7] and Dickson [8].

2.6. Residue analysis

2.6.1. Sampling

2.6.1.1. Soils

Triplicate samples (50 g) were taken for residue analysis. For lindane, sampling was made at the following times (crop growth stages): pre-treatment (4th leaf stage), day of first treatment, 1 day (ankle stage) and 14 days after first treatment (knee high stage). Samples were also taken on the day of second treatment, followed by 14, 30 and 60 days (post tusseling stage) after the second treatment. For the cowpea plots (including the endosulfan experiment), sampling was done on the pre treatment day (flowering stage), 1, 7 (pod formation stage), 14 (pod drying stage), and 22 (harvesting stage) days post-treatment.

2.6.1.2. Leaves

For the maize ecosystem, the youngest and oldest leaves were taken off randomly chosen plants from each of the sub-plots as follows: pre-treatment day of first treatment 1, 14 days after first treatment. Samples were also taken on the day of second treatment, followed by 14 and 30 days after second treatment. With cowpeas, leaves were randomly sampled from the various sub-plots as follows: 1, 7, 14 and 22 days post-treatment.

2.6.2. *Extraction and clean up*

2.6.2.1. Soils

The triplicate samples were accurately weighed and transferred into a single thickness cellulose extraction thimble, (43 mm i.d. × 123 mm long) and extracted in a Soxhlet through 10 cycles i.e. for ~5 h with methanol (250 mL). An aliquot (5 mL) of the methanol extract was evaporated to dryness over nitrogen. The residue was taken up in methanol (1 mL) and to this was added water (2.5 mL). Prior to this, a C-18 Bond Elut solid phase extraction column had been reconditioned by flushing through two volumes of methanol followed by two volumes of distilled water. The diluted extract was poured through the reconditioned column at a flow rate of 2 mL min⁻¹ and then washed with methanol + water (3+7 by volume, 1 mL) followed by distilled water (1 mL). The washed column was vacuum dried for 15 min and then the pesticide was eluted from the column with hexane (2 mL) into a glass vial. The eluent was made up to 2 mL with more hexane and then analysed by gas chromatography.

2.6.2.2. Leaves and seeds

Leaf tissue (20 g) and/or seeds were homogenised separately for 2 minutes in a Waring blender and mixed thoroughly with methanol (200 mL). The suspension was filtered and a volume (20 mL) of the methanol extract was dried over nitrogen and the residue was dissolved in methanol (1 mL) and diluted with water (2.5 mL). The aqueous methanol extract was passed through a preconditioned SPE column at a flow rate of 2 mL min⁻¹. The column was then washed with distilled water (1 mL). The washed column was vacuum dried for 15 min and then eluted with hexane (1.5 mL) into glass vials. The eluent was made up to 2 mL with hexane and analysed by gas chromatography.

2.6.3. *Recovery experiments*

2.6.3.1. Soils

Samples of pesticide-free soils (50 g) were air dried and sieved as before. Volumes (1 mL) of lindane and endosulfan standards (10 µg mL⁻¹) were added to the soils. The soil was then extracted using a Soxhlet for 5 hours (10 cycles) with methanol. The extracts were then cleaned up for GLC measurements as in Section 2.6.1.

2.6.3.2. Seed and leaves

Samples (10 g) of maize grains, cowpea seeds, maize leaves and cowpea leaves were homogenised using Waring blender. A volume (1 mL) of standard solutions (10 µg mL⁻¹) was added to the homogenised samples (maize leaves and grains, lindane; cowpea seeds and leaves, endosulfan). Methanol (100 mL) was added to each mixture and shaken mechanically for thorough mixing. The methanol extract was filtered and a volume (20 mL) of the extract cleaned up for GLC analysis as in Section 2.6.2.

3. RESULTS AND DISCUSSION

3.1. Recoveries and soil characteristics

Though the plot had been exposed to pesticides only a year before the investigation, initial analysis before application of any pesticide showed the absence of pesticide residues within the limit of detection of the gas chromatograph ($0.01 \mu\text{g g}^{-1}$). All blank samples were found to be below the detection limit. Spiked soil samples gave recoveries of 80.5 and 92.7% for lindane and endosulfan residues, respectively. Spiked leaf samples gave recoveries of 95.2 and 80.5% for lindane and endosulfan residues, respectively.

The persistence of pesticides in soils is known to depend on soil characteristics. The major ones include organic matter, moisture, texture and acidity. The soil analysis showed the soils to be slightly acidic with an average pH of 6.3 and had a sandy clay loam texture (Tables 2 and 3). The average moisture content of the soils from the four subplots during the sampling period ranges from 4.78-13.3 g kg^{-1} and 3.40-7.74 g kg^{-1} in the maize and cowpea soils respectively.

3.2. The maize ecosystem

The dissipation of lindane in maize soils was very fast (Table 5) with 46% of the initial residue detected on the day of treatment lost one day after the first application (6th leaf

Table 2 Soil characteristics of the maize plots.

Property ^{a)}	Plot	
	Control (SD)	Treated (SD)
pH	6.21 (0.17)	6.3 (0.15)
% sand	64.2 (5.4)	63.8 (7.3)
% silt	13.4 (2.8)	11.1 (3.9)
% clay	24.5 (4.7)	25.0 (3.8)
% organic matter	2.4 (0.8)	1.9 (0.8)
% organic carbon	1.4 (0.5)	1.1 (0.5)

a) texture class : sandy clay loam

Table 3. Soil characteristics of the cowpea plots

Property ^{a)}	Plot	
	Control (SD)	Treated (SD)
pH	6.4 (0.3)	6.3 (0.2)
% sand	57.0 (7.6)	63.0 (3.2)
% silt	12.2 (2.4)	11.0 (1.6)
% clay	29.0 (5.0)	26.2 (4.0)
% organic matter	2.0 (0.4)	2.2 (0.3)
% organic carbon	1.8 (0.7)	1.3 (0.2)

a) texture class : sandy clay loam

stage), and less than 5% remained after 14 days. This may be attributable to volatilisation under the hot windy and rainy conditions of the area (Table 4). Increase in temperature is known to increase pesticide loss from soils through volatilisation, chemical degradation and bacterial decomposition, all of which are temperature dependent [9].

Rainfall might also have contributed to the pesticide loss by causing surface run-off. Furthermore, water competes with pesticides for adsorption to the soil surface [10] and lindane will therefore be more easily volatilised from surfaces of soil that have high moisture content. The moisture content of 5.8 to 13.8 observed during the period of investigation were fairly high and could have been a contributing factor to the fast dissipation rate.

About 5% of the residue detected on the day of first treatment remained in the soils 14 days (knee high stage) after application (Table 5). This is less than the 13% of the residue detected in soils 14 days after the second application (above knee high stage). This relatively slower rate of lindane dissipation after the second application might have been due to the fact that the maize plants, which had matured at the time of the second spraying provided cover over the top soil and lowered temperatures and minimised loss through volatilisation. Only 2.8% of the initial residue of lindane remained 30 days after the second application.

Thirty percent of the initial lindane residues detected in the maize leaves on the day of first treatment remained one day after application and fell to 3% 14 days after the first application (Table 5). Similarly, 14 days after the second application of lindane, the residues had fallen to 1.7% of the initial level.

Table 4. Monthly total rainfall and mean temperatures

	May	June	July	August	September
total rain, mm	49	81	41	17	75
mean temp. °C	28.5	26.9	25.5	25.3	26.4

Table 5 Lindane residues in soil samples and plant samples of the maize ecosystem

Sampling time	Mean lindane residue, $\mu\text{g g}^{-1}$ (SD)	
	Soil	Plant leaves
pretreatment	nd	nd
day of treatment	0.22 (0.04)	0.22 (0.014)
1 day after treatment	0.12 (0.05)	0.067 (0.01)
14 days after treatment	0.011 (0.008)	0.006 (0.002)
day of second treatment	0.025 (0.013)	0.24 (0.02)
14 days after second treatment	0.032 (0.006)	0.004 (0.001)
30 days after second treatment	0.008 (0.003)	nd
60 days after second treatment	nd	-

nd = none detected

Mean lindane residues levels in the maize grains were $0.02 \mu\text{g g}^{-1}$ (Table 7), which is well below the acceptable maximum residue limit for grains of $1 \mu\text{g g}^{-1}$ [10].

Table 6. Endosulfan residues in soil samples and leaf samples of the cowpea ecosystem

Sampling time	Mean endosulfan residue, $\mu\text{g g}^{-1}$ (SD)	
	Soil	Plant leaves
pretreatment	nd	nd
day of treatment	0.29 (0.17)	0.33 (0.022)
7 days after treatment	0.022 (0.011)	0.011 (0.006)
14 days after treatment	0.008 (0.005)	0.002 (0.001)
22 days after treatment	nd	nd

nd = none detected

Table 7. Lindane and endosulfan residues in maize and cowpea grains

Crop	Mean residues, $\mu\text{g g}^{-1}$ (SD)
maize	0.020 (0.002)
cowpea	0.045 (0.002)

3.3 The cowpea ecosystem

The amount of endosulfan in cowpea soils at 7 days after spraying (i.e. pod formation stage) was found to be only 7.6% of the residue detected on the day of treatment, similar to that found with lindane in the maize soils (Table 6). The low persistence of endosulfan as explained in the section for the maize ecosystem can also be attributed to the low organic matter content of the soil (ranging from 1.91 to 2.18%), the fairly high temperature, the sandy clay loam texture of the soil as well as the fairly heavy rainfall (Tables 3, 4). The mean residue level of endosulfan in cowpea seeds was $0.045 \mu\text{g g}^{-1}$ (Table 7). This is higher than the residue levels of lindane in maize grains and could be due to the fact that endosulfan was applied close to the time of seed formation, making it possible for endosulfan to be quickly incorporated into the seeds before it could dissipate.

CONCLUSION

Lindane and endosulfan have been shown to dissipate quickly under Ghanaian climatic conditions which are characterised by heavy rainfall and fairly high temperatures. The residue levels of lindane in maize and endosulfan in cowpea were well below the acceptable maximum residue levels.

ACKNOWLEDGEMENTS

This work was supported by the SIDA through the International Atomic Energy Agency under Research Contract No. 6754. The authors also acknowledge the contribution of Messrs B.Q. Modzinuh and S. Afful of the Chemistry Dept. of the NNRI for their assistance during sampling.

REFERENCES

- [1] WARE, G. W., Pesticide, Theory and Applications. W. H. Freeman & Co., San Francisco (1978) 14.
- [2] FAO International Code of Conduct on the Distribution and Use of Pesticides (1986) FAO, Rome.
- [3] IAEA, Isotope Techniques for studying the Fate of Persistent Pesticides in the Tropics, IAEA-TECDOC-476, Vienna (1988).
- [4] FAO/IAEA Technical Report Series No 329 (1991) Vienna.
- [5] Personal Communication with people from Northern Ghana.
- [6] DAVIES, Soil Sci. **22** (1943) 405.
- [7] CHARMETSKI, W.A., LICHTENSTEIN, E.P., J. Econ. Entomol. **66** (1973) 344-349.
- [8] DICKSON, K.B., BENNEH, G., A New Geography of Ghana (1988) Longman, London 22-26.
- [9] EDWARDS, C.A., Residue Reviews (1966) **13** 83-132.
- [11] Codex Alimentarius Commission. Guide to Codex recommendations concerning pesticide residues (1986) 3rd prelim. issue FAO/CAC/VOL 8 Ed 2.

RESIDUES OF LINDANE AND ENDOSULFAN IN WATER AND FISH SAMPLES FROM RIVERS, FARMS IN BESEASE, AGOGO AND AKOMADAN IN THE ASHANTI REGION OF GHANA



XA9743720

S. OSAFO-ACQUAAH
Department of Chemistry,
University of Science and Technology

E. FRIMPONG
Department of Biological Sciences,
University of Science and Technology

Kumasi, Ghana

Abstract

Pesticide residue analyses were performed on water and fish samples from River Oda in Besese, River Afranso in Nobewam near Kumasi, River Atwetwe in Akomadan, and River Kowire at Agogo. Residues of lindane and endosulfan were found in water and fish (*Oreochromis niloticus*, *Tilapia zillii*, *Barbus trispulis*, *Heterobranchus sp.*, *Tilapia busumana*, *Ophiocephalus obscura* and *Chana obscura*) samples. The residues of lindane varied between the years and months in the year but were in the range of 0.3 - 15 ng L⁻¹ (1993-94) and 8.7-32.0 ng L⁻¹ (1995) for the water samples and 0.2 - 24 ng g⁻¹ (1993-93) and 8.4-120.4 ng g⁻¹ (1995) for the fish samples. Residues of endosulfan in the water and the fish samples were zero in 1993-1994 but, in 1995, were in the range of 6.4-35.2 ng L⁻¹ for the water samples and 5.0-267.5 ng g⁻¹ for the fish samples. In all cases the lindane and endosulfan concentrations in the water were 10,000-20,000 times lower than known toxic concentration levels and therefore unlikely to cause fish toxicity problems.

1. INTRODUCTION

Organochlorine pesticides have found wider application in agriculture for the control of pests and in public health for vector control [1]. Furthermore, there has been an increase in pesticide use in Ghana over the past ten years as a result of the efforts by the government to boost food production.

In agriculture, organochlorine pesticides are widely used by farmers due to their cost effectiveness and their broad spectrum of activity, with a neglect of their possible impact on the environment and man. Research in the temperate regions and some parts of Africa have found the pesticides to be very persistent in the environment and remain in the soil [2] for longer periods of time resulting in serious damage to soil flora and fauna [3].

Detailed information on the effects of organochlorine pesticide residues and the amounts present in water, food, soil, flora and fauna in Africa is however, very limited. In some African countries, organochlorine pesticides have been banned based on data obtained from the developed countries in the temperate regions, without any assessment of the effect of these pesticides in the tropical environment [4].

Lindane (Gammalin 20) is widely used in Ghana in the cocoa plantations for the control of capsids on vegetables and for the control of stemborers in maize [5]. Endosulfan is used in cotton-growing areas, as well as on vegetable farms and for the control of pests in coffee plantations.

Most of the vegetable farms in the commercial vegetable producing areas in Ghana like Akomadan and Agogo are situated along rivers which act as the source water supply for farming and for drinking purposes. Pollution of rivers is known to occur through run-off of pesticides from farms during heavy rain as well as through aerial spraying of pesticides and the improper disposal and handling of the chemicals during and after spraying.

Endosulfan and lindane are known to be toxic to fish. For example, lindane has an LC₅₀ (48h) of 0.16-0.3 mg L⁻¹ for guppies [6] and an LC₅₀ (96h) values of 1.2 mg L⁻¹ and 0.75 mg L⁻¹ for *Fundus heteroclitus* and *Oreochromis niloticus* respectively [7, 8].

This report is based on our work on the effects of organochlorine pesticide residues in water and fish, in the forest zone of Ghana, as part of the coordinated research work of the Joint IAEA/FAO research programme on the 'Adverse effects on flora and fauna from the use of organochlorine pesticides on the African continent'.

2. MATERIALS AND METHODS

2.1. Experimental materials

Pesticide grade hexane was obtained from Mallinckrodt Special Chemicals, Kentucky, USA. HPLC grade methanol and acetonitrile were obtained from Aldrich Chemical Company, USA. Bonded phase C-18 SPE columns were obtained from Varian, Sunnyvale, USA.

2.2. Experimental site

This general study covered about 500,000 hectares of land in the closed forest region of Ghana which is about 34.5% of the entire land area of Ghana. Three regions of intense farming activities were selected for the residue analysis work after the general survey work had established the nature of the agricultural activities undertaken in the areas under consideration and the extent of pesticide usage.

The farming community at Akomadan is involved in large-scale vegetable farming in addition to maize farming with tomato farming constituting about 80% of the farming activity in the area. Besease and Nobewam are used mainly for rice production and are the location of the Anum Valley Irrigation Project. Close to the rivers are over 200 rice fields covering more than 400 hectares of land. Agogo is noted for its vegetable farming, rice and maize production and has other surrounding villages involved in cocoa farming.

2.3. Sampling

The pH values of the river waters were in the range 7.0-8.2 and the average temperature was 28°C. Water samples were collected at a depth of 250mm from River Oda at Besease, River Aframso at Nobewam near Kumasi, River Kowire at Agogo and River Aframso and River Atwetwe at Akomadan in the Ashanti Region in one litre sterilized bottles closed with a stopper. The duplicate samples were collected between January 1993 and July 1994 and from June 1995 to October 1995. The rivers pass through regions which have intense cocoa production and other farming activities. The initial stage of work, during 1993-94, involved random sampling in the various rivers. In 1995, a much more systematic sampling was undertaken from June to October and involved the farming communities at

Agogo and Besease. No samples were taken from Nobewam because neither lindane nor endosulfan were being used in the farming in this area. The river Oda at Besease flows through several areas of cocoa farming where lindane is actively used. Endosulfan is used in the farming at Agogo while other nearby cocoa farms use lindane.

After collection the water samples were taken to the laboratory for analysis. Fish samples for pesticide analyses were collected from River Oda near the rice fields and River Aframso in Agogo where there is large scale vegetable production. The fish samples (seven species) were obtained by a hook and line method and also with cages and nets. The fish samples obtained were immediately washed with distilled water and stored in a freezer.

The samples were later taken to the Ghana Atomic Energy Commission laboratory and the Institute of Aquatic Biology in Accra, Ghana for pesticide residue analysis.

2.4. Sampling of other species

Although several varieties of birds, rodents (rats, squirrels, grasscutter) and reptiles (lizards) were identified, the project focused on residues in fish. The rodent (grasscutter) samples, which were collected from the area, are yet to be analysed.

2.5. Laboratory analytical procedures

2.5.1. Water residue analyses

Volumes (2 x 10 mL) of methanol followed by volumes (2 x 10 mL) of distilled water were passed through an SPE C-18 column via a reservoir mounted on top of the column before conditioning. A small pressure was applied to the top of the reservoir to help force each volume of liquid to flow through the column at a rate of at least 2 mL per minute.

A volume (800 mL) of each water sample was passed through a pre-conditioned column at a flow rate of 2 ml per min. Each column was then washed with water + methanol (1 mL, 3.3 + 1 by volume) followed by distilled water (1mL) and dried under vacuum for 20 minutes. The sample which was trapped in the column was eluted with hexane (5 x 0.5 mL) to recover the pesticide residues.

2.5.2. Fish residue analyses

A sample (10 g) of each fish was homogenized using a pestle and mortar. The sample was mixed with sodium sulphate (10 g) and then transferred to a separatory funnel (250 mL). Hexane (20 mL) was added followed by acetonitrile (100 mL) which had been saturated with hexane, and the mixture was shaken for about one minute. The acetonitrile was drained into a 1 litre separatory funnel. The mixture was further extracted with portions of acetonitrile (3 x 50 mL) and the extracts combined with those in the separatory funnel. These were washed with water (500 mL) followed by saturated sodium chloride solution (40 mL) and hexane (50 mL). The mixture was allowed to stand and the aqueous layer was drained into another 1 litre separatory funnel and extracted with a further volume (50 mL) of hexane. The hexane extracts were combined and passed through a plug of sodium sulphate (10 g) into a flask. The solvents were evaporated almost to dryness and the residue dissolved in methanol (10 mL) and diluted with water (25 mL). A volume (3.5 mL) of this solution was taken for pesticide residue analysis (equivalent to 1 g of the fish sample) and passed through a

previously pre-conditioned column at a flow rate of 2 ml per minute. The column was washed, dried and eluted with hexane as described for the water samples above. Volumes (1 µL) of this hexane eluent were taken for analysis by the gas chromatographic analysis method detailed in Section 2.5.3.

2.5.3. Gas chromatographic analysis

Gas chromatograph : Hewlett-Packard Series II model 5890; column : 30 m x 0.5 mm i.d.; stationary phase : SPB-5; carrier gas: nitrogen; flow rate : 30 ml min⁻¹; temperatures: column, 200°C, injector, 250°C, detector, 300°C; detector, ⁶³Ni electron capture. The sample injection volume was 1 µL. The limit of detection in this work was 1pg mL⁻¹. Recovery of lindane and endosulfan from the fish and water samples was 90%. Lindane and endosulfan in the residues were identified only by comparison of the retention times with those of their analytical standards and confirmation by an alternative method was not attempted.

3. RESULTS AND DISCUSSION

Seven species of fish caught in the various rivers were subjected to residue analysis. The species of fish caught depended on the time of the year when the trapping was done. It appeared that due to unknown adverse effects of some material, the fish populations had decreased considerably in these rivers. There were whole days when our team could not trap any fish in some of the rivers using a combination of a hook and line method, cages and nets.

Our initial survey work indicated that farms near River Oda at Besease were sprayed up to six times during the year. Farms near River Atwetwe and River Kowire which were along tomato-producing areas were subjected to pesticide applications of up to six times over either one or two seasons. Although lindane is not used widely in the spraying of tomatoes and rice, endosulfan is one of the commonly used insecticides in these regions. Despite this, the levels of endosulfan in the water samples were below the limits of detection (1 pg mL⁻¹). The concentrations of both lindane and endosulfan, when present, were found to be significantly higher in the fish than in the water (cf Tables 1, 2 with 3, 4) though the pesticide levels were well below their lethal levels, as discussed later.

Residues whose retention times corresponded with DDT and other organochlorines were observed in the chromatogram but were not quantified since the focus of this project was on lindane and endosulfan.

Table 1. Pesticide residues in river water samples during 1993-94

Sample		Residue, ng L ⁻¹	
river identity	date	lindane	endosulfan
Oda	15.01.93	14.2	0
Oda	21.12.93	15.0	0
Atwetwe	15.01.94	1.3	0
Aframso	12.03.94	3.9	0
Kowire	07.07.94	3.0	0
Aframso	22.07.94	0.3	0

Table 2. Pesticide residues in fish caught during 1993-94

Source of fish		Species	Residue, ng g ⁻¹	
river identity	date		lindane	endosulfan
Oda	15.01.93	<i>Heterobranchus sp</i>	14.2	0
Oda	21.12.93	<i>Heterobranchus sp</i>	24.3	0
Atwetwe	15.01.94	<i>Barbus trispulis</i>	23.2	0
		<i>Chana obscura</i>	0.8	0
Aframso	12.03.94	<i>Tilapia sp</i>	0.2	0
		<i>Barbus tripulis</i>	0.2	0
Kowire	07.07.94	<i>Ophiocephalus obscura</i>	1.5	0

Table 3. Pesticide residues in river water samples during 1995

Sample		Residue, ng L ⁻¹	
river	month	lindane	endosulfan
Oda	June	25.0	35.2
	July	24.0	30.0
	August	22.0	15.0
	September	32.0	14.0
	October	8.7	6.4
Kowire	September	10.0	15.4
	October	14.8	25.6
Atwetwe	September	12.5	28.2
	October	20.2	14.6

Table 4. Pesticide residues in fish samples caught during 1995

Source of fish		Species	Residue, ng g ⁻¹	
river	month		lindane	endosulfan
Oda	June	<i>Oreochromis niloticus</i>	20.0	80.2
		<i>Barbus trispulis</i>	120.4	158.0
		<i>Heterobranchus sp</i>	14.0	160.0
	July	<i>Oreochromis niloticus</i>	22.5	267.5
		<i>Tilapia zilli</i>	34.0	142.5
		<i>Heterobranchus sp</i>	35.2	64.0
	August	<i>Oreochromis niloticus</i>	36.4	20.2
		<i>Barbus trispulis</i>	16.8	11.4
		<i>Heterobranchus sp</i>	24.8	6.5
	September	<i>Oreochromis niloticus</i>	100.0	186.0
		<i>Tilapia zilli</i>	72.0	77.0
		<i>Heterobranchus sp</i>	80.0	66.0
Kowire	October	<i>Oreochromis niloticus</i>	38.2	5.0
		<i>Barbus trispulis</i>	22.0	10.0
	September	<i>Oreochromis niloticus</i>	8.4	12.4
	October	<i>Oreochromis niloticus</i>	13.4	22.8

In none of the samples of river water collected during 1993 and 1994 was endosulfan found while lindane was found at low concentrations up to 15 ng mL^{-1} (Table 1). During these two years no endosulfan residues were found in the fish samples but lindane was found at concentrations up to 24.3 ng g^{-1} . This represents over 1000 fold concentration of lindane by the fish and is in general accord with, though exceeding, the known bioconcentration factor (153) for lindane and fish in water. Although the toxic levels of lindane to these particular species of fish are not known, the fact that they were surviving is not too surprising since the actual concentrations of lindane found in the water were around 10,000 to 20,000 times below the toxic concentrations to another species of fish (guppy, LC50 (48 h), $160,000 - 30,000 \text{ ng L}^{-1}$) [6].

In the more complete studies of 1995, both lindane and endosulfan were found in the river water samples. Concentrations varied with the location and the month of sampling but were in the ranges $8.7-25.0 \text{ ng L}^{-1}$ and $6.4-35.2 \text{ ng L}^{-1}$ for lindane and endosulfan respectively. As in the previous two years, lindane residues were found in the fish samples, ranging from $8.4-120.4 \text{ ng g}^{-1}$. These levels were generally higher than those of the previous two years but still orders of magnitude below likely toxicity levels.

In contrast to previous years, endosulfan was found in the fish samples and often at concentrations higher than lindane. In fact this is in general accord with its known bioconcentration factor for fish (583) which is approximately four times higher than that for lindane. In the case of endosulfan the aqueous concentration toxicity levels against local species of fish are known (LC50 (96 h), $1,200,000 \text{ ng L}^{-1}$, *Fundus heteroclitus*; $750,000 \text{ ng L}^{-1}$, *Oreochromis niloticus*). It can be seen that the concentrations in the river water samples are over 20,000 times below these toxicity levels and therefore present no toxic hazard to the fish.

So, in conclusion, although both lindane and endosulfan were found in river waters and it could be seen that the fish in these rivers absorbed and concentrated these insecticides, the levels that were actually found were so far below known toxicity levels that it is unlikely that they were the cause of the perceived low populations of fish in the rivers and that must be attributed to another cause.

ACKNOWLEDGEMENTS

We are grateful to the International Atomic Energy Agency for their grant support for this project and Dr. C.A. Edwards for his comments and assistance. We also acknowledge with gratitude the assistance of the Ghana Atomic Energy Agency and the Institute of Aquatic Biology, Accra in performing the residue analysis.

REFERENCES

- [1] EDWARDS, C.A., "Pesticide residues in soil and water", Environmental pollution by pesticides (Edwards, C.A., Ed), pp 409-458. Plenum Publ. New York (1973).
- [2] WHO technical report series, Research into Environmental pollution, No. 406, Geneva (1968).
- [3] COPE, O.B., Contamination of the fresh water ecosystem by pesticides, J Appl. Ecol., 3 (1966) 33-34.

- [4] TALEKAR, N.S., CHEN, J.S., KAO, H.T., Long term persistence of selected insecticides in subtropical soil: their adsorption by crop plants, *J. Econ. Entomol.*, **76** (1983) 207.
- [5] Ministry of Agriculture Technical Bulletin No.GDB/M1 Ghana.
- [6] Pesticide Manual, 10th Edition (C. Tomlin, Ed) The British Crop Protection Council, Surrey, UK (1994).
- [7] FELTZ, A.R., National monitoring program for the assessment of pesticide residues in water, *Pesticide Monitoring Journal* **5** (1971)54-62.
- [8] GODZI, A., Undergraduate thesis, University of Science and Technology, Kumasi,Ghana.
- [9] NEGRO, J.J, DONAZAR, J.A., HIRALDO, F., HERNANDEZ L.M., FERNANDEZ, M.A., Organochlorine and heavy metal contamination in non-viable eggs and its relation to breeding success in a Spanish population of lesser Kestrels (*Falco naumanni*), *Environmental Pollution* **82** (1993) 201-205.
- [10] SAMUEL,T. PILLAI, M.K.K., Impact of repeated application on the binding and persistence of [¹⁴C]-DDT and [¹⁴C]-HCH in a tropical soil, *Environmental Pollution* **74** (1991) 205-216.
- [11] WALISZEWSKI, S. M., Residues of lindane, HCH isomers and HCB in the soil after lindane application, *Environmental Pollution* **82** (1993) 289-293.

**NEXT PAGE(S)
left BLANK**



STUDIES ON THE DISTRIBUTION, SALE, HANDLING AND USAGE OF AGROCHEMICALS IN ASHANTI, BRONG AHAFO AND EASTERN REGIONS OF GHANA

S. OSAFO-ACQUAAH

Department of Chemistry,
University of Science and Technology

E. FRIMPONG

Department of Biological Sciences, University of Science and Technology

Kumasi, Ghana

Abstract

Studies were carried on the distribution, sale, handling and usage of agrochemicals in 19 major towns located in three regions noted for their intensive agricultural activities and covering 500,000 hectares of land area in Ghana. The most prevalent of the agrochemicals were insecticides (52%) followed by herbicides (28%) and fungicides (16%). Of these the organophosphates constituted 24% and the organochlorines 6% in terms of the total numbers of pesticides sold, though lindane and endosulfan were among the insecticides most frequently used by farmers. The fungicide, copper oxychloride as "Kocide", and the insecticide, λ cyhalothrin as "Karate", with 39% and 22% respectively of the sales by volume, were found to be the most widely used agrochemicals. Most of the merchants of agrochemicals and the farmers had only primary school education and had very little knowledge on the handling and safety aspects of the use of agrochemicals. The Extension Service to the farmers was inadequate and the farmers depended largely on the agrochemicals merchants for information on pesticide usage.

1. INTRODUCTION

This study was undertaken to provide additional information on the application, handling and use of pesticides in Ghana. With the launching of the Economic Recovery Programme there has been considerable emphasis on the development of agriculture in Ghana and pesticides usage has been increasing during the past ten years. Owing to costs associated with these chemicals and the lack of adequate information on the usage and application of pesticides, deleterious effects on the environment have been largely ignored. Furthermore, major law enforcement agencies like the Environmental Protection Council, Plant Protection and Quarantine Division and the Custom, Excise and Preventive Service have very little information at their disposal for effective legislation on the use of pesticides and the management of pesticides in general. Most of the data available are based on studies in developed countries. The objective of the study is to increase our knowledge of the distribution, usage and handling of pesticides in Ghana as part of the coordinated research programme on the adverse effect on the use of organochlorines (lindane and endosulfan) on the African Continent.

The study covered about 500,000 hectares of land area in the Ashanti, Brong Ahafo and Eastern Regions of Ghana located within the closed forest vegetation zone of Ghana; this constitutes about 34.5% of the total land area of Ghana. Sixteen major towns, noted for their intensive agricultural activities, within these three regions were selected for the study.

These centres were Agogo, Juaso, Konongo, Ejisu, Kumasi and Mampong in the Ashanti Region; Akomadan, Ejura, Kintampo, Sunyani, Techiman and Wenchi in the Brong Ahafo Region; and Nkawkaw, Anyinam, Koforidua, Nsawam, Suhum, Asamankase and Oda in the eastern Region.

The major crops cultivated in these areas include cocoa, palm fruits, rice, maize, cowpea and vegetables such as tomatoes, okro, eggplant and pepper.

2. MATERIALS AND METHODS

Questionnaires specifically designed for the study were used to collect data on the distribution, sale and usage of agrochemicals from the selected towns within the three Regions. Seventy-eight agrochemical sellers and one hundred and ninety-five farmers were interviewed. Besides the interviews, visits were made to selected farms to obtain additional information on the application of the agrochemicals.

3. RESULTS

3.1. Types and distribution of agrochemicals

The agrochemicals on sale and used by the farmers in the three regions were made up of insecticides (56%), herbicides (29%) and fungicides (16%) (Table 1). Insecticides always constituted the highest percentage of agrochemicals used in each of the three regions.

On the basis of their chemical composition the available agrochemicals fall into the following groups of compounds: organochlorines, organophosphates, carbamates, synthetic pyrethroids, inorganic compounds (mainly fungicides) and a miscellaneous category with a variety of chemical structures (Table 2). The organophosphates and organochlorines make up 30% of the total number of agrochemicals while the newer pyrethroids form a surprisingly large proportion (15%) of the market taking over from the older organochlorines and, to some extent, the organophosphates. DDT was one of the agrochemicals also found to be in use, though it is supposed to be banned in Ghana.

In terms of the volume of purchases the most commonly bought agrochemical was found to be Kocide, a copper oxychloride fungicide, and one of the least commonly bought was Sumithion, the organophosphate insecticide, fenitrothion (Table 3).

3.2. Socio-economic and other characteristics of merchants of agrochemicals (retailers and local distributors)

3.2.1. Educational background

The highest proportion of the merchants were middle school leavers with the remainder having secondary or other type of final education (Table 4). None of the merchants had a tertiary education.

3.2.2. Distribution of merchants by sex and age

Males (76%) were, by far, the greater proportion of the 78 merchants in the survey. The majority of them were between the ages of 20 and 40 years.

Table 1. Types of Agrochemicals with respect to their use, by farmers from three regions in Ghana.

Insecticides	Herbicides	Fungicides
Actellic (pirimiphos methyl)	Gesaprim (atrazine)	Baygon (propoxur)
Cymbush (cypermethrin)	Bellator	Caocobre
Decis(deltamethrin)	Garlon (triclopyr)	Champion
Diazinon	Gramoxone(paraquat)	Cobox(copper oxychloride)
Dimethoate 40EC	Ronstar (oxadiazon)	Dithane (mancozeb)
Dursban (chlorpyrifos)	Roundup(glyphosate)	Kocide (copper hydroxide)
Endosulfan	Samppi No3	Gastoxin
Fenitrothion	Stam F34T(propanil)	
Fenom C	Basta 20 SL (glufosinate)	
Primigram	Paracol	
Grofol	Saturnmate (thiobencarb)	
Karate (1 - cyhalothrin)	Bladex (cyanazine)	
Nogos (dichlorvos)	Garex	
Ofunack (pyridaphenthion)		
Perfekthion (dimethoate)		
Phostoxin (aluminium phosphide)		
Ripcord (cypermethrin)		
Secto		
Sumithion (fenitrothion)		
Thiodan (endosulfan)		
Trebon (etofenprox)		
Gammalin 20 (lindane)		
Callifan 50CE		
Furadan (carbofuran)		
Unden (propoxur)		
Totals 25	13	7

Common names of the active ingredient of commercial products are given in parenthesis where known.

3.2.3. Sales experience and training of merchants

Most of the merchants had been in the agrochemical business for less than 5 years (Table 6). Less than 3% had been associated with the sale of agrochemicals for more than 20 years. Most (95%) claimed to have had some basic training in handling of such chemicals. Almost all of them acquired their knowledge on the job through seminars and workshops organised by the Ministry of Agriculture, Extension Services, Plant Quarantine and Protection Services and Agrochemical Companies.

Table 2. Types of agrochemicals on the basis of their chemical composition

Organo-chlorine	Organo-phosphate	Carbamate	Synthetic Pyrethroid	Inorganic Compound	Miscellaneous
Thiodan 35EC (endosulfan) Gammalin 20 (lindane)	Actellic (pirimiphos- methyl) diazinon dimethoate 40EC Dursban (chlorpyrifos) Sumithion (fenitrothion) Nogos (dichlorvos) Ofunack (pyridathion) Perfecthion (dimethoate)	Baygon (propoxur) Furadan (carbofuran) Saturnmate (thiobencarb?) Sebon (?)	Cymbush (cypermethrin) Decis (deltamethrin) Karate (λ-cyhalothrin) Ripcord (cypermethrin) Fenom C (?)	Champion (?) Caocobre (?) Cobox (copper oxychloride) Kocide (copper hydroxide) Dithsne (mancozeb)	Gesaprim (atrazine) Garlon (triclopyr) Ronstar (oxadiazon) Secto (?) Stam F34T (propanil) Callifan 50CE (?) Gramoxone (paraquat) Roundup (glyphosate) Basta 20SL glufosinate
2 (6%)	8 (24%)	4(12%)	5 (15%)	5 (15%)	9 (27%)

Table 3. Commonly used agrochemicals by volume of purchase

Chemical	Percentage
Kocide	39.0
Karate	22.0
Perfekthion	11.0
Actellic	11.0
Karate	11.0
Sumithion	6.0

Table 4. Educational background of agro-chemical sellers in the three regions.

Level of education	Number	Percentage
Middle School Leaving Certificate	41	52.6
Secondary School	30	38.4
Others	7	9
University	-	-

Table 5. Length of work experience in retail selling

Duration	Number	Percentage
0 - 5 years	36	46.2
6 - 10 years	31	39.8
11 - 20 years	9	11.5
> 20 years	2	2.5

Table 6. Source of Supply of agrochemicals to retailers and farmers in Ashanti, Brong Ahafo and Eastern Regions of Ghana

Source of agrochemical	Percentage of the total market
Chemico (GH) Limited	23.1
Ministry of Agriculture	18.0
Reiss and Co.	6.4
Imperial Chemical Company	12.8
Wienco	5.1
Retailers	23.1

3.2.4. Handling of agrochemicals by merchants

(a) **Record keeping**

Less than 30% of the merchants kept records of their business including financial records, expiry dates of chemicals and stock controls.

(b) **Storage and display conditions**

Chemicals were mostly stored in the shops in their containers and some packed in original boxes. A few of the sellers had storage rooms. Sellers in kiosks and those operating on table tops displayed some of their items in the open sun.

(c) **Other activities undertaken by merchants in shops**

Apart from the sale of agrochemicals over 90% of the merchants stocked other items including seeds, fertilizers, etc., in their shops. In a few instances, food items such as bread, vegetables, gari and rice were also sold in agrochemical shops.

(d) **Accidents**

There were some reported cases of accidents which occurred as a result of the improper repackaging of agrochemicals into smaller packages for sale to farmers. Reports of occasional headaches and other health problems were considered by the sellers to be related to the chemicals.

3.2.5. Major sources of supply of agrochemicals to retailers and farmers

The supply of agrochemicals to farmers was from commercial sales outlets except for a few cases in the cocoa industry where some farmers obtained chemicals directly from the Ministry of Agriculture. Most of the supply to farmers was from retailers in open market places throughout the regions.

3.2.6. Educational background of farmers

Nearly half (47%) of the farmers had Middle School Leaving Certificate though a reasonable proportion (17%) had no formal education at all (Table 7) and most (92%) were male.

Table 7. Educational background of farmers using agrochemicals.

Level of education	Number	Percentage
Middle School leaving Certificate	92	47.2
G.C.E. O'Level	39	20.0
Others - Post Sec, Agric. Colleges	26	13.3
University	4	2.1
None	34	17.4

3.2.7. Tenure and size of farm

Over one third (38%) of farmers owned their own lands with 18% based on the extended family system. The largest proportion (42%) of the land being farmed was rented but only a very small fraction (0.5%) was under the cooperative system. In general, the size of most farms for individuals was between 0.8 and 2.5 hectares. However there were a few cases where farmers owned about 15-20 hectares of land.

3.2.8. Experience of use, application and storage of pesticides by farmers

Most (66%) of the farmers had used pesticides actively over the past 6 years, though only about 7% had used pesticides for a period greater than 15 years (Table 9).

Most of the individual farms did not receive any Extension Service except those involved in cocoa farming. Some of the rented farms and cooperatives benefitted from those extension services provided by the Ministry of Agriculture with farms in the cocoa producing areas benefitting most from them.

Table 8. Type of farm tenure

Type of tenure	Number	Percentage of total
Owner	75	39
Family	36	19
Cooperative	1	0.5
Rented	83	43

Table 9 Duration of use of pesticides on farms

Years of Use of Pesticide on farm	Number	Percentage
0 - 5 years	128	65.6
6 - 10 years	47	24.1
11 - 15 years	6	3.1
> 15 years	14	7.2

Half (51%) of the farmers did not have any training in the use of agrochemicals although the remainder have had at least some information on the use of particular pesticides on their farm. Such information was from friends, Agricultural Extension Officers and agrochemical merchants.

Nearly all (98%) of the farmers used knapsack sprayers while a proportion (12%) of these used Mistblowers in addition to the knapsack sprayer. Less than 5% used other applicators and sprinklers.

Agrochemicals were mostly stored in the house before use though excess agrochemicals were stored under a variety of conditions. A third (31%) of the farmers used the agrochemicals immediately on the farm. A larger proportion (54%) stored them on the farm or buried the containers in the ground on the farm for future use, while a significant fraction (12%) poured them away on the farm.

A very small proportion (<5%) of the farmers used adequate protective clothing. The majority of the farmers used a variety of protective measures which appeared inadequate, though a quarter (24%) used nose protectors or cloths to cover their nose. Only a few (18%) either washed their hands after using the chemicals or used wellington boots or gloves (15%). A significant fraction (22%) indicated they did not use any safety measures at all.

There were reports of fainting (12%) after the spraying exercise and all the farmers complained of at least one ailment (fever, headache, stomach pains, chest pains) which occurred very frequently.

4. DISCUSSION

Pesticides have been in use in the Ashanti, Brong Ahafo and Eastern regions of Ghana for a relatively long time, and at least 3% of the agrochemical shops have been in operation for over 20 years whilst some farmers have used pesticides on their farms for more than 15 years.

Insecticides form the most prevalent type of agrochemical used in all three regions of Ghana and constitute about 52% of the total number of agrochemicals in use. Lindane and endosulfan are among the insecticides being used most extensively. These organochlorines can be persistent and their accumulation could pose a threat to man and the environment. Persistent insecticides, such as DDT, which have been banned are still available in shops and are used extensively on farms. This emphasises the need for a more strict enforcement of laws regarding the importation and use of such chemicals in Ghana. The danger posed by the low educational background of a large number (52%) of the agrochemical merchants needs to be seriously addressed. These merchants not only mishandle pesticides through lack of knowledge but farmers depend on them as their main source of information. There is the need to intensify training programmes to include not only the handling of such chemicals but also the disposal of stocks pesticides that have exceeded their expiration dates. The tendency for farmers to purchase such agrochemicals and even banned pesticides at relatively reduced prices is closely related to a lack of education coupled with the fact that the merchants who advise them may themselves lack the requisite knowledge about the chemicals. The relatively large number of farmers (51%) who have not had any previous training in the use of agrochemicals may further compound the problem of mishandling of these agrochemicals.

Improper record keeping also posed a big problem in the collection of data on volume of usage of these agrochemicals. The lack of an extension service to family and self-owned farms needs to be remedied. It deprives such farmers of the opportunity of benefitting from training programmes in the correct application and handling of agrochemicals.

Safety measures taken by the farmer were found to be grossly inadequate with less than 5% using adequate protective clothing while 22% did not use any safety measures. A programme of workshops, seminars and other training programmes, aimed at enhancing the farmer's capacity to handle and use agrochemicals to their full potential, needs to be set up. Furthermore, since pesticides can have adverse effects on the environment, a better knowledge of the proper handling of these chemicals from the manufacturers through retail merchants to the farmers could ensure for safer working conditions for the distributor and the farmer and reduce the potential for environmental contamination.

This study presents some of the problems related to the management of pesticides in Ghana which we believe could be mitigated through the coordinated effort of policymakers and scientists on the African Continent with the support of interested International funding agencies and organizations involved in the protection of the environment.

ACKNOWLEDGEMENT

We are grateful to the International Atomic Energy Agency for their grant support for this project.

PERSISTENCE AND FATE OF SOIL APPLIED ^{14}C - LINDANE IN A MAIZE ECOSYSTEM



XA9743722

P.O. YEBOAH
Department of Chemistry,
Ghana Atomic Energy Commission

K.G. MONTFORD
Biotechnology and Nuclear Agricultural Research Institute,
Ghana Atomic Energy Commission

W. APOH, S. DODOO
Department of Chemistry,
University of Cape-Coast

Legon, Ghana

Abstract

^{14}C -lindane applied to soil surface in a maize ecosystem (one month after planting) was taken up by the plant. Within the first 25 days of treatment, ^{14}C -lindane or its metabolites were found within the entire plant with the greatest concentration in lower leaves (from the ground level); and a sharp build up of lindane concentration towards the tip of each leaf. Radioactivity and hence pesticide concentration was uniformly distributed in the plant with time; to the extent that measurable levels of ^{14}C -compounds were detected in the tassel cob and the grain. This indicated that soil applied lindane was available to the maize plant. The persistence of ^{14}C -lindane in soils of variable organic matter content was also studied. Evidence is presented to show that ^{14}C -lindane dissipated faster in soils of lower organic matter content. Levels of surface applied pesticides that became bound in the soil increased with time after application and also with increasing organic matter content. ^{14}C -activity was mainly associated with the top soil layer (0-30 mm).

1 INTRODUCTION

The following study was designed to complement observations made in the previous paper [1].

2 MATERIALS AND METHODS

2.1. Chemicals

Uniformly labelled ^{14}C -lindane (specific activity $647 \text{ MBq mmole}^{-1}$) was obtained from Sigma Chemical Co. St Louis, USA, through the International Atomic Energy Agency (IAEA) in Vienna. The ^{14}C -concentration of the lindane (supplied as toluene solution) was found to be 1 kBq ml^{-1} . The scintillators 2,5-diphenyloxazole (PPO) and 2,2-phenylenebis-(4-methyl-5-phenyloxazole) (DMPOPOP) were purchased from Eastman Kodak Co., Rochester, N.Y., USA. All other chemicals were Analar grade.

2.2. Studies on the persistence of ^{14}C -lindane

Fifty-one PVC tubes (length, 300 mm; internal diameter, 70 mm) were buried in the soil in the rows between plants (in another plot), with 30 mm projecting above ground level. The tubes were arranged into groups of fifteen and in accordance with the organic matter content of the soils they contained. The soil in each of these group tubes were treated with ^{14}C -lindane (185 kBq) at the same time that plants on the field were sprayed with

commercial lindane. Two tubes in each soil were not treated with ^{14}C -lindane to serve as the control. The tubes were dug out carefully at intervals, wrapped in polythene bags and stored in a refrigerator pending extraction and determination of pesticides. The soil in each of these tubes was sprayed with ^{14}C -lindane (185 kBq) in hexane (100 mL).

2.3. Studies on the uptake of ^{14}C -lindane into plants

Twenty plastic buckets (length, 220 mm; internal diameter, 240 mm and open at both ends) were filled with soil from one of three sources, as described previously [1]. They were buried in the crop row with 30 mm projecting above ground level. Fifteen tubes of each soil type were grouped together. Maize was planted in each bucket on the same day that planting was done in the field. The soils in the buckets were treated with ^{14}C -lindane (370 kBq) on the same day that the plants on the experimental farm were sprayed with commercial lindane. Three plants were taken periodically and analysed for uptake of ^{14}C -activity. Both anatomical and sectional distribution of ^{14}C -lindane residues in the maize plants were studied.

2.4. Extraction and analysis

2.4.1. Soils

The soil contained in the PVC tubes was scooped out from top (end exposed to the atmosphere) in successive 150 mm layers. Each soil layer was weighed, dried, ground separately (using a pestle and mortar) and sieved through a 2 mm mesh sieve. Samples (40 g) of ground soil were weighed into cellulose extraction thimbles (size 43 × 123 mm) and soxhlet extracted for 2.5 h using methanol. The extracts were concentrated at $50 \pm 5^\circ\text{C}$ under vacuum to a volume of 5 mL which was then transferred into scintillation vials and analysed for ^{14}C -activity. Recovery was 80.2%.

2.4.2. Plants

Anatomical sections of maize plants above ground level (leaves, stem, cob, etc.) were removed at intervals and stored in a refrigerator for analysis. Stem and leaves were carefully removed from points of attachments cut into 30 mm segments and analysed for ^{14}C -activity using the dry combustion method.

2.5. ^{14}C -activity measurements

Total and bound residues in soils as well as in plant parts were determined by combusting weighed samples in a model 600 Biological Material oxidiser (R.J. Harvey Instrument Corporation, New Jersey, USA). The $^{14}\text{CO}_2$ was absorbed in absorber + cocktail (1+1 by volume, 2 mL). The absorber was made up of ethanolamine + methanol (2.5 + 17.5 by volume) and the cocktail consisting of PPO (5 g) + DMPOPOP (50 g) dissolved in toluene (1 L). ^{14}C -activity was measured by liquid scintillation counting (LSC) in a Tri-Carb liquid scintillation Analyser model 100 using the sample channel ratio method for quench correction.

3. DISCUSSION

3.1. Uptake and translocation in maize

Within 25 days of pesticide application to the soil, it was observed that the plants had taken up a measurable amount of ^{14}C -activity with the greatest amount of 0.08% of applied found in the lowest leaf, i.e. the first leaf from the ground level (Fig. 1). There was then a gradual decrease in pesticide concentration in the higher leaves and the stem. The upwards movement of ^{14}C -activity in the maize plant was further shown by the gradual build up of ^{14}C -activity towards the tip of lower leaves (Figs 3, 4). With time, i.e. within 50 days after




Fig 1. Percentage of initial dose recovered from leaves and stem.




Fig 2. Anatomical distribution of ^{14}C lindane in maize plant.

pesticide treatment, the ^{14}C -activity tended to spread out more uniformly throughout the plant though the greatest concentrations of ^{14}C -activity remained within the bottom three leaves with smaller ^{14}C -activity in the uppermost leaves and the stem (Fig. 2). Some ^{14}C -activity was also found in the tussel, the cob and the grain 50 days after treatment (Fig. 2) indicating that soil applied lindane or its metabolites were available for uptake by the plant. The translocation of soil applied lindane and its metabolites have been investigated in detail [2, 4, 5, 6]. Neither lindane nor its metabolites were evenly distributed within the plants. Comparatively high residue levels were always detected in the leaves whereas small amounts were translocated into the stems, leaves, and fruits. Paasivirta et al. [5] showed that in water plants, lindane concentrations were similar in both the roots and the stem. Differences in residue levels have been shown to be dependent on plant species. Of a series of edible crops grown in soils containing lindane, carrots were shown to have higher levels than beans, tomatoes and potatoes in that order. In soils with low organic matter contents, as in this study, insecticide residues are more available and hence susceptible to uptake by plants [7].




Fig 3. Distribution of ^{14}C lindane in maize leaves (25 days).




Fig 4. Distribution of ^{14}C lindane in maize leaves (50 days).




Fig 5. Bound residues of ^{14}C lindane in soils of different organic matter content.




Fig 6. Dissipation of ^{14}C lindane in soils of different organic matter content.




Fig. 7. Reciprocal value of activity against time.

3.2. Movement in soils

Lindane dissipated faster in soils of lower organic matter contents (Fig.5). In the three soils tested, there was a rapid rate of dissipation of ^{14}C -lindane within the first 10 days of application followed by a gradual decline in this rate. This rate of dissipation was attributed to the low organic matter content of the soils in this study. Although there were copious amounts of rainfall during the period after pesticide application (104 mm in 2 months), residues of ^{14}C -lindane were found only within the first 0-30 mm layer of the soil surface in agreement with Cliath and Spencer [7] who observed no mobility of lindane in clay soils.

ACKNOWLEDGEMENTS

This work was supported by SIDA through the International AtomicEnergy Agency under Research Contract No. 6754. The authors acknowledge the contributions of Messrs B.Q. Modzinuh and S. Afful of the Chemistry Department of the NNRI for their assistance during sample collection.

REFERENCES

- [1] YEBOAH, P.O., MONTFORD, K.G., KLAKE, R.K., AKPABLI, C.K., this TECDOC.
- [2] ITOKAWA, H., SCHALA, A., WEISBERGER, I., KLEIN, W., KORTE, W., Tetrahedron **26** (1970) 763-773.
- [3] BALBA, M.H., SABA, J., Environ Lett. **7** (1974) 181-194.
- [4] BRADBURY, F., Ann. Appl. Biol. **52** (1963) 361-370.
- [5] PAASIVIRTA, J., PALM, H., PAUKKA, R., AKHABUHAYA, J., LODENIUS M., Chemosphere **17** (1988) 2055-2062.
- [6] FUHREMANN, T., LICHTENSTEIN, E., J. Agric. Food Chem. **28** (1980) 446-452.
- [7] CLIATH, M., SPENCER, W., Environ Sci. Technol. **6** (1972) 910-914.

**NEXT PAGE(S)
left BLANK**

PERSISTENCE OF LINDANE IN RICE AND MAIZE ECOSYSTEMS IN NIGERIA



XA9743723

G.U. OKEREKE, C.C. ONOCHIE
Biotechnology Research Centre, Nnamdi Azikiwe University,
Awka, Anambra State, Nigeria

E. ONYEAGBA
Anambra College of Agriculture, Igbariam, Anambra State, Nigeria

Y. DJE
Department of Food Technology and Science, University of Tennessee,
Knoxville, Tennessee, United States of America

Abstract

A three year field study was undertaken to examine the fate of residues of lindane in soil and crops after repeated seasonal applications of lindane to soil in which maize and rice crops were grown. In the 1993 rice study, the lindane residues in the treated soil were 0.6 mg kg^{-1} (1day) and 0.04 mg kg^{-1} (2 months) after application respectively. In the soil from the maize plots, lindane residues were 0.19 mg kg^{-1} (1 day) and 0.16 mg kg^{-1} (1month) after application and not detectable after 6 months. In the 1994 maize study, the lindane residues in the soil were 0.12 (1 day), 0.10 (1 month) and 0.033 mg kg^{-1} (2 months) after application respectively while in the crop they were 0.19 (1 day), 0.105 (2 weeks) and 0.05 (4 weeks) mg kg^{-1} , respectively. In general, lindane residues were not detected in the soil one year after application indicating that there was no accumulation of lindane in the soil during the period 1992-1994 and therefore unlikely to be any long term effects on soil fauna. Lindane residues in rice and maize crops continued to decrease during the period up to harvest.

1. INTRODUCTION

Control of insect pests of rice and maize rely largely on the use of chemical pesticides. In addition to lowering the population levels of an insect, pesticides can cause changes within both the non-target organisms and co-inhabitants of its environment. Some pesticides can also accumulate in the soil and cause environmental pollution when they are leached to the underground water or runs off to streams and rivers.

Most pesticidal chemicals can be absorbed to varying degrees and in some cases can be translocated to the edible parts of plants growing in contaminated soils [1]. The presence of chlorinated hydrocarbon pesticide residues in human adipose tissue most often results from the consumption of contaminated food, either deliberately treated with chemicals for the control of pests or incidentally contaminated from environmental or industrial sources [2]. Despite the use of lindane in controlling insect pests, little is known about its fate in the soils and crops under conditions of Nigerian agriculture.

The publicity generated by the restrictions of certain pesticides, especially organochlorines, in developed countries has led to the ban of the use of these chemicals without sufficient information on their impact on other environments. Knowledge of the behaviour of pesticides and their residues in soils and crops in Nigeria is vital as it is not scientifically justifiable to make predictions of pesticidal residue accumulation using data generated elsewhere since pesticidal residue accumulation in will depend on factors such as the nature of the pesticide, soil type, moisture, soil pH, temperature, cultivation, mode of application and soil organisms [3]. The objective of the study was to determine lindane pesticide residues in soils and their absorption into crops in a field experiment after yearly applications of lindane.

2. MATERIALS AND METHODS

2.1. Materials

Lindane was obtained as the commercial formulation (200 g L⁻¹ emulsifiable concentrate, Gammalin 20, ICI Ltd, Nigeria). A solution containing standard lindane was obtained from Supelco, Baltenfonte, PA by the International Atomic Energy Agency, Vienna, Austria. The standard solution was stored in the refrigerator (4°C) until used. Analytical grade hexane, methanol and acetone were supplied by Aldrich Chemical Ltd, Gillingham, Dorset SP8 4JC. Sep Pak phase C-18 cartridges were obtained from Waters Chromatography Division, Millipore Cooperation, Milford, Massachusetts, 01757, USA.

2.2. Plot design and lindane application

The plot design and application procedures have been described elsewhere [see this Tecdoc, Umeh *et al.*].

2.3. Crop sampling

The aerial parts of 12 plants per subplot were collected 1 day and 2 months after lindane application from both the rice and maize fields in the 1992 and 1993 studies, while in 1994 the sampling intervals were 1 day, 2 weeks and 1 month after application. Amounts (~100 g) of plants were chopped into one mixture and stored in polyethylene bags in a freezer compartment until extracted.

2.4. Soil sampling

Soils were sampled at 1 day pre-treatment, 1 day, 2 and 6 months after lindane application. On each sampling occasion, 12 cores of soil were taken from a 5 m x 5 m area at the side of each subplot to a depth of 15 cm using a soil auger. The samples were ground to pass 0.25 mm mesh sieve and mixed to provide one bulk sample for each subplot. Duplicate soil samples were taken for measurement of moisture content and for residue analysis. All soil samples were stored in polyethylene bags in a freezer compartment until extracted.

2.5. Residue extraction and analysis

2.5.1. Extraction of plant tissue samples

The plant tissue samples were thawed at room temperature. Samples (20 g) were homogenised with of methanol (200 mL). These suspensions were shaken mechanically for 1 h in a screw-capped jar. The suspensions were filtered to remove plant debris. Volumes (10 mL, 1993; 50 mL, 1994) were dried by aspirating with nitrogen and redissolved in methanol + water (3.5 mL, 1+2.5 by volume).

2.5.2. Soil

Samples (20 g) of soil were placed in a screw-capped bottle, mixed with methanol (200 mL) and mechanically shaken for 1 h. The suspension was filtered and volumes (10 mL, 1993; 50 mL, 1994) were dried with nitrogen. Each residue was then redissolved in methanol + water (3.5 mL, 1 + 2.5 by volume).

2.5.3. Clean up procedure

Each aqueous methanol extract was passed through a preconditioned C-18 SPE column and then eluted with hexane. The hexane extract was dried with nitrogen and redissolved again in methanol and the clean up procedure repeated. The residue was dissolved in hexane (2 mL) and stored in a refrigerator until analyzed by the gas chromatographic method.

2.5.4. Gas chromatographic analysis

1992^{a)} : Instrument, Shimadzu 9; column, fused silica capillary, 30 m x 0.25 mm i.d.; temperatures, column, 240°C, injector, 250°C, detector, 350°C; carrier gas, nitrogen; detector, ⁶³Ni electron capture.

1993^{b)} : Instrument, Varian 3700; column, fused silica capillary; temperatures, column, 190°C; injector, 220°C; detector, 270°C; carrier gas, nitrogen; detector, ⁶³Ni electron capture.

1994^{a)} : Instrument, Hewlett-Packard 5890A; column, fused silica capillary, 30 m x 0.25 mm i.d.; temperatures, column, 240°C ; injector, 250°C; detector, 300°C; carrier gas, nitrogen; detector, ⁶³Ni electron capture.

^{a)} Dept. of Food Technology and Science, University of Tennessee, Knoxville, USA.

^{b)} Dept. of Chemistry, University of Ibadan, Nigeria.

3 RESULTS AND DISCUSSIONS

The limit of detection of lindane was 0.0044 mg kg⁻¹ for both crop and soil samples. The recovery from fortified soils after clean up was 93.5%. There were no detectable lindane residues before lindane application in 1992 for the rice plots and in 1992 and 1993 for the maize plots. Lindane residues were not detected in the untreated rice and maize soil because the field experimental design was oriented to avoid any contamination by erosion from treated to untreated plots. Absence of lindane residues at the beginning of the studies in each new site confirmed that lindane had not been used in the locations before the study was started.

Only data of the *extractable* lindane residues in the soil are reported. In the 1993 rice study, the lindane residues in the treated soil were 0.36 mg kg⁻¹ (1 day) and 0.04 mg kg⁻¹ (2 months) after application respectively. For the maize soil the residues were 0.19 mg kg⁻¹ (1 day) and 0.016 mg kg⁻¹ (1 month) after application and were either very low (< 0.09 mg kg⁻¹) or not detectable at 6 months after application. All these results suggest that there is no problem of long term residues of lindane building up in the soil from either rice or maize fields.

The 1994 maize study gave similar results to that of 1993 in that residues of lindane in the soil were not detectable after 6 months probably as a result of loss of lindane due to volatilization and microbial degradation. Some other studies have shown that organochlorine pesticides dissipate at higher rates under tropical and sub-tropical conditions than in temperate regions [4, 5]. The high lindane residues (0.25 mg kg⁻¹) found in maize foliage, 1 day after application, will have arisen from lindane sprayed onto the leaves, since it was applied as a surface treatment and not incorporated into the soil. The residues dissipated rapidly (0.094 mg kg⁻¹ at 2 months after application) but not as fast as they did in 1993 when

they were found to be as low as 0.0033 mg kg⁻¹ at the same interval after application. The cause of the differences between the years is unknown.

Table 1. Characteristics of the soils from the crop fields

Field Crop	% sand	%silt	%clay	% organic matter	pH	%N
rice ^{a)}	21	25	54	2.23	6.3	0.9
maize ^{a)}	25	20	55	1.85	6.1	0.08

a) classified as a clay

Table 2. Concentration of lindane samples taken from the rice field

Year	Crop	Sample	Lindane residues at period from spraying mgkg ⁻¹ (SD)				
			1 day before	1 day	1 month	2 months	6 months
1992	rice ^{a)}	soil ^{d)}	ND	NM	NM	0.17 (0.05)	0.09 (0.11)
		crop	-	0.53 (0.19)	NM	0.1 (0.04)	-
	maize ^{b)}	soil ^{d)}	ND	NM	NM	0.1 (0.02)	0.09 (0.01)
		crop	-	0.25 (0.19)	NM	0.094 (0.004)	-
1993	rice ^{a)}	soil ^{d)}	ND	0.36 (0.02)	NM	0.04 (0.05)	ND
		crop	-	NM	0.011 (0.01)	0.02 (0.002)	-
	maize ^{c)}	soil ^{d)}	ND	0.19 (0.013)	0.16 (0.013)	NM	ND
		crop	-	NM	0.0033 (0.003)	NM	-
1994	maize ^{c)}	soil ^{d)}	ND	0.12 (0.007)	0.103 (0.44)	0.033 (0.045)	NM
		crop	-	0.19 ^{c)} (0.06)	0.05 (0.03)	NM	-

ND = not detectable, NM = not measured

a) plot location, Igbariam; b) plot location, Ugwuoba; c) plot location, Enugu
d) there were no detectable residues of lindane in any of the samples of soils taken from the untreated control plots before and after application to treated plots on each occasion.

e) lindane residues at 2 weeks after application = 0.11 (SD,0.02) mgkg⁻¹.

4. CONCLUSIONS

In conclusion, our results indicate that lindane was not detected in the soil samples taken 6 months after application to rice and maize agro-ecosystems in Nigeria. The results also show that there was no accumulation of lindane or its metabolites in the soil during the study period thus giving little cause for concern as far as a build-up of lindane residues and long term effects on either soil fauna or transfer to the food chain is concerned. They also show that data from temperate countries should only be applied with caution to Nigerian conditions. The variability in the physical and biological properties of soil give rise to difficulties in the prediction of the fate of pesticides as their degradation is a complicated function of volatilisation loss by erosion, leaching, biotic and abiotic conversion [6].

ACKNOWLEDGEMENTS

The authors wish to thank the International Atomic Energy Agency for supporting the work under the funds provided by Swedish International Development Agency (SIDA). Thanks are also due to Professor O. Osibanjo of Department of Chemistry, University of Ibadan, Nigeria and Professor S.T. Melton of Department of Food Technology and Science, University of Tennessee, Knoxville, USA for use of their laboratory facilities and gas chromatograph to run the residue analyses.

REFERENCES

- [1] TALEKAR, N.S., SUM, L.T., LEE, E.M., CHAN, J.S.J., Persistence of some insecticides in the Tropics, Kenya. *J. Sci. Technol. (Ser A)* **5** (1977) 31-44.
- [2] SANBORN, J.R., CHILDOR, W.F., HANSEN, L.G., Uptake and elimination of ¹⁴C hexachlorobenzene (HBC) by the green sunfish (*Leponis cyanellus* Raf.) after feeding contaminated food, *J. Agric. Food Chem.* **25** (1977) 551-553.
- [3] LICHTENSTEIN, E.P., SCHULTZ, K.R., Persistence of insecticidal residues in soils, *J. Econ Ent.* **54** (1961) 617-622.
- [4] SLEICHER, C.A., HOPCRAFT, J., Persistence of pesticides in surface soil and relation to sublimation, *Environ. Sci. Technol.* **18** (1984) 514-518.
- [5] PERFECT, T.J., Environmental impact of DDT in a tropical agro-ecosystem, *Ambio*, **16**(1980) 16-21.
- [6] HORVATH, L., KLING, F., SIMON, L.P., Fate of pesticides in soilsof various organic matter, *Pesticides, Food and Environmental Implications. Proceedings of a symposium, Neuherberg 24-27 (Nov. 1987)* 177-183.

**NEXT PAGE(S)
left BLANK**

LINDANE RESIDUES IN FISH INHABITING NIGERIAN RIVERS

G.U. OKEREKE

Biotechnology Research Centre, Nnamdi Azikiwe University,
Awka, Anambra State, Nigeria



XA9743724

J.CHIDODBEM

Enugu State University of Science and Technology,
Enugu, Enugu State, Nigeria

Y. DJE

Department of Food and Technology and Science, University of Tennessee,
Knoxville, Tennessee United States of America

Abstract

Analysis for residues of lindane in fish collected from various rivers close to rice agroecosystems showed that the concentrations of lindane ranged from none detectable to 3.4 mg kg⁻¹. Fish from rivers where strict regulations prohibits its use had no detectable lindane residues while appreciable amounts of lindane were found in fish where such restriction was not enforced with the variation attributed to the extent of use of lindane in the area of contamination. The investigation confirms that the use of lindane in rice production in Nigeria can cause the contamination of fish in nearby rivers.

1. INTRODUCTION

Pesticides have benefitted man by reducing loss and increasing food production, but unfortunately have been a major source of pollution to fish [1, 2]. Lindane residues can be stored in tissues of birds, and mammals including man [3] and can cause death to organisms at the end of the food chain [4]. Pollution of the environment and its effects on the health of man, animals, fish, wildlife and the ecosystem in general have become a subject of concern[5]. Broad conclusions regarding lindane levels can be drawn from comparison of their distribution in biological materials from a given area. Because of the widespread distribution of fish, their opportunity for exposure to pesticides applied in rice agroecosystems, their role in Nigerian economy and their importance in the web of life, fish was chosen as a logical organism for monitoring lindane residues.

Reports show that the use of pesticides requires close monitoring of the effects on non-target organisms especially on the economically important fishery resources in rivers close to farm lands (6). In developed countries in the temperate regions there is a continuous effort to monitor pesticide residues in air, water, soil, man, wildlife and fish. On the other hand, the effects of pesticide residues on non-target organisms in Nigerian agriculture and its environment is still speculative because of a lack of quantitative data. The purpose of the present study was to monitor certain rivers in Anambra State to evaluate the level of lindane contamination in fish.

2. EXPERIMENTAL

2.1. Sample collection and storage

In 1991, fish samples of different species were collected from different rivers close to rice fields that were suspected to have received lindane applications (Table 1). Fish from locations thought to be untreated were also collected. The fish were collected from the river

with a cast net and transported to the laboratory in ice buckets where they were sorted out into species, wrapped in aluminum foil and frozen until analysed.

2.2. Sample preparation, extraction and clean-up

Each frozen fish sample was ground and homogenized in a meat grinder. Samples (20 g) of the homogenized fish were blended and mixed with anhydrous sodium sulphate (80 g). The pesticide was extracted for 1 hour in a screw-capped jar with acetone + hexane (1+1 by volume, 200 mL). After tumbling, the solids were allowed to settle and the solvent extract filtered into a separating funnel containing sodium sulphate solution (20 gL⁻¹, 200 mL) and shaken for about one minute. The lower aqueous layer was discarded and the hexane extract washed twice more with sodium sulphate solution (20 gL⁻¹, 50 mL). Quadruple hexane extracts were combined and evaporated to about 10mL and then

Table 1. Residues of lindane found in fish caught at the sampling stations in Anambra State, Nigeria

River	Location	Fish species	Lindane residues, mgkg ⁻¹
Ani	Ufuma ^{a)}	<i>Heterotis niloticus</i> ^{d)}	ND ⁱ⁾
		<i>Clarias zariipinus</i> ^{e)}	ND
		<i>Heterobranchus bidorsalis</i> ^{f)}	ND
		<i>Oreochromia niloticus</i> ^{g)}	ND
Ozzi	Enuguabo ^{a)}	<i>Heterotis niloticus</i>	ND
		<i>Clarias zariipinus</i>	ND
		<i>Heterobranchus bidorsalis</i>	ND
		<i>Oreochromia niloticus</i>	ND
Ozalla	Oyi ^{b)}	<i>Heterotis niloticus</i>	<1 ^{j)}
		<i>Clarias zariipinus</i>	<1
		<i>Heterobranchus bidorsalis</i>	<1
Omoh	Oyi ^{b)}	<i>Heterotis niloticus</i>	<1
		<i>Clarias zariipinus</i>	<1
		<i>Oreochromia niloticus</i>	<1
Anambra	Oyi ^{b)}	<i>Heterotis niloticus</i>	3.1
		<i>Clarias zariipinus</i>	2.1
		<i>Heterobranchus bidorsalis</i>	3.1
		<i>Oreochromia niloticus</i>	2.2
Nkissi	Onitsha ^{c)}	<i>Heterotis niloticus</i>	3.4
		<i>Cyprinus carpio</i> ^{h)}	1.9
		<i>Heterobranchus niloticus</i>	2.0

Local Government area :

a) Orumba North

b) Oyi

c) Onitsha

i) none detectable

j) detectable, but not quantifiable

Local name : d) African boy tongue (Okpo)

e) Mudfish

f) Catfish (Azu isi)

g) Tilapia (Ikpokpo)

h) Carp

transferred into vials and evaporated using a N₂ sparge. The residue was redissolved in hexane (2mL). This extract was passed through a column containing Florisil to remove non-pesticidal extracts by the procedure reported previously [7].

2.3. Gas chromatographic analysis

The hexane extracts were analysed by gas chromatography at the Department of Food Technology and Science, University of Tennessee, Knoxville, USA using a Shimadzu 9 AM; column: 30 m x 0.25 mm i.d. SPB-5 fused silica column (Supelco Inc. Belafonte, PA); temperatures : column, 70°C for 1.5min, increased to 200°C at 10°C min⁻¹ and then to 240°C for 20 min; injector, 250°C; detector, 250°C; carrier gas, helium; flowrate, 2 mL min⁻¹; detector, ⁶³Ni electron capture interfaced with a Shimadzu QP 1000 mass spectrometer. At the beginning of the analysis each day, several consecutive injections were done to prime the column [8].

3. RESULTS AND DISCUSSION

3.1. Questionnaire survey of lindane use in Anambra State

The preliminary questionnaire survey involved extensive visits and interviews with officials of the State Ministry of Agricultural Development Agencies, Pest Control Department, Office of Statistics as well as on-the-spot interviews of rice farmers in various locations in Anambra State, Nigeria. The survey revealed that there was no information on the occurrence, distribution, present usage, effects and the impact of lindane in the rice agroecosystem in Nigeria. It was observed that lindane, commonly called gammalin, is still sold in local markets even though the Federal Government of Nigeria has banned its use. Farmers still use it for pest control in various crops and also for killing fish. In some areas the Government ban is strictly observed while, in most other areas, farmers use it for pest control because it is effective and relatively cheap compared with other pesticides. The unavailability of authentic data on the usage and distribution of lindane in Nigeria as evidenced by our survey, justified the collection of fish from various rivers close to rice growing areas to begin the collection of baseline data of lindane residue levels in fish.

3.2. Lindane residues in fish

Comparison of lindane residues in fish from different rivers show that Anambra and Nkissi rivers were more exposed to pesticide contamination than the other rivers (Table 1) with the residues in fish ranging from none detectable to 3.4 mg kg⁻¹. Only small amounts of lindane were detected in fish from the Omoh and Ozalla rivers and no detectable residues in the fish from the Ani and Ozzi rivers. The absence of lindane residues in fish collected from the Ani and Ozzi rivers reflects the strong restrictions on the use of lindane for farming and fishing in these areas, while the low level of lindane found in fish from the Omoh and Ozalla rivers indicate that only small amounts of lindane, that is applied to the rice in this area, entered the rivers. These low levels of lindane are comparable to those found in areas subjected only to airborne contamination [9]. Such levels may not be harmless since it has been reported that DDT at low levels can interfere with the reproduction of certain fish species [2].

Fish obtained from Anambra and Nkissi river contained appreciable amounts of lindane probably because the river flowed through areas where lindane was frequently used

by farmers. Reports show that under certain conditions pesticide residues arising from normal agricultural use may contribute to diffuse (non-point) pollution of the aquatic environment [11]. In an experiment in Zimbabwe, all the fish sampled contained insecticide residues ranging from 0.055mg kg⁻¹ in the liver of living species up to 5.1mg kg⁻¹ in the liver of dead specimens, the latter presumably being the cause of death [12]. Individual fish vary in residue levels and this can be attributed to differences in fish movement, size, food habits and fat content [13]. Other field and laboratory studies showing vast differences in residue levels in fish have also been reported in Wisconsin [14]. Although a very limited number of fish samples were collected in this study, when present the levels of lindane were similar to those found in fish in the midwest of the USA, where total pesticide residue of 2.10 µgg⁻¹ were recorded [15] but higher than those in Utah where only 0.05-0.096 µgg⁻¹ of organochlorine residue was detected [16]. Because some fish continue to accumulate residues over a period of years, the levels observed in the fish from the Nkissi and Anambra rivers reflect the integrated history of exposure. Interviews with villagers indicate a direct contamination as the farmers in the Oyi Local Government areas, where Anambra and Nkissi rivers flow through, reported non-restricted use of the lindane in agriculture and in fish farming.

4. CONCLUSION

In the rivers studied, the extent of contamination of fish by lindane varied. Lindane which was found in fish collected from the Anambra, Nkissi and Ozalla rivers that may have originated from run-off from the rice fields or from direct contamination of the rivers by fishermen. The considerable variation in lindane residue levels in different fish samples, could be caused by variation of fish movement, food habits, species size, age and fat content. Because of these variations, there should be caution in interpretation and application of these data.

ACKNOWLEDGEMENTS

We wish to thank the International Atomic Energy Agency, Vienna and Swedish International Development Agency for providing the fund to support this study under the Coordinated Research Project on Adverse Effect on Flora and Fauna from the use of Organochlorine Pesticides in African Continent, IAEA Research Contract No. 6585 SD.

REFERENCES

- [1] INGLIS, A., HENDERSON, C., JOHNSON, W.L., Expanded program for pesticide monitoring of fish, *Monitoring Journal* **5** (1) (1971) 47-49.
- [2] JOHNSON, D.W., Pesticides in fish: a review of selected literature, *Trans. Amer. Fish. Soc.* **97** (1968) 398-424.
- [3] EDWARDS, C.A., *Environmental pollution by pesticides*. Plenum Press, London. (1973).
- [4] AN de WAERDT, J., *Drins and related insecticides: toxicology, use and alternatives with special reference to Latin America and South East Asia*, *Econoscript* **23** (1983) 1-176.
- [5] KLASSES, H.E., KADDOUM, A.M., Pesticide residues in natural fish population of the Smoky River of Western Kansas, 1967-69. *Pesticide Monit. J.* **7** (1973) 53-61.

- [6] ANTWI, L.A.E., "Studies on the effect of some insecticides on the brain acetylcholinesterase activity of *Tilapia zilli* in two treated tropical rivers", Pesticides, Food and Environmental Implications. IAEA, Vienna 247-256.
- [7] MELTON, S.L., "Pesticide report on Bush Brothers and Company, Vegetables, October1, 1990 - October30, 1991", Chemical fall out 2nd Edition (Muller, N.W., Berg. G.G., Eds) Thomas, C.C. Illinois USA (1991) 119.
- [8] WANDIGA, S.O., MUTERE, A., Determination of γ -BHC in breast milk of Kenyan women., Bull. Chem. Soc. Ethiop.2 (1988) 39-44.
- [9] SODERGREN, A., Transport, distribution and degradation of organochlorine residue in a south Swedish lake ecosystem, Vatten (1973) 108-290.
- [10] SODERGREN, A., DSIRARAI, R., CHARIBZADEU N., MOIMPOUR, A., Organochlorine residues in aquatic environments in Iran, Pest Moni J. 12 (1978) 81- and 87.
- [11] KREUGER, J.K., BRINK, N., Losses of pesticides from agriculture, Pesticides: Food Environmental Implications IAEA, Vienna (1988) 101-112.
- [12] MULLER, P., Effects of pesticides on flora and fauna, Pesticides: Food and Environmental Implications. IAEA, Vienna (1988) 11-27.
- [13] HENDERSON, C., JOHNSON, W.L., IGLIS, A., Organochlorine insecticide residues in fish (National Pesticides Monitoring Program). Pest. Mon. J. 3 (1969) 145-171.
- [14] KLWINWER, A.J. DEGURSE, P.E., WIRTH, T.L., Occurrence and significance of DDT and dieldrin residues in Wisconsin fish, Tech Bull (1968) 41- 43. Dept of Natural Resources, Madison, Wisconsin.
- [15] HENDERSON, C., INGLIS, A., JOHNSON W.L., Organochlorine insecticide residue in fish. National Pesticide Monitoring Program, Pesticides Mon. J. 5 1971).
- [16] HAW, J.S., LOVERRIDGE, E.L. RIPPEE, K.P., PETERSON, J.L., WHITE, D.A., BARTON, J.R., FUHRIMAN, D.K., Seasonal variations in residues of chlorinated hydrocarbon pesticides in the water of the Utah lake drainage system, 1970-1971. Pest Mon. J. 6 (1972) 167-170.

**NEXT PAGE(S)
left BLANK**



ADVERSE EFFECTS ON FLORA AND FAUNA FROM THE USE OF ORGANOCHLORINE PESTICIDES ON THE AFRICAN CONTINENT: THE NIGERIAN CASE

E.D.N. UMEH

Enugu State University of Science and Technology, Enugu

R.I. EGWUATU, E.N. NWAKEZE

Nnamdi Azikiwe University, Awka

P.E. AMULU

Institute of Management and Technology, Enugu

Nigeria

Abstract

Lindane was found to reduce stem borer damage significantly in both rice and maize plots in most years. However, a significant difference in yield was recorded only for maize in 1993 and 1995. There was a slight but steady decrease in the yield of maize over the years in the lindane treated plots although yields in these plots remained higher than in the control plots. Significant differences in the mean dry weight of retrieved leaf discs were recorded in the maize plots in 1994 and 1995. Neither insects nor spiders, were caught from plants in the treatment plots 1 day after lindane application, although almost equal numbers of insects and spiders were recovered from D-vac ground samples of treatment and control plots on the same day. Data from pitfall traps showed that lindane significantly affected the population of *Aranae*, *Formicidae*, *Collembola* and *Acarina* for up to 6 weeks after application. Its effect on *Carabidae* was inconsistent.

1. INTRODUCTION

Organochlorine pesticides are still widely used in Africa for agricultural and public health purposes. Their cost effectiveness and broad spectrum of activity have continued to encourage their demand and use, in spite of international pressures calling for their ban. Presently, there is no pesticide legislation in Nigeria and perhaps also, in some of the African countries resulting in unrestricted importation and use of various kinds of pesticides including organochlorines. Many organochlorine pesticides are persistent in the environment for a long time. They are also known to leave residues in the soil [1,2] resulting in the disruption of soil fauna and flora [3] in the temperate regions, and thus affecting organic matter breakdown and nutrient cycling [4,5]. Treatment of a cultivated forest soil in Nigeria with DDT resulted in a steady decline in yield of cowpea over a four year period [6] and affected activity of certain crickets, lycosid spiders and millipedes [7].

The objectives of this investigation are to determine the effects of lindane (γ -1,2,3,4,5,6 - hexachlorohexane) on:

- (i) the key pests of maize and upland rice, i.e stem borers,
- (ii) predatory and other non-target species in maize and upland rice agro-ecosystems,
- (iii) soil biology and biological activity.

2. MATERIALS AND METHODS

The experiments were carried out from 1992 to 1995 in maize (*Zea mays*) plots, and in 1992 and 1993 in rice (*Oryza sativa*) plots. In 1992, the maize and rice plots were located at Ugwuoba, Enugu State, and Igbariam, Anambra State, respectively, the two towns being about 20 km from each other. Both towns are located in the derived rain forest zone of

Nigeria. In 1993 to 1995, the maize plots were located in Enugu, Enugu State and is at the northern fringe of the derived rain forest. These towns fall within the zone whose annual rainfall does not fall below 1,500 mm and temperature ranges between 27 and 32°C during most of the year.

2.1. Experimental plots

The plots used in Enugu and Ugwuoba were under fallow for 1 and 2 years, respectively, before the experiments began in 1993 and 1992. The plots used in Igbariam were cropped to yam (*Dioscorea* sp.), maize, egusi melon (*Colocynthis vulgaris*), cassava (*Manihot esculenta*), and various leafy vegetables in a mixed farming system. These plots had no previous history of pesticide use.

At the beginning of each experimental season, all the vegetation was ploughed into the soil which was later harrowed twice before mapping out the plots. Soil samples were taken in the three locations and found to be clayey (48-57%), acidic (pH ranges from 6.2 to 6.8), with high organic matter content (Ugwuoba: 8.0%; Igbariam: 3.8% and Enugu: 2.1%).

In 1992 (for maize and rice) and 1993 (for rice), the plot size was 50 m x 50 m each, for treatment and controls, and separated by a 3-metre row of grass. From 1993, the plot sizes for maize were 32 m x 80 m for both treatment and controls, separated by a 1-metre row of grass. In these years (1992 and 1993) there were 4 sub-plots (5 m x 5 m) located within each of the two plots 10 m away from the edge of the plots, and used for sampling. In 1994 and 1995, there was randomised block design with 4 blocks and two treatments. Within each plot was a sub-plot (10 m x 10 m). Because of the possible interference of the various sampling activities within the sub-plots with yield, a yield area plot (2 m x 2 m) was located outside but close to each sub-plot.

2.2. Planting and fertiliser application

The maize variety used throughout these experiments was TZSR yellow while the rice varieties used were FARO 43 (1992), and ITA 306 (1993). Maize was planted in rows, 1 m apart and 50 cm between hills at 2 seeds/hill in 1992. In the subsequent years, the inter-row distance was reduced to 75 cm. Rice was planted in rows by dibbling seeds, 15 cm apart both between rows and hills at the rate of 3-4 seeds/hill. A compound fertiliser (NPK 15.15.15) was applied to rice and maize soils at the rate of 200 kg ha⁻¹, 21 days after planting. A second fertiliser application of urea was made at the tasselling stage (for maize) and the booting stage (for rice), at the rate of 50 kg ha⁻¹. Weeding was done twice manually during each experimental season.

2.3. Pesticide applications

Lindane (200 g L⁻¹ EC) was applied to the treatment plots in rice (1992 and 1993) and maize (1992) in a single dose of 1 kg AI ha⁻¹. From 1993 to 1995, 1 kg AI ha⁻¹ lindane was applied to maize in split doses of 0.5kg AI ha⁻¹. The first lindane application corresponded to the 3-leaf stage in the maize development (stage 13 in the decimal code [8]). The second lindane application was made two weeks later. High volume sprays (120-200 L of water ha⁻¹), using a knapsack sprayer(CP15) with a red nozzle were used in the lindane application. There was no lindane application to the control plots.

2.4. Sampling

Pre-treatment samples of stem borer damaged plants, insects, other arthropods, and soil fauna were taken from the sub-plots a day before lindane application. Thereafter, samples were taken at two weekly intervals for 2 months. In 1994 and 1995, insect and other arthropod samples were also taken a day, and up to week 10 after lindane application.

2.4.1. Plant damage by stem borers

Stem borer damage was assessed by counting all the affected plants in the maize sub-plots. Because of the high plant population in the rice sub-plots, only half the plants were randomly sampled for stem borer damage. At the early stages, window panes and dead hearts were used as indices for stem borer damage. At the later stages, holes in the stem, lodging, and white heads (dried up tassels and panicles) arising from stem borer damage were also used in assessing plant damage. In 1993, the populations of stem borer larvae was estimated 4 weeks after the last lindane application by dissecting 50 stalks, randomly selected outside the sub-plots, and counting all the larvae and tunnels found. Cob damage was also estimated by counting all the damaged cobs in the yield area plots.

2.4.2. Sweep netting, D-vac sampling and pitfall traps for arthropods

Insect, spider and mite populations were sampled using sweep net and pitfall traps in 1992 and 1993. In 1994 and 1995, sweep net sampling was replaced with use of a D-vac sampler which was used to sample both plants and ground. Sweep netting was carried out along two diagonal lines in each sub-plot at the rate of 5 sweeps/diagonal line. After 5 sweeps, the collections were removed and dropped into 70% alcohol until counted. Collections from the sub-plots were stored separately.

D-vac samples for plants and soil surface were taken at the rate of 4 samples per sub-plot. Each sample consisted of 5 sub-samples. For the ground samples, each sub-sample consisted of a 5 second suck with the sampling cone lightly pressed to the ground. For the plant samples, each hill containing 1 or 2 plants, chosen at random, represented one sub-sample. At the earlier stages, the D-vac cone was placed over the plants while sampling. Later, as the plants became taller, the cone was used to sweep the plants from all sides. D-vac sampling of plants was carried out up to 6 week after lindane application while ground sampling continued up to the 10th week.

Four pitfall traps were set up at random, in each sub-plot. The plastic containers used (10 cm in diameter and 10 cm deep) were buried so that their rims were flush with the soil surface. The traps were half filled with soapy water. In 1992, 1993 and 1994, and 1995, the traps were left in the field for 1 day, 2 days and 1 week, respectively, before collection. In 1995, the traps were emptied twice during the sampling week.

In 1992, the target taxa included numbers of adult stem borers, *Carabidae*, *Staphylinidae*, *Coccinellidae*, *Chrysopidae*, *Formicidae*, *Dermaptera* and *Aranae*. In 1993, stem borers, *Carabidae* and *Aranae* were sampled, while in 1994 and 1995, stem borers, *Carabidae*, *Coccinellidae*, *Formicidae*, *Aranae*, *Acarina* and *Collembola* were sampled.

2.4.3. Breakdown of organic matter

Fifty leaf discs (each measuring 25 mm in diameter), cut from mango leaves (*Mangifera indica*), were put in each 12 cm² litter bag with 10 mm apertures. The litter bag was attached to a marker peg and buried to a depth of 50 mm. Four litter bags were buried randomly in each sub-plot prior to lindane application. The litter bags were retrieved three months later and the contents of each litter bag were washed separately, oven dried at 60°C for 24 hrs, and weighed.

2.4.4. Soil fauna

Four soil cores, 150 mm deep and 50 mm in diameter were taken at random from each sub-plot before pesticide application in 1992 and 1993. Each soil core was cut into 6 pieces and the soil fauna present were extracted using Tullgren type extractors. A second set of soil core samples were taken 2 months after lindane application and treated similarly.

3. ANALYSIS OF DATA

The various data obtained were analysed using a two-way analysis of variance incorporating effects of treatment and time. The sub-plots in 1992 and 1993 were regarded as real plots and the data generated were analysed similarly.

4. RESULTS

4.1 Maize 1992

There was heavy rainfall (31.6 mm), which lasted several hours on June 9, 1 hr after lindane application and this seemed to have affected the experiment. The important stem borer species of maize encountered in this work were *Busseola fusca* Fuller, *Sesamia calamistis* Hamp, and *Eldana saccharina* Wlk. Although damage caused by these stem borers was low in 1992, there was an increase in their damage with time in both treatment and control plots (Fig. 1) but the differences were only significant ($P < 0.05$) at one interval (6 weeks after treatment).

A few insects and spiders were caught with the sweep net (Table I). These included specimens from *Coccinellidae*, *Formicidae* and *Aranae*. The *Coccinellids* caught were the phytophagous species, *Epilachna* sp., known to scrape leaf tissues from young maize plants. More insects and spiders were caught in the control than in the treatment plots but their differences were not statistically significant ($P < 0.05$).

Pitfall trap collections included specimens from *Carabidae*, *Staphylinidae*, *Formicidae* and *Aranae* (Table II and Fig. 2). For *Carabidae* and *Staphylinidae* (Table II), there was no significant difference in the mean catches made on any of sampling dates in both treatment and control plots at a 5% level. For *Formicidae* (Fig. 2), only the second post-treatment sample, with mean catches of 3.5 and 8.0 for treatment and control plots, respectively, was found to be statistically significant. Analysis of the mean spider catches showed that significantly more spiders were trapped in the control than in the treatment plots in the 4th, 6th and 10th week after treatment. These results (*Formicidae* and *Aranae*), do not show any clear trends in their response to lindane application. *Chrysopidae* was caught neither in sweep nets




FIG. 1. Stem borer damage in maize and rice plots.




FIG. 1. (cont.).

nor trapped in pitfalls in this experiment. Overall, it would appear that the heavy rain which occurred soon after lindane application may have affected some of these results.

The soil fauna in the samples included *Collembola* (springtails), *Acarina* (mites) and other soil mesofauna (Table IV). Although the pre-treatment samples showed no significant differences in the mean numbers of the various soil fauna extracted from both treatment and

control plots, post-treatment samples showed significant differences in the mean numbers of *Collembola* and *Acarina* extracted from the lindane-treated compared with the control plots. Both *Collembola* and *Acarina* are micro-arthropods inhabiting the soil and leaf litter. They would have received lindane sprays directly and may have been killed before commencement of the rain on the treatment day. Their population seemed unable to recover from the impact of lindane, even at two months after application.

Mean dry weight of leaf discs retrieved (Table V) were 4.7g and 3.8g for treatment and control plots, respectively, and this difference was not statistically significant. Yields obtained from the lindane treated plot (5.12 t ha⁻¹) was not significantly different (P < 0.05) from the yields obtained from control plots (4.99 t ha⁻¹). This result may be due partly to low pest infestation on maize this year but also due to the effect of rainfall which may have leached or washed away most of the lindane deposits on the treatment day.

Table I. Sweep net collections in maize and rice

a) Maize

Weeks after treatment	Mean number of specimens caught ±SE							
	<i>Coccinellidae</i>		<i>Aranae</i>		<i>Formicidae</i>		Stem borers	
	T	C	T	C	T	C	T	C
Pretreatment	0.00	0.00	0.00	0.00	0.00	0.00	-	-
2	0.75 ±0.5	1.50 ±1.3	0.50 ±0.6	0.50 ±0.6	0.50 ±0.5	1.25 ±0.5		
4	0.00	0.25 ±0.5	0.00	0.00	0.00	0.00		
6	0.00	0.50 ±0.7	0.00	0.00	0.25 ±0.5	0.75 ±0.5		
8	0.25 ±0.5	0.75 ±0.5	0.00	0.00	0.00	0.00		
10	0.75 ±0.5	0.75 ±0.5	0.25 ±0.5	0.25 ±0.5	6.75 ±1.0	6.50 ±1.3		

T = treated plot, C = untreated control plot

b) Rice

Weeks after treatment	Mean number of specimens caught ±SE							
	<i>Coccinellidae</i>		<i>Aranae</i>		<i>Formicidae</i>		Stem borers	
	T	C	T	C	T	C	T	C
Pretreatment	0.25 ±0.5	1.25 ±0.5	0.00	0.25 ±0.5	1.25 ±0.5	1.25 ±0.5	1.50 ±0.6	1.25 ±0.5
2	0.25 ±0.5	1.00 ±0.0	0.00	0.00	0.25 ±0.5	4.25 ±1.3*	1.00 ±0.8	6.25 ±1.7*
4	0.00	0.50 ±0.5	0.00	0.75 ±0.5	2.25 ±1.3	0.00	0.75 ±1.0	3.00 ±1.4*
6	0.50 ±0.6	0.50 ±0.6	0.25 ±0.5	0.75 ±0.5	1.25 ±0.5	3.00 ±0.8*	1.00 ±0.8	0.75 ±0.5
8	0.25 ±0.5	0.50 ±0.5	0.50 ±0.6	1.00 ±0.8	1.00 ±0.0	1.00 ±0.0	0.25 ±0.5	0.25 ±0.5
10	0.25 ±0.5	0.25 ±0.5	0.50 ±0.6	1.50 ±0.6	0.50 ±0.6	0.25 ±0.5	0.25 ±0.5	0.50 ±0.6

T = treated plot, C = untreated control plot

* significantly different from the treated plot at P = 0.05

Table II. Pitfall trap collections from the maize trials

Year	Weeks after treatment	Mean number of specimens caught \pm SE			
		<i>Carabidae</i>		<i>Staphylinidae</i>	
		T	C	T	C
1992	pretreatment	2.50 \pm 0.6	2.25 \pm 1.0	1.75 \pm 0.5	2.25 \pm 0.9
	2	1.25 \pm 1.0	2.25 \pm 0.5	0.00	1.00 \pm 0.8
	4	1.25 \pm 0.5	2.75 \pm 0.5	0.00	0.75 \pm 0.8
	6	1.25 \pm 0.5	1.25 \pm 1.0	0.50 \pm 0.6	1.00 \pm 0.8
	8	0.25 \pm 0.5	0.50 \pm 0.6	0.25 \pm 0.5	0.25 \pm 0.5
	10	2.00 \pm 1.1	2.00 \pm 1.2	0.50 \pm 0.6	0.75 \pm 1.0
1993	pretreatment	0.75 \pm 0.5	0.75 \pm 1.0		
	2	0.25 \pm 0.5	1.75 \pm 1.0		
	4	0.25 \pm 0.5	2.00 \pm 1.0		
	6	0.50 \pm 1.0	2.25 \pm 0.5		
	8	1.00 \pm 0.5	2.50 \pm 0.6		
	10				
1994	pretreatment	1.25 \pm 1.3	1.50 \pm 0.6		
	1 day	1.75 \pm 0.5	1.75 \pm 1.0		
	2	2.25 \pm 1.3	3.25 \pm 1.3		
	4	1.50 \pm 0.6	2.25 \pm 0.5		
	6	1.25 \pm 0.5	1.25 \pm 0.5		
	8	1.75 \pm 0.5	1.50 \pm 0.6		
1995	pretreatment	7.25 \pm 5.3	7.5 \pm 5.1		
	1 day	0.00	8.25 \pm 1.5		
	2	1.00 \pm 0.8	12.75 \pm 3.8		
	4	2.75 \pm 1.0	15.00 \pm 2.5		
	6	7.00 \pm 2.7	17.00 \pm 1.7		
	8	11.00 \pm 2.6	12.25 \pm 2.2		
10	7.75 \pm 3.9	10.25 \pm 2.9			

T = treated plot, C = untreated control plot


Table III. Pitfall trap collections from the rice trials

Year	Weeks after treatment	Mean number of specimens caught \pm SE			
		<i>Carabidae</i>		<i>Staphylinidae</i>	
		T	C	T	C
1992	pretreatment	1.50 \pm 0.6	1.50 \pm 0.6	0.75 \pm 0.5	0.50 \pm 0.6
	2	1.00 \pm 0.0	1.75 \pm 0.5	0.00	0.25 \pm 0.5
	4	2.50 \pm 0.6	1.75 \pm 1.0	0.00	0.75 \pm 1.0
	6	2.00 \pm 0.8	2.75 \pm 1.0	0.50 \pm 0.5	1.50 \pm 0.6
	8	3.75 \pm 1.0	5.00 \pm 0.8	0.25 \pm 0.5	1.25 \pm 0.5
	10	4.50 \pm 1.3	6.75 \pm 3.1	2.25 \pm 0.5	2.00 \pm 0.8
1993	pretreatment	4.00 \pm 1.6	3.50 \pm 1.3		
	2	1.50 \pm 1.0	4.50 \pm 1.3*		
	4	2.50 \pm 1.0	4.75 \pm 1.5*		
	6	4.00 \pm 0.8	4.75 \pm 1.3		
	8	3.25 \pm 0.5	3.00 \pm 0.8		
	10				


T = treated plot, C = untreated control plot

* significantly different from the treated plot at P = 0.05

1992 spiders



1994 spiders



1995 spiders






FIG. 2. Pitfall trap collections in maize plots.

1992, ants



1994 ants



1995 ants




FIG. 2. (cont.)





FIG. 2. (cont.)

4.2. Rice 1992

The major stem borers of upland rice are *Diopsis macrophthalma* Dalm., *Chilo zacconius* Bleszynsky, *S. calamistis* and *Scirpophaga* sp. During the vegetative stage of rice, plant damage by stem borers showed up as dead hearts while damage during the reproductive phase resulted in white heads. There was no significant difference in the mean number of dead hearts obtained in the pre-treatment, and in the week 2 samples (Fig. 1). Week 2 sample results may have been because damage attributable to the larvae arising from eggs laid in control subplots between lindane application in the treatment plots and the week 2 sample may not have been manifested. It takes about 12-20 days from egg deposition to appearance of dead hearts. Subsequent samples showed significant differences in the mean number of damaged plants recorded in the treatment compared to the control plots at 5% level.

1992 ants



1992 spider




FIG. 2. (cont.)

Table IV. Fauna found in the soil cores taken from the maize and rice fields

Year	Crop	Species	Mean number of species found \pm SE			
			pretreatment		posttreatment	
			T	C	T	C
1992	maize	Collembola	150.50 \pm 16.7	151.00 \pm 8.7	69.00 \pm 6.5	155.00 \pm 19.6*
		Acarina	16.50 \pm 5.2	16.75 \pm 2.5	10.25 \pm 1.9	11.75 \pm 1.7*
		others	9.75 \pm 3.1	10.50 \pm 3.9	9.75 \pm 3.1	11.00 \pm 3.4
1992	rice	Collembola	74.25 \pm 14.1	60.00 \pm 15.2	27.00 \pm 4.2	80.75 \pm 19.9*
		Acarina	10.00 \pm 3.2	11.00 \pm 1.4	4.75 \pm 3.3	13.50 \pm 6.6
		others	7.75 \pm 4.0	8.75 \pm 1.7	3.00 \pm 1.6	12.00 \pm 2.2*
1993	maize	Collembola	10.00 \pm 3.7	9.25 \pm 1.7	8.00 \pm 2.4	7.25 \pm 1.7
		Acarina	5.50 \pm 1.7	3.75 \pm 1.0	5.00 \pm 1.8	4.50 \pm 1.0
1993	rice	Collembola	52.50 \pm 5.1	47.25 \pm 3.3*	43.00 \pm 7.1	51.25 \pm 5.5*
		Acarina	12.75 \pm 1.7	13.50 \pm 1.7	18.25 \pm 4.1	27.00 \pm 3.7*

T = treated plot, C = untreated control plot

* significantly different from the treated plot at $P = 0.05$

Table V. Retrieval after 3 months of litter buried in the control and treated plots of maize and rice

Crop	Year	Mean retrieved weight, g \pm SE	
		Treated	Control
maize	1992	4.70 \pm 0.5	3.76 \pm 0.6
	1993	5.42 \pm 2.4	4.64 \pm 0.8
	1994	1.33 \pm 0.3	0.67 \pm 0.2*
	1995	2.40 \pm 0.0	1.71 \pm 0.2*
rice	1992	5.49 \pm 1.7	4.46 \pm 1.0
	1993	5.93 \pm 2.1	3.81 \pm 1.4*

* significantly different from the treated plot at P = 0.05

The rice plots had a richer faunal composition than the maize plots. Adult stem borers, *Coccinellidae*, *Formicidae* and *Aranae* were the main groups caught with the sweep net during the vegetative phase. At this stage, most coccinellids caught were the phytophagous species. In the later stages of rice development, a few *Carabidae*, *Dermaptera* and predaceous *Coccinellidae* were also caught in the sweep net. Statistical analyses of sweep net catches of adult stem borers showed significant differences in the mean catches from the lindane treated and the control plots during week 2 and week 4 sampling (Table I). Lindane affected stem borers of rice for up to one month after application. For all the other groups of insects and spiders, except the *Formicidae*, lindane application had no effect. The effects of lindane on *Formicidae* appeared inconsistent.

Pitfall trap collections (Table III and Fig. 2) showed that the important families trapped were *Carabidae*, *Staphylinidae*, *Formicidae* and *Aranae*. Stem borers, particularly, *D. macrophthalma* were also trapped in the pitfalls. In terms of numbers of arthropods caught in both pitfall traps and sweep nets, *D. macrophthalma* was the dominant stem borer species. Carabid beetles were trapped throughout the sample periods but no significant difference was found in their mean numbers in the lindane treated and the control plots. Similarly *Staphylinidae*, though caught on fewer occasions and in lower numbers than *Carabidae*, was also found not to be statistically significant in treatment and control plots at the 5% level of probability.

For *Formicidae* and *Aranae* (Fig. 2) significant differences in their means were recorded in treatment and control plots in all post treatment samples. Thus lindane remained effective against ants and spiders in the rice plots for more than two months after its application.

Soil fauna data (Table IV) showed a mean of 81 *Collembola* in the control plots against 27 in the lindane treated plots. These data were found to be statistically significant at the 5% level of probability. Mite populations, as well as rates of leaf disc disintegration, were unaffected by lindane in the rice plots. The difference in rice yield from treatment (2.48 t ha⁻¹) and control (2.10 t ha⁻¹) plots was not statistically significant.

4.3. Maize 1993

Stem borer damage increased steadily with time and ranged from 1.90% in the pre-treatment to 14.90% in the post-treatment control plots (Fig. 1). In lindane treated plots, plant

damage ranged from 1.35% before treatment to 3.53% after treatment. These differences between treated and control were statistically significant at a 5% level. Thus, the split dose application of 1 kg AI ha⁻¹ of lindane provided protection to the plants against stem borer attack for up to 6 weeks. Larval populations determined by stem dissection showed that more larvae ($X = 17$) were recovered from the control plots than from the lindane treated plots ($X = 4$) and these were found to be statistically significant at the 5% level. Similarly stem borer damage (stem tunnelling), also determined by stem dissection, showed that a mean of 27 and 18 stems were damaged in the control and the lindane treated plots, respectively, and these were found to be statistically significant. On the other hand, more cobs were damaged in treatment plots ($X = 12$) than in control plots ($X = 6.3$) and these data were also statistically significant. None of the target families were caught in the sweep net in 1993.

Generally, pitfall trap catches were lower in 1993. There was no significant difference in the mean numbers of *Carabidae* and *Aranae* caught in the pre-treatment samples. Post-treatment samples of *Carabidae* showed inconsistent trends thus making interpretation difficult. Except the first post treatment sample, lindane application affected spider catches significantly in the treatment plots throughout the sampling period.

Mean dry weights of leaf discs (Table V) retrieved from the lindane treated plots (5.42 g) were not statistically significant from the dry weight of leaf discs retrieved from control plots (4.64 g) at a 5% level. *Collembola* and *Acarina* populations extracted from the treatment and control soil cores two months after lindane application were not significantly different.

Maize yield was found to be significantly higher (5.03 t ha⁻¹) in the treatment than in the control plot (3.61 t ha⁻¹).

4.4. Rice 1993

The major stem borer species monitored was *D. macrophthalma*. However, stem borer damage reported here resulted from the combined effects of *D. macrophthalma* and other stem borers. Plant damage was found to be significantly lower in the treatment than the control plots, in all but the last two post-treatment samples. Sweep net samples showed low populations of *D. macrophthalma* in the treatment plots at 2 and 4 week after lindane application. Differences in the mean numbers of *D. macrophthalma* found in the samples from both the treated and the control plots were not statistically significant throughout.

Pitfall trap collections of *Carabidae* showed that the lindane application reduced carabid populations significantly, for up to 4 weeks after application. Post-treatment soil fauna samples showed that significantly fewer *Collembola* and *Acarina* were extracted from the lindane treated plots, and the mean dry weight of leaf discs retrieved was also significant ($P < 0.05$) in treatment compared with control plots. The yield obtained from treatment plots (2.45 t ha⁻¹) was not statistically different from that obtained from control plots (2.08 t ha⁻¹).

4.5. Maize 1994

Stem borer damage increased steadily with time although damage was lower in treatment than control plots. Damage caused by stem borers was significantly less in the lindane-treated compared with the control plots during the first 6 weeks after lindane application (Fig. 1). Another type of plant damage which became important this year was that

caused by the coccinellid, *Epilachna* sp. which feeds on the leaf tissues of young maize plants, leaving large strips of papery epidermis. Although this damage was not quantified, visual observation showed far greater damage in control than treatment plots.

D-vac samples of plants showed that *Coccinellidae*, *Formicidae* and *Aranae* were the main families caught. Pre-treatment samples showed that in these 3 families, there were no significant differences in the mean numbers caught in the lindane-treated and the control plots. Samples taken 1 d after lindane application showed that no insects or spiders were caught on plants in the lindane treated plots. Subsequent samples also showed significant differences in the numbers of insects and spiders caught in treatment compared with control plots for between 1 and 2 months after treatment (Table VI).

More faunal specimens were collected in D-vac ground samples than from plants. Specimens collected included those of *Collembola*, *Acarina*, *Carabidae*, *Formicidae* and *Aranae*. The mean number of insects, spiders, and mites collected in the pre-treatment and 1d post-treatment samples were statistically similar in all plots. Similarity in these results occurred because specimens killed by lindane the previous day were also picked up by the D-vac. Subsequent soil surface samples showed that significantly fewer *Collembola* and *Acarina* were caught in the treatment than control plots (Table VII) at the 5% level, throughout the sampling period. Carabid catches in D-vac samples were low and not statistically significant between the treatment and the control plots. Lindane application caused significantly fewer spiders and ants in the lindane treated than the control plots during the first 6 weeks of the initial lindane application.

Pitfall trap collection showed that lindane affected *Collembola*, *Aranae* and *Formicidae* populations adversely (Fig. 2). Spider and ant populations recovered by the 8th week. Springtail populations built up steadily in the treatment plots throughout the experiment, but on a significantly lower level than in the control plots. Pitfall catches of *Carabidae* were statistically similar in both plots throughout the experiment (Table II).

Table VI. Insect species caught from the maize plants by the D-VAC sampler

Year	Weeks after treatment	Mean number of specimens caught \pm SE					
		<i>Aranae</i>		<i>Coccinellidae</i>		<i>Formicidae</i>	
		T	C	T	C	T	C
1994	pretreatment	2.50 \pm 0.6	2.25 \pm 0.5	4.00 \pm 0.8	4.00 \pm 0.8	1.50 \pm 1.0	1.25 \pm 0.5
	1day	0.00	2.50 \pm 0.6*	0.00	3.00 \pm 0.8*	0.25 \pm 0.5	3.50 \pm 1.3*
	2	0.00	1.25 \pm 0.5*	2.75 \pm 0.5	5.00 \pm 0.8*	0.25 \pm 0.5	2.25 \pm 1.0*
	4	1.00 \pm 0.8	3.00 \pm 0.8*	4.75 \pm 1.0	6.75 \pm 1.3*	3.00 \pm 1.2	5.25 \pm 2.2
	6	1.00 \pm 0.8	2.75 \pm 0.5*	2.50 \pm 0.6	3.00 \pm 0.8	7.00 \pm 1.8	11.75 \pm 2.4
1995	pretreatment	0.75 \pm 0.5	0.50 \pm 0.6	1.25 \pm 0.6	1.50 \pm 0.6	0.00	0.25 \pm 0.5
	1day	0.00	0.75 \pm 0.5	0.00	1.50 \pm 0.6*	0.00	0.00
	2	0.00	0.75 \pm 0.5	0.00	2.00 \pm 0.8*	0.00	2.00 \pm 0.8*
	4	0.00	1.00 \pm 0.5	0.75 \pm 0.5	3.25 \pm 0.5*	0.50 \pm 0.6	5.25 \pm 2.2*
	6	1.50 \pm 1.0	1.25 \pm 0.5	4.25 \pm 1.0	3.75 \pm 1.7	2.00 \pm 1.4	9.75 \pm 1.3*

T = treated plot, C = untreated control plot

* significantly different from the treated plot at P = 0.05

Table VII. Insect species caught from the soil in the maize plots by the D-VAC sampler

Year	Weeks after treatment	Mean number of specimens caught \pm SE							
		<i>Acarina</i>		<i>Aranae</i>		<i>Carabidae</i>		<i>Collembola</i>	
		T	C	T	C	T	C	T	C
1994	pretreatment	46.8 \pm 10.1	40.2 \pm 1.8	1.0 \pm 0.8	1.50 \pm 1.3	1.25 \pm 1.0	1.00 \pm 0.8	469 \pm 100.9	4
	1 day	43.5 \pm 5.8	41.5 \pm 4.8	0.5 \pm 0.6	1.00 \pm 0.8	0.5 \pm 0.6	2.00 \pm 0.8	424 \pm 83.3	4
	2	12.8 \pm 3.6	43.8 \pm 4.3*	0.5 \pm 0.6	1.25 \pm 0.5*	0.25 \pm 0.6	1.00 \pm 1.0	41.8 \pm 3.2	5
	4	18.8 \pm 6.7	49.5 \pm 5.2*	0.5 \pm 0.6	1.75 \pm 0.5*	0.25 \pm 0.5	0.5 \pm 0.6	127 \pm 23.2	4
	6	30.2 \pm 5.8	51.8 \pm 2.1*	1.0 \pm 0.6	3.00 \pm 1.4*	0.5 \pm 0.6	0.75 \pm 0.5	141 \pm 37.2	5
	8	40.2 \pm 7.1	73.0 \pm 4.1*	2.0 \pm 0.8	3.00 \pm 0.8	0.75 \pm 0.5	1.75 \pm 0.5	160 \pm 43.1	5
	10	58.2 \pm 6.5	41.2 \pm 2.2*	3.5 \pm 1.3	2.75 \pm 0.5	1.25 \pm 0.5	1.5 \pm 0.6	295 \pm 34.2	5
1995	pretreatment	38.5 \pm 2.5	39.8 \pm 3.1	1.25 \pm 0.5	0.75 \pm 0.5	3.00 \pm 5.4	0.50 \pm 0.6	276 \pm 20.4	2
	1 day	39.3 \pm 2.6	34.5 \pm 3.0*	1.00 \pm 0.8	0.75 \pm 0.5	0.00	0.50 \pm 0.6	276 \pm 19.3	2
	2	8.8 \pm 1.8	38.0 \pm 2.2*	0.00	2.25 \pm 1.3*	0.00	0.00	21.8 \pm 2.2	2
	4	8.5 \pm 1.3	40.5 \pm 1.3*	0.25 \pm 0.5	1.25 \pm 1.0	0.25 \pm 0.5	0.50 \pm 0.6	43.0 \pm 13.0	2
	6	20.5 \pm 4.1	44.0 \pm 5.7*	1.00 \pm 0.8	0.75 \pm 1.0	0.25 \pm 0.5	0.25 \pm 0.5	78.0 \pm 14.8	2
	8	35.5 \pm 4.1	41.8 \pm 3.9*	0.75 \pm 1.0	1.00 \pm 0.8	0.00	0.00	245.5 \pm 23.6	3
	10	40.3 \pm 2.4	41.0 \pm 4.1	1.25 \pm 0.5	1.25 \pm 0.5	0.50 \pm 0.6	0.00	280.5 \pm 27.1	3

T = treated plots, C = untreated control plots

* significantly different from the treated plot at P = 0.05

Rates of leaf disc degradation were significantly lower in the lindane treated than in the control plots. Yields, though higher in the treatment (4.53 t ha⁻¹) than in the control plots (3.51 t ha⁻¹) were not statistically different. There was a drop in yield in the lindane treated plots from 5.03 t ha⁻¹ in 1993 to 4.53 t ha⁻¹ in 1994.

4.6. Maize 1995

As in the previous years, stem borer damage increased steadily with time in both treatment and control plots, although at a significantly lower level in treatment than control plots (Fig. 1).

Arthropod samples (Table VI) had no significant difference in the mean numbers of specimens of the various families obtained in the pre-treatment samples from both control and treatment plots. Similar to the results of the previous year, no insects or spiders were caught on the plants in treatment plots 1 day after treatment. *Epilachna* sp. was found to be an important pest of maize seedlings this year, but was not found on the plants in the treated plots two weeks after lindane application. By the 4th week, coccinellids were recorded on plants in treated plots but their number ($X = 0.75$) was significantly lower than control plots ($X = 3.25$) at 5% level.

Like in the previous year, there were no significant differences in the mean numbers of insects and spiders obtained in D-vac ground samples of the treated and the control plots in the pre-treatment and 1 day post treatment samples (Table VI). Subsequent samples showed that the lindane application significantly reduced *Formicidae*, and *Collembola* and *Acarina* populations for about 4 and 8 weeks, respectively, after application. The effects of lindane on carabid populations was not significant. In general, few *Aranae* were caught in the D-vac this year and analysis showed no significant difference in numbers caught in treated and control plots on most sampling dates. This result may be misleading because of the few numbers involved.

Pitfall traps were left in the field for 1 week and this resulted in richer collections of *Collembola* and *Formicidae* than in the previous year. *Collembola* trapped in pitfalls in the pre-treatment samples were enormous ($X = 5629$ and 5555 for control and treatment plots, respectively) compared to the previous years. Samples 1 day after treatment showed that the *Collembola* population in the lindane treated plots had crashed ($X=173$) compared with the controls ($X = 5651$). The *Collembola* population remained significantly lower in the lindane-treated than in the control plots throughout the experiment. For *Formicidae*, the mean catches in the treatment plots remained significantly lower up to 4 weeks after lindane application.

The mean dry weight of leaf discs retrieved from treatment and control plots were statistically different. Higher yields (4.03 t ha^{-1}) were obtained in treatment than control plots (3.60 t ha^{-1}), and their differences were statistically significant. Decrease in yield in treatment plot was also recorded from 4.53 t ha^{-1} in 1994 to 4.03 t ha^{-1} in 1995.

5. DISCUSSION

Lindane is used extensively in Nigeria on cocoa and sometimes also in rice farms. However, it is not recommended for use on maize. Throughout, the maize crop showed phytotoxicity after lindane application and this tended to retard plant growth for a while.

Lindane protected maize and rice plants effectively from stem borer damage in most years. However, this protection was not always reflected in crop yields. Indeed, significant differences in maize yields in the lindane-treated compared with the control plots were recorded only in 1993 and 1995. Rice yield in 1992 and 1993 were not statistically significant. This would suggest that stem borers and perhaps other pests controlled by lindane were not the only determining factors in yield in these experiments. It is possible that lindane may have had an indirect negative effect on yields which declined in the treatment plots over the years. Perfect et al. [6] recorded similar decline in yields of cowpea grown on soils treated with DDT in Nigeria over a 4-year period. They postulated that DDT adversely affected the numbers and structure of soil fauna as well as the decomposition of organic matter which affected some soil properties. Their postulation was based on an earlier paper [9] which showed that elimination of soil fauna by fumigation adversely affected plant growth. The indirect effects of lindane on yield, speculated in this work, may be the result of its adverse effects on some aspects of soil biology which in turn affected yield.

The target soil fauna in these experiments were *Collembola* (springtails) and *Acarina* (mites). The mites included both predatory and detritivorous species. These soil invertebrates are micro-arthropods which, by virtue of their sizes should easily be affected by pesticides. The degree of susceptibility of these micro-arthropods to lindane in these experiments appeared to depend on the concentration of the spray solution, the quantity of the spray solution that reached the soil and perhaps also, their intrinsic ability to resist adverse effects of lindane. In the maize plots (1992) with plant distances of 1m and 500mm between rows and hills, respectively, large areas of soil surface were exposed and received lindane (1 kg AI ha^{-1}) sprays directly, and this decimated both *Collembola* and *Acarina* populations, resulting in significantly fewer numbers recovered in the lindane treated than the control plots in the post-treatment samples. In 1993, with reduced inter-row distances, a closer plant canopy was formed, which reduced the exposed soil surface thereby giving greater protection to the soil fauna. The split dose application of 1 kg AI lindane per hectare this year may also have helped these micro-fauna to resist lindane toxicity resulting in no significant differences in the mean numbers of both *Collembola* and *Acarina* recorded in treatment and control plots.

In rice (1992) with plant distances of 150 mm both between rows and hills, a close plant canopy was formed, leaving limited exposed soil surfaces which received lindane sprays (1 kg AI ha⁻¹ in a single dose) directly. In this situation, the *Acarina* which probably had greater intrinsic resistance to lindane than the *Collembola* were unaffected. However in 1993, significantly fewer *Acarina* was recorded in the lindane treated than in the control plots. It is possible that rainfall in the season had spread lindane residues from where the lindane was deposited to other areas in the plot in 1992. Igbariam had a total annual rainfall of 2181 mm in 1992. Ploughing and harrowing in 1993 may have also helped to redistribute residual lindane in the farm. Samuel and Pillai [10] reported that about 15% of the initial concentration of lindane applied to an Indian soil remained as extractible and bound residues one year after application. Yeadon and Perfect [11] found that after 1 year of soil treatment with DDT, 4.2% of the initial concentration was detected as residues (chiefly DDE) and after 4 years the percentage of the detectable DDE rose to 24%. With 1993 spray, the level of lindane in the soil may have become high enough to also affect *Acarina*.

Micro-arthropods, particularly *Collembola*, which feed on leaf litter, are regarded as important in the fragmentation and incorporation of leaf litter into the soil. Surprisingly, in these experiments (1992 and 1993), there was no significant difference in the mean dry weight of leaf discs retrieved from treatment and control plots in spite of the significant differences in collembola population in treated and control plots. On retrieval of litter bags, they were found to be infested with young earthworms, nematodes, other unidentified mesofauna, roots of weeds and in a few cases, ants. These agents, together with some soil microflora (bacteria and fungi) probably played more important role in litter fragmentation inside the soil than collembola.

The effects of lindane on the agents of litter fragmentation when the same plot was used over the years appeared to be cumulative. For example, in 1992 (on rice) and 1993 (on maize), the effects of lindane on litter disintegration was not significant. In subsequent years however, the effect became progressively more significant ($P < 0.024$ in 1994 and $P < 0.005$ in 1995). This may suggest a cumulative effect due probably to lindane residue which may have increasingly accumulated over the years to a level that drastically affected agents of litter fragmentation in sprayed plots in 1995.

Throughout these experiments, spiders and ants were consistently adversely affected by lindane. Spiders are generalist predators known to range freely over sprayed vegetation and soil surfaces in search of their preys. Critchley et al. [5] also recorded adverse effects of DDT on spiders and ants.

In general, a sweep net appeared inappropriate for sampling maize plants as very few specimens were caught in sweep net compared with the numbers caught with the D-vac sampler.

REFERENCES

- [1] EDWARDS, C.A., Insecticide residue in soils, *Residue Rev.* **13** (1966) 83-132.
- [2] EDWARDS, C.A., 'Pesticide residues in soil and water', in *Environmental Pollution by Pesticides* (EDWARDS, C.A., Ed.), Plenum, London (1973) 1-7.
- [3] EDWARDS, C.A., THOMPSON, A.R., Residues and the soil fauna, *Residue Rev.* **45** (1973) 1-79.

- [4] KELSEY, J.M., ARLIDGE, E. Z., Effects of isobenzan on soil fauna and soil structure, New Zealand Jour. Agric. Res. **11** (1968) 245-260.
- [5] PALISA, A., Soil animals and soil fertility, Urania **20** (1668) 3- 33.[6] PERFECT, T.J., et.al., The effects of crop protection with DDT on some elements of the subterranean and surface active arthropod fauna of a cultivated forest soil in the humid tropics, Pedobiologia **20** (1980) 31-38.
- [7] CRITCHLEY, B.R., et al., The effects of crop protection with DDT on some elements of the subterranean and surface active arthropod fauna of a cultivated forest soil in the humid tropics, Pedobiologia **20** (1980) 31-38.
- [8] ZADOKS, J.C., *et al.*, A Decimal code for the growth stages of cereals, Weed Research **14** (1994) 415-421.
- [9] EDWARDS, C.A., LOFTY, J.R., The influence of arthropods and earthworms upon root growth of direct drilled cereals, Jour. Applied Ecol. **15** (1978) 789-795.
- [10] SAMUEL, T., PILLAI, M.K.K., "Persistence and fate of ^{14}C -gamma-HCH in an Indian sandy loam soil under field and laboratory conditions", Isotope Techniques for Studying the Fate of Persistent Pesticides in the Tropics, IAEA-TECDOC-476, IAEA,Vienna (1988) 93-102.
- [11] YEADON, R., PERFECT, T.J., DDT residues in crop and soil resuting from application of cowpea *Vigna unguiculata* (L.) Walp.in the sub-humid tropics, Environ. Pollution **B2** (1981) 275-294.

**NEXT PAGE(S)
left BLANK**

THE DEVELOPMENT OF LABORATORY AND SEMI-FIELD METHODS TO TEST THE EFFECTS OF PESTICIDES ON PREDATORY BEETLES

P. CHIVERTON, H. WALLIN*
Department of Entomology,
Swedish University of Agricultural Sciences,
Uppsala, Sweden



XA9743726

Abstract

Following the sequential testing procedure adopted by the IOBC/WPRS Working Group Pesticides and Beneficial Organisms, two simple, robust methods are presented which were designed for testing the effects of pesticides on predatory beetles. In a laboratory initial toxicity test both DDT and lindane were found harmful to the carabid *Pterostichus cupreus*, whereas α -endosulfan was 'harmless'. DDT was found harmless to *P. melanarius*. Sub-lethal doses of both DDT and lindane incorporated in prey caused *P. cupreus* females to produce smaller eggs. In a semi-field test it was demonstrated that Lindane reduced the beneficial capacity of *P. cupreus*. Climatic conditions at the time of the test however were such that the majority of test animals in control treatments escaped. Caution was therefore advised in the choice of test animal and test design for the semi-field test.

1. INTRODUCTION

Testing or screening for the side-effects of pesticides on beneficial arthropods is gaining increasing attention by research workers in different parts of the world [1]. Results from such tests enable agronomists to choose safe preparations for use in Integrated Pest Management schemes. They can also be of use to authorities concerned with pesticide registration.

Recognising the need for standardised testing methods and greater international co-operation, a Working Group Pesticides and Beneficial Organisms was formed in 1974 within the framework of the International Organisation of Biological Control (IOBC), Western Palaearctic Regional Section (WPRS) [2]. Aware of the fact that no single test method would provide sufficient information to show the side-effects of a pesticide on a beneficial organism, this group devised a sequential procedure that includes laboratory, semi-field and field methods. These methods are carried out in a particular sequence progressing from laboratory to semi-field and, ultimately, to a field test only if a preparation under examination is found to be harmful to the test organism. The reasoning behind this is that pesticides found harmless to a particular beneficial in a stringent laboratory initial toxicity test are most likely to be harmless to the same organism in the field.

In this paper we adapt a laboratory and a semi-field test method, originally designed for and used in the IOBC Working Group, to examine the direct effects of organochlorine pesticides on predatory arthropods. We also examine whether any sub-lethal effects occur when predatory beetles consume organochlorine-tainted prey. The aim has been to produce protocols for robust test methods that could be used by African colleagues participating in the FAO/IAEA/SIDA Co-ordinated Research Programme on Adverse Effects on Flora and Fauna from the Use of Organochlorine Pesticides on the African Continent.

* Present address: Anticimex AB, Box 47025, S 100 74, Stockholm, Sweden.

2. MATERIALS AND METHODS

2.1. Selection and treatment of test animals

The beetles chosen to illustrate our test methods were two medium-sized carabids, *Pterostichus cupreus* L. and *Pterostichus melanarius* (Illiger) (Carabidae, Coleoptera). Both are common in agricultural fields in Sweden and both have been shown to prey upon the most serious aphid pest in Swedish cereals, *Rhopalosiphum padi* L.[3]. Large numbers of beetles were caught in May and June 1991 in dry pitfall traps. The latter were placed on uncultivated land where no previous pesticide treatments had been made. Beetles were held in plastic containers lined with moistened tissue paper, in a refrigerator (5°C) and fed dog biscuits. All beetles were acclimatised to test conditions (15°C, 16 h light, 8 h dark, 70% r.h.) at least 72 h prior to testing.

2.2. Determination of insecticidal activity in laboratory tests

The basic test method was as follows: Individual beetles were transferred to small plastic beakers (height, 60 mm; diameter, 60 mm) containing 20 mm of even-grained sterile sand. Twenty such beakers were then placed in four rows of five on a plastic seeding tray (400 x 600 mm x 80 mm deep) in a fume cupboard. The beakers (+ beetles) were then sprayed with a hand-held sprayer connected to a compressor set at 35 kPa. The spray nozzle was passed over the beakers at a constant speed and height, calculated to deliver the recommended concentration at the IOBC standard amount (for initial toxicity tests in laboratories) of 6 mg fluid per 100 mm² sand. Two test runs of 20 replicates per run were performed for each pesticide/dose (see below). Controls were sprayed with water + alcohol (19 + 1 by volume), as this was used to dissolve the organochlorines tested. Between sprays, the sprayer was thoroughly washed with repeated dosings of acetone followed by water and then dried.

Treated groups were transferred to a large, well ventilated controlled climate room set at the above conditions. The small beakers were then inserted into larger ones of the same diameter (i.e. height, 60 mm; diameter, 60 mm). This ensured that the beetles a) could not escape and b) did not come into contact with excess pesticide on the beaker walls. Each beetle received one dog biscuit as food, and this was moistened daily. Beetles were examined at 1, 6 and 24 h post treatment, and thereafter daily until the sixth day whereupon the test was terminated. Numbers of beetles found dead or moribund were recorded.

Two toxicity tests were performed using this basic method. Different dilutions of DDT, starting with a 'field dose' equivalent to 1000g AI ha⁻¹ (as this was the most common dose found in the literature) were tested on *P. melanarius*, and several organochlorines (i.e. DDT at a rate equivalent to 1000g AI ha⁻¹, α - and β -endosulfan : both at 1100g AI ha⁻¹ and lindane : at 350g AI ha⁻¹) were screened for toxicity to *P. cupreus*. In both tests water containing alcohol (19 + 1 by volume) was used as a control treatment.

2.3. Determination of sub-lethal effects in laboratory tests

In June 1991, 50 female *P. cupreus* were selected from our field-collected stock. These were placed individually in petri dishes containing moistened pulverised peat, and divided into five groups of ten. Every third day one group of ten beetles were each fed with

one (initially two) blowfly larvae (maggots) which had previously been sprayed with a 1:32 dilution of the 'field dose' DDT used above. The other four groups of ten beetles were fed maggots that had been sprayed with dilutions (1:32) of α - and β -endosulfan, lindane, and water + alcohol (19 + 1 by volume) as control, respectively. After spraying, all but two of the maggots were frozen for later use. At subsequent feeding events, maggots were thawed and then almost cut in half prior to feeding in order to ensure that the beetles could penetrate the larval cuticle. Twice a week beetles were temporarily removed from the petri dishes and the peat substrate was replaced. The used substrate was washed through a fine-meshed sieve. Any eggs the beetles had laid in the substrate were subsequently caught in the sieve. The number of eggs found per female were recorded. Eggs were placed briefly on a filter paper to dry and then weighed.

2.4. Determination of insecticidal activity in semi-field tests

The species of beetle used in this test (*P. cupreus*) was collected and held under the same conditions as above. In June 1992, 16 plastic seeding trays (400 x 600 x 80 mm) were half-filled with potting soil and sown with three rows of spring barley at normal field seed rates (4-500 per square meter, in this case ~50 plants per 600 mm row) and at a row width of 125 mm. Trays were then placed in a greenhouse to hasten germination. The inner edges of each tray, from the soil surface to the rim, were painted with Fluon (polytetra-fluoroethylene) to discourage beetles from climbing the tray walls and escaping. When the plants had reached growth stage 32 4 four of the trays were taken outdoors and sprayed with Lindane at a 'field' dose equivalent to 300 g AI ha⁻¹. Four trays were sprayed with a dilution (1:4) of the field dose, and a further four trays were sprayed with a second dilution (1:16). The remaining four trays were sprayed with water + alcohol (19 + 1 by volume) containing several drops of a wetting agent (Triton X-100) as controls. The trays were allowed to dry for 24 h, after which four *P. cupreus* were released onto each tray. The trays were then placed in a large, outdoor bird-proof cage and the beetles allowed to acclimatise for 48 h.

At 72 h post-treatment, each tray was provided with 100 freshly-frozen *R. padi* arranged on four moistened filter papers with 25 aphids per paper. The filter papers were placed on the soil surface and covered with a plastic roof to protect the aphids from the rain and wind. The number of aphids removed after 24 h was recorded. This procedure was repeated at weekly intervals for 3 weeks. Four fresh beetles were released onto each tray 24 h before each feeding occasion. The mean total number of aphids eaten per treatment and per feeding occasion was noted and the results examined by analysis of variance (ANOVA).

Dead beetles found on the soil surface were removed during the course of the experiment and numbers recorded. After the last feeding period, plants were cut at the stem base and removed from the trays. The soil surface was then carefully searched and surviving beetles removed with a pooter. Thereafter, the soil was sieved onto a larger tray and the beetles that had buried into the soil were recovered. The mean number of survivors per treatment was noted and the results analysed by ANOVA.

3. RESULTS

3.1. Insecticidal activities in laboratory tests

The chosen 'field' dose of 1000g AI ha⁻¹ DDT caused 40% mortality in *P. melanarius* with dilutions of this field dose giving a typical dose response curve (Fig. 1). In contrast, the

same field dose of DDT caused 100% mortality in *P. cupreus*, as did the chosen field dose of lindane, whereas α -endosulfan only caused 30% mortality even at the relatively high dose of 1100 g AI ha⁻¹ (Fig. 2).




Fig 1. Initial toxicity of DDT to the carabid beetle *Pterostichus melanarius*




Fig 2. Selectivity of organochlorine insecticides to the carabid beetle *Pterostichus cupreus* in a laboratory initial toxicity test

3.2. Sub-lethal effects in laboratory studies

No significant differences were found between the number of eggs laid by female *P. cupreus* fed on organochlorine tainted prey or fed on water treated control prey ($F = 0.28$, d.f. = 50, n.s.). Eggs from female beetles fed sub-lethal doses of DDT and lindane, however, were significantly smaller (Table 1).

3.3. Insecticidal activities in a semi-field test.

The mean number of aphids eaten per treatment for each feeding occasion are given in Table 2. Mean numbers of aphids eaten in the field dose treatment were always significantly

lower than numbers in the diluted (1:4) field dose treatment. Surprisingly, mean numbers eaten in the control were not significantly different from those in either the field dose or from the different dilutions of the field dose. An explanation was found at the end of the experiment when beetles were recovered. Only one beetle of a treatment total of 48 was recovered from the control trays. There was no significant difference in the number of survivors in the treated plots (Table 3), although dead beetles were only recovered from the field dose (1:1) treatment.

Table 1. Sub-lethal effects of organochlorine tainted prey on egg weights of the carabid beetle *Pterostichus cupreus*.

Pesticide	Egg wt. (mg)	N	P
DDT	0.65	291	0.0001
endosulfan	0.70	288	n.s.
endosulfan	0.69	261	n.s.
lindane	0.63	315	0.0001
water control	0.71	254	

n.s. = not significant in a Student's T-test against control

Table 2. Consumption of aphids by the carabid *Pterostichus cupreus* in arenas sprayed either with different dilutions of lindane or with water treated controls.

Week after treatment	Control	lindane (300g AI ha ⁻¹)		
		1:1	1:4	1:16
1	12.9 ab	4.3 b	20.9 a	16.4 ab
2	14.3 ab	4.5 b	20.8 a	20.0 a
3	14.6 bc	11.1 c	24.4 a	24.4 ab

Numbers with the same letter are not significantly different at $P < 0.05$ (Duncan's multiple range test)

Table 3. Recovery of alive and dead *Pterostichus cupreus* from arenas sprayed either with lindane dilutions or water treated controls.

Insect status	Control	lindane (300 g AI ha ⁻¹)		
		1:1	1:4	1:16
live	0.3 b	1.8 ab	3.5 a	2.5 a
dead	0	41	0	0

Numbers with the same letter are not significantly different from the control at $P < 0.05$

4. DISCUSSION

The carabid species chosen for the semi-field test, *P. cupreus*, is macropterous and can therefore fly. However, extensive behavioural studies of this species (in which beetles confined on similar arenas were video filmed at different temperatures in a controlled climate chamber 5) had revealed that the beetle rarely flies unless the soil surface temperature exceeds ~ 30°C, whereupon they climb to the top of the plants and fly away. Unfortunately the summer of 1992 was one of the hottest and driest on record in Sweden, with ambient temperatures often exceeding 26-28°C throughout the duration of the experiment! Soil surface temperatures, usually 10°C higher than ambient temperature, would therefore have exceeded the 30°C upper threshold for this species and caused it to flee. Interestingly, fewer beetles fled, or were capable of fleeing, from the 1:4 and 1:16 dilution treatments of lindane as significantly more were recovered from the arenas compared with controls. We have previously used this method successfully to screen the effects of pesticides on a small, brachypterous carabid, *Bembidion lampros* Herbst., with good results and would therefore encourage the use of the method, though with a caution that the beneficial insect should be carefully selected.

This laboratory test has proved useful, and is robust enough to be carried out in most laboratories. The basic method can even be adapted and used as a persistence test, in which beetles are placed in the beakers at intervals post-treatment. This would be a useful exercise particularly regarding the organochlorines, and might reveal differences between, for example, DDT and lindane regarding the persistence of harmful effects. Should a macropterous species be chosen for the laboratory test, we suggest that a simple muslin cloth coated with Fluon and held in place by a rubber band over the top of the beaker will suffice to prevent escape. This was demonstrated at the CRP Workshop in Arusha where the method to screen the effects of organochlorines was used on a coccinellid species.

The increase in egg production following ingestion of a carbamate (pirimicarb) has previously been observed in *P. cupreus* [6] and is thought to be due to a hormoligosis effect [7]. Egg size has been found to affect the survival of carabid larvae [6], and first instars hatching from large eggs can withstand starvation better than those hatching from small eggs. Therefore ingestion of DDT or lindane tainted prey by female ground beetles could have significant effects on the survival of their progeny. These and other sub-lethal effects are not exclusive to the organochlorines and need to be considered when describing the overall side effects of pesticides.

ACKNOWLEDGEMENTS

We thank Carol Högfeldt for technical assistance. This work was carried out within the project "The effects of organochlorine pesticides on the non-target fauna" and was supported by grants from the Swedish International Development Authority (SIDA).

REFERENCES

- [1] HASSAN, S.A., Procedure for testing the side effects of pesticides on beneficial arthropods as being considered by the International Working Group Pesticides and

- Beneficial Arthropods, Mitteilungen der Deutschen Gesellschaft fuer Allegemeine und Angewandte Entomologie **4** (1983) 86-88.
- [2] FRANZ, J.M., Pesticides and Beneficial Arthropods, IOBC/WPRS Bulletin **1** (1975)147-152.
- [3] CHIVERTON, P.A., Predation on Rhopalosiphum padi by polyphagous predatory arthropods during the aphids pre-peak period in spring barley, Ann. Appl. Biol. **111** (1987) 257-269.
- [4] ZADOKS, J.C., CHANG, T.T., KONZAK, D.E., A decimal code for the growthstages of cereals, Weed Research **14** (1974) 415-421.
- [5] CHIVERTON, P.A., Searching behaviour and cereal aphid consumption by Bembidion lampros and Pterostichus cupreus, in relation to temperature and prey density, Entomol. Exp. Appl. **47** (1988) 173-182.
- [6] WALLIN, H., CHIVERTON, P.A., EKBOM, B.S., BORG, A., Diet, fecundity and egg size in some polyphagous predatory beetles, Entomol. Exp. Appl. **65** (1992) 129-140.
- [7] LUCKEY, T.D., Insecticide hormoligosis, J. Econ. Entomol. **61** (1968) 7 - 12.

**NEXT PAGE(S)
left BLANK**



EFFECTS OF LINDANE ON A MAIZE ECOSYSTEM AT TPRI, ARUSHA, TANZANIA

E.M. MINJA, R. MAKUSI, F. TESHA
Tropical Pesticides Research Institute,
Arusha, Tanzania

Abstract

The effects of the organochlorine insecticide, lindane, on target and non-target organisms was studied in field plots of maize at Arusha, Tanzania for 4 growing seasons. A single application of $1 \text{ kg}\cdot\text{ha}^{-1}$ caused leaf scorch to the crop but two applications of $0.5 \text{ kg}\cdot\text{ha}^{-1}$ at an interval of 2 weeks were tolerated. The insecticide reduced damage by the stem borers *Busseola fusca* and *Sesamia calamistis* in all seasons and maize yields were higher on the treated plots although the differences were not always significant ($P > 0.05$). Collembola and sometimes ant and spider numbers were lower in treated plots early in the season but the differences did not persist. Differences in rates of decomposition of buried leaf litter were not significant.

1. INTRODUCTION

The use of synthetic pesticides in crop and forestry production as well as in the control of both animal and human disease vectors has been a common practice worldwide. These pesticides are harmful to the environment particularly those which persist for long periods. Persistent pesticides include insecticides in the organochlorine group (e.g. DDT, benzene hexachloride-BHC, chlordane, heptachlor, toxaphene, methoxychlor, aldrin, dieldrin, endrin and endosulfan) which are lipophilic. While most of these have been banned from use in many countries on the basis of their persistence and accumulation in animal and human tissues, some of them are still registered for use in Tanzania [1].

Lindane (gamma benzene hexachloride) is among the organochlorine insecticides registered for use in crop protection in Tanzania. In most developed countries the environmental impact of organochlorine pesticides has been monitored and documented [2]. It has been calculated that as much as 50% of pesticide sprays applied to crops miss their target and fall on to the soil surface. Also the air can easily become contaminated with pesticides during spraying operations [3]. In most developing countries including Tanzania, very little information is available on pesticide effects on the environment. The maize stem borer *Busseola fusca* Fuller is one of the major insect pests limiting maize production in Tanzania. DDT dust and high volume sprays were recommended and commonly used by farmers but after the banning of DDT due to its persistence and hazardous effects on the environment, farmers have been using various other insecticides including lindane. The work being presented was therefore proposed to investigate the effects of lindane on field maize arthropod fauna.

2. MATERIALS AND METHODS

The studies were conducted on a 0.7 ha field at the Tropical Pesticides Research Institute, Arusha for four planting seasons starting in 1991/92 short rain season. Planting dates were based on the on-set of the rains which was November/December (this applied to 1991/92, 1992/93 and 1994/95) for the short rains and March/April for the long rains (1994 trial). Planting of the trial was done in the 1994 long rain season (which is cooler and wetter, and therefore with lower insect populations compared with the warm short rain season) due to failure of the 1993/94 short rain season crop from a severe drought in January/February 1994. The field was first ploughed and for the first two seasons animal manure (cow dung) at 3 tons

per ha was broadcasted and harrowed into the volcanic sand loamy soil which is known to be low in organic matter. The cow dung was applied to the soil a month before planting maize. The field was then divided into four equal blocks which were each further divided into two 25 m x 25 m plots. The plots in a block were each randomly assigned two treatments i.e. sprayed and unsprayed.

Maize hybrid seed CG 4141 was sown at 90 cm x 30 cm at two seeds per hole and the seedlings were thinned to one per hill at three weeks after emergence (3WAE). All field operations apart from tractor ploughing and harrowing were carried out by hand. Fertilizer top dressing using urea at 25 kg N/ha was done twice at 3WAE and 6WAE.

In each of the two plots per block, 5 m x 5 m subplots were marked out permanently for different investigations. Peripheral subplots acted as guard rows. In one subplot 4 standard pitfall traps were installed in the ground for monitoring ground fauna population activity. The traps were set for three days with a weak soap solution. They were then emptied and arthropod fauna sorted out and recorded. A second subplot was used for aerial fauna monitoring in which arthropods were swept from plants using a sweep net (three seasons) and a D-Vac in the fourth season. A third subplot was used for visual fresh plant damage assessment where all plants with fresh borer damage were counted and converted into a proportion of total subplot plant population. A fourth subplot was left undisturbed until harvest for grain yield assessment. A fifth subplot was used for organic matter decomposition studies in which four litter bags containing *Grevillea robusta* A. Cunn leaves were randomly buried for three months. Twenty plants were randomly uprooted from the remaining four subplots at each assessment day for two seasons to assess stem borer larval population and parasite incidence. The stems were each dissected for stem borer larval counts.

The crop was constantly inspected for stem borer eggs and first instar larvae to help in setting out the date for initial insecticide sprays. Spraying was carried out at 4WAE when either eggs and or first instar stem borer larvae were recorded on 0.5-1.0% of the plants. Lindane 20 EC. (Gammalin 20% from Imperial Chemical Industries) was sprayed once at 1.0 kg AI.ha⁻¹ in 1991/92 and 1992/93. Because this caused severe leaf scorching the application was split and sprayed twice at 0.5 kg AI.ha⁻¹ at two week intervals starting at 4WAE in 1994 and 1994/95. CP15 and CP20 knapsack hand sprayers with polijet cone type of nozzles and operated at a pressure of 3 x 10⁵ Pa (3 bar) were used at high volume rate (300 L water ha⁻¹). Arthropod populations and plant damage were monitored just before the first spray, 24 hours later and then every two weeks until harvest. All data was subjected to analysis of variance.

3. RESULTS

Field observations showed that the maize stem borer, *B. fusca* was the dominant pest early in the four seasons starting towards the end of the 3WAE. Towards the middle of each growing season, however, minor infestations of *Sesamia calamistis* Hamp. occurred and the two borers cohabited on the plants until harvest. Damage assessments were based on *B. fusca* which feeds in the leaf whorl, in the stems and on maturing seed on the cobs. *S. calamistis* normally rings the lower part of the stems and migrate to other stems or to the maturing seeds on the cobs.

The total monthly rainfall and mean temperatures for the institute where the experiments were carried out are shown in Figures 1-4 corresponding to the four seasons. The results on beneficial arthropod population groups represented by ants, spiders, and collembola,

and plant damage for 1991/92, 1992/93, 1994 and 1994/95 are shown in Tables 1-4, respectively. Buried leaf litter weight losses are shown in Table 5 and maize grain yields in Table 6.

The results for 1991/2 show that arthropod populations were very low in that season. The one lindane application was sufficient to check borer infestations for 7 weeks by which time the plants were old enough to withstand a degree of pest attack.

In 1992/3 lindane significantly reduced Collembola numbers initially but the population recovered to a similar level to that in the control plots by the end of the season. Ant and spider numbers were essentially the same in treated and control plots but treated plots showed significantly less crop damage.

The 1993/4 short season crop failed because of drought so a crop was grown in the 1994 long rainy season. The long rainy season is normally cooler and wetter with relatively low pest insect activity but more abundant beneficial soil arthropods. Splitting the lindane treatment into two applications prolonged the depression of arthropod populations but, as in the previous seasons, they recovered by harvest time. Again, pest damage to the crop was significantly lower on the treated plots. These trends were repeated in 1994/5. In addition a gregarious *Apanteles* spp. which is a parasite of *B. fusca* larvae was observed in similar numbers in treated and untreated plots.

Buried leaf litter decomposition (Table 5) was slightly, but, not significantly, lower in treated than control plots in all three years that litter bags were buried. Grain yields (Table 6) were 31% higher on average in the treated plots over the 4 seasons although the variability was such that the differences were not significant ($p > 0.05$).

4. DISCUSSION

The results indicate that lindane was effective in reducing *B. fusca* damage on maize. On the other hand it had adverse effects on most of the early season beneficial arthropod fauna monitored in the four seasons. It was especially so with Collembola, and in some seasons it also affected ants and spiders. Similar reductions in population activity for ants and spiders were observed in the humid tropical zone of Nigeria in plots treated with DDT [4].

The mid- and late-season beneficial arthropods including spiders (Araneae) and ants (Formicidae) were not significantly affected by lindane. Collembola populations were the most abundant beneficial soil microfauna at the beginning of each season but their numbers would normally decline gradually even in the controls as soil moisture decreased towards crop maturity. These results indicate that lindane could have short-lived adverse effects on the maize environment at the rate in which it was used in the warm and semi-humid climatic conditions in which the experiments were conducted. Edwards [5] observed that in a combination of wet and warm weather, the pesticides are released and may break down and disperse into the atmosphere much more rapidly than from dry soils; and that there was evidence that pesticides which persist for more than 10 years in temperate climates may disappear almost entirely in a year in the tropics. The results in this programme from Ghana, Nigeria and Zambia confirm this.

Lindane sprays in the present study seem to have reduced buried leaf litter decomposition rate (though not significantly) in the three seasons. Although there was no

Text cont. on p. 179.




Figure 1: Total monthly rainfall (mm) at T.P.R.I. for July 1991 - June 1992




Figure 2: Total monthly rainfall (mm) and mean monthly temperature at T.P.R.I. during 1992/1993




Figure 3: Total monthly rainfall (mm) at T.P.R.I. for July 1993 - June 1994




Figure 4: Total monthly rainfall (mm) at T.P.R.I. for July 1994 - June 1995

Table 1. Mean number of ants (Formicidae) populations in sprayed (lindane) and unsprayed plots

Sampling occasion	1991/92		1992/93		1994		1994/95	
	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed
1	0	0	36	16.75	56.25	65.75	7.25	11.75
2*	0	0	27.25	18.50	18.75	65.00	10.50	5.25
3	0	1.5	21.25	16.00	18.50	55.75	64.75	92.50
4**	1.75	6.0	28.25	33.25	8.00	53.50	44.75	53.00
5	3.25	0.75	52.50	47.00	27.50	47.25	22.25	25.50
6	8.75	12.00	30.00	82.25	37.00	41.75	23.75	23.00
7	0.75	11.75	61.00	54.25	65.75	52.75	0	0
8	0.0	13.00	66.00	74.25	98.25	220.00	0	0
9					113.50	238.75		
10					144.25	236.50		
Mean±SE	1.8±0.30	5.6±1.20	40.3±5.62	43.5±6.89	58.8±7.54	107.7±10.7	28.9±5.22	35.2±6.80

- Sampling occasion 2 corresponds to 24 hours after the first spray for all four seasons

** Sampling occasion 4 corresponds to two weeks after the second spray in 1994 and 1994/95

Mean values for 1992/92 are significantly different ($p < 0.05$), values for other years are not

Table 2. Mean number of spider populations in sprayed and unsprayed plots

Sampling occasion	1991/92		1992/93		1994		1994/95	
	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed
1	0	0	3.00	3.75	4.00	8.25	10.5	12.00
2*	0	0	3.75	2.00	1.25	6.75	1.75	5.00
3	0	0.75	1.25	5.50	1.25	8.25	3.50	4.00
4**	0.75	2.50	2.5	3.75	1.25	12.25	8.00	11.75
5	0.50	1.50	4.25	3.00	2.25	12.00	0.50	3.00
6	2.75	1.50	7.00	6.75	5.25	11.00	3.25	3.00
7	0.75	1.50	3.25	5.50	5.25	10.25	0	0
8	2.75	2.00	3.75	6.00	7.75	11.00	0	0
9					9.25	10.25		
10					10.00	11.00		
Mean±SE	0.94±0.21	1.2±0.40	3.6±0.84	4.5±0.67	4.8±0.96	10.1±2.90	4.6±1.10	6.5±1.21

Mean values for 1994 are significantly different ($p < 0.05$), values for other years are not

Table 3. Mean number of Collembola populations in sprayed and unsprayed plots

Sampling occasion	1992/93		1994		1994/95	
	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed
1	268.75	275.75	594.75	528.25	787.50	790.00
2*	190.00	262.25	228.00	603.50	17.50	12.50
3	189.25	247.50	204.50	688.00	275.00	430.00
4**	145.00	249.00	175.25	366.00	282.50	787.50
5	143.75	187.00	82.75	101.50	166.50	294.00
6	162.00	175.50	61.25	74.75	91.25	108.50
7	180.00	193.75	85.00	70.75	45.00	73.75
8	148.50	173.00	87.25	69.00	0	0
9			87.50	89.00		
10			94.50	98.50		
Mean \pm SE	178.4 \pm 12.8	220.5 \pm 20.6	170.1 \pm 18.7	288.05 \pm 40.63	237.9 \pm 17.5	356.6 \pm 20.6

Mean values for treated and control plots are significantly different ($p < 0.05$) for all years

Table 4. Fresh borer damage (% mean) on maize plants in sprayed and unsprayed plots

Sampling occasion	1991/92		1992/93		1994		1994/95	
	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed	Sprayed	Unsprayed
1	1.5	0.7	1.2	1.4	0.25	0.37	1.10	1.00
2*	0	0.7	2.9	6.8	0	0.75	0.40	1.42
3	0	5.0	2.7	5.1	0	1.12	0	2.30
4**	0	6.2	4.4	9.0	0.25	1.37	1.12	3.55
5	2.2	8.0	0.6	3.0	0.62	1.62	1.92	7.72
6	1.1	6.0	0	0.5	1.12	1.87	2.70	12.40
7	1.0	3.2	0	0	1.25	2.50	0	0
8	1.0	3.4	0	0	1.50	2.62	0	0
9					2.00	2.75		
10					2.00	3.05		
Mean \pm SE	0.85 \pm 0.12	4.2 \pm 0.58	2.0 \pm 0.04	4.3 \pm 0.61	0.90 \pm 0.03	1.8 \pm 0.06	0.9 \pm 0.17	3.5 \pm 0.71

Mean values for treated and control plots are significantly different ($p < 0.05$) for all years

TABLE 5. MEAN PERCENTAGE BURIED LEAF LITTER WEIGHT LOSS IN SPRAYED AND UNSPRAYED PLOTS

Season	Sprayed	Unsprayed	P value
1992/93	34.4 ± 2.6	37.6 ± 3.2	Non significant
1994	25.8 ± 2.5	27.7 ± 1.9	Non significant
1994/95	9.4 ± 2.4	10.1 ± 1.6	Non significant

TABLE 6. MAIZE GRAIN YIELD (T HA⁻¹) IN SPRAYED AND UNSPRAYED PLOTS (MEANS ± SE)

Season	Sprayed	Unsprayed	%yield increase	P value
1991/92	2.20 ± 0.06	1.90 ± 0.03	15.79	Non significant
1992/93	2.40 ± 0.05	1.50 ± 0.02	60.00	Non significant
1994	2.56 ± 0.04	1.80 ± 0.04	42.22	Non significant
1994/95	2.99 ± 0.03	2.79 ± 0.04	7.00	Non significant

microarthropod extraction from the buried leaf litter in this experiment, the reduced Collembola activity in pitfall traps reported above could help to explain this slow-down in decomposition. Cook *et al.* [6] showed that Collembola and Acari account for 90-100% of fauna responsible for buried leaf litter decomposition and that their numbers were significantly reduced by DDT sprays.

The results further show that lindane reduced early season stem borer infestations thereby protecting the young maize plants at their critical stage of vegetative growth (4-8WAE). This protection is revealed by low plant damage. Protection of young maize from borer damage is important in that early infestations lead to direct plant loss due to lack of compensatory growth. Timing of spray application before borer larvae tunnel the stems is equally crucial. Delayed applications will not reach the protected larvae in the stems. Furthermore, many larvae which will have hatched from an egg batch on a single plant will have migrated to other plants if sprays are delayed until the larvae reach their third instar stage. Plant damage can be related to pest populations and insecticide applications. Severely damaged plants would certainly give poor grain yields. The insecticide sprays used in this study contributed substantially to grain yield increase in maize for the four seasons. These yield increases were however, not statistically significant in any of the seasons. Grain yield increases have been reported on cowpea after DDT applications [7]. The above results therefore show that although Lindane 20 e.c. reduced stem borer population and damage on maize thereby improving grain yields slightly, it also showed short-lived adverse effects on most of the early season beneficial arthropods which highlights the importance of detailed studies to monitor residues in the soils and on some of the beneficial arthropod fauna groups.

ACKNOWLEDGEMENTS

The authors are grateful to the Swedish International Development Agency (SIDA) for financial support and to the Director General, International Atomic Energy Agency (IAEA) Vienna, Austria for availing the funds. Sincere thanks to the Director, Tropical Pesticides Research Institute (TPRI), Arusha, Tanzania for providing facilities for the field work at the Institute. The assistance extended to us by Dr. P. Chiverton and other staff of the Swedish University of Agricultural Sciences (SUAS), Uppsala, Sweden is highly appreciated. Lastly but not least we are thankful to Mr. G. Kayange and Ms. G. Shayo for their keen supervision of all the field work.

REFERENCES

- [1] TANZANIA GAZETTE, 18th January 1991 pp. 29-63.
- [2] CAREY, A.E., GARVEN, J.A., TAI, H., MITCHELL, W.G., WIERSMA, G.B., Pesticide residue levels in soils and crops from 37 states - National Soils Monitoring Program (iv). Pesticide Monitoring Journal, **12** (1979) 209-233.
- [3] TABOR, E.C. . Contamination of urban air through the use of insecticides. Transactions of the New York Academy of Sciences, Series 2. **28** (1966); 569.
- [4] CRITCHLEY, B.R., COOK, A.G., CRITCHLEY, U., PERFECT, T.J., RUSSELL-SMITH, A.. The effects of crop protection with DDT on some elements of the subterranean and surface active arthropod fauna of a cultivated forest soil in the humid tropics., *Pedobiologia* **20** (1980) 31-38.
- [5] EDWARDS, C.A.. Environmental aspects of the usage of pesticides in developing countries. Med. Fac. landbouw. Rijksuniv. Gent 42/2. (1977) pp. 853-68.

- [6] COOK, A.G., CRITCHLEY, B.R., CRITCHLEY, U., PERFECT, T.J., RUSSELL-SMITH, A., YEADON, R., The effects of soil treatment with DDT on the biology of a cultivated forest soil in the sub-humid tropics., *Pedobiologia* **19** (1979) 279-292.
- [7] PERFECT, T.J., COOK, A.G., CRITCHLEY, B.R., CRITCHLEY, U., DAVIES, A.L., SWIFT, M.J., RUSSELL-SMITH, A., YEADON, R.. The effect of DDT contamination on the productivity of a cultivated forest soil in the semi-humid tropics. *Journal of Applied Ecology* **16** (1979) 705-719.



PRELIMINARY STUDIES ON THE EFFECT OF ORGANOCHLORINE PESTICIDES ON BIRDS IN TANZANIA

A.S.M. IJANI, J.M. KATONDO, J.M. MALULU

Tropical Pesticides Research Institute,
Arusha, Tanzania

Abstract

Preliminary studies to investigate the effects of organochlorine pesticides on birds was conducted in Lower Moshi, NAFCO West Kilimanjaro, Arusha seed farm, Tropical Pesticides Research Institute (TPRI) farms, Manyara ranch and areas around Lake Victoria as well as in the TPRI laboratory in Tanzania. Large quantities of the pesticides particularly DDT, endosulfan, dieldrin, lindane and toxaphene are still being applied against pests of cotton, coffee, maize, beans and other crops as well as disease vectors in the country. Several groups of birds including waterbirds, African Fish Eagles, Marabou storks, Oxpecker, ducks, etc. were found feeding, roosting and swimming in the water and exposed to other substances that were contaminated with organochlorine pesticides and were presumably at risk. Analytical results from the tissues of the African Fish Eagles collected from Lake Victoria areas showed that the kidneys were contaminated with p,p'-DDE and o,p'-DDE at levels of 0.4 ng g⁻¹ and 1.45 ng g⁻¹ respectively. These organochlorine insecticides as well as β -HCH were also present in the brain and liver tissues. The levels of the organochlorine residues were well below the lethal and sublethal levels for bird raptors reported in the literature.

1. INTRODUCTION

Organochlorine pesticides, including DDT, dieldrin, endosulfan, lindane and toxaphene, are still being used to control agricultural pests and vectors of human and animal diseases in developing countries in the tropical and subtropical regions. Their use is greatly restricted in the developed temperate countries. This is mainly due to their high persistence in the environment and tendency to accumulate in living tissues and to move through different trophic levels in the food chains. Scientists have indicated that, compared with the temperate regions, the chemicals may behave in a different manner in tropical environments due to high temperatures, humidity, ultra-violet light, microbiological complexes etc. Information on the amount and effects of organochlorine pesticide residues present in water, food, soil, animals and birds in the tropical and subtropical countries is scanty. Talekar et al. stated that organochlorine pesticides were banned in some tropical countries without sufficient information about their impact on the local environment [1].

A number of organochlorine pesticides have been used in Tanzania for many years and are still registered for use against insect pests of coffee, especially in the Kilimanjaro and Arusha regions, and against pests of cotton particularly in the Lake Victoria zone, and those of maize and beans in many other regions as well as pests of domesticated animals and man. However, no research has been conducted to evaluate their effects on the higher fauna, particularly birds, despite the fact that in Tanzania there is a vast number of birds species including some endangered species. Therefore, studies on both the short and long-term effects of the organochlorine pesticides on the birds is very important. In the Lake Victoria zone the piscivorous and raptorial birds, particularly the African Fish Eagles, seem to be most affected by the pesticides. This is because some fishermen have been reported to pour endosulfan and other pesticides into the lake to kill fish and these pesticides, together with those washed into the lake from the crop fields during the heavy rains, accumulate in the lake so that when fish, birds and other animals that use the water they absorb the pesticides into their bodies. The African Fish Eagle (*Haliaeetus vocifer*) feeds on fish and swims in the contaminated water and hence it can be expected that high concentrations of the pesticides could accumulate in their tissues. The

population of these birds around Lake Victoria has been declining for the past few years (Mgallah, Pers. Com.). Probably the pesticides accumulate in the birds tissues and consequently can interfere with reproductive and other metabolic activities of the birds.

This study was conducted firstly, to identify the various organochlorine pesticides used in the Lake Victoria zone, Kilimanjaro and Arusha regions; secondly, to assess utilization of crop fields by avifauna in these areas; thirdly, to assess the distribution and abundance of the birds species particularly African Fish Eagles in the areas; fourthly, to investigate their feeding habits, breeding habits and success; fifthly, to determine the levels of organochlorine residues present in different tissues of the African Fish Eagles.

2. MATERIALS AND METHODS

2.1. Experimental site

Surveys were conducted in lower Moshi rice and sugarcane farms and NAFCO West Kilimanjaro (Kilimanjaro region); Arusha seed farm, Tropical Pesticides Research Institute (TPRI) farms, Manyara ranch, and Manyara National Park (Arusha region) and areas around Lake Victoria namely Bunda, Magu, Igoma, Mwanza and Butimba (Mwanza region) to investigate the pesticides used and birds species present. The experiment was started in March 1992 and took about three years to investigate the pesticides used and birds species present.

2.2. Surveys and ecological study

Pesticide shop keepers, cattle keepers, crop farmers, agricultural extension and health officers were visited and interviewed to find out the types and quantities of organochlorine pesticides used against crop pests and disease vectors in their areas. The time and persistence of the various pesticides used were recorded. The results obtained were then used to demarcate the areas into "contaminated areas", that is, those with regular use of organochlorine pesticides and "control areas" which are those areas with no such use.

Observations were made according to respective crops and pesticides spraying calendars and the numbers of dead birds and their behavioural changes during and immediately after the sprayings were recorded. The birds found in the different study sites were monitored once every month apart from those in TPRI which were monitored once every week. The birds found were sampled and their distribution and abundance, feeding habits, breeding habits and success and population dynamics were investigated. The movement behaviour of the birds was used to group them into sedentary and migrant groups. The results obtained from the ecological studies of the birds were used to determine suitable indicator species for further intensive studies.

2.3. Sampling and laboratory analysis

During the different surveys a small number of birds were sampled for residue analyses. However, because of some technical problems with GLC, only a few samples of the African Fish Eagles collected from the Lake Victoria areas could be analyzed. These birds were selected from different spots around the Lake Victoria area and shot during two different periods. The first sampling was effected at the beginning of the short rains in October 1994 and the second one in February 1995 during the heavy rainfalls. The killed birds were taken to the TPRI laboratory where liver, brain, kidney and fat tissues were removed from their bodies. The removed tissues and water which was also sampled from the same areas were then analyzed by

following the procedure outlined in the protocol provided by the IAEA but with slight modifications as shown in the technique given previously in order to suit circumstantial needs [2].

Some tissue samples were prepared for residue analysis but without using the SPE column. In this case an amount (20 g) of tissue was mixed with anhydrous sodium sulfate (four times the weight of the tissues) and ground. This mixture was tumbled in acetone + hexane (200 mL, 1 + 1 by volume) and left to settle for one hour. The suspension was filtered into a separating funnel containing sodium sulfate solution (200 mL). The hexane extract was separated and washed three times with sodium sulfate solution (50 mL, 20 g L⁻¹).

The hexane extract was concentrated to 2.0 mL and passed through a microcolumn (containing 1 g of 60 to 80 mesh silica gel for chromatography and 0.5 g anhydrous sodium sulphate) previously washed with hexane (50 mL) followed by methanol (50 mL) and dried at 75°C overnight. The column was eluted with a further volume of hexane (10 mL). The hexane extract was concentrated to 10 mL and 5 µL analysed by gas chromatography. The conditions were: column, length, 1.5 m, diameter, 6.35 mm; packing, 50 g kg⁻¹ QF1 on Chromosorb WHP; temperatures, column, 220°C, injector, 260°C, detector, 260°C; carrier gas, nitrogen; flowrate, 60 mL min⁻¹; detector, electron capture detector (⁶³Ni).

When the sample was extracted without using the SPE column broad unresolved peaks were observed. Consequently, the extracts were cleaned further by using microcolumn (or pipette column) as mentioned above and resolved peaks were obtained. The peaks were then quantitated using analytical standards after determining the retention time of the individual peaks and concentrations of the corresponding compounds. The pesticide residue concentrations were calculated as follows:

$$R = \frac{A (\text{weight std injected}) \times B (\text{sample peak ht/area}) \times C (\text{final extract vol.}) \times P (\text{std})}{D \text{ mL (inj. vol.)} \times E (\text{sample tissue weight}) \times F (\text{std peak height/area}) \times 100}$$

A = weight of standard used (ng/mL)

B = Area or height of sample peak (mm)

C = Final volume of extract (mL)

D = Microlitres injected (mL)

E = weight of tissue sample taken (g)

F = Area or height of standard peak (mm)

R = Residue concentration (ng/mL)

P = Purity of standard, %

3. RESULTS AND DISCUSSION

Endosulfan, dieldrin, lindane, DDT and several cocktails of DDT and other pesticides are still recommended as cheap and effective pesticides for controlling cotton and coffee insect pests as well as disease vectors in Tanzania and were found to be sold in large quantities in the Tanganyika Farmers Association (TFA) shops and other shops. Vegetables, maize, beans, rice, sugarcane and other crop producers were buying and using the various organochlorine pesticides against their crop pests, since there were few pesticides specifically recommended for control of pests of such crops. Water which was being used to irrigate sugarcane in the lower

Moshi crop contained high concentrations of dieldrin, endosulfan and DDT which were presumably washed from the coffee farms in Kilimanjaro and Arusha regions. Endosulfan which was being sold with the trade name Thiodan 35 EC, dieldrin and DDT were being used extensively against coffee, maize, vegetables and bean pests in the Arusha seed and TPRI farms. Toxaphene was extensively used against ticks and other animal disease vectors in the Manyara ranch. Endosulfan in ULV and EC formulations was being used in large quantities to control cotton and maize insect pests in most of the areas around Lake Victoria (Mwanza region). Large quantities of lindane and DDT had also been used in the lake zone in previous years. Endosulfan and other pesticides were being used to kill fish in Lake Victoria.

The species of birds which were observed in the lower Moshi, NAFCO West Kilimanjaro, TPRI, Arusha seed farm, Manyara Ranch and Manyara National Park are listed in Table 1. The lower Moshi rice irrigation Scheme (1200 ha) and sugarcane plantation are inhabited by numerous species of birds most of which are permanent residents, and pests, of rice. Queleas visit the farms in millions between July and February and can devastate the rice crop if not managed. The majority of the waterbird species (Egrets, Herons, Ibises, Storks) were observed eating rice grains, drinking and swimming in standing water which was probably contaminated with endosulfan, dieldrin, DDT and other pesticides. During the survey conducted in the area in May 1994, immediately after extensive spraying of the sugarcane and rice fields with endosulfan and other pesticides, some birds particularly the weaver birds and fiscal shrikes were found dead and some failed to fly for long distances. The dead birds were collected for analysis of pesticides residues in the laboratory but the analyses were not done owing to problems we had with the gas chromatograph.

NAFCO West Kilimanjaro is an open Woodland area inhabited by a number of bird species including the Grey Crowned Cranes and Cattle Egrets. The birds are likely to be affected by DDT, dieldrin, endosulfan and other pesticides which are frequently applied against pests of coffee, beans, maize, vegetables and other crops. TPRI and Arusha seed farms have relatively fewer species of birds but only a few of them, including the Abdimn's and stork, are sedentary hence are likely to be affected by the organochlorine pesticides which are being used in the areas. Yellow weavers which feed on bollworms in maize crops in Arusha seed farm and TPRI experimental fields are at risk. Fiscal shrikes may be exposed to organochlorine pesticides, when feeding on insects which survive after spraying in maize, coffee and bean farms. Manyara ranch, a wooded grassland, has a high diversity of bird species and shares many bird species with lower Moshi. Several waterbirds were drinking and swimming in the five ponds which were being used for washing cattle after the water had been treated with DDT, dieldrin, endosulfan and particularly toxaphene. Scavengers, such as Marabou storks, are permanent residents; birds of prey, especially vultures, also visit the ranch. These birds as well as oxpecker, doves, weavers and other species drink the contaminated water and some may also eat cattle meat contaminated with the pesticides. The five ponds are the only sites in the area where the birds can get water. Other birds found in the ranch include ducks, geese, ibises etc., which appear to be local immigrants from nearby Manyara National Park. Queleas also utilized the acacia trees in the area as nesting and breeding sites and later migrated to feed on wheat, rice and barley in the nearby farms. The birds which are permanent inhabitants in the Manyara National Park and do not visit the areas contaminated with the organochlorine pesticides are considered to be in the "control" area. The amount of organochlorine residues obtained from the tissues of these birds (if any) can be compared with that obtained from the similar species of bird which live in the contaminated areas. The Marabou stork a raptorial and sedentary species, is a good indicator species for this kind of study as it is found in both the National Park where no pesticide is being used and the ranch where several pesticides are applied.

Table 1. Species, distribution and abundance of birds.

Species	Manyara Ranch	Lower Moshi	NAFCO-West Kilimanjaro	TPRI-Arusha seed farm	Manyara National Park
Ostrich	+				++
Little Grebe	+			+	
Long-tailed Cormorant	+	+			
Squacco heron	+	+			+
Great white Egret	+	+			
Black Heron	+	+			
Little Egret					+
Black-headed heron		+		+	
Purple Heron		+			
Cattle Egret		++	++	+	+++
Yellow billed egret		++			+
Woolly beaked stork		+			+
Marabou stork	+	+			
Yellow billed stork		+			+
Hadada Ibis		+		+	+
Glossy Ibis		+			
Sacred Ibis	+	+		+	+
African Spoonbill	+	+			+
White faced Tree Duck	+	+			+
Egyptian Goose	+	+			+
Red-billed Duck	+	+			+
Knob-billed Duck	+	+			+
Macawa Duck					+
Augur Buzzard	+	+			+
Black shouldered Kite		+			+
Hilderbrandt's Francolin	+				+
Helmeted Guinea fowl	+				+
Button Quail	+	+			+
Grey Crowned Crane	+	+	++		+
Blacksmith Plover	+	+		+	
Black Crake		+			
Two-banded Courser		+			
Chestnut-bellied Sandgrouse	+			+	
Namagua Dove	+	+			
Ring-necked Dove	+	+		+	
Laughing Dove	++	+		+	
Red eyed Dove	+	+			+
White-bellied Go-away bird	+				+
White-browed Coucal	+	+			+
African Jacana		+			+
Sandpipers		+			+
Speckled Mousebird	+	+	+	+	+
Little Boo-eater	+			+	
African Hoopoe	+				
Red-billed Hornbill	+				
Long tailed fiscal	+	+		+	
Fiscal shrike	+	+	+	+	
White Crowned Shrike	+	+			
Hilderbrandt's Starling	++		+		
Superb Starling	++		+		
Ashy Starling	++				
Yellow billed oxpecker	++				
Queleas	+++	+++	+++	+	
YellowWeavers (<i>Ploceus</i> ssp)	+	+++		++	
Pelicans					+
White storks				+	
Black kites				+	
Abdim's				+	

number of birds : + <100

++ 100 - 1000

+++ >1000

In Mwanza region, the African Fish Eagle, a raptorial bird was found in all the areas that were surveyed around Lake Victoria. The areas included Butimba, Igoma, Nyamahanga, Tahamu near Rubara River and Guta where the number of birds observed ranged between four and eight. The Eagles were found together with the Marabou stork, Black kite, Great Egrets, Yellow billed duck, Yellow billed stork, Black headed herons and other species of birds.

Possibly, these birds have succeeded in staying in the same place because their ecological niches do not quite overlap. The African Fish Eagle feeds on fish and small animals such as frogs living in shallow water along the lake shore and river streams. They feed during day times especially between 10.0 a.m. and 4.0 p.m. It appears that the African Fish Eagles build their nests and lay their eggs during the dry season or towards the end of the rainy season because no nests with eggs were found during the survey, which was conducted at the beginning of the rainy season. Only two young birds were spotted. These birds laid two eggs which succeeded in hatching. Their nests were 30 cm in diameter and built by using small twigs or tree branches. These were found on the highest tree tops and were not easily viewed. Such behaviour of building nests in high positions and the great parental care which the African Fish Eagles have towards their young ones, enable all their chicks to survive normally. However, because of the small number of eggs which the birds lay, their population is usually small in most areas.

The analytical results from the different tissues of the African Fish Eagles collected from Mwanza region showed that the kidney contained what is probably o,p'DDE and p,p'DDE and was more heavily contaminated with residues of DDT type than the liver and brain tissues (Table 2). Possible traces of lindane and β -HCH were also observed in the brain and liver tissues. The work on organochlorines in Africa has generally concentrated on residue levels

Table 2. Residues of organochlorine insecticides in tissues of African Fish Eagle and in samples of water taken from lakes

Sampling		Sample identity	Residues	
Time	Location		Concentration, $\mu\text{g g}^{-1}$	Identity
October 1994	Magu	liver	0.002	o,p'- DDE
		brain	0.024	o,p'- DDE
		kidney	0.430	o,p'- DDE
			0.022	p,p'- DDE
	Igoma (MZ)	liver	0.009	o,p'- DDE
			0.007	p,p'- DDE
		kidney	0.200	o,p'- DDE
			0.460	p,p'- DDE
	Mwanza	brain	0.010	o,p'- DDE
		0.006	p,p'- DDE	
		0.019	lindane	
February 1995	Mwanza	kidney	1.45	o,p'- DDE
		liver	0.001	β - HCH
		brain	0.001	β - HCH
	Kayenyeze(MZ)	water	0.001	o,p'- DDE
		water	0.114	o,p'- DDT
		Bunda	0.004	o,p'- DDE
		Magu	0.154	o,p'- DDT

without investigating the associated effects except for egg shell thinning [3]. Therefore, the side effects on birds due to the organochlorines, have been estimated by comparison with the results of studies made in other countries. This has also been difficult mainly because sensitivities to pesticides vary widely between birds species [4]. However, diurnal raptors have broadly similar sensitivities [5]. Mortalities of raptors due to DDT and its residues were reported to occur when levels of DDT exceeded 30 mg kg⁻¹ [6]. In addition to raptors, DDT and its residues are reported to cause eggshell thinning and breakage as well as addling and reduction of hatchability [7]. Several studies of organochlorine pesticides in Africa have indicated that the residue levels are generally lower than those found in Europe and North America [4]. This is mainly due to a lower proportion of agricultural land in Africa and the shorter persistence of organochlorines in tropical as compared to temperate climates. Assuming that the amounts of residues obtained from tissues of the African Fish Eagles during this study can be confirmed they are well below lethal and sublethal levels calculated by other scientists [6]. However, the data obtained cannot be considered as representative results for organochlorine residues present in the African Fish Eagles found in Mwanza areas. This is because other important tissues of the bird including muscles and fats have not been analyzed for the residues. In addition, the birds were sampled only at the beginning of the rainy season, when most of the pesticides applied in the cotton fields had not been washed into the lake where the birds were feeding and swimming. Furthermore, the number of samples analyzed was small hence could not provide consistent and reliable results and the identity of the peaks has not been confirmed by GC-MS.

However, together with studying the feeding habits, breeding habits and home ranges of the indicator bird species in relation to contamination with organochlorines, the tentative data on the African Fish Eagles indicate that it is necessary to determine the levels of organochlorine residues that are present in tissues and or eggs of the representative samples of the exposed birds and to investigate their associated effects. The preliminary results presented in this paper show a need for the further study of the effects of organochlorine pesticides on birds in Tanzania such as the African Fish Eagles and Marabou storks found around the Lake Victoria areas where lindane and endosulfan are extensively used against crop pests and on the Oxpecker birds in the Manyara areas where large quantities of toxaphene are applied in cattle dips [7].

ACKNOWLEDGEMENTS

We are grateful to the International Atomic Energy Agency/FAO/SIDA and the Director of TPRI for their financial support which enabled us to carry out this project.

REFERENCES

- [1] TALEKAR, N.S., CHEN, J.S, KAO, H.T., Long term persistence of selected insecticides in subtropical soil: their absorption by crop plants, *J.Econ. Entomol.* **76** (1983) 207.
- [2] WATTS, R.R., Analysis of pesticide residues in human and environmental samples. A compilation of methods selected for use in pesticide monitoring programs, US Environmental Protection Agency, Health Effects Research Laboratory, Environmental Toxicology Division Research Triangle Park, North Carolina, USA (1979).

- [3] CRICK, H.Q.P., Organochlorine pesticides and birds of prey in Africa, Proc. VII Pan. Afr. Orn. Congr. (1992) 171-189.
- [4] de KOCK, A.C., LORD, D.A., Chlorinated hydrocarbon residues in African Fish Eagle (*Haliaeetus vocifer*) eggs, Institute for Coastal Research Report No.10 (1986).
- [5] NEWTON, I., Effects of organochlorine pesticides on birds, Proc. 2nd Symp. Afr. Predatory Birds. (1984) 151-159. Durban: Natal Bird Club.
- [6] BLUS, L.J., HENNY, C.J., STAFFORD, C.J., GROVE, R.A., Persistence of DDT and metabolites in wildlife from Washington State Orchards, Arch. Environ. Contam. Toxicol. **16** (1987) 467-476.
- [7] NEWTON, I., Population ecology of raptors, Edited by T. Berkhamsted and A.D. Poyser (1979).

ADVERSE EFFECTS OF LINDANE IN A MAIZE AGRO-ECOSYSTEM IN UGANDA



E.M. TUKAHIRWA , W.TINZAARA, B.T.KIREMIRE
Institute of Environment and Natural Resource, Makerere University,
Kampala, Uganda

Abstract

The impact of lindane, at commercial rates of application, on invertebrate fauna, soil microbial activity, earthworm populations, crop damage and yields in a maize agro-ecosystem was studied and compared with unsprayed control plots of maize using a pitfall trap, D-Vac suction, litter bag, the earthworm formalin expulsion and crop assessment methods. The findings of the study generally portrayed lindane as having very few effects on the maize agro-ecosystem.

1. INTRODUCTION

To increase food production to cope with the ever rising demands by increasing populations in developing countries, agricultural practices must become increasingly intensive and, for the foreseeable future, pesticides will continue to be used.

However, the dangers to individuals and the environment of unqualified dependence on pesticides are very well documented [1-3]. There is no doubt therefore that Third World farmers too need to be made aware of the ills of overdependence on these chemicals. Hence chemicals used in developing countries should be of the type that does not pose risks that are too serious to farmers and the environment. Many of the pesticides presently in use in developing countries are both expensive and lacking in the persistence necessary for them to be effective on tropical pests. The ones that are can be toxic and pose a hazard to the often poorly trained applicators of the Third World.

Until its use became restricted worldwide, lindane was an acceptable pesticide for the Third World mainly on account of its relative cheapness, combined with its low toxicity to applicators. However because of its persistence, its residues tend to accumulate in soil and aquatic sediments, and in plant and animal tissue which led to the restriction on its usage.

The economic viability of lindane in the tropics is however, not fully documented. Whereas its persistence as demonstrated in researches done in temperate environments is an undesirable characteristic, the extent of these negative attributes in tropical environments has not been fully investigated. In any case, some persistence can be a desirable attribute for tropical farmers.

It is well known that temperature, among other factors, influences the dissipation rates of pesticides in the environment [4]. In that case, it is possible that under tropical conditions, lindane will dissipate at rates which preclude local accumulation of residues in the soil and wildlife. This implies that its use may still be acceptable, and not cause significant environmental problems in the tropics.

The research effort presented in this report is part of a the FAO/IAEA co-ordinated programme to provide more information on the merits and demerits of using lindane in the tropics. It is hoped that the data so generated will contribute to the resolution of the argument as to whether lindane should or should not be used by Third World farmers, especially given that it

possesses attributes that they probably would welcome. The following account is based on a field study that was carried out at Kawanda Research Station, Uganda, each year during the months of April-July of the years 1992-1995. The principle objective of the study was to assess the impact of lindane on the invertebrate fauna and to determine its impact on yields.

2. MATERIALS AND METHODS

2.1. Field trial design and preparation

In the study periods of April-July 1992 and 1993, two parallel fields of about 0.7 hectares were used for the experiments. The experiments were designed to have four sprayed plots in one field and four control plots in another field. The fields were at least 50 m apart, while the plots within the fields measured 10 m × 10 m and were 10 m apart and in the centre of the fields, leaving 10 m from the edges. In the study periods of April-July 1994 and 1995, one field of about 0.8 hectares was used for each of the experiments. In this case the experiments were designed to have four plots (i.e. measuring 10 m × 10 m and 10 m apart) sprayed with lindane, with another four plots left as controls. The positions of the sprayed and unsprayed plots were randomised, so as to give a randomised block experiment with four blocks and two treatments in each block. The soil properties for the years 1994 and 1995 for each of the blocks are given in Tables 8 and 9.

In all cases, the fields were ploughed by tractor and planted with dressed maize seed (*Zea Mays* var. Namulonge) at a spacing of 60 cm between rows and 30 cm within rows. Initially two seeds were placed in each hole, but these were later thinned to single plants after germination. All weeding was done manually, and the NPK fertilizer was added at the rate of 200 kg/ha when the crop was four weeks old.

The first application of lindane (bought locally as Gammalin), using a 20L knapsack sprayer at a rate of 0.5 kg AI ha⁻¹, was normally made four weeks after germination and two weeks after the laying of pitfall traps.

2.2. Pitfall traps

Following the demarcation of plots (i.e two weeks after germination), four pitfall traps (10 cm rim diameter and 10 cm deep) were randomly placed in each plot to trap any surface-active invertebrates that came by. The traps contained formalin (20 g L⁻¹) as the trap agent and preservative. Traps were emptied once a week.

2.3. D-vac sampling

The D-vac suction apparatus was used for both plant and ground sampling in the experiments conducted in the study periods of 1994 and 1995. It was normally first done three weeks after germination, and subsequently every after two weeks. For ground sampling, one sample was taken per sampling occasion per subplot, each sample consisting of four sub-samples taken from four different places in each subplot. Each sub-sample consisted of five-seconds suck with the sampling cone tightly pressed to the ground. Similarly for plant sampling, one sample was also taken per sampling occasion per subplot, each sample consisting of four sub-samples taken randomly by sucking the maize plant inserted into the D-vac cone. At the earlier stages, the D-vac cone was placed over the plants while sampling. Later, as the plants became taller, the cone was used to sweep the plants from all sides. In all cases, the

invertebrates caught were killed with chloroform and kept in formalin (50 g L⁻¹) in sample bottles for later identification and counting.

2.4. Assessment of microbial activity

In order to compare the level of microbial activity in the different plots, leaf discs of a Musanga tree (*Cecropioides* sp) were buried to a depth of about 5 cm. The discs were buried in bags made of fine mesh plastic netting, with each bag containing 20 discs of 1.5 cm diameter. The discs were oven-dried at 70°C for at least 24 hour, before weighing both at the beginning of the experiment and after 2 months when the experiment was terminated. The loss in disc weight represents the proportion of the disc tissue that was consumed by microbes in the soil.

2.5. Earthworm sampling

The standard earthworm expulsion method was used [4]. A half-metre quadrat was placed on the soil surface in each of the fields, and 9 litres of a dilute formalin solution (50 mL of 40% formaldehyde in 9 litres of water) poured slowly within the quadrat. Earthworms would then emerge from the soil during the following 15 minutes. Two expulsion samples were taken per study period, once at the beginning of the experiment when the maize was two weeks old, and the other towards the end of the experiment when the maize had matured. Earthworm abundances at these two stages of the maize phenology were compared.

2.6. Pest damage

The maize crop was monitored by physical inspection for pest damage in all subplots throughout the experimental period. However, more intensive work was done at harvest to assess damage by *Busseola* spp.

2.7. Yield

Maize yield was estimated from four subplots (2 m × 2 m) in both sprayed and unsprayed subplots. These subplots for estimation of yield were usually located outside, but close to the subplots where sampling activities had taken place during crop growth.

2.8. Statistical methods

Students t-test was used for analysis of the data collected in the experiments conducted in the study periods 1992 and 1993. A two-way analysis of variance was employed using a statistical package (Minitab for Windows) for the data collected in the experiments conducted in 1994 and 1995.

3. RESULTS

3.1. Pitfall trap catches

Overall, there were no significant differences between the catches from the sprayed fields and those from the control fields except for fauna groups of Gryllidae and Blattidae (P<0.05) and Collembola (P<0.01) (Table 1) indicating that lindane application generally did not have a dramatic effect on the invertebrate populations in the sprayed plots.

TABLE 1. The mean catch per plot (\pm S.E) of selected key organisms caught using pitfall traps from sprayed and unsprayed plots during the years 1992-1995.

PRE-SPRAYING:

YEAR	FAUNA GROUP	SPRAYED (\pm S.E)	CONTROL (\pm S.E)	t-test Value	ANOVA P-value
1994	Araneida	9.8 \pm 2.2	6.8 \pm 1.4	-	0.207NS
	Acarina	5.0 \pm 1.5	7.3 \pm 3.8	-	0.516NS
	Gryllidae	4.5 \pm 0.9	4.3 \pm 1.3	-	0.817NS
	Formicidae	166.3 \pm 56.1	233.8 \pm 127.0	-	0.517NS
	Collembola	102.3 \pm 54.1	57.5 \pm 36.2	-	0.309NS
1995	Araneida	29.5 \pm 8.1	39.3 \pm 4.3	-	0.226NS
	Acrididae	17.5 \pm 5.6	12.3 \pm 0.5	-	0.697NS
	Acarina	3.0 \pm 1.5	2.3 \pm 1.9	-	0.769NS
	Gryllidae	12.5 \pm 1.9	17.5 \pm 1.9	-	0.164NS
	Formicidae	290.3 \pm 40.1	304.0 \pm 61.5	-	0.783NS
	Blattidae	1.8 \pm 1.4	3.0 \pm 1.2	-	0.43NS
	Collembola	0.8 \pm 0.8	0.5 \pm 0.5	-	0.391NS

POST-SPRAYING:

1992	Araneida	21.3 \pm 4.8	17.5 \pm 3.8	0.42 NS	-
	Gryllidae	29.5 \pm 5.5	53.8 \pm 9.1	2.56 *	-
	Formicidae	577.8 \pm 91.9	719.8 \pm 80.1	0.88 NS	-
	Carabidae	21.3 \pm 4.3	37.5 \pm 7.5	0.43 NS	-
1993	Acrididae	48.0 \pm 6.1	55.5 \pm 7.8	0.79 NS	-
	Gryllidae	39.3 \pm 8.1	67.0 \pm 7.0	2.17 NS	-
	Formicidae	1056.0 \pm 488.0	707.5 \pm 126.1	1.77 NS	-
	Blattidae	14.0 \pm 5.1	29.3 \pm 3.4	2.35 NS	-
	Collembola	80.8 \pm 30.2	7.3 \pm 2.1	5.24 **	-
1994	Araneida	57.5 \pm 11.4	46.3 \pm 6.3	-	0.191NS
	Acarina	48.8 \pm 21.6	47.5 \pm 11.4	-	0.855NS
	Gryllidae	39.5 \pm 2.2	34.3 \pm 5.6	-	0.325NS
	Formicidae	1186.5 \pm 205.0	1452.8 \pm 500.1	-	0.528NS
	Collembola	2428.0 \pm 993.0	1507.3 \pm 171.8	-	0.064NS
1995	Araneida	66.0 \pm 6.7	89.8 \pm 38.8	-	0.694NS
	Acrididae	38.6 \pm 11.6	28.3 \pm 1.4	-	0.783NS
	Acarina	44.8 \pm 26.0	31.0 \pm 21.4	-	0.131NS
	Gryllidae	58.0 \pm 10.2	76.3 \pm 6.7	-	0.225NS
	Formicidae	1180.3 \pm 324.6	2382.0 \pm 761.1	-	0.372NS
	Blattidae	48.3 \pm 8.1	60.5 \pm 9.5	-	0.008*
	Collembola	17.0 \pm 1.4	27.0 \pm 5.9	-	0.288NS

NS= Not significant ($P > 0.05$) *= Significant ($P < 0.05$)

**= Significant ($P < 0.01$)

TABLE 2. Mean number (\pm S.E) of selected organisms caught using a D-vac sampler from the sprayed and unsprayed control plots.

A: GROUND SAMPLES:

YEAR	ORDER	SPRAYED(\pm S.E)	CONTROL(\pm S.E)	P-Value
1994	Araneida	2.0 \pm 0.4	6.5 \pm 0.6	0.002 *
	Homoptera	21.8 \pm 3.2	30.8 \pm 4.9	0.149 NS
	Hymnoptera	14.5 \pm 0.6	14.5 \pm 1.3	0.995 NS
	Diptera	98.0 \pm 12.3	114.5 \pm 18.5	0.138 NS
	Coleoptera	6.0 \pm 1.3	8.8 \pm 1.9	0.241 NS
1995	Araneida	5.5 \pm 0.9	8.5 \pm 2.3	0.289 NS
	Homoptera	17.0 \pm 5.3	20.5 \pm 4.3	0.546 NS
	Hymnoptera	17.8 \pm 3.7	18.0 \pm 2.0	0.212 NS
	Diptera	68.8 \pm 12.2	55.0 \pm 15.7	0.042 *
	Coleoptera	7.5 \pm 2.5	4.5 \pm 1.4	0.181 NS

B: PLANT SAMPLES:

YEAR	ORDER	SPRAYED(\pm S.E)	CONTROL(\pm S.E)	P-Value
1994	Araneida	2.0 \pm 0.7	3.0 \pm 0.7	0.091 NS
	Homoptera	8.8 \pm 2.7	9.3 \pm 3.8	0.940 NS
	Hymnoptera	13.0 \pm 2.3	18.8 \pm 1.9	0.267 NS
	Diptera	27.5 \pm 1.4	46.3 \pm 5.3	0.037 *
	Coleoptera	1.0 \pm 0.4	1.5 \pm 0.6	0.606 NS
1995	Araneida	1.8 \pm 0.9	4.0 \pm 0.7	0.029 *
	Homoptera	7.5 \pm 1.3	8.0 \pm 1.8	0.229 NS
	Hymnoptera	15.0 \pm 9.3	9.8 \pm 2.9	0.628 NS
	Diptera	81.0 \pm 32.0	82.8 \pm 27.1	0.855 NS
	Coleoptera	4.3 \pm 2.5	3.5 \pm 1.3	0.252 NS

NS=Not Significant ($P>0.05$). *=Significant ($P<0.05$)
 **=Significant ($P<0.01$)

TABLE 3. Mean percentage(\pm S.E) loss in weight of buried leaf discs of a Musanga tree over a period of two months in sprayed and unsprayed control plots.

YEAR	SPRAYED(\pm S.E)	CONTROL(\pm S.E)	P-Value
1992	63.6 \pm 2.0	66.2 \pm 1.5	0.383 NS
1993	24.8 \pm 4.7	41.2 \pm 5.6	0.001 **
1994	16.4 \pm 2.6	20.7 \pm 1.4	0.233 NS
1995	36.1 \pm 5.5	52.3 \pm 8.5	0.045

*NS= Not Significant ($P>0.05$) *= Significant ($P<0.05$)
 **= Significant ($P<0.01$)

TABLE 4. Total number of earthworms expelled from sprayed and unsprayed control plots.

YEAR	SPRAYED	CONTROL
1992	0.0	0.0
1994	10.0	9.0
1994	6.0	5.0
1995	2.0	7.0

TABLE 5. Mean percentage cob damage by *Busseola* spp assessed at harvest in sprayed and unsprayed control plots.

YEAR	SPRAYED(±S.E)	CONTROL(±S.E)	P-Value
1992	4.5±0.7	5.2±1.3	0.623 N
1993	3.6±1.0	2.7±0.9	0.519 N
1994	8.4±3.3	11.5±4.5	0.224 NS
1995	6.0±1.1	9.3±3.9	0.869 NS

NS=Not Significant (P>0.05)

TABLE 6. Mean weight (Kg) of maize yield per hectare from sprayed and unsprayed control plots.

YEAR	SPRAYED x 10 ⁴	CONTROL x 10 ⁴	P-Value
1992	#	#	#
1993	#	#	#
1994	0.90	0.88	0.594 NS
1995	1.00	0.90	0.173 NS

#= No Data Available, NS= Not Significant, P>0.05

TABLE 7. Comparison of rainfall (mm) availability between different years for the study period January-July

	JAN		FEB		MAR		APR		MAY		JUN		JUL	
	RD	TOT	RD	TOT	RD	TOT	RD	TOT	RD	TOT	RD	TOT	RD	TOT
1990	#	#	#	#	12	284.4	14	235.7	15	304.2	5	16.6	6	1.7
1991	10	39.2	11	93.8	13	230.4	14	256.0	17	116.7	9	110.3	6	27.3
1992	7	56.9	5	30.0	9	73.5	17	97.6	14	128.3	8	113.4	7	3.8
1994	5	57.6	9	69.0	5	83.4	8	109.4	19	103.3	9	140.8	9	4.8
1995	3	18.4	5	14.3	12	223.6	14	148.7	17	150.0	8	80.6	10	4.4

= Equipment was dysfunctional; RD= Rainfall days; TOT=Rainfall total

SOURCE: Kawanda Agricultural Research Institute

Table 8. Soil characteristics (1994)

Composition	Block 1	Block 2	Block 3	Block 4
Sand	58.36	64.24	62.28	60.32
Clay	27.49	27.49	25.53	31.41
Silt	14.15	8.27	12.19	8.27
% org. mat.	5.45	7.73	6.82	8.82
% nitrogen	0.09	0.10	0.11	0.12
pH	5.0	4.8	5.2	5.4

Table 9. Soil characteristics (1995)

Composition	Block 1	Block 2	Block 3	Block 4
Sand	43.0	47.0	55.0	49.0
Clay	40.0	36.0	34.0	34.0
Silt	17.0	17.0	11.0	17.0
% org. mat.	4.09	4.24	3.94	3.79
% nitrogen	0.14	0.16	0.13	0.13
pH	5.4	5.5	5.6	5.6

The overall numbers of individual groups throughout the experimental periods are presented in Table 1. The figures show that none of the invertebrate groups was ever significantly affected at any time during the cropping season. Most groups decline in numbers as the season progresses, and this happens equally in the sprayed field as in the unsprayed field.

3.2. D-vac samples

Although the individual number of invertebrates caught by both ground and plant sampling are generally higher in the control plots than in the sprayed ones, most differences were not significant ($P < 0.05$) except for Araneida and Diptera (Table 2).

3.3. Assessment of microbial activity

Soil microbial activity in control plots was generally higher than in the sprayed plots with differences significant in 1993 ($P < 0.01$) and in 1995 ($P < 0.05$) (Table 5).

3.4. Earthworm sampling

In general, few earthworms were expelled from the soil in either sprayed or unsprayed control plots on any occasion and no significant were recorded (Table 4). Soil moisture seems to have been the principal limiting factor, as the earthworms were only be expelled during the first sampling occasions when rainfall was high (Table 7). No earthworms were expelled during the second sampling occasions when the climatic conditions were rather dry.

3.5. Pest damage

The maize crop was monitored regularly to assess pest damage, being done by regular inspection of the individual plants within the demarcated plots. There were no pests on the maize during its vegetative growth period in all years. The maize stalk borer (*Busseola* sp) was the only significant pest which, however, appeared once the maize had formed cobs very late in the season. The damage recorded in the sprayed plots was not significantly different from that recorded in the control plots in all study periods (Table 5).

3.6. Maize yield

The sprayed plots gave a slightly higher, not statistically significant ($P < 0.05$), mean cropping yield compared with the control plots. In both cases, the yields were quite low, mainly due to insufficient rainfall during the cob-forming period.

4. DISCUSSION

The results of the experiments do not show great differences between the lindane sprayed and unsprayed control plots. Focusing on the macro-invertebrate fauna, it is clear that although there were some numerical differences, these were mostly not statistically significant, implying a merely moderate impact by the pesticide. Such a low impact may have been due to rapid dissipation or adsorption of the pesticide onto soil particles [4]. But it must be pointed out also that in all study periods, a few weeks of the growing season were characterised by high rainfall. This could have quickly leached away the pesticide from the maize and soil surfaces although organochlorine leaching is not considered to constitute a significant mode of loss. Whatever the cause, however, the pesticide impact was low.

The microbial activity seems to have been less in sprayed as compared to control plots. The low mean percentage weight loss in both sprayed and control plots could be explained by the rather tough leaves used, suggesting that the two months period was not long enough to allow for higher total decomposition. Although not conclusive, this observation is in line with those of other workers [5], who registered reduced microbial activity following application of DDT.

A meaningful measure of pesticide impact on an agro-ecosystem must relate to the capacity of the system in the short term to maintain crop yield [3]. In this experiment, cob damage was slightly less and the yields slightly higher, though in no case were the differences statistically significant between sprayed and unsprayed control plots. The mean maize yields were higher in sprayed than in control plots. This implied that lindane was, as expected, providing some measure of control, albeit limited, probably owing to the low pest load.

Generally, any differences in invertebrate populations, loss in weight of leaf discs, plant damage and grain yields could be attributed to the application of lindane since the inter- and intra-field soil factors at the Research Station, which are known to be small, could not have been responsible for such differences. The major observation however, is that the effects of lindane application were small on the population levels of macro-invertebrates or the functional capabilities of micro-organisms in the soil.

The findings of the experiments generally portray lindane as not being highly damaging to the agro-ecosystem, and therefore an acceptable option for insecticide application in the tropics.

ACKNOWLEDGEMENT

The authors wish to thank FAO/IAEA/SIDA for supporting the work.

REFERENCES

- [1] EDWARDS, C.A., Impact of herbicides on soil ecosystems, *Critical Rev. in Plant Sci*, **8** (1989) 221-257.
- [2] EDWARDS, C.A., Methods of assessing populations of soil inhabiting invertebrates, *Agric. Ecosys. and Environ.* **34** (1991) 145-176.
- [3] PERFECT, T.J., A.G. COOK, B.R. CRITCHLEY, U. CRITCHLEY, A.L. DAVIES, M.J. SWIFT, A.RUSSELL-SMITH, R. YEADON., The effect of DDT contamination on the productivity of a cultivated forest soil in the sub-humid tropics, *J. Appl. Ecol.*, **16** (1979) 705-719.
- [4] EDWARDS, C.A., The environmental impact of pesticides *Parasitis* **86** (1987) 309-329.
- [5] COOK, A.G., B.R. CRITCHLEY, U. CRITCHLEY, T.J. PERFECT, A. RUSSELL-SMITH, R.YEADON, The effects of soil treatment with DDT on the biology of a cultivated soil in the sub-humid tropics, *Pedobiologia* **19** (1979) 279-292.

**NEXT PAGE(S)
left BLANK**



THE EFFECT OF LINDANE ON NON-TARGET FAUNA IN A MAIZE AGRO-ECOSYSTEM IN ZAMBIA

Y.D. DEEDAT, S. CHANDA, A M. CHIVUNDU, G. KALEMBE, C.D. MECHA
School of Agriculture, University of Zambia,
Lusaka, Zambia

Abstract

The effect of lindane on non-target fauna in a maize agro-ecosystem was studied in Zambia in 1992 and 1993. While lindane was effective against the stalk borers, a target pest, it also affected other non-target fauna. Ants, spiders and springtails were significantly reduced. However the effect was only transient and lasted for approximately two months. Lindane appeared to have no real effect on aerial non-target fauna or on soil inhabiting microorganisms

1. INTRODUCTION

Organochlorine pesticides such as DDT, endosulfan, lindane and toxophene have been widely used for the control of crop and animal pests in Zambia. These pesticides are highly persistent and may affect many of the non-target organisms. Acquiring quantitative information about the effects of pesticides on non-target fauna is fundamental in the determination of the role of insecticides in integrated pest management programmes.

In Zambia some studies have been made to determine the side effects from the use of organochlorine pesticides. However, these studies have been conducted solely in relation to tsetse fly eradication programmes [1,2]. There is no information on the effect of organochlorine pesticides within the agro-ecosystem. Thus this study was undertaken to determine the non-target effects of lindane in a maize agro-ecosystem.

2. MATERIALS AND METHODS

2.1 Experimental layout

This study was conducted at UNZA farm, 25km east of Lusaka during the 1991/92 and 1992/93 growing season. The field was cultivated by a tractor and planted with maize (Table I) at the rate of 25 kg ha⁻¹ with a row to row spacing of 90 cm and a plant to plant distance of 20 cm. fertiliser, at the rate of 400 kg ha⁻¹ was applied as basal dressing at planting. Urea was top dressed at the rate of 200 kg ha⁻¹ when the crop was at knee height. Soil characteristics (Table I) as well as weather data (Table II) for the experimental site were recorded during each cropping season.

Two plots, each 100 m × 50 m, were marked out and separated by a distance of 100 m. About three weeks after plant emergence one plot was sprayed with Gammaxane 20 (Table I) at the rate of 1 kg AI ha⁻¹ using a CP15 knapsack sprayer. The other plot was untreated and served as a control. Within each plot, four sub-plots (5 m × 5 m) were demarcated with an isolation distance of 5 m. These sub-plots were used for all sampling activities. All data were analyzed using F-test at P = 0.05 level of significance.

Table I. Crop, plot and soil characteristics of the experimental site at the University of Zambia farm for the 1992 and 1993 growing seasons.

Component	Characteristic	1992	1993
crop	variety	MM 602	MM 603
	seed	undressed	undressed
	planting date	25 November, 1991	30 December, 1992
	spraying date	7 January, 1992	20 January, 1993
	harvest date	6 May, 1992	5 June, 1993
Soil	sand	55.8 ± 1.3 %	65.6 ± 2.8 %
	silt	10.0 ± 0.4 %	14.3 ± 6.4 %
	clay	34.7 ± 1.4%	20.1 ± 3.6 %
	organic carbon	0.8 ± 0.1 %	0.3 ± 0.2 %
	organic matter	1.36 ± 0.1 %	0.5 ± 0.4 %
	pH (soil : water=1:2.5)	6.3 ± 0.1	9.7 ± 0.5
plot	cultivation	tractor	tractor
	weeding	by hand	by hand
	previous crop	fallow	maize (MM 602)
	neighbouring crop	maize	maize
	pesticide history	unknown	Gammoxane

Table II. Mean temperature and rainfall for the 1992 and 1993 growing seasons

Month	1992		1993	
	temperature, °C	rainfall, mm	temperature, °C	rainfall, mm
January	22.4	168	24.4	167.8
February	23.5	44.9	22.1	179.6
March	23.5	166.4	21.4	83.5
April	21.9	5.0	22.1	4.6
May	19.1	6.0	19.6	0.0
June	16.9	0	16.0	0.0

2.2. Sampling activities

Sampling for arthropods was done immediately pre-treatment and thereafter every two weeks for two months. Aerial fauna were swept using a standard net at two opposite diagonals across each sub-plot. Pitfall traps were used to assess fauna that inhabit the soil surface. Four pitfall traps (diameter, 50 mm; depth, 100 mm) were randomly placed in each sub-plot. Each trap was filled with water to which a few drops of liquid detergent was added to reduce the surface tension. The traps were emptied after 48 hours.

To assess soil inhabiting fauna, four soil cores (diameter, 50 mm; depth, 150 mm) were randomly taken from each sub-plot prior to application of lindane and once after two months. The arthropods were extracted from the soil using Berlese funnels. Light bulbs serving as heat source were placed above the samples to obtain a temperature gradient. The soil samples were left in the funnels for 72 hours.

2.3. Determination of soil microbial activity

Prior to the application of lindane, four nylon bags (120 mm × 120 mm), each containing 50 leaf discs (25 mm in diameter) were randomly buried to a depth of 50 mm in each of the sub-plots. Three months later the bags were retrieved and the contents dried (at 60°C for 24 hours) and weighed.

2.4. Assessment of crop damage and yield

To assess damage due to stalk borers, all the plants in each sub-plot were searched and the proportion of infested plants recorded.

In each plot, maize yields were recorded from four sub-plots (2 m × 2 m) which were earmarked from the start of the experiment and left undisturbed until harvest. The cobs were dried, shelled and weighed at a moisture content of 13.8%.

3. RESULTS

3.1. Sweepnet

Due to low numbers of insects caught in the sweepnet, the data were grouped into two insect orders. In general there were no significant differences in the number of Coleoptera caught either before or after application of lindane (Table III). Prior to insecticide application, there was an apparently significantly higher number of Diptera in the "treated" plot in 1992 (Table III), for reasons unknown. After treatment, there were no significant differences between the treated and untreated plots. Overall there were no significant differences in the number of Diptera between the treated and untreated plots in 1993 (Table III).

Table III. Mean (\pm SE) number of insects caught using a sweepnet in the trials in 1992 and 1993.

Species	1992			1993		
	interval after spraying, days	untreated	treated	interval after spraying, days	untreated	treated
Coleoptera	pretreatment	0.8 \pm 0.3	0.6 \pm 0.3	pretreatment	0.5 \pm 0.4	0.3 \pm 0.3
	13	0.8 \pm 0.3	0.0 \pm 0.0*	15	0.5 \pm 0.4	0.5 \pm 0.5
	27	0.1 \pm 0.1	0.5 \pm 0.2	29	1.5 \pm 1.0	1.3 \pm 0.6
	41	1.3 \pm 0.3	0.6 \pm 0.1	43	1.8 \pm 0.4	3.3 \pm 1.6
	56	2.1 \pm 0.4	2.0 \pm 0.4	57	1.5 \pm 0.8	1.0 \pm 0.7
	69	1.6 \pm 0.6	0.6 \pm 0.3	71	2.0 \pm 0.7	1.5 \pm 0.5
Diptera	pretreatment	0.3 \pm 0.3	1.3 \pm 0.1*	pretreatment	0.8 \pm 0.4	0.5 \pm 0.4
	13	2.3 \pm 1.0	4.2 \pm 2.0	13	2.7 \pm 0.4	2.7 \pm 0.1
	27	3.1 \pm 0.9	0.8 \pm 0.3*	27	14.8 \pm 5.1	18.0 \pm 9.5
	41	1.6 \pm 1.0	2.1 \pm 0.6	41	2.3 \pm 1.2	5.8 \pm 1.1*
	56	1.8 \pm 0.6	2.3 \pm 1.0	56	3.0 \pm 1.1	7.5 \pm 5.1
	69	0.5 \pm 0.0	1.1 \pm 0.1*	69	2.8 \pm 1.9	1.5 \pm 1.0

* Means within rows significantly different at P = 0.05 level.

3.2. Pitfall

Overall the number of spiders caught in 1993 were higher than in 1992. Both in 1992 and 1993 the number of spiders were significantly reduced in the treated plots shortly after spraying (Table IV). Thereafter there were no significant differences between the treated and untreated plots.

In 1992 the number of ants, although not significantly different, was higher in the treated plots before the application of lindane. After spraying, the number of ants was significantly lower in the treated plots (Table IV). In 1993 the number of ants was significantly higher in the treated plots, before the application of lindane. After spraying there were no significant differences between the treated and untreated plots.

3.3. Soil core

In both years, neither springtails nor mites were recorded from the soil core samples. However, that springtails were present in the field, was confirmed by the numbers caught in pitfall traps. The number of springtails recorded in 1993 was much lower than in 1992. Spraying of lindane did significantly reduce the number of springtails in both years. The effect lasted for almost two months (Table V).

3.4. Organic matter breakdown

The losses in weight of the leaf litter bags owing to soil microbial activity were moderate with around 50% losses during the period of burial. In 1992 the organic matter breakdown was significantly higher in the treated plots (0.74 ± 0.33 g) than in the untreated

Table IV. Mean (\pm SE) number of arthropods caught using pitfall traps in the trials in 1992 and 1993.

Species	1992			1993		
	Interval after spraying, days	untreated	treated	interval after spraying, days	untreated	treated
Araneida	pretreated	0.8 \pm 0.2	0.5 \pm 0.2	pretreated	35.7 \pm 30.5	4.5 \pm 1.9
	14	1.6 \pm 0.4	0.2 \pm 0.2*	15	3.5 \pm 1.2	1.0 \pm 1.0*
	28	0.5 \pm 0.1	0.4 \pm 0.1	29	8.5 \pm 4.6	3.3 \pm 1.4
	42	0.4 \pm 0.2	1.4 \pm 1.2	43	4.8 \pm 1.4	3.3 \pm 1.4
	57	0.9 \pm 0.4	0.6 \pm 0.2	57	2.5 \pm 1.4	3.8 \pm 1.4
	70	0.6 \pm 0.1	0.2 \pm 0.1	71	4.3 \pm 1.4	7.0 \pm 4.3
	Formicidae	pretreated	4.6 \pm 3.3	14.3 \pm 7.6	pretreated	10.3 \pm 10.2
14		23.7 \pm 3.3	9.8 \pm 0.6*	15	7.8 \pm 3.8	15.3 \pm 4.7
28		9.0 \pm 2.7	6.2 \pm 2.4	29	7.5 \pm 7.0	11.0 \pm 2.8
42		13.8 \pm 3.2	5.3 \pm 2.7	43	17.5 \pm 17.5	21.3 \pm 8.9
57		7.9 \pm 1.2	4.1 \pm 1.0*	57	42.8 \pm 25.5	34.3 \pm 12.1
70		16.4 \pm 2.0	8.1 \pm 2.6*	71	38.5 \pm 5.4	33.3 \pm 25.9

* Means within rows significantly different at P = 0.05 level.

Table V. Mean (+SE) number of Collembola caught in pitfall traps in the trials in 1992 and 1993.

1992			1993		
interval after spraying, days	untreated	treated	interval after spraying, days	untreated	treated
pretreatment	38.6±14.2	216.0±68.6*	pretreatment	18.0±4.7	5.8±3.8
14	75.3±17.1	0.3±0.2*	15	9.8±2.2	1.8±2.1*
28	18.4±1.7	3.1±1.3*	29	5.8±1.5	0.3±0.4*
42	14.4±9.1	1.5±0.7	43	9.5±5.6	3.3±1.9*
57	46.5±19.1	10.3±2.4	57	7.5±2.2	2.3±0.8*
70	31.2±10.6	22.7±1.5	71	17.8±2.4	10.8±4.2

* Means within rows significantly different at $P < 0.05$ level.

plots (0.89 ± 0.03). In 1993, this situation was reversed with a greater recovery of leaf material from the bags buried in the treated plots (0.89 ± 0.19 g) than in the untreated plots (0.67 ± 0.38 g) though with the higher experimental variation in this, the difference was not statistically significant.

3.5. Plant damage

In both years a single application of lindane significantly reduced damage due to stalk borers. However, significantly higher yield of maize was only obtained in 1993 (Table VI). The results showed that about 10-12% damage could be tolerated without significantly losing yield in 1992. Whereas, in 1993 about 8-10% damage due to stalk borers could not be tolerated without significantly affecting the yield.

4. DISCUSSION AND CONCLUSIONS

In general, the activity of insects was low probably due to the drought conditions that prevailed during the experimental period. Data from soil core samples and organic matter breakdown also indicated low faunal activity. This was not surprising considering the highly alkaline soils having poor organic matter.

Table VI. Proportion of plants damaged by stalk borers and maize yields in the trials in 1992 and 1993.

Observation	1992			1993		
	interval after spraying, days	untreated	treated	interval after spraying, days	untreated	treated
% damage	pretreat	13.6±1.0	10.4±0.6	pretreat	8.0±2.2	7.1±2.4
	13	8.4±0.9	2.8±0.6*	15	13.3±3.7	2.0±0.6*
	27	15.5±2.5	2.0±0.4*	29	13.5±1.9	1.3±0.5*
	41	12.2±2.1	2.5±0.3*	43	7.5±2.2	1.3±0.4*
	-	-	-	57	2.9±1.0	0.9±0.3
yield, kg ha ⁻¹		5988±1037	8041±1016		5799±782	7395±1504*

* Means within rows significantly different at $P < 0.05$ level.

Application of lindane appeared to have no real effect on aerial fauna. Although lindane is known to be toxic to Coleoptera and Diptera [3], the lack of significance may have been due to the low activity of these insects as well as emigration and immigration. Furthermore, the effectiveness of lindane is reduced under low rainfall conditions [4] which were experienced during the present studies. In addition, because of its relative high vapour pressure, the action of lindane under tropical conditions is not as persistent when compared to lower temperatures under temperate conditions [5].

Lindane significantly reduced the number of ants, springtails and spiders, recovered from pitfall traps. These findings are in agreement with those of other reports [6]. However in the present study the impact of lindane on these arthropods lasted for about two months. These findings agree well with other concurrent studies which have shown that lindane dissipates rapidly from the soil under Zambian field conditions. About 75% of the initial concentration of lindane is lost during the first 60 days after treatment [7].

Organochlorine insecticides are known to reduce microorganisms in the soil [8] and in turn affect the breakdown of leaf litter (organic matter). Contrary to the expected findings the present study showed that lindane did not significantly affect these microorganisms and our data are difficult to harmonize with those of other studies. Probably greater breakdown of organic matter would have occurred in the untreated plots had the litter bags been left in the soil for a longer period.

In both years lindane was highly effective against the target pest (stalk borers) as indicated by a reduction in the proportion of plants damaged. The yields were relatively higher in the treated plots when compared with the untreated plots although a clear relationship between damage and yield could not be established by the results in this study.

ACKNOWLEDGEMENTS

Financial support provided by the joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture is gratefully acknowledged. We are also appreciative of the technical assistance provided by Alfred Phiri.

REFERENCES

- [1] GRAHAM, P., Destruction of birds and other wildlife, by dieldrin spraying against tsetse fly in Bechuanaland. *Arnoldia* **10** 1-4 (1964).
- [2] WILSON, V.J., Observation on the effect of dieldrin on wildlife during tsetse fly *Glossina morsitans* control operations in Eastern Zambia, *Arnoldia* **5** 16-20 (1972).
- [3] ZINCHENKO, V.A., KALININ, V.A., ISLOVTSOV, R., Chemical Protection of Plants. R. Islovtsov (Ed.). Mir Publishers, Russia (1988).
- [4] SHEETS, T.J., Pesticides; Contemporary Ideas in Agriculture Health and Environment, T.J.Sheets, D. Pimentall (Eds), Clifton, New Jersey (1979).
- [5] KUMAR, R.,(Ed.), Insect Pest Control with Special Reference to African Agriculture, Edward Arnold, London (1984).
- [6] BROWN, A.W.A. (Ed.), Ecology of Pesticides, John Wiley and Sons, New York (1978).
- [7] MWANGALA, et al., this TECDOC.
- [8] PERFECT, T.J., Environmental impact of DDT in a tropical agro-ecosystem, *Ambio* **9** 16-21 (1980).

EFFECTS OF ENDOSULFAN ON A MAIZE AGRO-ECOSYSTEM IN ZAMBIA

Y.D. DEEDAT, G.C.MAIMBO, A.S.PHIRI
School of Agriculture, University of Zambia,
Lusaka, Zambia



Abstract

The effect of endosulfan on non-target fauna in a maize agro-ecosystem was studied in Zambia in 1994 and 1995. Endosulfan was well tolerated by a number of beneficial arthropods such as spiders, coccinellids, carabids and ants. Springtails were significantly reduced. However the effect was only transient and lasted for at most eight weeks. While endosulfan was effective against the target pest (stalk borers) it appeared to have no real effect on the soil inhabiting microorganisms.

1. INTRODUCTION

Maize is an important cash crop, both for commercial and small scale farmers in Zambia. Over the years production of this crop has increased significantly. However, successful cultivation of the crop depends upon, among other things, the use of insecticides to control major insect pests such as the stalk borers. In Zambia [1] as, in many other African countries [2], these insecticides are applied on a prophylactic basis irrespective of pest infestation level.

Endosulfan is among the organochlorine pesticides that is still being used in Zambia for the control of maize as well as cotton, groundnuts and soyabean pests. However, its use in other African countries and the world over is being questioned because of environmental concerns. To date, there is no information on the effect of endosulfan on either the maize or the cotton agro-ecosystem in Zambia. Hence, this study was conducted to determine the non-target effects of endosulfan in a maize agro-ecosystem.

2. MATERIALS AND METHODS

2.1. Experimental layout

This study was conducted at Chalimbana (N.C.S.R.) farm, 30 km east of Lusaka during the 1993/94 and 1994/95 growing season. The field was cultivated by a tractor and planted with maize (Table I) at a rate of 25 kg ha⁻¹ with a row to row spacing of 900 mm and a plant to plant distance of 200 mm. Fertiliser, at the rate of 400 kg ha⁻¹, was applied as basal dressing at planting. Urea was top dressed at the rate of 200 kg ha⁻¹ when the crop was at knee height. Soil characteristics (Table I) as well as weather data (Table II) for the experimental site were recorded during each cropping season.

The experimental site was divided into four blocks, each 50 m x 50 m. Each block was further subdivided into two 25 m x 25 m plots. Within each block one plot was randomly assigned and sprayed with Thiodan 350 g L⁻¹ at the rate of 1 kg AI ha⁻¹, using a CP15 knapsack sprayer. The other plot was left untreated. Thus, the experiment was conducted as a randomized complete block design with two treatments within each of the blocks. Within the centre of each plot, a subplot of 10 m x 10 m was demarcated for sampling activities.

Two-way analysis of variance was used to analyze the data at $P = 0.05$ level of significance. Where appropriate the data were transformed using the square-root transformation ($\sqrt{Y + 1}$).

2.2. Sampling activities

Sampling for arthropods was done one day before and after treatment and every two weeks thereafter for two months. The D-vac was used to sample arthropods on the plant as well as on the ground. A total of four samples were taken from each sub-plot during each sampling occasion. Each sample consisted of 5 sub-samples. In the case of plant sampling one plant represented one sub-sample. Sampling was done by either placing the cone of the D-vac over the entire plant when plants were small or by sweeping the plant with the cone from all sides when the plants were big. With ground sampling, each sub-sample consisted of lightly placing the D-vac cone on the ground for five seconds.

Table I. Crop, plot and soil characteristics of the experimental site at Chalimbana farm for 1994 and 1995 season.

Component	Characteristic	1994	1995
Crop	variety	PHB 3402	CG4144
	seed	dressed	dressed
	planting date	5/2/94	1/12/94
	spraying date	15/3/94	11/1/95
	harvesting date	- - -	7/4/95
Soil	sand, %	52.9	52.9
	silt, %	4.9	4.9
	clay, %	42.2	42.2
	organic matter, %	2.14	2.14
	pH (in CaCl_2)	5.1	5.1
Plot	cultivation	tractor	tractor
	weeding	by hand	by hand
	previous crop	fallow	maize
	neighbouring crop	maize	maize
	pesticide history	unknown	Thiodan

Table II. Mean temperature and rainfall for 1994 and 1995 the growing seasons.

Month	1994		1995	
	temperature, °C	rainfall, mm	temperature, °C	rainfall, mm
January	22.8	244	23.3	109
February	22.3	141	22.1	131
March	22.2	3	22.3	83
April	21.6	22	21.8	0.3
May	18.7	0	16.0	0
June	16.2	0	16.0	0

Pitfall traps were used to assess fauna that abound (dwell) on the soil surface. Four pitfall traps (diameter, 50 mm; depth, 100 mm) were randomly placed in each sub-plot. Each trap was filled with water to which a few drops of liquid detergent was added to reduce the surface tension. The traps were emptied after 72 hours.

To assess soil inhabiting fauna, four soil cores (diameter, 50 mm ; depth, 150 mm) were randomly taken from each sub-plot prior to application of endosulfan and once, two months later. The arthropods were extracted from the soil using Berlese funnels. Light bulbs serving as heat source were placed above the samples to obtain a temperature gradient. The soil samples were left in the funnels for 72 hours.

The breakdown of organic matter was used to assess the effect of endosulfan on microfauna and mesofauna. Prior to spraying, four nylon bags (12 cm x 12 cm), each containing 50 leaf discs (2.5 cm in diameter) were randomly buried to a depth of 50 mm in each of the sub-plots. Three months later the bags were retrieved and the contents dried (at 60°C for 24 hours) and weighed.

To determine damage due to stalk borers, all the plants in each sub-plot were examined and the proportion of the infested plants recorded. In each plot maize yield was recorded from another sub-plot (2 m x 2 m) which was earmarked from the start of the experiment and left undisturbed until harvest. The cobs were dried, shelled and weighed at a moisture content of 14.1%.

3. RESULTS

3.1. D-vac plant sampling

Both in 1994 and 1995 there were no significant differences in the number of spiders caught on the plants either before or after application of endosulfan (Table III). In both years before treatment, the mean number of coccinellids caught in the treated plots were relatively higher, but were lower following treatment. Only in 1994 was the mean number of coccinellids significantly lower in treated plots immediately after treatment.

The number of ants recorded in 1995 were relatively higher than in 1994. In both years the number of ants, although not significantly different were higher in the untreated plots before the application of endosulfan. After treatment the number of ants were slightly lower in the treated plots than in the untreated ones though the differences were small and not significant between the treated and untreated plots.

3.2. D-vac ground sampling

Generally the number of spiders, carabids and ants recovered from the ground were relatively higher in 1995 when compared with 1994 (Table IV). Before the application of endosulfan there were no significant differences in the number of spiders, carabids and ants caught in the treated and untreated plots. After spraying, although there were no significant differences, the number of these arthropods were slightly reduced in the treated plots. Only in 1995 was the mean number of spiders and carabids significantly reduced at 4 and 6 weeks post-treatment, respectively.

Table III. Mean number of arthropods caught using D-vac plant sampling in the trial plots in 1994 and 1995.

Arthropods	1994				1995			
	interval after spraying, days	untreated	treated	CV, %	interval after spraying, days	untreated	treated	CV, %
Araneida	pretreat	0.16	0.28	7.0	pretreat	1.34	1.53	3.1
	1	0.41	0.47	16.1	1	0.69	0.63	9.6
	16	0.09	0.28	13.2	14	0	0	0
	31	0.72	0.47	14.3	27	1.66	1.66	15.7
	52	0.56	0.72	14.5	41	1.94	1.94	7.4
Coccinellidae	pretreat	0	0.09	5.4	pretreat	0.16	0.22	12.0
	1	0.56	0.19*	4.5	1	0.69	0.47	9.9
	16	0.09	0.22	12.3	14	0.06	0.19	8.7
	31	0.78	0.69	7.6	27	0.59	0.38	14.2
	52	0.44	0.25	16.5	41	0.69	0.78	11.7
Formicidae	pretreat	0.75	0.66	10.5	pretreat	1.31	1.16	6.4
	1	0.34	0.22	8.0	1	0.94	0.34	9.6
	16	0.84	0.34	16.0	14	0.03	0.06	4.5
	31	0.28	0.41	10.3	27	2.03	2.20	4.4
	52	0.63	0.72	11.6	41	1.50	2.00*	2.9

* Means within rows significantly different at $P=0.05$.

3.3. Pitfall trap results

Both in 1994 and 1995, there were no significant differences in the number of spiders, carabids and ants caught in the treated and untreated plots either before or after the application of endosulfan (Table V). The mean number of springtails recovered from the pitfall traps were relatively higher in 1994 when compared with 1995. Both in 1994 and 1995 the mean number of springtails were significantly reduced in the treated plots but only at 6 and 4 weeks post-treatment for the two years, respectively.

3.4. Soil core fauna results

In both years the number of mites and springtails recovered from the soil were too low to carry out statistical analysis. In 1994 only 7 mites and 6 springtails were recovered from the soil core samples. Whereas in 1995, 18 mites and 18 springtails were collected from the soil. Most of these arthropods were collected from soil core samples taken after the application of endosulfan.

3.5. Organic matter breakdown

Both in 1994 and 1995 the organic matter breakdown, although not significantly different was slightly lower in the sprayed plots when compared with the unsprayed plots. In 1994 the organic matter breakdown over a period of three months was 0.39 g in the untreated

Table IV. Mean number of arthropods caught using D-vac ground sampling in the trial plots in 1994 and 1995.

Arthropods	1994				1995			
	interval after spraying, days	untreated	treated	CV, %	interval after spraying, days	untreated	treated	CV, %
Araneida	pretreat	0.06	0.16	8.8	pretreat	1.53	1.53	5.2
	1	0.38	0.03	10.8	1	0.94	0.34	12.3
	16	0.22	0.25	9.4	14	0.50	0.03	10.2
	31	0.44	0.28	5.8	27	0.47	0.22*	3.5
	52	0.81	0.63	18.7	41	3.28	3.47	12.5
	59	0.34	0.03	8.6	55	1.44	1.09	10.6
	72	0	0.16	3.2	71	2.81	2.63	6.5
Carabidae	pretreat	0	0	0	pretreat	1.47	1.63	11.0
	1	0.03	0.41	13.1	1	0.88	0.22	12.4
	16	0.13	0.13	0.4	14	0.28	0.03	12.6
	31	0.44	0.06	11.4	27	0.53	0.31	19.8
	52	0.13	0.22	11.1	41	5.00	3.56*	6.5
	59	0	0.16	6.7	55	1.56	1.84	8.1
	72	0.03	0.03	3.0	71	2.19	2.84	4.7
Formicidae	pretreat	0.44	0.06	9.7	pretreat	1.75	1.63	7.4
	1	0.44	0.84	18.3	1	0.84	0.44	22.4
	16	0.94	0.47	21.8	14	0.69	1.66	24.2
	31	0.06	0.03	4.5	27	1.22	1.47	19.1
	52	0.34	0.69	12.6	41	4.63	3.81	4.1
	59	0.91	1.19	18.8	55	2.16	2.63	13.5
	72	0.31	0.25	8.7	71	3.19	3.69	9.2

* Means within rows significantly different at $P=0.05$.

plots and 0.34 g in the treated plots. This represented a loss in weight of leaf material of 69.9 and 62.2% (CV 21.3%) for the two treatments, respectively. For 1995, the actual breakdown of organic matter over the period of four months was 0.61 g and 0.60 g for the unsprayed and sprayed plots, representing 72.0% and 70.5% (CV 27.0%) for the two treatments, respectively.

3.6. Plant damage

In 1994 there were no significant differences in the proportion of plants damaged between the sprayed and unsprayed plots (Table VI). Sampling to assess plant damage could not be continued beyond 4 weeks post-treatment because of the effects of drought on the plants. The severity of drought also negated collection of data on maize yield. In 1995 a single application of endosulfan significantly reduced damage due to stalk borers. Consequently, significantly higher yields were obtained in the treated plots. Endosulfan appeared to take effect about three weeks post-treatment and thereafter.

Table V. Mean number of arthropods caught using pitfall traps in the trial plots in 1994 and 1995.

Arthropods	1994				1995			
	interval after spraying, days	untreated	treated	CV, %	interval after spraying, days	untreated	treated	CV, %
Araneida	pretreat	1.94	5.53	40.9	pretreat	2.38	2.16	17.9
	14	4.72	4.88	14.6	14	4.44	3.25	12.7
	29	1.56	0.72	23.4	28	4.36	4.53	2.0
	43	3.06	1.72	19.9	42	4.13	3.34	13.1
	56	3.59	2.72	9.0	56	5.97	5.50	12.8
	70	1.97	2.00	12.4	69	3.59	3.44	20.9
	-	-	-	-	84	2.19	2.72	10.1
Carabidae	pretreat	0.41	0.47	15.7	pretreat	0.56	0.75	3.1
	14	2.31	1.38	10.7	14	1.84	2.03	12.4
	29	1.44	0.50	23.7	28	2.13	2.50	9.5
	43	5.03	3.47	12.9	42	2.38	2.69	14.1
	56	6.13	2.75	13.5	56	2.28	1.53	15.4
	70	1.88	2.63	9.6	69	1.06	1.22	12.5
	-	-	-	-	84	0.69	0.88	8.1
Formicidae	pretreat	11.25	9.50	22.6	pretreat	15.34	15.56	18.4
	14	22.91	20.03	9.8	14	10.34	10.66	16.0
	29	13.72	13.94	28.7	28	12.56	10.88	12.1
	43	24.34	27.13	16.9	42	4.13	4.31	23.4
	56	18.22	28.31	31.0	56	8.47	2.78	31.3
	70	6.09	9.66	13.0	69	13.22	11.84	23.2
	-	-	-	-	84	15.34	12.22	22.8
Collembola	pretreat	11.56	13.47	11.7	pretreat	3.44	3.44	18.0
	14	18.72	7.56	34.7	14	10.69	8.81	12.9
	29	14.94	8.09	22.3	28	12.66	5.28*	18.0
	43	23.41	3.97*	22.8	42	5.34	2.69	21.0
	56	82.91	30.03	34.9	56	11.22	4.31	26.8
	70	58.38	21.81	28.2	69	12.84	3.56	28.8
	-	-	-	-	84	6.97	6.00	11.8

* Means within rows significantly different at P=0.05.

4. DISCUSSION AND CONCLUSIONS

Overall the activity of arthropods was lower in 1994 compared with 1995. Although there were no significant differences between the treated and untreated plots, the number of arthropods recovered from the treated plots after spraying were relatively low. In all the experiments the number of arthropods recovered from the treated plots appeared to reach pre-treatment levels within a period of six to eight weeks.

Data from D-vac sampling and pitfall traps showed no significant effect of endosulfan on spiders as well as insects such as coccinellids and carabids. The results obtained from sampling Formicidae also showed no significant effect of endosulfan. This is contrary to the

Table VI. Proportion of plants damaged by stalk borers and maize yield from the field trials in 1994 and 1995

Component	1994				1995			
	interval after spraying, days	untreated	treated	CV, %	interval after spraying, days	untreated	treated	CV, %
Damage, %	pretreat	7.3	8.0	23.7	pretreat	18.8	21.1	9.2
	14	10.1	8.0	24.4	7	20.9	22.0	4.3
	35	9.2	7.4	15.0	21	31.2	23.3*	9.0
	-	-	-	-	35	33.6	24.3*	8.6
	-	-	-	-	48	34.7	24.9*	9.7
Yield, kg ha ⁻¹	-	-	-	-	-	4893	6995*	5.1

* Means within rows significantly different at P = 0.05.

report of Brooks who found endosulfan to be toxic to ants [3]. The rate of application of the chemical as well as immigration and emigration between plots may account for the discrepancy.

Data from pitfall traps showed that endosulfan significantly reduced the springtail populations. These findings agree well with the paper of Youdeowi and Service who stated that endosulfan is used to protect soil inhabiting arthropods such as mites and springtails [4]. However, the effect of endosulfan on springtails in the present study was transient and lasted for approximately 6 weeks. These data fit well with the findings of Mwangala et al. [5] who found that endosulfan dissipated rapidly under tropical conditions. About 65% of α -endosulfan and 70% of β -endosulfan was lost within 60 days.

There were no significant differences in the breakdown of organic matter between the treated and untreated plots as indicated by the loss of weight of the leaf litter. This was despite keeping the litter bags for longer than the usual two months period. The results indicate that endosulfan had no effect on the microorganisms that are important agents of organic matter breakdown in the soil. The data from the present study agree well with those of Gorback and Knauf who also found endosulfan to have no significant effect on soil microorganisms [6].

Data from 1995 showed that a single application of endosulfan was effective against the target pest as indicated by the proportion of plants damaged in agreement with Youdeowi and Service who have shown that endosulfan was especially effective against the maize stem borers [4]. The present study showed that one application of endosulfan was adequate to obtain a significant increase in yield.

ACKNOWLEDGEMENT

Financial support provided by the joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture is gratefully acknowledged.

REFERENCES

- [1] JAVIAD, I., Application of insecticides on cotton in Zambia; timing of spray applications. *Trop. Pest. Mgmt.*36:1-9 (1990).
- [2] MATTHEWS, G.A., *Pesticide Application Methods*, Longman, London (1979).
- [3] BROOKS, G.T., *Chlorinated Insecticides, Technology and Application*, CRC Press, Vol. 1 (1974).
- [4] YOUDEOWEI, A., SERVICE, M.W., *Pest and Vector Management in the Tropics*. ELBS (1986).
- [5] MWANGALA, et al., these proceedings.
- [6] GORBACH, S., KNAUF, W., *Endosulfan and the Environment, Environmental Quality and Safety*, Vol.1. (1972).

PERSISTENCE OF LINDANE AND ENDOSULFAN UNDER FIELD CONDITIONS IN ZAMBIA



XA9743732

F.S. MWANGALA, P.M. MUNDIA, J.C. NONDO, R. BANDA, C. MANGOYE
National Council for Scientific Research, Livestock and Pest Research Centre,
Chilanga, Zambia

Abstract

The persistence of lindane and endosulfan was studied under field conditions in Zambia in 1992 to 1994. Both pesticides dissipated rapidly under field conditions. About 29% and 73% of initial concentration was lost during the first 30 and 60 days after treatment, respectively in 1992. After 180 days, about 11% of the initial concentration was recovered from the soil. In 1993, 40% of initial residues were lost during the first 30 days. At 180 days after spraying, slightly more residues (25% of the initial values) were recovered at this time than in 1992. This indicated a change in the longer term behaviour of lindane in the soil since the calculated half-lives of lindane, covering the shorter term behaviour, were 55-80 days in 1992 and ~17 days in 1993. In 1994, losses of α -Endosulfan and β -Endosulfan were 40% and 37% respectively during the initial 30 days after treatment. A further 25% of α -Endosulfan and 33% of β -Endosulfan were lost during the following 30 days. These data allow estimates of the half-lives of α - and β -Endosulfan (40 and 38 days) under the field conditions pertaining in Zambia at the time of the trials showing that this compound has only moderate persistence and unlikely to cause long term environmental problems.

1. INTRODUCTION

Pesticides are still a major component of the inputs used in agriculture in Zambia. The newer pesticides such as organophosphorus insecticides, carbamates and pyrethroids, which are more environmental friendly, are more expensive and resource-poor farmers, who are the majority, cannot afford them. Organochlorine pesticides, on the other hand, are relatively cheap and some are still effective against most pests. However, their use in Africa and the world wide is being restricted because of environmental concerns. These concerns have been based on data generated from temperate regions where organochlorine pesticides have been found to persist for longer periods [1]. In recent years, there have been indications that organochlorine pesticides may be less persistent in tropical environments than in temperate zones [2-6].

Lindane and endosulfan are among the organochlorine pesticides that are still being marketed in Zambia. Use of lindane for general agricultural purposes in Zambia was withdrawn in 1994 by the agrochemical companies, after the start of this project, but it is still being used for seed dressing. Endosulfan is still being used for control of cotton, potato, maize, soyabean and groundnut pests in Zambia. This study was undertaken to determine the persistence of lindane and endosulfan in a Zambian agro-ecosystem.

2. MATERIALS AND METHODS

2.1. Field experiments

In 1992 and 1993, a field of maize at the University of Zambia farm (east of Lusaka) was divided into two plots, 100m apart. One plot was treated with lindane (Gammexane 200 g⁻¹) at a rate of 1 kg ha⁻¹. One plot was untreated and served as a control. Each plot was divided into 5 m x 5 m subplots. The subplots were 5 m apart and at least 10 m from the edge of the large plot.

In 1994, a field of maize at National Council for Scientific Research, Chalimbana farm (east of Lusaka) was divided into four 50 m x 50 m plots. Each plot was further subdivided into two 25 m x 25 m plots. One sub-plot was treated with endosulfan (Thiodan 350 g L⁻¹) from Zambia Cooperative Federation Agrochemical Division) at 1 kg ha⁻¹.

Soil samples were taken at random over the whole sub-plots using soil core samplers (diameter, 55 mm; Eikelkamp Giesbeek soil equipment) at depth of 0-50, 50-100 and 100-150 mm on day 1, 30, 60, and 180 in 1992 and on day 0, 1, 30, and 180 in 1993 after treatment. In 1994, soil cores were collected at a depth of 0-50 mm only on day 0, 1, 30, and 60 after treatment. A minimum of nine cores were collected at each depth. Cores from each subplot, depth and sampling date were mixed together after sieving in 0.25 mm sieve. The samples were stored in a deep freezer before analysis.

2.2. Moisture content

Moisture content of soil samples was determined gravimetrically by oven drying 10 g sub-samples at 110°C for 24 hours and expressing the moisture content as a percentage of water lost over weight of oven-dried soil. Soil characteristics were determined by the Department of Soil Science, University of Zambia in 1992 and 1993, and Soil Survey Section, Mt Makulu Research Station, Zambia in 1994.

2.3. Extraction and analysis

In 1992, two 20 g soil samples from each subplot were extracted with acetone+hexane (1+1 by volume, 100 mL) by tumbling for 1 hour on a wrist action shaker. The extract was filtered into a separating funnel containing sodium sulphate (20 g L⁻¹, 100 mL). The funnel was shaken for 1 minute. The lower layer was discarded after the two layers separated. The hexane extract was washed twice with sodium sulphate solution (20 g L⁻¹, 50 mL). The hexane extract was dried with anhydrous sodium sulphate (5 g) and analysed by the gc method.

In 1993 and 1994, duplicate soil samples (50 g) were extracted with methanol (200 mL) in a Soxhlet apparatus for 3 hours. Aliquots (4 mL) were blown to near dryness with a stream of nitrogen and the residues were redissolved in 1 mL methanol and diluted with 2.5 mL water. The samples were cleaned up on SEP-PAK C-18 cartridge columns (Waters, Milford, Massachusetts, USA). The columns were conditioned before the samples were added using methanol (3 mL), followed by distilled water (3 mL). The columns were washed with water + methanol (3 + 1 by volume, 1 mL), followed by distilled water (1 mL) before elution of the column with of hexane (2.5 mL) and analysed by the gc method.

The gas liquid chromatography method was as follows - column : 1 m x 4.5 mm silanised glass; packing, OV17 (16 g kg⁻¹) + OV210 (64 g kg⁻¹) on Chromosorb WHP; temperatures: column, 200°C, injector, 200°C, detector, 250°C; detector type : ⁶³Ni electron capture; carrier gas: argon. The limit of determination of lindane and endosulfan was 1 µg kg⁻¹ soil by the overall methods described. Recoveries from fortified samples were 83-98% for lindane and 86-96% for endosulfan and its sulfate.

3. RESULTS AND DISCUSSION

The environmental conditions that prevailed during the experimental period are presented in Table 1. The properties of the soil at the farms in 1992 to 1994 are presented in

Table 2. Moisture content of the soil on each sampling day is presented in Tables 3-5. There were no significant differences in the moisture content of soil from treated and untreated plots.

No residues were detected from the control plots in 1992 and 1993. Lindane dissipated relatively rapidly in both years (Tables 6, 7). In 1992, approximately 29% and 73% of the amount extracted on day 1 after treatment was lost during the initial 30 and 60 days, respectively from the 50 mm top layer (Table 6). After 180 days, the amount of lindane extracted from the soil was about 11% of the initial concentrations in the upper 50 mm layer. There were few variations in residue levels at lower depths and no pattern was discernible.

Table 1. Environmental conditions that prevailed during the experimental period.

Month	Year	Max. Temp.	Mean Temp.	Min. Temp.	Rainfall
January	1992	28.7	22.4	16.0	168.0
	1993	27.6	24.4	17.2	167.8
	1994	27.9	22.8	17.6	244.1
	1995	28.9	23.3	17.7	108.7
			(27)*	(21.6)	(17.0)
February	1992	31.1	23.5	16.0	44.9
	1993	26.7	22.1	17.4	179.6
	1994	27.9	22.3	16.6	140.7
	1995	28.1	22.1	17.1	131.2
			(27)	(21.5)	(17.0)
March	1992	29.9	23.5	17.2	166.4
	1993	26.3	21.4	16.4	83.5
	1994	29.7	22.2	14.6	2.7
	1995	29.1	22.3	15.5	82.7
			(27)	(21)	(16.0)
April	1992	29.1	21.9	14.8	5.0
	1993	28.6	22.1	15.6	4.6
	1994	28.7	21.6	14.4	21.9
	1995	28.5	21.8	15.1	0.3
			(27)	(19.9)	(14.0)
May	1992	26.7	19.1	11.5	6.0
	1993	28.1	19.6	11.0	0.0
	1994	26.7	18.7	10.7	0.0
	1995	24.3	16.0	7.7	0.0
			(26)	(15.1)	(10.0)
June	1992	24	16.9	8.5	0.0
	1993	23.4	16.0	8.6	0.0
	1994	24.2	16.2	8.1	0.0
	1995	24.3	16.0	7.7	0.0
			(25.3)	(17.0)	(8.0)

a) numbers in brackets = 30 year mean values

Table 2. Characteristics of soil in the experimental plots.

	1992	1993	1994
Sand (%)	55.8 ± 1.3	65.6 ± 2.8	73
Clay (%)	34.7 ± 1.4	20.1 ± 3.6	8
Silt (%)	10.0 ± 0.4	14.3 ± 6.4	19
Organic Matter (%)	1.60	0.51 ± 0.4	0.3
Organic Carbon (%)	0.81 ± 0.08	0.26 ± 0.2	0.15
pH (1:2.5 Soil:Water Ratio)	6.3 ± 0.1	9.7 ± 0.5	5.8

Table 3. Moisture content (% , ± SD) of soil in experimental plots at various depths in 1992.

DAY	CONTROL			TREATED		
	0-50mm	50-100mm	100-150mm	0-50mm	50-100mm	100-150mm
1	9.2 (± 1.1)	8.8 (± 0.5)	10.8 (± 0.9)	10.1 (± 0.9)	10.4 (± 0.1)	9.3 (± 0.3)
30	7.1 (± 0.3)	8.3 (± 0.3)	8.6 (± 0.2)	8.5 (± 0.3)	9.4 (± 0.2)	8.6 (± 0.2)
60	5.7 (± 0.1)	8.3 (± 0.2)	8.8 (± 0.2)	5.4 (± 0.2)	7.8 (± 0.1)	8.1 (± 0.2)
180	4.8 (± 0.1)	7.0 (± 0.6)	6.9 (± 0.3)	5.0 (± 0.1)	6.9 (± 0.2)	6.3 (± 0.3)

Table 4. Moisture content (% , ± SD) of soil in experimental plots at various depths in 1993.

DAY	CONTROL			TREATED		
	0-50mm	50-100mm	100-150mm	0-50mm	50-100mm	100-150mm
0	10.3(± 0.9)	12.4(± 0.7)	12.8(± 0.8)	8.5(± 1.1)	10.5(± 0.5)	10.8(± 1.1)
1	13.5(± 0.6)	13.5(± 1.3)	12.2(± 1.1)	11.7(± 1.5)	10.7(± 0.5)	11.1(± 0.9)
30	14.5(± 2.5)	15.2(± 1.8)	14.1(± 0.8)	13.9(± 1.2)	14.9(± 1.5)	14.6(± 3.5)
180	4.6(± 0.6)	6.4(± 0.6)	7.1(± 0.6)	4.5(± 0.8)	6.2(± 0.7)	5.9(± 1.7)

Table 5. Moisture content (% , ± SD) of soil from Chalimbana farm on each sampling day in 1994.

TREATMENT	DAY AFTER TREATMENT			
	0	1	30	60
Treated	8.15 ± 0.45	5.02 ± 0.72	2.50 ± 1.10	0.90 ± 0.76
Untreated	8.54 ± 0.72	5.00 ± 0.67	2.58 ± 1.27	0.86 ± 0.78

Table 6. Lindane levels ($\mu\text{gg}^{-1} \pm \text{SE}$) in soil from University of Zambia farm during 1991/1992 season.

Depth (mm)	Day After Treatment			
	1	30	60	180
50	0.76 ± 0.02	0.54 ± 0.03	0.20 ± 0.10	0.090 ± 0.006
100	0.06 ± 0	0.05 ± 0.007	0.04 ± 0.03	0.070 ± 0.006
150	0.01 ± 0	0.11 ± 0.01	0.012 ± 0.004	0.020 ± 0

Table 7. Lindane levels (ngg^{-1}) in soil from the University of Zambia farm during the 1992/1993 season.

Depth (mm)	Day			
	0	1	30	180
50	0.95 ± 0.07	13.5 ± 0.7	5.8 ± 0.07	3.4 ± 0.8
100	0.6 ± 0.1	4.8 ± 0.2	0.6 ± 0.2	0.8 ± 0.1
150	0.6 ± 0.1	4.0 ± 0.3	0.2 ± 0.02	0.8 ± 0.1

Table 8: α - and β -Endosulfan residues ($\mu\text{gg}^{-1} \pm \text{SE}$) in soil from the Chalimbana farm after treatment with Thiodan 350gL^{-1} during the 1993/94 season.

Day	Endosulfan I		Endosulfan II	
	Treated	Untreated	Treated	Untreated
0	0.70 ± 0.12	0.52 ± 0.15	0.16 ± 0.07	0.18 ± 0.08
1	16.34 ± 1.61	1.79 ± 0.10	6.50 ± 0.37	0.85 ± 0.11
30	9.87 ± 0.69	1.31 ± 0.15	4.13 ± 0.42	0.60 ± 0.05
60	5.80 ± 0.87	1.03 ± 0.12	2.00 ± 0.23	0.47 ± 0.03

In 1993, lindane declined by more than 40% during the first 30 days after treatment in the upper 50 mm layer (Table 7). After 30 days, lindane levels declined to pre-treatment levels at 100 and 150 mm depths but the level was still high in the upper 50mm. After 180 days, 25% of the residues detected on day 1 after treatment still persisted in the upper 50 mm layer of the soil. The residues at 100 and 150 mm depths did not change and were at pre-treatment levels. The calculated half-lives of lindane in the upper 50 mm depth, during the period immediately

after application, were 55-80 and 17 days in 1992 and 1993, respectively. At later periods (> 60 days), although the data are limited, the rate of loss of lindane appeared to decline probably owing to the drier conditions of both the weather and the soil (Tables 1, 3-5) giving rise to stronger sorption of lindane and also lower microbial activity pertaining in the soil, during this period.

In the trials with Thiodan in 1994, only α - and β -endosulfan were detected from the soil and not endosulfan sulphate. During the initial 30 days, 40% and 37% of the initial amounts of α - and β -endosulfan, recorded on the day of application, were lost. Another 25% of α -endosulfan and 33% of β -endosulfan were lost after 60 days leaving residues remaining in the soil after 60 days of 36% and 31% of their initial concentrations. Both α - and β -endosulfan dissipated at the same rate with estimated half-lives of 38-40 days. Small amounts of residues were detected in the untreated sub-blocks (Table 8) possibly attributable to drift during spraying or contamination during sampling and sample handling.

The data indicate that both lindane and endosulfan dissipated rapidly under Zambian conditions. This is in agreement with other studies done in Kenya and Tanzania with DDT [3, 7]. Most of the residues were restricted to the upper 50 mm layer. There was little evidence to suggest that lindane was leached downwards to lower layers. The amount of lindane detected at the 50-150 mm depth could be due to contamination by soil particles from the top layer. The dissipation of lindane and endosulfan was possibly biphasic in nature and dependent on the varying meteorological and ensuing soil conditions during the trials. During the earlier, wetter period of the trials, particularly the first 30 days (Tables 1, 3-5), the rate of disappearance of lindane and endosulfan was fast but slowed down considerably thereafter.

It is concluded that lindane and endosulfan dissipate rapidly under Zambian field conditions. The use of lindane and endosulfan will not result in accumulation of residues. This is in agreement with other studies that have shown that the residence period of organochlorine pesticides in the soil is much shorter under tropical field conditions than under temperate zones [2, 3, 5, 7].

REFERENCES

- [1] EDWARDS, C.A., *Persistent Pesticides in the Environment*, CRC Press, Cleveland, Ohio (1973).
- [2] EL-ZORGANI, G.A., Persistence of organochlorine insecticides in the field in the Gezira soil under cotton, *Bull. Environ. Contam. Toxicol.* **15** (1976) 378-382.
- [3] LALAH, J.O., ACHOLLA, F.V, WANDINGA, S.O., Fate of ^{14}C -p,p'-DDT in Kenyan Tropical Soils, *J. Environ. Sci. Health* **B29** (1994) 57-64.
- [4] PERFECT, T.J., Environmental impact of DDT in a tropical agro-ecosystem, *Ambio* **9** (1980) 16-21.
- [5] SLEICHER, C.A., HOPCRAFT, J., Persistence of pesticides in surface soils and relation to sublimation, *Environ. Sci. Technol.* **18** (1984) 514-518.
- [6] TALEKA, N.S, SUN, L.T., LEE, E.M., CHEN, J.S., Persistence of some insecticides in subtropical soil, *J. Agric. Food. Chem.* **25** (1977) 348-352.
- [7] STEVENS, J., MAENDA, D. N., NGOWI, A.V., MOSHI, A.O., MUSHY, P., MAUSA, E., Dissipation and degradation of ^{14}C -p,p'-DDT and ^{14}C -p,p'-DDE in Tanzania Soils under field conditions, *J. Environ. Sci. Health*, **B29** (1994) 65-71.

SOIL PERSISTENCE, PLANT AND NON-TARGET INSECT UPTAKE OF ENDOSULFAN AND LINDANE APPLIED TO SOYA BEAN AND MAIZE IN FIELD TRIALS IN ZIMBABWE

M.F. ZARANYIKA, P. MUGARI
Chemistry Department, University of Zimbabwe,
Harare, Zimbabwe



Abstract

The persistence of lindane and endosulfan in the soil, uptake by, and distribution in plants, and effects on and absorption by non-target insects, following application of the insecticides for the control of maize pests and soya bean respectively were determined under Zimbabwean weather conditions. No large scale effects on the non-target insects were observed though some small effects on the populations of semiloopers and orthoptera in the endosulfan treated soya bean plot were noted. Concentrations of the insecticides in spiders declined during the trial though those in grasshoppers and crickets appeared to increase. Concentrations of both insecticides in soil fell rapidly during the first 7 weeks after application but, after that, the rates of loss were much slower possibly owing to the drier conditions prevailing during this later period, reducing both physicochemical and microbial loss processes. Initial concentrations of both insecticides in all the vegetative parts of plants examined after spray application declined systematically to low levels during the 10 weeks of observation, probably owing to both metabolism within the plants and to crop volume dilution effects and will have declined to even lower levels by harvest time. Surprisingly, low concentrations of lindane and endosulfan were found in the harvested maize and soya bean seeds. At early stages after application, there were also traces of both insecticides in the vegetative parts of the plants from the untreated, control plots probably arising from uptake of soil residues from the previous year and/or spray drift but these became undetectable at later stages of growth.

1. INTRODUCTION

1.1. Background

By 1989 it was estimated that Africa accounted for 5% of the world pesticide use [1]. Most of these pesticides are used for pest control in agriculture. Although the use of pesticides in agriculture has been reported to have adverse effects on both man and other non-target organisms in the environment, agricultural development and the need to increase food production demands the use of pesticides. It is however necessary to assess the impact of the use of pesticides on the environment.

This impact will depend on several factors including temperature, rainfall, soil type, biotic activity, light intensity, land cultivation and other agricultural practices. These factors determine the persistence of the pesticide in a specific environment, and in this respect organochlorine pesticides have been found to be the most persistent as a group. Thus although the impact of pesticides on the environment has been studied extensively in the developed countries, there is still need to assess their impact in developing countries as agricultural practices in these countries may differ from those used in the developed countries.

Of major concern in Zimbabwe has been the use of organochlorine pesticides to control tsetse fly and malaria vectors. Pesticide sprays for tsetse fly control began in the early 1960s. Pesticides which have been used include dieldrin (1962-1967) and DDT (1968 to present) [2, 3]. Endosulfan and deltamethrin are also used, especially in aerial sprays [4]. In addition to its use in the control of tsetse fly and malaria vectors, DDT was used extensively in agriculture prior to 1983 when the use of DDT in agriculture was banned. Dieldrin,

endosulfan and deltamethrin are used extensively in agriculture. Table 1 shows the list of organochlorine pesticides registered for use in agriculture in Zimbabwe [5]. Contamination of non-target invertebrates and vertebrates like birds, reptiles, amphibia, fish, mice and sometimes large mammals as a result of aerial spraying of riverine and lake shore vegetation with DDT and endosulfan to kill tsetseflies in the Zambezi Valley area has been reported [6], but direct study of the effects of organochlorine pesticides on the agrosystem have yet to be undertaken in this country.

The aims of this study were: (a) To determine the persistence of organochlorine pesticides in the agrosystem, and (b) to monitor pesticide residues in the different compartments of the agrosystem. The pesticides selected for the study are endosulfan and Lindane. Endosulfan is used extensively for the control of the heliothis boll-worm, the semi-looper caterpillar and aphids in soya bean and groundnut crops, aphids in potatoes and the cutworm and red mite in maize [1] while lindane is used as for seed dressing [5]. A number of organochlorine formulations registered for use in Zimbabwe is listed in Table 1.

1.2. Persistence of lindane and endosulfan in soil

The soil is the major sink for chemicals applied to crops. Persistence patterns will vary from one climate to another and also depending on the chemical nature of the pesticide and the type of soil [7]. In cold climates degradation was generally found to be slower than in tropical climates [7, 8]. Lindane undergoes microbial degradation by dechlorination to pentachlorocyclohexane [9]. The bacteria, *Clostridium sporogenies* and *Bacillus coli*, produce benzene and monochlorobenzene from lindane [10]. Other possible transformations of lindane which occur especially in wet and submerged conditions include isomerization into the α - and β -HCH isomers of lindane [11]. The isomers were shown to be rapidly degraded

Table 1. Some organochlorine pesticides registered for use in agriculture in Zimbabwe

Trade Name	Active Ingredient	Toxicity Class*
Aldrin 40% w.p.	aldrin 40%	P
Anti-Kil	chlordan 30%	A
Razor	chlorthal-dimethyl 36%	G
Dicofol 40 E.D.	dicofol 40%	A
Kelthane	dicofol 18,5%	G
Dieldrex 50 w.p.	dieldrin 50%	P
Thionex 1% granules	endosulfan 1%	G
Thiodan 1% granules	endosulfan 1%	G
Thiodan 20 e.c.	endosulfan 20%	P
Multi Benhex	γ -BHC 12% +total BHC 75%	A
Gamatox House spray	γ -BHC 5,0%	A
Bexadust (L)	γ -B.H.C. 0,6%	G
Agri seed Dress 75%	lindane 1%	G

*P = extremely toxic; A = toxic; G= non-toxic.

in soil [12]. Endosulfan was found to undergo epoxidation which is enzyme catalyzed [13] and other transformations which include hydrolysis, reduction and hydroxylation [14]. Mechanical processes such as volatilization and run off also deplete the levels of these chemicals in soil.

The persistence of endosulfan in the soil has attracted attention from several workers. Burns studied the degradation of the insecticide in soils [15]. Kathpal et al. [16] and Martens [17] studied the kinetics of its degradation in soils and reported rates of 63% loss in 2-3 months and 5.4% in 15 weeks, respectively. Martens also studied the non-microbial degradation of the insecticide in soils under varying conditions of pH, sample wetness and temperature, and found that degradation was faster in wet samples with a higher pH. El Beit studied the kinetics of leaching of endosulfan in sediments and found that the rate of leaching was slow compared to the rate of degradation by micro-organisms [18].

1.3. Uptake of the pesticides by non-target fauna

Insects were selected for study as examples of non-target lower fauna. The uptake of organochlorine pesticides by insects following application for the control of agricultural pests in Europe and the American continents has been reported by several workers [19]. Gish found organochlorine pesticide residues of up to 0.60-0.65 $\mu\text{g g}^{-1}$ in beetle larvae from two agricultural fields in the USA [20]. Korschgen reported values as high as 8 ppm for aldrin and dieldrin residues in ground beetles from aldrin treated fields [21]. In Africa, DDE, DDT and dieldrin were detected in *Chiron. L.*, *Corixidae*, and *Chironomidae* from Kenya by Lincer [22] and Greichus [23], while Muller reported on the uptake of dieldrin by ants, lepidoptera and isoptera following dieldrin application for the control of tsetse fly in Cameroon [24]. Dieldrin, DDT and DDE were also detected in various insects by Greichus et al. in Zimbabwe [23].

1.4. Pesticide residues in crops

Several workers have studied the occurrence of pesticide residues in crops. Beestman et al. studied the translocation of dieldrin into corn from the soil and found that although 70-90% of the residues were found in the stems, appreciable quantities, up to 2%, eventually reached the seed [25]. These workers concluded further that residues measured at harvest time do not represent the total uptake by the plant during its life time because some of the absorbed insecticides are metabolized. Dormal et al. [26] and Brett and Bowery [27] found 0.3-1.5 $\mu\text{g g}^{-1}$ DDT residues and 0.4 $\mu\text{g g}^{-1}$ lindane residues respectively in beans and McCaskill et al. found 6 ng g^{-1} lindane residues in soybean [28], while Bruce et al. reported levels as high as 0.11 $\mu\text{g g}^{-1}$ heptachlor in soybean [29].

1.5. Aims of the present study

The work reported in this paper was carried out between 1991 and 1995, and was conducted as part of the FAO/IAEA Co-ordinated Research Program on "Adverse Side Effects on Flora and Fauna from the Use of Organochlorine Pesticides on the African Continent". The trials conducted during the 1991/92, 1993/94 and 1994/95 growing seasons, November to March in Zimbabwe, were adversely affected by severe droughts during these periods. As a result, meaningful data were only collected during the 1992/93 growing season.

2. MATERIALS AND METHODS

2.1. Experimental field site preparation

The 1992/93 field experiments were conducted on specially designated land in the experimental section of the University of Zimbabwe Farm approximately 10 km from the University. After mechanical land preparation, the trial field of ~2 hectare was divided into 4 plots of approximately 0.25 ha each and alternate plots planted with commercially dressed soya and maize seed on 23 December 1992. The experimental plot was bordered by cattle fodder to the north and east, while plot A was bordered by a maize crop, and plot B was bordered by a roadway. The crops were separated by a 1.5 m buffer zone. This was done to simulate actual agricultural practice as closely as possible.

2.2. Pesticide application

The spraying of both soya and maize was carried out on 20 February, 1993. Thiodan (endosulfan) 50 WP (300 g) and Multi-Benhex (lindane) 75 WP (270g) were diluted in water (120 L) and applied using 15 L knapsack sprayers (Taurus Spraying Systems, Zimbabwe) to the respective plots, equivalent to application rates of 600 and 810 g AI ha⁻¹, respectively. To ensure the pesticide was applied evenly, preliminary spraying tests were carried out using water.

2.3. Sample collection

2.3.1. Soil samples

In each plot 4 subplots, each 5 m x 5 m and 10 m from the edge of the plot, were marked out. Four soil samples were collected from each of the four subplots in each plot. Each plot was divided into 4 quadrants and one soil core was collected from each quadrant. Sampling was done by means of a cylindrical soil corer 50 mm x 150 mm. Each soil core was placed into a clean polyethylene bag which was then sealed and labelled with the subplot number and the sampling point number. The samples from each plot were then placed into a single bag which was sealed and labelled with the plot letter, for transportation to the laboratory where they were stored in a deep freezer until they were taken for analysis.

2.3.2. Insect samples

Insect samples were collected using 4 pitfall traps per subplot after Critchley et. al. [30]. The traps were set 2, 4 and 10 weeks after the experimental plots were sprayed with the pesticides. Pretreatment traps were set 2 weeks before spraying. On each occasion the traps were left overnight for collection the next day. The inner cups of the traps containing the trapped insects were transported to the laboratory where the insects were classified. Each order of insects was transferred to a separate plastic vial and stored in a deep freezer without a preservative. Because of the low numbers of insects trapped, all insects of the same order from each plot were pooled together to form a single composite sample as shown in Table 4.

2.3.3. Plant samples

For residue analysis, 12 plants per plot were randomly taken. For each plot the plants were divided into stems and leaves, and roots at the field. Each of the two samples per plot

Table 2. Recovery of lindane and α - and β - endosulfan from soil samples spiked at three levels

lindane			α - endosulfan			β - endosulfan		
amount added ngg ⁻¹	amount found ngg ⁻¹	% recovery	amount added ngg ⁻¹	amount found ngg ⁻¹	% recovery	amount added ngg ⁻¹	amount found ngg ⁻¹	% recovery
3.01	2.50	83	2.48	2.10	85	2.53	2.20	87
6.32	5.00	80	5.00	4.20	84	5.04	4.40	87
9.32	9.04	97	9.00	8.93	99	9.83	9.77	99

Table 3. Mean residues of lindane and α - and β - endosulfan found in soil cores from the treated and untreated plots in the 1992/93 field trial

Interval after application, weeks	Mean residue concentration, ngg ⁻¹ (\pm SE)					
	treated plots			untreated plots		
	lindane	α -endosulfan	β -endosulfan	lindane	α -endosulfan	β -endosulfan
maize plots						
pretreatment	nd ^{a)}	20 \pm 7	6 \pm 2	nd	nd	nd
2	227 \pm 71	29 \pm 13	17 \pm 10	1 \pm 2	1 \pm 2	nd
5	137 \pm 33	4 \pm 3	7 \pm 6	nd	nd	nd
7	53 \pm 13	8 \pm 6	3 \pm 5	- ^{b)}	-	-
10	37 \pm 8	5 \pm 4	0.4 \pm 0.8	nd	nd	nd
25	24 \pm 4	3 \pm 2	1 \pm 2	-	-	-
soya plots						
pretreatment	16 \pm 7	nd	nd	-	-	-
2	3 \pm 2	185 \pm 5	119 \pm 121	3 \pm 3	1 \pm 1	nd
5	6 \pm 5	83 \pm 10	40 \pm 7	4 \pm 4	2 \pm 1	nd
7	7 \pm 8	71 \pm 1	23 \pm 2	2 \pm 3	4 \pm 4	nd
10	-	-	-	1 \pm 1	1 \pm 2	nd
25	1 \pm 2	40 \pm 3	13 \pm 2	-	-	-

a) nd = not detected b) - = analysis not conducted

was then placed in a polyethylene bag which was then sealed, labelled with the plot number and sampling date. The seed samples were randomly taken from each of the four plots at harvest and placed into polyethylene bags which were labelled as described above. The samples were then transported to the laboratory where they were stored in the deep freezer prior to residue analysis.

2.4. Residue analysis

2.4.1. Preparation of Florisil cleanup columns

Florisil (residue grade, Supelco, USA, ~5 g) was packed into a Pyrex glass column (10 mm i.d. x 110 mm) followed by anhydrous sodium sulphate (Merck, Germany, ~5 g).

Table 4. Total insect pitfall catches over 24 hours

Plot	Crop	Pesticide treatment	Insect order	Number of insects caught				
				Week / Date				
				pretreat-ment	0 20Feb	2 6Mar	3 11Mar	11 11May
A	maize	lindane	arachnida	40	43	48	41	62
			coleoptera	45	63	53	59	69
			orthoptera	17	23	15	15	28
B	soya	endosulfan	arachnida	23	24	30	21	26
			coleoptera	4	4	3	3	6
			orthoptera	62	50	58	12	29
C	maize	untreated control	arachnida	30	38	41	49	23
			coleoptera	11	21	31	49	42
			orthoptera	10	13	11	8	7
D	soya	untreated control	arachnida	48	41	38	42	49
			coleoptera	44	62	51	40	44
			orthoptera	19	15	14	18	23

The column was washed with hexane (20 mL) followed by methanol + hexane (0.5 + 9.5 by volume, 20 mL) and then dried at 180°C overnight. The column was cooled and then prewetted with hexane (10 mL).

2.4.2. Extraction and clean up of soil samples

Each soil core was taken for residue analysis. The frozen sample was thawed, ground and passed through a 2.5 mm sieve. A mass (equivalent to 1 g dry weight of the soil) was taken from the core and weighed into a beaker (10 mL). Acetic acid (0.5 mL) was pipetted into the beaker and the mixture was stirred using a glass rod. Nonane (0.5 mL) was added to the slurry and the mixture was again stirred with the glass rod. The resulting slurry was ultrasonicated for 30 minutes. The mixture was then allowed to stand and pesticide grade silica gel (Merk, 5 g) was added and the mixture stirred. The finely divided powder was transferred into a cellulose extraction thimble (Merk) containing silica gel (5 g). The thimble was then placed into a Soxhlet extraction apparatus and extracted with hexane + benzene (2 + 1 by volume) for 4 hours. The crude extract was concentrated in a Kuderna-Danish flask to 1 mL. This concentrate was quantitatively transferred to a Florisil clean up column and eluted with hexane (20 mL - first fraction) followed by methanol + hexane (0.5 + 9.5 by volume, 20 mL - second fraction). The fractions were separately concentrated in the Kuderna-Danish flask (to 1 mL) and stored in glass vials (1.8 mL) with Teflon lined screw caps (Supelco, USA) at 4°C until analysed by the GC method described in section 2.4.4.

In preliminary studies soil samples (1 g) were spiked with a solution containing a mixture of lindane, α -endosulfan and β -endosulfan standards. Each spiked sample was stirred to ensure even distribution of the pesticides in the sample and then stored in the dark for 24 hours to equilibrate before extraction, clean-up, concentration and analysis as described above.

Table 5. Mean residues of lindane, α - and β -endosulfan and endosulfan sulfate in insects caught in the pitfall traps within the sprayed plots at intervals after spray application

Interval after application, weeks	Insect species	Residue concentrations, ngg ⁻¹			
		lindane	α -endosulfan	β -endosulfan	endosulfan sulfate
<u>maize</u> pretreatment	spiders	nd ^{a)}	nd	nd	nd
	crickets	nd	nd	nd	nd
	semiloopers	nd	nd	nd	nd
2	spiders	5.1	0.5	nd	nd
	crickets	2.87	0.42	0.002	nd
	semiloopers	nd	0.045	nd	nd
4	spiders	1.75	nd	nd	nd
	crickets	- ^{b)}	-	-	-
	semiloopers	-	-	-	-
10	spiders	-	-	-	-
	crickets	20.06	6.16	nd	nd
	semiloopers	-	-	-	-
<u>soya</u> pretreatment	spiders	nd	nd	nd	nd
	crickets	nd	nd	nd	nd
	grasshoppers	nd	nd	nd	nd
2	spiders	nd	6.91	nd	nd
	crickets	nd	1.56	2.36	nd
	grasshoppers	0.002	nd	nd	nd
4	spiders	0.53	3.86	nd	nd
	crickets	-	-	-	-
	grasshoppers	nd	4.29	nd	nd
10	spiders	-	-	-	-
	crickets	nd	1.84	nd	nd
	grasshoppers	nd	28.16	nd	nd

a) nd = none detectable b) - = not analysed

Table 6. Recovery of lindane, α - and β -endosulfan and endosulfan sulfate from spiked plant samples

Plant sample	Spiked concentration, ngg ⁻¹ ; % Recovery							
	lindane		α -endosulfan		β -endosulfan		endosulfan sulfate	
	spike	% rec.	spike	% rec.	spike	% rec.	spike	% rec.
maize stems+roots	5.02	76.1	3.08	93.3	3.88	96.9	4.62	105.2
soya stems+roots	7.32	84.6	14.3	70.3	5.78	70.3	6.00	77.5
maize seed	16.0	90.3	18.0	97.1	13.2	97.8	18.4	95.7
soya seed	10.0	82.0	5.76	81.6	13.2	106.2	14.1	88.7

Table 7. Residues of lindane, α - and β -endosulfan and endosulfan sulfate in plant samples at intervals after spraying

Plant sample	Interval after spraying, weeks	Residue concentration, ngg ⁻¹							
		Sprayed plots				Unsprayed control plots			
		lindane	α -endo	β -endo	endo sulfate	lindane	α -endo	β -endo	endo sulfate
maize roots	pretreat	nd ^{a)}	nd	nd	nd	nd	nd	nd	nd
	2	13.8	0.4	nd	nd	nd	0.2	nd	nd
	5	8.8	0.1	0.5	nd	nd	nd	nd	nd
	10	8.2	1.0	nd	nd	nd	nd	nd	nd
maize stems +leaves	pretreat	nd	nd	nd	nd	nd	nd	nd	nd
	2	19.3	0.3	nd	nd	nd	0.2	nd	nd
	5	9.5	0.2	0.15	nd	nd	nd	nd	nd
	10	1.06	0.07	nd	nd	nd	nd	nd	nd
maize seed		1.32	0.62	nd	nd	nd	nd	nd	nd
soya roots	pretreat	nd	nd	nd	nd	nd	nd	nd	nd
	2	1.72	12.7	3.31	nd	0.8	0.2	nd	nd
	5	0.13	6.55	0.4	nd	0.79	nd	nd	nd
	10	0.14	1.53	0.19	nd	nd	nd	nd	nd
soya stems +leaves	pretreat	nd	nd	nd	nd	nd	nd	nd	nd
	2	0.01	2.22	0.10	nd	0.77	0.40	0.18	nd
	5	0.07	6.57	2.02	nd	0.80	2.91	nd	nd
	10	nd	1.50	0.10	nd	nd	nd	nd	nd
soya seed		0.28	2.21	0.06	nd	0.04	0.02	nd	nd

a) nd = none detectable

2.4.2. Extraction and clean up of insect samples

A composite sample of insect species was crushed and divided into two approximately equal portions. The portions were weighed into a beaker (20 mL) and anhydrous sodium sulphate (~10 g) added. The mixture was ground using a glass rod until it was finely divided. Acetonitrile (15 mL) was added and the mixture stirred for about 10 minutes using a sonicator. The mixture was centrifuged and the supernatant transferred to a Kuderna-Danish flask. The residue after centrifugation was re-extracted with a further portion of acetonitrile and the supernatant combined with the first extract. The combined extract was then concentrated (to 1 mL), quantitatively transferred to a Florisil clean up column and eluted with hexane (20 mL, - fraction 1) followed acetonitrile + hexane (0.5 + 9.5 by volume, 20 mL - fraction 2). The two fractions were concentrated and analyzed separately as described above for soil samples.

In preliminary studies a composite sample of insects from one pitfall trap was crushed and divided into 6 approximately equal portions three of which were spiked with 18.3 ng g⁻¹, 12.1 ng g⁻¹, 1.62 ng g⁻¹ and 18.2 ng g⁻¹ of lindane, α -endosulfan, β -endosulfan and endosulfan sulphate respectively. The fortified samples were left to stand for 24 hours to allow the

samples to equilibrate before being extracted, concentrated and analysed as described above. The unspiked portions were used as the blanks.

2.4.3. Crop samples

The samples were ground using a Phillips grinder (Phillips, Mexico). Duplicate aliquots (5 g) were accurately weighed into a Phillips homogenizer (Phillips, Mexico) containing acetonitrile (50 mL) and homogenized at high speed until the sample was reduced to a slurry. The slurry was quantitatively transferred into 6 centrifuge tubes and then centrifuged at ~5000 rpm. The supernatants were combined into a Kuderna-Danish concentration tube. The sediment in each centrifuge tube was washed with further acetonitrile (3 mL) and recentrifuged. The rinsings were then transferred to the concentration tube containing the first supernatants. The extracts were then concentrated (to 1 mL) and transferred to a prepared Florisil clean up column and allowed to stand for 2 minutes to equilibrate before being eluted first with hexane (15 mL - fraction 1) and then with acetonitrile + hexane (0.5 + 9.5 by volume, 15 mL - fraction 2). The fractions were each concentrated in a Kuderna-Danish flask (to 1 mL) and either analysed immediately or stored in Teflon stoppered glass vials at 4°C until analysed by the GC method described in section 2.4.4.

In preliminary studies the ground root, stem, leaf, and seed samples were spiked with an aliquot (1 mL) of a solution containing a mixture of lindane, endosulfan and endosulfan sulphate as shown in Table 8. Each sample was stirred to ensure even distribution of the spike in the sample and allowed to equilibrate for 24 h before homogenizing, extraction, clean up and analysis as described above.

2.4.4. Details of the gas chromatographic analysis method

Gas chromatograph : Varian Model 3300 GC (Varian AB, Solna, Sweden) fitted with a microprocessor, a split/splitless capillary injector, a Varian Model 4400 integrator and a ⁶³Ni electron capture detector (ECD); column : 30 m x 0.25 mm refined silica capillary column coated with DB-1701 (J and W Scientific, CA, USA); carrier gas: ultra pure nitrogen: flow rate : 5 mL min⁻¹; make-up gas : 25 mL min⁻¹ ; temperatures : column, programmed from 100°C held for 2min and then heated to 250°C at 10°C min⁻¹ and held for 3min; injector, 150°C and operated in the splitless mode during the first 30 seconds only; detector, 300°C. Quantitation was by the external standard calibration method.

3. RESULTS AND DISCUSSION

3.1. Persistence of lindane and endosulfan in the soil

Recoveries of the insecticides from the spiked soil samples were reasonable for all compounds varying between 80-97% (lindane), 84-99% (α -endosulfan) and 87-99% (β -endosulfan) (Table 2) suggesting that extraction and analysis of the field core samples would be satisfactory.

Lindane, α - and β -endosulfan disappeared rapidly during the first 7 weeks after application but declined to much slower rates in the period up to harvest at 25 weeks after application (Table 3). Losses suggested pseudo first order degradation rates over the initial 7 week stage. Similar results were obtained by Liechtenstein and Schultz following a single

application of 25 lbs of dieldrin per acre of Wisconsin soil, USA [31]. The rapid initial rate of loss may be attributed both to microbial degradation during the weeks of high rainfall when the soil was moist and to physicochemical losses by volatilisation and photodegradation. Loss of insecticide by run-off was probably minimal as shown by the extremely low levels of the insecticides detected in the untreated plots.

Lindane was also detected in plot B, while endosulfan was also detected in plot A, the plots which were not treated with that specific pesticide (Table 3). This was attributed to the fact that these plots had been treated by these insecticides in the previous year and indicates the likely extent of carry-over each year.

3.2. Variation of non-target insect populations

At the intervals examined, there were only a few obvious effects on the populations of the species of insects caught in the pitfall traps (Table 4). Firstly, there was a very low population of coleoptera in the endosulfan-treated soya bean plot (B) throughout the study, including the pretreatment period. Secondly, in this plot, the population of semiloopers declined systematically during the 10 week period of monitoring and, thirdly, the population of orthoptera was also lower at the last two intervals of observation though perhaps not outside statistical confidence limits.

It is possible, of course, that at intervals immediately following spray application there could have been larger reductions in the populations of all the species but at the time of the first assessment (2 weeks) their populations had recovered and remained unchanged during the trial, except for the coleoptera, and orthoptera in plot B.

3.3. Uptake of lindane and endosulfan by non-target insects

No insecticide was detected in any of the insects caught in the pitfall traps before spray application. At an interval of two weeks after application, lindane was detected in the insects caught in the traps from the maize plot treated with lindane but also traces of endosulfan (Table 5), presumably from cross-contamination. Similarly, both endosulfan isomers were detected in the insects trapped in the plot of soya beans treated with the endosulfan but, in this case, cross-contamination with lindane was hardly apparent (Table 5). Initially, spiders absorbed more than the other insects but the levels of both lindane and endosulfan (mainly the α -isomer) seemed to decline at later stages, though the data is limited. Spiders are known to be very susceptible to pesticides and can die from very low insecticide doses. Thus the levels in new generations or invasions of spiders decrease as the levels of the insecticides in the plot decrease with time.

Interestingly, the concentration of lindane in crickets and of endosulfan in grasshoppers appeared to increase with time reaching 20 ng g⁻¹ and 28 ng g⁻¹, respectively. The reason for this is unknown but may be related to their diet (Table 5).

In contrast, no detectable residues were found in most of the insects caught in the traps in the untreated, control plots C and D (data not shown). The only exceptions were that traces of lindane were detected in grasshoppers in both plot C (0.09 ng g⁻¹) and plot D (0.003 ng g⁻¹) and α -endosulfan was detected, also in grasshoppers, in plot D (0.004 ng g⁻¹). Grasshoppers are much more mobile and these results may be due to the migration of these insects from the treated plots.

3.3. Distribution of residues in crops

In the preliminary studies, recoveries of all compounds from the spiked samples of the various parts of the crop plants were reasonable with recoveries of 76-90% (lindane), 70-97% (α -endosulfan), 70-106% (β -endosulfan) and 78-105% (endosulfan sulfate) (Table 6), indicating that the overall extraction and analysis procedures were satisfactory for examination of the field crop samples.

Lindane and endosulfan were detected in all components of the maize and soya bean plants from the treated plots. For both maize and soya bean roots, insecticide levels were initially high then declined with time (Table 7). For example, lindane concentration in maize roots was initially found to be 13.8 ng g⁻¹ at 2 weeks after spraying but then declined to 8.2 ng g⁻¹ at 10 weeks after spraying. In soya bean roots, endosulfan concentration was initially at 12.7 ng g⁻¹ at 2 weeks after spraying and declined to 1.53 ng g⁻¹ at 10 weeks after spraying. Lindane concentrations in maize stems and leaves was initially higher than in the roots at 2 weeks after spraying but, as the volume of crop foliage increased its concentration declined to low levels and below the level in the roots at 10 weeks after spraying.

The trend for endosulfan in soya bean stems and leaves is different to that of lindane in maize stems and leaves. Endosulfan in soya stems and leaves is initially low at 2.22 ng g⁻¹ at 2 weeks after spraying then increased to 6.55 ng g⁻¹ at 5 weeks after spraying before dropping to a low concentration of 1.53 ng g⁻¹ at 10 weeks after spraying. The decline in pesticide levels with time is a complex combination of absorption, metabolism and crop volume dilution effects. The concentrations obtained for lindane in maize and soya bean grain at harvesting time were 1.32 and 2.21 ng g⁻¹ respectively. These concentrations, although low, are still surprising since neither compound is expected to be transported to the storage sinks of plants.

Occasionally, trace levels of insecticide were detected from plants growing in the untreated control plots in the interval shortly after application, possibly as a result of drift during spraying (Table 7). The plant samples from the treated plots show higher levels of alien insecticides than the untreated plots (Table 7). This may be because the crops were rotated from the previous year's trial and so any higher level of an alien insecticide from plants in these plots could have resulted from both drift and carry-over from the previous year's pesticide application.

4. CONCLUSIONS

It has been shown that applications of lindane and endosulfan to maize and soya bean crops cause few medium-term changes in the populations of non-target insect species with only possible cause/effect relationships between endosulfan and coleoptera in soyabean. Analysis showed that insecticide concentrations in these species declined for spiders but increased for grasshoppers over the period of observation with both trends probably dependent on the feeding habits of these insects.

Concentrations of the insecticides in the soil initially fell rapidly but this rate of loss declined at later intervals probably as the soil dried out reducing both physicochemical and microbial mechanisms of degradation. Concentrations of both lindane and endosulfan in crop plants also declined to low levels during the monitoring period up to 10 weeks and will have declined even further in the period up to harvest. Surprisingly, low concentrations of both lindane and endosulfan were found in the harvested seeds of the treated crops.

ACKNOWLEDGEMENTS

This work was carried out as part of the FAO/IAEA Co-ordinated Research Programme on "Adverse Side Effects on Flora and Fauna from the use of Organochlorine Pesticides on the African Continent", and was supported in part by a grant from the International Atomic Energy Agency, in part by a grant from the Swedish International Development Agency (SIDA), and in part by a grant from the Research Board of the University of Zimbabwe. We wish to thank the University of Zimbabwe Farm Board for the use of the farm in carrying out this work.

REFERENCES

- [1] MHLANGA, A.A., MADZIVA, J.J., Pesticides Residues in Lake McIlwaine, *Ambio*, **19** (1990) 368-372.
- [2] WHITWELL, A.C., PHELPS, R.J., THOMPSON, W.R., Further records of chlorinated hydrocarbon pesticide residues in Rhodesia, *Arnoldia (Rhodesia)* **6** (1974) 1-7.
- [3] MPOFU, S.M., DDT and its use in Zimbabwe, *Zimbabwe Sci. News* **21** (1987) 31-36.
- [4] CHAPMAN, N.G., Aerial spraying of tsetse flies (*Glossina* spp) in Rhodesia with ultra low volumes of endosulfan, *Trans. Rhod. Sci. Assoc.* **57** (1976) 12-21.
- [5] MATHUTHU, A.S., Feasibility Study on the use of Agro-chemical Pesticides in Zimbabwe: SADC-ELMS sub-programme 4.4 Final report (August 1993).
- [6] ZARANYIKA, M.F., MAKHUBALO, J.M., "Organochlorine Pesticide Residues in Inland Waters in Zimbabwe", in *Pesticides in Zimbabwe: Toxicity and Health Implications*, University of Zimbabwe Publications (1994), in press.
- [7] EDWARDS, C.A., Environmental impact of pesticides, *Protection, Parasitism*, **86** (1987) 309-329.
- [8] EDWARDS, C.A., "Persistent pesticides in the environment" 2nd Ed., CRC Press, Ohio (1976).
- [9] CHIBA, M., MORLEY, H.V., Factors influencing extraction of aldrin and dieldrin residues from different soil types, *J. Agric. Food Chem.*, **16** (1968) 916-922.
- [10] ALLAN, J., "Loss of Biological efficiency of cattle dipping wash containing benzene hexachloride", *Nature (London)* **175** (1955) 1130-1132.
- [11] NEWLANDS, L.W., CHESTERS, G., LEE, G.B., Degradation of γ -BHC in simulated lake impoundments as affected by aeration. *J. Water Pollut. Contr. Fed.* **41**(1969) R174-R188.
- [12] McRAE, I.C., RAGHU, K., CASTRO, T.F., Persistence and biodegradation four isomers of benzene hexachloride in submerged soils, *J. Agric. Food Chem.* **15** (1967) 911.
- [13] MENZI, C.M., Metabolism of Pesticides, U.S. Fish Wildl. Serv., Spec. Sci. Rep (1969) pp 127.
- [14] MILES, J.R.W., MOY, P., Degradation of endosulfan and its metabolites by a mixed culture of soil microorganisms, *Bull. Environ. Contam. Toxicol.* **23** (1978) 1.
- [15] BURNS, R.G., Degradation of thiodan in soil, *Bull. Environ. Contam. Toxicol.* **6** (1971) 361-322.
- [16] KATHPAL, T.S. *et al.*, Degradation of thiodan in soil under Indian conditions, *Indian J. Entomol.* **43** (1981) 420-427.
- [17] MARTENS, R., Degradation of endosulfan-8,9-¹⁴C in soil under different conditions, *Bull. Environ. Contam. Toxicol.* **17** (1977) 438-446.

- [18] EL BEIT, I.O.D., WHEELOCK, J.V., COTTON, E.D., Factors involved in the dynamics of pesticides in the soil: The effect of pesticide concentration on leachability, *Int. J. Environ. Stud.* **16** (1981) 187-7.
- [19] EDWARDS, C.A., THOMPSON, A.R., Pesticides and the soil fauna. *Residue Rev.*, **45** (1973) 1-79.
- [20] GISH, C.D., Organochlorine insecticides in soils and soil invertebrates from agricultural lands, *Pest. Monit. J.* **3** (1970) 241.
- [21] KORSCHGEN, L. J., Soil-food-chain-wild-life relationships in aldrin treated fields, *J. Wildl. Manage.* **34** (1970) 186.
- [22] LINCER, J.L., Organochlorine insecticide residues in Kenya Rift Valley lakes, *J. Appl. Ecol.* **18** (1981) 157 -171.
- [23] GREICHUS, Y.A., GREICHUS, A., AMMANN, B., HOPCRAFT, J., Insecticides, polychlorinated biphenyls and metals in African lake ecosystems.III. Lake Nakuru, Kenya, *Bull. Environ. Contam. Toxicol.* **19** (1978) 454-461.
- [24] MULLER, P., NAGEL, P., FLACKE, W., Ecological side effects of dieldrin application against tsetse flies in Adamaoua, Cameroon, *Ecologia* **50** (1981) 187-194.
- [25] BEESTMAN, G.B., KEENEY, D.R., CHESTERS, K., Dieldrin translocation and accumulation in corn, *Agron. J.* **61** (1969) 390.
- [26] DORMAL, S., MARTENS, P.H., DECLEIRE, M., DE FAESTRAETS, L., Study of the persistence of insecticide residues in various vegetables, fresh, blanched and sterilized., *Bull. Inst. Agron. Stat. Rech. Gembox*, **27** (1959) 137.
- [27] BRETT, C.H., BOWERY, T.G., Insecticide residues on vegetables, *J. Econ. Entomol.* **51** (1958) 818.
- [28] McCASKILL, W.R., PHILLIPS, B.H, Jr., THOMAS, C.A., Residues of chlorinated insecticides in soybean and surface soils from selected counties of South Carolina, *Pest. Monit. J.* **4** (1970) 42.
- [29] BRUCE, W. N., DECKER, G.C., WILSON, J.G., The relationship of the levels of pesticide contamination of crop seed to fat content and soil concentrations of aldrin, heptachlor and their epoxides, *J. Econ. Entomol.* **59** (1966) 179.
- [30] CRITCHLEY, B.R., COOK, A.G., CRITCHLEY, U., PERFECT, T.J., RUSSELL-SMITH, A., The effects of crop protection with DDT on some elements of the subterranean and surface active arthropod fauna of a cultivated forest in humid tropics, *Pedobiologia*, **20** (1980) 31-38.
- [31] LICHTENSTEIN, E.P., SCHULTZ, K.R., Residues of aldrin and heptachlor and their translocation into various crops, *J. Agric. Food Chem.* **13** (1965) 57.

**NEXT PAGE(S)
left BLANK**

Appendix I

EXPERIMENTAL GUIDELINES

1. FIELD EXPERIMENT

This experiment was carried out in maize as a common crop in all participating countries and (optionally) in one other crop chosen according to local conditions. The two crops are hereafter called maize and alternative crop respectively. If the crop is not specified in the following the comments are valid for both crops. The term pre-treatment applies only to the first treatment.

1.1. Experimental layout

1.1.1. Experimental design

Use a randomised block design with four blocks and two treatments (sprayed and unsprayed) and hence eight plots. A plot size of 25 m by 25 m. gives an experimental area of 0.5 ha. Randomise the position of sprayed and unsprayed plots within each block. Where a uniform field is not available establish the experiment so that most of the variation is between blocks and as little as possible between treatments within the same block. If there is a known gradient in the field, (e.g. soil type or slope), place the blocks perpendicular to the gradient (see Fig. 1).




Fig. 1. Experimental design. Two examples of the layout are shown in the figure. The actual layout can be modified slightly to suit local conditions. The thick lines indicate demarcation between blocks. Blocks are marked with roman numerals. s=sprayed plot; u=unsprayed plot. The actual position of sprayed and unsprayed plots within each block is not fixed but must be randomised.

Mark out one subplot (10 m x 10 m) in the centre of each plot for sampling activities (see Fig. 2). Take samples for yield outside this subplot. Mark out other subplots for destructive sampling, e.g. plant and soil samples for residue analyses. Do not take samples within the outer 5 metres of the plot.

Analyse results by two-way analysis of variance.

1.1.2. Pesticides

1.1.2.1. Compounds

Lindane in maize and an appropriate pesticide in the alternative crop.

1.1.2.2. Rate of pesticide application

1 kg (AI) lindane divided in two applications in maize and according to local recommendations in the alternative crop.

1.1.2.3. Method of pesticide application

As a high volume (200-400 L ha⁻¹) spray in maize and according to local recommendations in the alternative crop. Choose the time of spraying to minimize spray drift (usually early in the morning) and keep the nozzle as low as possible.




Fig. 2. Plot layout. Do not sample in the hatched outer area. The 2.5 metres surrounding the 10m by 10 m subplot can be used for destructive sampling, e.g. soil samples and yield.

1.1.2.4. Timing

Maize: The first application of lindane at the three leaf stage and the second two weeks later at a rate of 0.5 kg AI ha⁻¹ on each occasion.

Alternative crop: according to local recommendations.

1.1.3. Field maintenance

1.1.3.1. Weeding

Mechanical

1.1.3.2. Seed dressing:

Accepted if necessary to secure the crop.

1.1.4. Background information to be recorded

- | | |
|---------------------|--|
| 1.1.4.1. Pesticide: | Formulation
Manufacturer (including trade name)
Dose rate
Application details (type of sprayer nozzle type and identification letters/numbers, concentration of spray solution, volume rate of application) |
| 1.1.4.2. Soil: | Mechanical composition (% sand, silt & clay)
organic carbon content
pH (specify soil/water ratio of slurry) |
| 1.1.4.3. Plot: | Size
Method of cultivation
Previous cropping
Neighbouring crops
Previous pesticide history (if known)
Fertilizers applied |
| 1.1.4.4. Crop: | Variety
If seed dressed specify dressing
Seed rate
Sowing date
Density of established crop
Spraying date
Harvest date |
| 1.1.4.5. Weather: | As much as is available regarding rainfall and temperature. Ideally on a daily basis. At least monthly rainfall and mean, maximum and minimum temperatures. |

1.2. Monitoring pests

1.2.1. Species

Stem borers in maize and one chosen key pest in the alternative crop.

1.2.2. Sampling

- (i) Maize crop: search all plants in the subplots for plant damage and record the observations as proportion of damaged plants in each subplot. An estimate of the population of stem borers in maize is optional, made 2-4 weeks after the last spray by dissecting 50 stalks, chosen randomly outside the subplots, and counting all larvae and tunnels.
- (ii) Alternative crop: according to the standard method for crop and pest.

1.2.3. Sampling intervals

One immediately pre-treatment and thereafter every two weeks for two months. Dissection of stalks to be done only once.

1.3. Monitoring non target arthropods

The objectives are to get good estimates of both ground living and foliar living predators. It is, therefore important that appropriate methods for these categories are used. The ground living fauna can be assessed with pitfall trapping or ground search and the foliar living fauna with sweep netting or D-vac sampling. None of these methods is perfect. Pitfall traps are rather easy to use but the catches are dependent on both number of insects and their activity. Ground searches are more tedious but give a more reliable estimate of predator density if performed correctly. Pitfall traps, because they are relatively easy to use, are suggested as the standard method for ground living predators.

The best method for foliar living arthropods is D-vac sampling. However, the apparatus is expensive and not readily available. Sweep netting is not an exact method but should give a reasonably accurate estimate of predator density. Sweep netting is suggested as the standard method for this study. However, those institutes that can use a D-vac sampler can substitute sweep netting with D-vac sampling.

It is very important that time and place of catch of each sample is identified. Each sample must, for statistical reasons, be counted and stored separately. It is not necessary to identify the catches beyond the indicated categories. However, it is recommended that all the listed categories are identified. If possible, store the samples in 50 - 70 % alcohol in airtight containers for future identification of important species.

1.3.1. Soil living insects

The following taxa should be identified and quantified where possible:

- Order Collembola (spring tails)
- Class Araneae (spiders)
- Fam. Coccinellidae (lady bird beetles)

Fam. Formicidae (ants)
Fam. Carabidae (ground beetles).

Other groups (e.g. parasitoids, crickets, etc.) could be added if they are considered important natural enemies and/or are caught in substantial numbers.

1.3.1.1. Pitfall traps

Populations of ground living arthropods can be assessed periodically using simple pitfall traps. Plastic pots or beakers (7~10 cm in diameter; 10- 5 cm deep) buried with the upper rim at soil surface level can be used. It is important that the rim does not protrude above the soil surface (see Fig 3). Four traps are placed in each subplot (see Fig 3). The traps should normally be kept covered with lids or boards. The traps are uncovered and filled to half volume with water with a few drops of detergent (e.g. washing up liquid) at each sampling occasion. The traps are emptied after 48 hours.

Do not use coloured traps (they may attract or repel certain species). Use white or grey traps.

The best way to collect the catches from pitfall traps is to flush the whole contents of the trap through a sieve (ca 1 mm mesh). The catch is then washed down with 70% alcohol (use a wash bottle) through a funnel into a container (this can be done in the field with some training). The arthropods are kept in 70% alcohol for later identification.

1.3.1.2. Ground search

Half metre square, wooden or metal quadrats are used for sampling. The quadrat is placed at random locations in the subplot. The vegetation and soil surface is carefully inspected




Fig 3. Arrangement of pitfall traps.

and any arthropods are caught (use a soft forceps or a pooter) and stored in 70% alcohol for later identification. Ground search should be done in the morning a few hours after sunrise, and always at the same time of the day.

NOTE! Ground search should ideally involve removal of the vegetation and inspection of the soil down to 5 cm to be of real value. However, in view of the small subplots, this would be too destructive to other sampling activities.

1.3.1.3. Pooter

The pooter or aspirator is convenient for the rapid collection of small insects in sweep nets, during ground search, etc. (Fig. 4). The catching end is placed above the insect and, by sucking the mouthpiece, the operator can catch the insects inside the bigger glass tube. By removing one of the corks the pooter can be emptied into a container.




Fig. 4. The pooter.

NOTE! It is important to place a piece of gauze at the end of the glass tube leading into the mouth in order to avoid the risk of inhaling small insects or other particles.

1.3.1.4. D-Vac

The D-vac can be used for both plant and ground sampling. The cone with the biggest opening (1 sq. ft.) is for ground sampling and the smaller (1/3 sq. ft. opening) is for plant sampling.

Take ground samples at 4 different places in each subplot. A sample consists of five subsamples. Each subsample consists of a 5 second suck with the sampling cone lightly pressed to the ground. Take subsamples at intervals of two or three metres until 5 subsamples have been taken without emptying the collecting bag or switching off the

engine in between. After the 5th subsample, remove the collecting cone and empty the collecting bag into a plastic bag. Leave the engine running until the catch is secured in the plastic bag to prevent the insects flying away. Two persons are needed and prior training is necessary. It is recommended to carry a screw driver so that the air duct can be loosened from the engine to recover the bag in the event it is dropped. Repeat the whole procedure of 5 subsamples at 4 different places (e.g. near each pitfall trap) in each subplot.

1.3.1.5. Soil cores

A soil corer 5 cm in diameter and at least 20 cm long is recommended. Ideally it should be possible to divide the sample into sections, for example using a design like that of Tanton [1] (Fig. 5) but this is not essential. Four soil cores (5 cm in diameter 15 cm deep) should be taken from each subplot prior to treatment and again after two months.

The arthropods in the soil sample can be extracted in a Berlese-Tulgren funnel (see Fig. 6). The diameter of the funnel can be between 15-40 cm but the angle should be steep. Laboratory glass funnels are suitable but plastic should be avoided because of potential problems caused by static electricity. The mesh size of the metal gauze should be about 3mm. Some kind of heat source (e.g. a light bulb) should be placed above the sample or the extraction should be done in a warm room. The purpose is to obtain a temperature gradient of 10-20°C from top to bottom of the sample. The sample should be broken into




Fig. 5. Soil corer with split liner.




Fig. 6. The Berlese-Tullgren funnel.

smaller sections before extraction. The sample should, however, be at least 1 cm thick to avoid it drying out. Leave the sample in the funnel for 72 h. Extraction should be done as soon as possible after sampling but it is still possible for up to 4 months if the sample is stored at 4-6°C.

1.3.2. Foliar living insects

1.3.2.1. Sweep net

A standard sweep net consists of a cloth bag fitted into a wire loop (ca 40 cm in diameter) and attached to a 50-60 cm long handle. The net bag is cone shaped and approximately 60 cm deep. As you sweep, tilt the lower rim of the net so it is ca. five cm in advance of the upper rim to catch insects which drop from the plant. Be careful to keep the net below the tops of the plants until the end of the sweep. A single sweep consists of one 180° arc as you step forward. Raise the net at the end of each sweep and reverse the direction of your swing alternately. Take five sweeps for each diagonal. Two diagonals (= two samples) is taken per subplot (see Fig. 7). The catch can be collected with a pooter or put directly in a container with 70% alcohol.

1.3.2.2. D-Vac

As with the ground search, plant samples consist of 5 subsamples. In this case each subsample consists of one plant. Take a total of 4 samples in each plot. To make the sampling as exhaustive as possible place the cone over the whole plants when they are small; when larger, sweep the plants with the cone from all sides. Since the method is impractical on large plants, sample plants only in the first 6 weeks.




Fig. 7. Use of the sweep net.

1.3.3. Soil biology

1.3.3.1. Categories

Breakdown of organic matter (compulsory), earthworms and soil fauna (optional).

1.3.3.2. Sampling methods

(i) *Breakdown of organic material*: Cut disks (2.5 cm diameter), using a cork borer or similar tool, from suitable leaves collected from trees of a single, identified spp growing nearby the experimental plots. The leaves should be rather tough, uniform and growing on the same side of the tree. Place batches of fifty disks in nylon mesh (7-10 mm aperture) bags (ca. 12 cm square), closed with staples and attached to a marker peg with thread. Bury four bags per subplot at random to a depth of ca. 5 cm prior to treatment. Dig up the bags after 2-5 months and wash the contents, then dry in an oven (ca. 600 C for 24 hours) and weigh. Dry and weigh two additional batches of fifty leaves prior to treatment. Present the results in two ways: (a) percent loss in weight (the freshly dried disks are 100%) and (b) the actual weight of the content.

Note! It is important that the bags are left in the soil for an "appropriate", time that is when approximately 50% of the organic matter has been broken down. It is therefore recommended to bury two extra bags in two control plots to be used to decide when to dig up the rest of the bags. Thus one extra bag can be dug up after 1-2 months and the second after 2-3 months.

(ii) *Earthworms*: Use half metre square, wooden or metal quadrats for sampling. Lay them on the soil surface and apply nine litres of dilute formalin (50 mL of 40% in 9 litre water) gradually with a fine spray watering can so that the soil does not become flooded. Most of the worms come to the surface within 15 minutes, and can be picked off and collected into jars containing 5% formalin for storage and eventual identification to species. Take two samples per subplot prior to treatment and again after two months.

1.3.4. Yield

1.3.4.1. Sampling

Estimate yield from 4 subplots (2 m × 2 m) in each plot. Because sampling activities may have an effect on the yield, locate the subplots for yield outside, but close to, the subplots for sampling. Note the moisture content in the crop at harvest.

1.3.5. Residues

1.3.5.1. Categories

Crop (compulsory), soil, earthworms and insects (optional).

(i) *Crop sampling*: For basic analysis a sample of 12 plants per plot is suggested. Collect the leaves only 1 day pre-treatment, 1 day post-treatment, after 2 weeks and after 4 weeks. Weigh, chop, preferably mechanically, and thoroughly mix. Take duplicate subsamples for dry weight and residue analyses.

(ii) *Soil sampling*: Take 12 cores at random from each subplot. Cores 2.5 diameter and 15 cm deep are advised. Weigh the cores before and after removal of any stones and grind to pass a 0.25 cm sieve. Mix the ground cores thoroughly to provide one bulk sample for each plot by successive quartering of the soil on a plastic sheet or in a mechanical mixing device. Take subsamples in duplicate for measurement of moisture content and for residue analysis pre-treatment, one day post-treatment and after 2 and 6 months from both treated and untreated plots.

(iii) *Earthworms*: Worms can be obtained for residue analyses either by the formalin method or by digging. If formalin is used, drop the earthworms into a bucket of clean water immediately they emerge on to the soil surface. Then leave for one day on a damp filter paper in covered petri dishes to allow them to empty their intestines. Then wash, remove surplus moisture with clean tissue paper, and place in clean petri dishes in a deep-freeze and leave until brittle. Take samples pre-treatment and after two months.

1.4. Higher fauna

It is not possible to make up detailed protocols for this area at present. The methods must be adjusted to local conditions and to the range of species that are chosen.

Basically two different approaches can be adopted:

1. Field trial
2. Extensive survey.

In both cases there must be a choice of suitable "indicator" species to study. The criteria could be:

species that are particularly likely to be exposed to the chemical;
species thought to be particularly sensitive to exposure;
species for which an adverse effect would be especially damaging;
abundant species permitting large samples for data collection;
relatively sedentary species;
species whose ecology and behaviour provide easy opportunities to measure effects.

It is unlikely that a single species will fulfill all these roles, and it may be desirable to study a range of species from several categories, at least initially.

1.4.1. Field trial

This type of study requires rather large areas of land (several hectares) where the use of pesticides can be supervised. There must also be some kind of control data to enable comparison. This could be done in two ways: to look for changes before and after treatment on the same site or to compare the treated area with an untreated. A combination of both approaches is recommended.

The measurement should include:

estimation of population numbers (e.g. trapping for mammals and nest counts for birds);
reproductive success;
corpse counts;
residue analyses (corpses, blood, eggs);
biomarkers (if applicable).

Although dead animals give the clearest evidence of a damaging effect this may not always be the most useful approach to detect adverse effects from pesticide use. Carcass searching should, therefore, be complemented with a measure of other effects, e.g. estimation on population changes and reproductive success. If possible a biomarker should be used such as egg shell thinning or repression of enzymatic processes.

1.4.2. Extensive survey

This survey should be done as a comparative study between an area with regular use of organochlorine pesticides (contaminated area) and an area with no such use (control area).

This should include:

background survey, i.e. information on pesticide use, information on residues in higher fauna and identification of indicator species;
identification of contaminated area and control area;
collection of samples for residue analyses.

One crucial step is the choice of indicator species (see above). For this type of study it is important that relatively sedentary species are chosen. For birds, it is recommended to choose one or a few predatory birds e.g. falcons and one or a few insect eating birds, e.g. shrikes.

1.4.3. Sampling for residue analyses

1.4.3.1. Carcasses

Carcasses may have been shot or simply found dead in the field. They should be put in plastic bags, labelled and taken to the laboratory (preferably at 0-4°C) where they should be dissected as soon as possible. Carcass weights and fresh organ and/or tissue weights should be recorded. Liver samples should always be analysed and where appropriate brain, breast muscle and fat should also be sampled. Whole tissues or organs should be homogenised and subsamples taken as needed. If storage is necessary samples or subsamples should be kept at -20°C (or lower) in glass tubes or vessels fitted with tight sealing caps.

1.4.3.2. Eggs

After weighing, eggs should be carefully blown to remove all the contents. The content should be weighed and stored, if necessary, at -20°C. The shells should be retained to assess shell thickness either by the weight/volume ratio method or by direct measurement of sections.

A very useful publication for this area is Sommerville, L. & Walker, C.H. [2].

2. RESIDUE ANALYSES

2.1. Residue analyses (unlabelled pesticides)

If chlorinated hydrocarbons are analysed routinely, follow routine procedures. If not, the following notes can be used as a basis to develop a procedure. No extraction and cleanup procedure can be guaranteed to work first time. Always check the recovery through the procedure by 'spiking' samples (at least 3) of untreated substrate with a known quantity of analyte. The easiest way to do this is to use ¹⁴C labelled pesticides, 3.7 mBq (0.1 g Ci) per sample is enough.

Always run control (untreated) samples of substrate through the procedure to establish the chromatographic background. This is necessary to determine the limit of determination. There are a number of statistical procedures that have been proposed but for most purposes the 'rule of thumb' approach of setting the limit of determination at twice the blank value is satisfactory.

2.1.1. Extraction

2.1.1.1. Plant and non-fatty animal tissue

Homogenise 10-20 g tissue in 200 mL methanol. Filter if necessary. Take a known volume of filtrate (up to 142 mL) and add 2.5 times that volume of water to give a solution containing not more than 40% methanol. The final volume should not exceed 500 mL.

2.1.1.2. Fatty animal tissue and fish

Homogenise 10 g tissue, add 20 g Na₂SO₄ and mix thoroughly. Transfer to a 250 mL separating funnel and add 50 mL hexane or petroleum ether followed by 100 mL acetonitrile saturated with hexane or petroleum ether and shake 1 minute. Drain the acetonitrile into a 1 L separating funnel. Repeat with 3 further portions of 50 mL acetonitrile and combine the extracts. (For reasons of economy you could check the recovery using smaller quantities of acetonitrile.) Add 500 mL water, 40 mL of saturated NaCl solution and 50 mL petroleum ether or hexane. Shake 30 seconds, allow layers to separate and drain aqueous layer into a 2nd 1 L separating funnel. Shake aqueous layer with 50 mL petroleum ether or hexane. Discard aqueous layer and combine the petroleum ether/hexane extracts. Drain through a plug of 10 g Na₂SO (held in a funnel for example) into an evaporating basin or a rotary evaporator. Evaporate just to dryness and dissolve residue in 10 mL methanol. Dilute with 25 mL water.

2.1.1.3. Soil

Extract 50 g soil with methanol in a Soxhlet apparatus for 10 cycles (2-4 hours). Dilute the methanol extract with 2.5 volumes of water as above.

2.1.2. Cleanup using solid phase extraction (SPE) columns

A number of suppliers produce SPE columns. They vary slightly and different batches from the same manufacturer may also vary, so the procedure needs to be checked. Recovery of standard from the column should be at least 90%. For this analysis 6 mL C-18 SPE columns are needed.

2.1.2.1. Column conditioning

Pass two column volumes of methanol, followed by two column volumes of distilled water through the column.

2.1.2.2. Sample addition

Add 200 mL of water diluted methanol extract to the extraction column(s). The flow rate through the column(s) should be 30-40 mL/min. Residual particulates can be removed by placing a 6 mL filtration column and adaptor between the extraction column and reservoir.

2.1.2.1. Column wash

Remove reservoir(s) and wash with one column volume of distilled water. Air dry column under vacuum for 10 minutes.

2.1.2.3. Sample elution

Elute with three 500 µL aliquots of hexane and make up eluate to 2 mL with hexane.

2.1.3. Gas chromatography

There are many sets of GLC conditions cited in the literature. Examples include:

Glass column 5 feet long, 1/4 in diameter packed with 5% QF I on 60-80 mesh Chromosorb G. Column temp. 220°C, detector temp 235°C carrier gas 20 mL/min [3].

Glass column 165 cm × 2 mm packed with 1: 1 mixture of 10% DC-200 and 15% QF-1 on Gas Chrom G. Injector temp 235°C, column 220°C, detector 275°C. Carrier gas 70 mL/min [4].

Glass column 1 m × 3 mm packed with 5% SE30 on Chromosorb W. Column temperature 200°C [5].

2.2. Analysis of (¹⁴C) organochlorine pesticides in maize

(i) Homogenize the maize sample with methanol as in 2.1.1.1.

(ii) Take three replicate samples of not more than 300 mg dry weight and combust in a biological material oxidizer, trapping evolved ¹⁴C₂ in Carbosorb. Analyze the trapped ¹⁴C₂ in LSC to give total radioactivity in the sample.

(iii) Weigh a 5g sample of maize homogenate into a flask and add 20mL methanol. Shake for 30 min. then sonicate for 5 min. in an ultra sonic bath. Transfer the sample to a centrifuge tube, centrifuge for 10 min. at 3000 RPM and remove supernatant. Determine total volume and count 2 × 1 aliquots in LSC for calculation of total extractable radioactivity. Dry the residue and combust to determine unextractable radioactivity. Determine total recovery.

SUMMARY OF SAMPLING PROGRAMME

Type of sampling	Compulsory/Optional	No. of pre-treatment samples	Intervals of post-treatment samples
Plant damage	Comp	1	2, 4, 6, 8, 10 weeks
Dissection of stalks	Opt	0	ca. 8 weeks
Pitfall traps	Comp	1	0*, 2, 4, 6, 8, 10 weeks
D-vac (ground)	Comp	1	1 day, 2, 4, 6, 8, 10 weeks
D-vac (plant)	Comp	1	1 day, 2, 4, 6 weeks
Breakdown organic matter	Comp	-	2-5 months
Earthworms	Opt	1	ca. 2 months
Soil fauna	Opt	1	ca. 2 months
Yield	Comp	-	-
Residues in crop	Comp	1	1 day, 2, 4 weeks
Residues in soil	Opt	11	1 day, 2, 6 months
Residues in earthworms	Opt	1	ca. 2 months

REFERENCES

- [1] TANTON, M.T., A corer for sampling soil and litter arthropods, *Ecology* **50** (1969) 134.
- [2] SOMERVILLE, L. , WALKER, C.H. (Eds), *Pesticide Effects on Terrestrial Wildlife*. Taylor & Francis (1990) pp 404.
- [3] TOOBY, T.E., DURBIN, F.J., Lindane residue accumulation and elimination in rainbow trout (*Salmo gairdinerii* Richardson) and roach (*Rutilus rutilus* Linnaeus), *Environmental Pollution* **8**: (1975) 79-89).
- [4] CARO, J.H., TAYLOR, A.W., Pathways of loss of dieldrin from soils under field conditions, *Journal of Agricultural and Food Chemistry* **19** (1971) 379-384).
- [5] EDWARDS, C.A., 1979. *Proceedings Environmental Seminar on Ecological Tests Relevant to Environmental Chemicals: Evaluation and Research Needs* Berlin December 1977:240-253.

LIST OF PARTICIPANTS

- Benhura, M.A.N. Department of Biochemistry,
University of Zimbabwe,
P.O. Box MP 167, Mount Pleasant,
Harare, ZIMBABWE
- Bennaceur, M. Centre de Développement des Techniques Nucléaires,
Boulevard Frantz Fanon,
B.P. 1017 - Alger Gare, ALGERIA
- Chiverton, P. Department of Plant & Forest Protection,
Swedish University of Agricultural Sciences,
P.O. Box 7044,
S-750 07 Uppsala, SWEDEN
- Deedat, Y. University of Zambia,
Department of Crop Science,
P.O. Box 32379,
Lusaka, ZAMBIA
- Edwards, C.A. Department of Entomology,
The Ohio State University,
103 Botany & Zoology Bldg.,
1735 Neil Avenue,
Columbus, OH 43210-1220, USA
- Farghaly, M.M. M.E.R. Radioisotope Centre for the Arab Countries,
Pesticide Toxicology Laboratory,
Sh. Malaeb El-Gamaa,
Dokki, Cairo 12311, EGYPT
- Hance, R.J. (*Scientific Secretary*) Agrochemicals and Residues Section,
Joint FAO/IAEA Division,
International Atomic Energy Agency,
Box 100, Wagramerstrasse 5,
A-1400 Vienna, Austria
- Ijani, A. Faculty of Agriculture,
Sokoine University of Agriculture,
Department of Crop Science & Production,
P.O. Box 3005,
Morogoro, TANZANIA
- Kiremire, B.T. Department of Chemistry,
Makerere University,
P.O. Box 7062,
Kampala, UGANDA
- Maeda, D. Tropical Pesticides Research Institute,
P.O. Box 3024,
Arusha, TANZANIA
- Meguenni, H. Centre de Développement des Techniques Nucléaires,
2 Boulevard Frantz Fanon, BP 1017
Alger Gare, ALGERIE

- Uronu, B. Agricultural Research Division,
Tropical Pesticides Research Institute,
P.O. Box 3024,
Arusha, TANZANIA
- Minja, E. INCRISAT Office,
P.O. Box 39063,
Nairobi, KENYA
- Montford, K. Department of Biology, Food & Agriculture,
National Nuclear Research Institute,
Ghana Atomic Energy Commission,
P.O. Box 80,
Legon-Accra, GHANA
- Mwangala, F.S. National Council for Scientific Research,
Livestock & Pest Research Centre,
Pesticide Toxicology Laboratory,
P.O. Box CH 310158,
Chelston, Lusaka, ZAMBIA
- Okereke, G.U. Biotechnology Research Center,
Nnamdi Azikiwe University,
P.M.B. 5025, Awka,
Anambra State, NIGERIA
- Osafo-Acquaah, S. Department of Chemistry and Biology,
University of Science and Technology,
University Post Office,
Kumasi, GHANA
- Tukahirwa, E.M. Institute of Environment & Natural Resources,
Makerere University,
P.O. Box 7062,
Kampala, UGANDA
- Umeh, E-D.N. Enugu State University of Science & Technology,
P.M.B. 01660,
Enugu, NIGERIA
- Witkelius, C. S. Department of Entomology,
Swedish University of Agricultural Sciences,
P.O. Box 7044,
S-750 07 Uppsala, SWEDEN
- Yeboah, P. Department of Chemistry,
Ghana Atomic Energy Commission,
P.O. Box 80,
Legon, GHANA
- Zaranyika, M.F. Chemistry Department,
University of Zimbabwe,
P.O. Box MP 167, Mount Pleasant,
Harare, ZIMBABWE