

ANNUAL REPORT
TO THE
DEPARTMENT OF THE ARMY
ARMED FORCES SPECIAL WEAPONS PROJECT
Contract No. DA-49-007-MD-755

STUDY OF THE POST-IRRADIATION SYNDEROME IN HUMANS

Report Period: September 1, 1956-August 31, 1957

Date of Report: October 1, 1957

Investigator: James J. Nickson, M.D., Head, Section of Experimental Radiation, Division of Experimental Pathology, Member, Sloan-Kettering Institute; Chief, Department of Radiation Therapy, Memorial Center

Address: Sloan-Kettering Institute for Cancer Research
410 East 68th Street, New York 21, New York

Not for publication or publication reference without approval of author and Chief, Research and Development Division, Office of the Surgeon General, Department of the Army, Washington 25, D. C.

Amended

Washington National Record Center
Office of the Army Surgeon General
Record Group 112
Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - MD - 755 (OI - 41 - 57) Sloan Kettering Inst

Introduction:

This report will be divided into four sections: I- Hematology and clinical findings, II- Lipoprotein studies, III- Electrolyte and metabolic studies, IV- Future plans.

Selection of Patients

All patients given total body irradiation had disseminated cancer; however, the criteria for the selection of suitable cases were severe so that the patients were as near as possible to the physiological norm. A detailed list of these criteria was included in last year's report.

Method of Irradiation

All external irradiations were accomplished with the 1 Mevp G.E. Resonant generator using exposure ports: upper and lower halves of the body, anterior and posterior. The upper and lower fields were separated by 5 cm. to compensate for the penumbra and the divergence of the beam. This results in a fairly uniform dose throughout the body. All doses are expressed as air doses, measured at the mid-plane of the body. The mid-plane was 290-300 cm from the x-ray target. The dose rate was 5-6 r/min.

In collaboration with Drs. Bond, Cronkhite, and Robertson at the Brookhaven National Laboratory, one patient was given 18 mc of Na^{24} by mouth for an estimated mid-plane body dose of 50 r.

I. Hematological data and case reports

Hematological data on all patients are presented in Table I and Figures 1-13.

Case Reports

Patient M. W. received 18 mc of Na^{24} for a calculated mid-plane dose of 50 r. This 50 year old white female had had a cancer of the base of the tongue with no evidence of residual disease and also had a carcinoma.

Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - UD - 755 (OI - 41 - 57) Sloan Kettering Inst

of the esophagus with regional metastases. Liver function studies and blood counts were within normal limits. No symptoms resulted from the total body irradiation. Pre-irradiation platelet counts averaged about 225,000 and showed very little change until the 20th day post-irradiation when they began to fall, reaching a minimum on the 27th day post-irradiation when the count was 116,000. By the 30th day post-irradiation the platelet count had risen to 177,000 and the patient was discharged from the Brookhaven hospital. Subsequent counts in the laboratory were somewhat lower and probably result from differences in techniques of counting. By the 169th day post-irradiation, the platelet count had returned to 222,000. No pronounced leukopenia was observed. There was a slight anemia of 10.0 gms of Hgb around the 40th day (control 13.0 gms) with a gradual increase to 11.4 gm % on the 141st day. Hematological data are summarized in Table I. The patient showed no adverse effects from the procedure. There has been little advance in her neoplastic process. There was no clinical evidence of bleeding. It is anticipated that the patient will be given a second course of radiation in the near future. She will be re-admitted to Brookhaven National Laboratory Hospital, October 21, 1957 for a second course of total body irradiation.

Patient H.S. had a primary carcinoma of the cervix with pulmonary and pleural metastases. Prior to total body irradiation with 150 r in air at the mid-plane with 1 Mevp x-rays, liver function studies and blood counts were within normal limits. The control white blood count was about $11.3 \times 10^3/\text{cu. mm.}$ and it fell steadily until the 34th day when it reached a minimum of $3.3 \times 10^3/\text{cu. mm.}$ From this time on it rose slowly to $7.0 \times 10^3/\text{cu. mm.}$, but it never returned to control values.

Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - WD - 755 (OI - 41 - 57) Sloan Kettering Inst

By the third day, the lymphocytes fell to about one-half of the control value. Platelet counts showed little change from the control values of 3.5×10^5 /cu. mm. until about the 14th day when they began to drop, reaching a minimum of $.435 \times 10^5$ /cu. mm. on the 27th day. By the 59th day they had returned to 2.84×10^5 /cu. mm. There was evidence of clinical bleeding in the form of petechiae and ecchymoses. The patient had some transitory nausea following irradiation but no vomiting or diarrhea. Graphs of the hematological response are attached as figures 10 to 13.

Patient L.G. had a malignant melanoma of the left leg with widespread metastases in that extremity only. She received 150 r in air to the mid-plane of the body delivered by the 1 Mevp x-ray machine. She experienced no nausea, vomiting or diarrhea. The hematological response followed the same pattern as that of the previously reported cases. The platelet count was 2.57×10^5 /cu. mm. prior to irradiation and began to fall about the 12th day post-irradiation when it was 2.3×10^5 /cu. mm., and it reached a minimum of 0.20×10^5 /cu. mm. on the 28th day. It increased steadily from this point on, reaching 3×10^5 /cu. mm. on the 56th day. Granulocyte counts paralleled the total white count and the lymphocytes showed the usual prompt drop to about one-half of the control value with a slow and incomplete recovery. Details are shown in the attached graphs, figures 1 to 5.

One patient (C.H. previously reported as K.H.) deserves special attention. She had a primary carcinoma of the stomach which was not resectable as proven by exploratory laparotomy. On December 19, 1955 she received 150 r total body irradiation. Radiation treatment of the

Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - MD - 755 (OI - 41 - 57) Sloan Kettering Inst.

primary lesion was started on February 7, 1956 with the cobalt 60 machine. She received a tumor dose of 5000 r in 5 weeks. The patient did extremely well until June 5, 1957 when pelvic metastases were found. The patient had minimal symptoms from these lesions and was in excellent general health. On July 1, 1957 she received a second course of total body irradiation up to a dose of 75 r in air. Following this, she received a total of 4400 r to the pelvis in 32 days. This treatment was completed on September 20, 1957. (Patient included in this report even though some of data obtained after August 31, 1957.) The attached graphs figures 6 to 9 summarize the hematological findings following the second course of total body irradiation.

Table II

Maximum Changes During the First and Second Courses
of Total Body Irradiations

Patient, C.H.	<u>First Irradiation 150 r</u>		<u>Second Irradiation 75 r</u>		<u>Prior to Pelvic Irradiation</u>
	<u>Control</u>	<u>Maximum Change</u>	<u>Control</u>	<u>Maximum Change</u>	
Hgb	12.3	no change	12.3	no change	12.8
RBC	5.6	_____	4.4	_____	3.7
WBC	7.9	2.9	5.4	2.4	5.4
Granu- %70 locytes		53	57	71	
Lym- %27 phocytes		40	17	12	
Plate- 225 lets		50.4	154	117	204

Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - MD - 755 (OI - 41 - 57) Sloan Kettering Inst

There was no evidence of bleeding at any time and bleeding and clotting time remained within normal limits. Bone marrow aspiration before the first total body irradiation was reported as rather hypocellular and appeared to be diluted with peripheral blood. What marrow elements were seen appeared "normal" and prior to the second course, the report stated: "marrow appears fatty and hypoplastic in some areas, in others it is adequately cellular with all cell elements represented, with increased numbers of cells of the erythroid series."

Since institution of pelvic irradiation there has been a progressive anemia and thrombocytopenia. This patient has been re-admitted to the hospital for investigation of possible sites of bleeding, and to evaluate the status of the bone marrow.

II. Lipoprotein studies

Lipoprotein studies as previously described were carried out on one patient (L.G.). At the time of the maximum thrombocytopenia there was a marked increase (+++) in the amount of clearing factor. This persisted, even after the platelet count had risen above 200,000. At the time of the greatest thrombocytopenia there were petechiae. This observation was consistent with earlier ones, viz. that there has been an increase in clearing factor in all patients that showed clinical evidence of bleeding. Patients who have not had clinical evidence of bleeding have shown no increase in clearing factor; however they have had severe thrombocytopenias.

Shifts of emphasis in the lipoprotein research program within the Institute have resulted in the discontinuation of this facet of the investigation. The professional personnel and equipment formerly available to this phase of our project are now being fully utilized in other studies.

Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - MD - 755 (OI - 41 - 57) Sloan Kettering Inst

III. Electrolyte studies

The common electrolyte abnormalities occurring after irradiation have been outlined, and appropriate therapies have been established (1). These are potassium depletion alkalosis from inanition, hypochloremic alkalosis from vomiting, and hyperchloremic acidosis from diarrhea. In addition to these, adrenal insufficiency, ammonia toxicity, and hyperkalemia may be seen on occasion.

Potassium spaces have been studied pre- and post-irradiation. Such data are being analyzed by an analogue computer to see if there is any difference in the various body compartments into which potassium goes. Preliminary data suggest that there may be differences in the very early portion of the fall-off curve.

The CNS death modality in dogs has been studied from the physiological point of view. It has been shown that the convulsive phase of this syndrome is associated with a respiratory alkalosis (2). This work is being continued. Pathological examination of the irradiated brains is being done in collaboration with the Pathology Department of Cornell University Medical College.

IV. Future plans

Careful and detailed dosimetry will be done utilizing total body phantoms. These will be designed to measure integral dose by means of chemical detectors. A phantom has been designed which contains a complete skeleton. This will permit measurement of dose at any point that is considered to be of biological importance.

Hematological studies will be continued, and additional work will be done on enzymes associated with the clotting mechanism.

Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - WD - 755 (OI - 41 - 57) Sloan Kettering Inst

Potassium spaces will be studied pre- and post-irradiation with particular emphasis on the immediate fall-off curve. Glucose tolerance curves will also be determined. Further work is anticipated on mechanisms of the super-lethal death in dogs.

Some patients will receive total body irradiation by means of internal isotopes to see if there is any difference in response to a more protected exposure. It is hoped that more patients can be given two courses of total body irradiation to see if there is a difference in response to the second exposure.

BIBLIOGRAPHY

- (1) Bane, H. N., Glicksman, A. S., and Nickson, J. J. Common electrolyte problems in radiation therapy. Am. Journal of Roentgenology, in press.
- (2) Bane, H. N., Whitehouse, W. L., and Nickson, J. J. Physiological changes in the acute radiation syndrome in dogs. Radiation Research Meeting, 1956.

Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - MD - 755 (OE - 41 - 57) Sloan Kettering Inst

Table I

PATIENT M.W.: HEMATOLOGICAL DATA, PRE- AND POST-IRRADIATION

	Days, Pre- and Post-Irradiation									
	-15	-12	-9	-5	-2	2	5	8	12	
Hemoglobin	13.0									
RBC (1,000,000)										
WBC (1,000)	4.1	5.4	4.9	3.8	6.4	4.7	5.2	3.9	6.9	
Polynuclear	61									
<u>fil.</u>										
non-										
<u>fil.</u>										
Eosinophiles	6									
Basophiles										
Monocytes	5									
Lymphocytes	28									
Hematocrit	33.8	35.5	34.8	35.5	37.2	34.3	36.2	35.0	35.0	
Platelets (1,000)	253.8	212.5	227.5	250.0	261.3	296.3	225.0	332.5	252.5	

Days 16 - 33 (continued)

	Days, Post-Irradiation									
	16	20	23	26	27	28	29	30	33	
Hemoglobin										10.8
RBC(1,000,000)										3.7
WBC (1,000)	6.0	5.4	4.8	4.9						4.7
<u>fil.</u>	68									72
Polynuclear										
<u>non-</u>										
<u>fil.</u>	4									8
Eosinophiles	5									6
Basophiles	2									1
Monocytes	11									4
Lymphocytes	10									9
Hematocrit	35.5	32.6	37.0	36.0				34.4		
Platelets (1,000)	225.0	231.2	130.0	128.8	116.3	142.5	147.5	177.5	125.8	

(Continued on next page)

Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - UD - 755 (OI - 41 - 57) Sloan Kettering Insts

Table I
(Continued)

PATIENT M.W.: HEMATOLOGICAL DATA, PRE- AND POST-IRRADIATION

	Days, Post-Irradiation							
	36	40	43	47	54	57	61	71
Hemoglobin	10.2	9.9	10.8	10.6	10.8	10.8	10.9	10.8
RBC (1,000,000)	3.5	3.3	3.3	3.1	3.8	3.6	3.6	3.6
WBC (1,000)	4.7	4.4	3.1	2.9	3.4	4.3	4.5	4.0
fil.	77	67		77	77	79	81	80
Polynuclear	-----							
non-								
fil.	5	2		2	3	2	2	3
Eosinophiles	1	2		1	1		1	3
Basophiles	1	1		1				
Monocytes	6	8		3	5	6	8	7
Lymphocytes	10	20		16	14	13	8	7
Hematocrit								
Platelets (1,000)	123.0	125.4		124.0	133.0	126.0	133.2	148.0

Days 85 - 169 (continued)

	Days, Post-Irradiation				
	85	99	127	141	169
Hemoglobin	11.2	11.1	11.3	11.4	10.9
RBC (1,000,000)	3.7	3.7	3.8	3.5	3.7
WBC (1,000)	4.5	5.0	4.9	4.2	6.3
fil.	84	82	81	76	78
Polynuclear	-----				
non-					
fil.	1	2	3	5	4
Eosinophiles	3	2	1	1	2
Basophiles			2		
Monocytes	2	4	6	8	7
Lymphocytes	10	10	7	10	8
Hematocrit					
Platelets (1,000)	140.4	162.8	167.2	196.0	222.0


Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583


Box #: 165

File: DA - 49 - 007 - MD - 755 (OI - 41 - 57) Sloan Kettering Inst

L.G. MALIGNANT MELANOMA LEFT LEG



L.G. MALIGNANT MELANOMA LEFT LEG




Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165


File: DA - 49 - 007 - UD - 755 (OI - 41 - 57) Sloan Kettering Insts

L.G. MALIGNANT MELANOMA LEFT LEG



4

L.G. MALIGNANT MELANOMA LEFT LEG




Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583


Box #: 165

File: DA - 49 - 007 - MD - 755 (OE - 41 - 57) Sloan Kettering Insts

L.G. MALIGNANT MELANOMA LEFT LEG



C.H. CANCER OF GASTRO-ESOPHAGEAL JUNCTURE




Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583


Box #: 165

File: DA - 49-007-UD-755 (OI-41-57) Sloan Kettering Inst

C.H. CANCER OF GASTRO-ESOPHAGEAL JUNCTURE



C.H. CANCER OF THE GASTRO-ESOPHAGEAL JUNCTURE




Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - UD - 755 (OI - 41 - 57) Sloan Kettering Inst

C.H. CANCER OF GASTRO-ESOPHAGEAL JUNCTURE




Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165


File: DA - 49 - 007 - MD - 755 (OI - 41 - 57) Sloan Kettering Inst

H.S., CA. CERVIX & PULMON. METAS.



//

H.S., CA. CERVIX & PULMON. METAS.




Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583


Box #: 165

File: DA - 49 - 007 - MD - 755 (OI - 41 - 57) Sloan Kettering Insts

H. S., CA. CERVIX & PULMON. METAS.



H. S., CA. CERVIX & PULMON. METAS.



Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - MD - 755 (OI - 41 - 57) Sloan Kettering Inst.

Initial Distribution

Copies

5 Armed Forces Special Weapons Project (SWPTI-3)
5 Armed Services Technical Information Agency (TIOSCP-2)
4 The Surgeon General, Department of the Army (MEDDH-SP)
1 Army Environmental Health Laboratory
1 Army Medical Research Laboratory (MEDEA-XD)
1 Walter Reed Army Institute of Research
1 Office of Naval Research (Code 454)
1 Bureau of Medicine and Surgery, Department of Navy
1 Air Research and Development Command
1 Dr. V. P. Collins, Baylor University
1 Dr. J. K. Hampton, Jr., Tulane University

22

Washington National Record Center
Office of the Army Surgeon General
Record Group 112

Accession #: 62A 1583

Box #: 165

File: DA - 49 - 007 - UD - 755 (OI - 41 - 57) Sloan Kettering Inst.