

MARSHALL ISLANDS FILE TRACKING DOCUMENT

Record Number: 295

File Name (TITLE): Uptake of Iodine-131 by the
Red Alga

Document Number (ID): UWPR-44

DATE: 11/1955 (?)

Previous Location (FROM): CR

AUTHOR: R. F. Palumbo

Additional Information: _____

OrMIbox: 17

CyMIbox: 11

**UPTAKE OF IODINE -131 BY THE RED ALGA
ASPARACOPSIS TAXIFORMIS**



Ralph F. Palumbo

**Applied Fisheries Laboratory
University of Washington
Seattle, Washington**

**Radiochemical analyses
Dorothy South**

November 20, 1958

**Operated by the University of Washington under Contract No
AT(40-1)1840 with the United States Atomic Energy Commission**

**BEST AVAILABLE
COPY**

UNCLASSIFIED

UWFL

44

1 OF 1

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, express or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission to the extent that such employee or contractor prepares, handles or distributes, or provides access to, any information pursuant to his employment or contract with the Commission.

ABSTRACT

Specific activity measurements, radiochemical analyses, and half-life studies were made of the red alga, Asparagopsis taxiformis, collected at Eniwetok Atoll in March, April and May, 1954. The results of these studies show that this alga more than any other tested, concentrated the 131 produced by the nuclear weapons tests at Bikini and Eniwetok Atolls during this period. This alga may prove useful as an indicator of radioactive fallout from nuclear detonations.

Introduction

During the weapons testing program of Operation Castle at Bikini and Eniwetok Atolls in the spring of 1954 the Applied Fisheries Laboratory of the University of Washington carried out studies on the uptake of radioactive materials in a variety of aquatic and terrestrial plants and animals. In addition to the programmatic studies, exploratory studies also were continued, including a study of the reef algae.

Collections of algae from near the Marine Biological Laboratory on Parry Island, Eniwetok Atoll, indicate that a small red alga (*Asparagopsis*) has a remarkable affinity for iodine as measured by the uptake of I^{131} immediately following weapon detonations at the Pacific Proving Ground.

Collection of Material

Immediately following the March 1, 1954 experiment at Bikini Atoll, it was observed that the background level of radioactivity on the reef of the seaward side of Parry Island had increased. Most of the algae growing on this reef had only slightly higher readings than the background level. However, the level for *Asparagopsis taxiformis* (Delile) Collins and Hervey, a delicate, branching red alga, was unexpectedly high.

Radioassays were made of algae collected at Parry and Bngambngo Islands during March, April, and May, 1954. The results of these studies are given in Table I.

Table I Radioactivity of Algae Collected at Eniwetok Atoll
March, April, May, 1954

Values expressed in thousands of d/m/g of wet weight

Species	Date Collected (1954)	Location	d/m/g (Thousands)
<u>Asparagopsis taxiformis</u>	3/6	Parry I. (seaward reef)	3.92 2.03 1.50
<u>Euteromorpha</u> sp	"	" "	195
<u>Halimeda tuna</u>	"	" "	113
<u>Padina Compressalis</u>	"	" "	307
<u>Asparagopsis taxiformis</u>	4/3	" "	12.4 11.4
<u>Dictyota planatifida</u>	"	" "	12
<u>Asparagopsis taxiformis</u>	4/25	Bogombogo (lagoon reef)	2.4 4.42
<u>Spyridia filamentosa</u>	"	" "	5.6 5.02
<u>Asparagopsis taxiformis</u>	5/19	" "	39,100 21,000
<u>Caulerpa racemosa</u>	"	" "	4,540 2,420 918
<u>Halimeda opuntia</u>	"	" "	8,750 5,610

It was thought that the radioactivity in the samples of Asparagopsis collected in early March might be due to a short half-life isotope resulting from the test of March 1, 1954 at Bikini. In order to verify this assumption series of counts for determining radioactive half life were initiated at the Eniwetok laboratory. Duplicate samples were sent to the home laboratory at the University of Washington for more complete analyses. The results of tests conducted on samples collected at Eniwetok and Rongelap Atolls from March through June are given in Table II.

Plots of the radioactive decay of three samples of Asparagopsis are shown in Figure 1.

Discussion

The results of the half-life studies and radiochemical analyses* indicate that at least 90 percent of the radioactivity in Asparagopsis was due to 131 . The half life of the isotope contributing 90 percent of the radioactivity in Asparagopsis was found to be 7.6 to 8.5 days. This compares well with the accepted value for 131 , which is given as 8.14 days (3). The lowest values obtained at Eniwetok may have been due to the presence of

* The radiochemical method for determination of 131 was adapted from Olenchuk et al (2) and is as follows. The dried algae were heated several minutes with NaClO and 0.1 N NaOH containing iodine carrier (added as KI), filtered, and discarded. The iodine in the solution was reduced with hydroxylamine hydrochloride, extracted into CCl_4 , and re-extracted into water containing a little bisulfite. The aqueous solution was acidified and treated with sodium nitrate. The extraction into carbon tetrachloride and into water was repeated. The iodine was precipitated from the resulting solution as silver iodide and counted in an internal gas flow counter.


Table II. Results of Radiochemical and Half-life Studies of Samples Collected March, April, May, 1954

Species	Date Collected (1954)	Location, Atoll - Island	Half Life	¹³¹ I percent of total activity *
<u>Asparagopsis taxiformis</u>	3/10	Eniwetok - Parry	7.8	
" "	3/19	" "	8.2	
" "	4/3	" "	8.5	90
<u>Caulerpa racemosa</u>	3/28	Rongelap - Kabelle		0.18
<u>Ulotea indica</u>	"	" "		0.20
<u>Boerhaavia tetrandra</u> **	"	" Labaredj		0.25
<u>Morinda citrifolia</u>	"	" "		0.38
<u>Goettardia speciosa</u>	"	" "		3.5
<u>Asparagopsis taxiformis</u>	5/19	Eniwetok - Bogaubogo	7.8	90.
" "	"	" "	8.2	

* Rare earths, 2.8, Zr⁹⁵, 1.0; unidentified, 6.2 percent of total activity

** Leaves

Figure 1 Decay curves of Asperagopsis collected at Eniwetok Atoll (A) Whole sample from Parry Island, 3/10/54 (n 1), (B) whole sample from Bugumbogo Island, 3/19/54 (n 100), (C) extractable 131 from sample "B" (n 10).



132 , 133 , and 135 isotopes whose faster decay rates would tend to lower the value

It was evident from the short half life and high level of radioactivity of two samples of Asparagopsis collected on April 3, 1954 at Parry Island that these algae were concentrating a recently formed isotope. The values for the counts of these samples were 12, 000 and 11, 000 d/m/g as opposed to an average value of 200 d/m/g based on radioactive decay of samples collected in the same locality on March 8, 1954. The increase in specific activity might have been due to material resulting from the March 27 test at Bikini Atoll, since it was observed that the background count at Parry Island increased a few days after this date. Changes in the activity levels in the algae would be expected to follow closely those of the environment, for it has been shown in laboratory tests that 131 uptake reaches equilibrium after the first hour in Acrophyllium (4) and after three hours in Laminaria (7). As laboratory tests have not been conducted with Asparagopsis to determine this matter, it cannot be stated with certainty that it would behave similarly.

It has been known for many years that iodine is present in the sea and in a majority of the algae (8). In Asparagopsis iodine constitutes 0.002 percent of the dry matter (2). Laboratory experiments have indicated that certain algae will concentrate 131 from sea water (3, 4, 7). The observations made at Eniwetok during the present investigations support these findings since the 131 concentration in Asparagopsis was approximately 10, 000 times that of the surrounding water. (The total activity of

the water was 24,700 d/m/g and the activity due to 131 in the algae was 30,000,000 d/m/g. The contribution of 131 to the total activity of a fission products mixture at 10 days is 6.8 percent, therefore, assuming the ratio to be the same, the activity in the water due to 131 was 1,800 d/m/g. Thus the concentration factor for 131 by the algae was 17,000. If other iodine isotopes are considered, this figure would probably be higher.

The role of iodine in the physiology of the algae is uncertain, but it has been detected as mineral iodide (5, 6) as well as the iodine amino acid, iodo-tryptane (5, 6). It is now generally accepted that iodine occurs in both forms and that the ratio of the two varies in the different species of algae (3). The absorption of 131 has been shown to be associated with respiration (4), since its uptake in Acrophyllum, a brown alga, decreases when nitrogen is used in place of oxygen. The specimens of Asparagopsis from Eilat were collected in a region of pounding surf and active water movement, conditions favoring maximum aeration and maximum uptake of 131 .

Conclusions

It has been observed from the investigations following the weapons tests at the Pacific Proving Ground that a red alga, Asparagopsis taxiformis, has a remarkable affinity for 131 , an isotope with a half life of 8.14 days. It is believed that the alga may prove useful as an indicator of radioactive fallout from nuclear detonations.

REFERENCES

- 1 Dixit, S C 1930 A note on the percentage of iodine in certain algae
J Indian Chem. Soc. 7(12) 959.
- 2 O'endenis, L E, R P Metcalf, T. B. Novey and C D Coryell. 1951
Interchange of radioactive iodine with carrier iodine. in Radiochem-
ical Studies The Plaston Products. Book 3 McGraw Hill, N.Y.
pp 1629-1634
- 3 Institute of Seaweed Research Annual Report 1953
- 4 Kelly, S 1953. Respiration and iodine uptake in Acicophyllum Biol
Bull. 104 130.
- 5 Kineman, S. 1954 Radiological Health Handbook Cincinnati, Ohio
p 136
- 6 Kyla, H 1928 Ueber das Vorkommen von Iodiden, Bromiden und
Iodidoxydosen bei den Meeresalgen Hoppe-Seyl Z 166 29
- 7 Rucka, J and Y Yagi. 1952 Sur la fixation de l'iode radioactif par
les algues et sur les constituants iodés des Laminaires Compte
Rend Soc Biol, Paris. 146 642
- 8 Vinogradov A P 1953 The elementary chemical composition of
marine organisms Sears Foundation for Marine Research, Mem-
oir No II, Yale Univ. New Haven Conn.

END
