

~~CONFIDENTIAL~~

410914

NRDL MARSHALL ISLAND RESURVEY - 1956

RESULTS OF ANALYSES PERFORMED AT HASL

Laboratory Report 56-7

CONFIRMED TO BE UNCLASSIFIED  
BY AUTHORITY OF DOE/OC

Carl Wilson 2/1/85  
REVIEWED BY DATE

by

E. P. Hardy

W. R. Collins

US DOE ARCHIVES	
326 U.S. ATOMIC ENERGY	
RG	COMMISSION
Collection	<u>DCS McCraw</u>
Box	<u>9 Job 1320</u>
Folder	<u>Radiological Survey</u>

~~CONFIDENTIAL~~

49  
51

~~CONFIDENTIAL~~

in polyethylene containers. Concentrations are based on the weight of the material at the time it was received at HASL. Consequently, all radiochemical and analytical results are reported here in terms of d/m or grams, per gram of material as received at HASL. Dr. H. Weiss has stated via letter<sup>(2)</sup> that the wet weights of the plant specimens were not recorded at the time of collection. He has proposed that the results be expressed in terms of d/m/kg of material as received at NRD.

For determination of total beta activity an aliquot of the solution of wet-ashed material was transferred to a glass planchet, evaporated and dried under an infra-red lamp. Counts were converted to disintegrations by applying a geometry factor based on  $K^{40}$  as a standard. A self-absorption correction was also applied in each case.\* Under the wet ashing and plating conditions used at HASL possible loss of volatile fission products such as  $Ru^{106}$  -  $Rh^{106}$  is avoided.<sup>(3)</sup> For practical reasons, the coconut outer and inner shells were dry ashed at 500°C prior to dissolution.

\* For the particular specimen type under consideration, several values of activity vs. dry weight of the plated aliquot were plotted and a smooth curve fitted for the points. Another curve (based on extrapolation to zero mass) of activity ratio ( $A/A_0$ ) was drawn and used for determining the self-absorption correction. See Figures 2 through 8.

DOE ARCHIVES

The procedure outlined in NYO-4617 was followed for the radiochemical analysis of Sr<sup>90</sup>. The Cs<sup>137</sup> analyses were performed by S. Tarras using a method which to date has not been documented. It involves the coprecipitation of cesium with ammonium alum to eliminate mixed fission products as well as potassium, then a final precipitation as the chloroplatinate. Radiochemical and gravimetric yields of 95% are attainable. The samples were analyzed for calcium by C. Baxter employing the oxalate-permanganate titration method<sup>(4)</sup>.

As a check on radiochemical purity, beta absorption analyses were carried out by N. Hallden<sup>(5)</sup> on the Cs<sup>137</sup> fractions of two pooled urine samples (specimens collected at Utirik and Likiep), one water sample (HASL #3457), and one soil (HASL #3462). In each case Cs<sup>137</sup> was positively identified and there was no evidence of other interfering isotopes. The radioactive decay of the Y<sup>90</sup> fractions of the urine samples was followed over a period of one hundred hours. Within statistical limits concurrence with the theoretical half-life was observed.

Analytical results are shown in Tables 2 through 6. The error term accompanying each absolute result represents one standard deviation due to the error in counting. The only available interlaboratory cross-check data are given in Table 7. These

DOE ARCHIVES

results were obtained from Dr. S. Cohn by phone on April 24, 1956. A follow-up letter from Dr. Cohn<sup>(6)</sup> expressed his idea that the discrepancy in the beta count probably lies in the conversion from c/m to d/m. NRDL used Sr<sup>90</sup> - Y<sup>90</sup> as a standard in this case and for purposes of comparison, the HASL results were also standardized against Sr<sup>90</sup> - Y<sup>90</sup>. It is felt that the use of K<sup>40</sup> as a standard allows the best approximation of the energy for mixed fission products among the available long-lived isotopes<sup>(7)</sup>.

As an aid in evaluating these data, Figure 1 and Table 8 are included.

DOE ARCHIVES

REFERENCES

1. Memorandum from Dr. J. H. Harley to Mr. M. Eisenbud, "Rongelap Resurvey Samples from NFDL", April 17, 1956.
2. Letter of May 9, 1956 from Dr. H. Weiss to Edward Hardy.
3. Memorandum from G. Hamada to Dr. J. H. Harley, "The Effect of the Wet Ashing Technique Used at HASL on Ruthenium Volatilization", March 29, 1956.
4. Private communication, Mr. I. B. Whitney.
5. "Analyzing Beta Absorption Graphically to Identify Emitters", J. H. Harley and N. Hallden, Nucleonics, 13, 1, January 55, pp 32-35.
6. Letter of May 2, 1956 from Dr. S. Cohn to Mr. I. B. Whitney.
7. HASL Laboratory Report 56-1, "Standardization and Operation of Fallout Counters", N. A. Hallden and J. H. Harley.

DOE ARCHIVES

TABLE 1

## NNDL MARSHALL ISLAND RESURVEY - 1956

## Samples Received at HASL

(samples analyzed at HASL shown in parenthesis)

MARINE ORGANISMS - 65 (23), received 3/6/56

	<u>Rongelap</u>	<u>Oejen</u>	<u>Eniaetok</u>	<u>Eniwetak</u>	<u>Sifo</u>	<u>Utirik</u>	<u>Likiep</u>	<u>Kabell</u>
<u>Fish</u> - 37 (13)								
Unicorn	1			1				
Mullet	1							
Surgeon	1 (1)	1 (1)		1 (1)		2 (2)		
Damsel	1 (1)			1 (1)		2 (2)	1 (1)	1 (1)
Sea Cucumber	1							
Bl. Tip Shark								1
Trigger								1
Siganus					1		1	1
Butterfly					1 (1)		1 (1)	1 (1)
Snapper		1		1	1			
Squirrel		1			1	1		
Parrot			1		1	1		
Angel			1		1			
Goat			1					
Sergeus					1			
Sea Bass							1	
<u>Crab</u> - 11	3	1	1		3	3		
<u>Clam</u> - 2	2							
<u>Snail</u> 9 (4)	2	4 (4)*				2		1
<u>Coral</u> 6 (6)	1 (1)	1 (1)	1 (1)			2 (2)	1 (1)	

LAND PLANTS - 77 (14), received 4/3/56

<u>Coconuts</u> - 26 (5)	3 (3)	3	4	4	4	4 (1)*	4 (1)*	
<u>Portulaca</u> 6	1	1	1	1		1	1	
<u>Pandanus</u> - 18 (2)	3 (2)*	2	3	2	2	3	3	
<u>Papaya</u> - 9	3					3	3	
<u>Arrowroot</u> 14 (7)	2 (1)*	2 (1)*	2 (1)*	2 (1)*	2 (1)*	2 (1)*	2 (1)*	
<u>Banana</u> - 2							2	
<u>Taro</u> - 2							2	

SOIL - 21 (13), received 4/3/56

	3	3 (2)*	3 (3)*	3 (2)*	3 (2)*	3 (2)*	3 (2)*	
--	---	--------	--------	--------	--------	--------	--------	--

LAND WATER - 7 (6), received 4/3/56

<u>Well</u> - 4 (4)						2 (2)	2 (2)	
<u>Cistern</u> - 2 (1)	1 (1)					1 (1)*		
<u>Lens</u> - 1 (1)			1 (1)					

SEA WATER - 14 (14), received 4/3/56

<u>Ocean</u> - 7 (7)	1 (1)*	1 (1)*	1 (1)*	1 (1)*	1 (1)*	1 (1)*	1 (1)*	
<u>Lagoon</u> - 7 (7)	1 (1)*	1 (1)*	1 (1)*	1 (1)*	1 (1)*	1 (1)*	1 (1)*	

URINE - 24 (24), received 3/29/56

	5 (5)					10 (10)	9 (9)	
	(Majuro)							

\* Interlaboratory cross-check samples.

INCOMPLETE DOCUMENT REFERENCE SHEET

The archive copy of this document is incomplete.

Pages missing 7-11

Enclosures missing \_\_\_\_\_

Attachments missing \_\_\_\_\_

Other \_\_\_\_\_

W Teach  
signature

9/17/87  
date

US DOE ARCHIVES
326 U.S. ATOMIC ENERGY
RG COMMISSION
Collection <u>DOS McCraw</u>
Box <u>9</u> Job <u>1320</u>
Folder <u>Radiological Survey</u>

DOE ARCHIVES

56  
58

TABLE 7

NRDL MARSHALL ISLAND RESURVEY - 1956

INTERLABORATORY COMPARISON

(Snail Solutions Prepared at NRDL from Specimens Collected on Gejen Island)

<u>HASL #</u>	<u>NRDL #</u>	<u>Type</u>	<u>HASL*</u>	Total $\beta$ Activity (d/m/gram-wet) <u>HASL*</u>	<u>NRDL<sup>o</sup></u>	Total $\gamma$ Activity (d/m/gram-wet) <u>NRDL</u>	<u>HASL</u> Sr <sup>90</sup> (d/m/gram-wet) <u>NRDL</u>
3326	1636	Spider	520 $\pm$ 10	570 $\pm$ 11	877	378	4.4 $\pm$ 0.39
3327	1637	Spider	2180 $\pm$ 29	2400 $\pm$ 32	2965	1605	1.3 $\pm$ 0.34
3328	1638	Scorpion	23310 $\pm$ 290	25600 $\pm$ 300	29700	9150	1.1 $\pm$ 0.44
3329	1639	Scorpion	9800 $\pm$ 120	10800 $\pm$ 125	14250	4640	1.5 $\pm$ 0.58

\* Standardized against K<sup>40</sup>

† Standardized against Sr<sup>90</sup>- Y<sup>90</sup>

o Standardized against Sr<sup>90</sup>- Y<sup>90</sup>

DOE ARCHIVE

NOTE: Net weights furnished by NRDL  
NRDL results forwarded by phone to I. B. Whitney  
from S. Cohn on April 24, 1956.

TABLE 8

NRDL MARSHALL ISLAND RESURVEY - 1956

TOTAL  $\beta$  ACTIVITY - d/m/gram\*

	<u>Rongelap</u>	<u>Enlaketok</u>	<u>Kaballe</u>	<u>Gejen</u>	<u>Enlaketak</u>	<u>Sifo</u>	<u>Utirik</u>	<u>Likiep</u>
<u>FISH</u>								
Surgeon	52				34		22; 18	11
Damsel	37		120	230	20		14; 22	51
Butterfly						95		
<u>CORAL</u>	35	200		310			18; 21	115
<u>SNAILS</u>								
Spider				520; 2200				
Scorpion				23,000; 9800				
<u>LAND PLANTS</u>								
Coconuts								
Outer Husk	71;66 } 26							
Inner Shell	26;35 }							
Meat and Milk	98;87							
Milk	43							
Pandanus	42; 30			300	67	59	26	7.3
Arrowroot	lost	180		69,000; 120	3000; 461	620; 457	1600; 173	153; 165
<u>SOIL</u>								
<u>LAND WATER**</u>								
Well	1500						1600; 173	153; 165
Cistern							20; 19; 28	120
Lens		560					43	
<u>SEA WATER**</u>								
Ocean	49	123		118	25	119	121	45
Lagoon	126	120		121	119	120	119	120





\* Weight of material as received at HASL  
 \*\* Samples scavenged with Fe(OH)<sub>3</sub>

DOE ARCHIVES



SELF-ABSORPTION CURVES

FIGURE




Residue Weights in Milligrams

60  
62


~~CONFIDENTIAL USE ONLY~~  
NFDL MARSHALL ISLAND RESURVEY - 1956

SELF-ABSORPTION CURVES


FIGURE



6



7



8

Residue Weights in Milligrams

DOE ARCHIVE

~~CONFIDENTIAL USE ONLY~~

67  
63