

A 41

UWFL-47

HEALTH AND SAFETY

UNITED STATES ATOMIC ENERGY COMMISSION


SURVEY OF RADIOACTIVITY IN THE SEA AND  
IN PELAGIC MARINE LIFE WEST OF THE  
MARSHALL ISLANDS, SEPTEMBER 1-20, 1956

By  
Allyn H. Seymour  
Edward E. Held  
Frank G. Lowman  
John R. Donaldson  
Dorothy J. South

	US DOE ARCHIVES
	326 U.S. ATOMIC ENERGY
RG _____	COMMISSION
Collection _____	<del>SECRET</del> <i>McClaw</i>
Box _____	<i>9</i>
Folder _____	<i>SURVEY OF RADIOACTIVITY IN THE SEA + IN PELAGIC MARINE LIFE WEST OF THE MARSHALL ISLANDS, 9-1 THRU 20, 1956</i>

March 15, 1957

Applied Fisheries Laboratory  
University of Washington  
Seattle, Washington



Technical Information Service Extension Office, Tenn.

Folder	<i>1320</i>
Box	<i>9</i>
Page	<i>1320</i>
File	<i>1320</i>

INVENTORY CONTROL

UWFL-47

SURVEY OF RADIOACTIVITY IN THE SEA AND IN PELAGIC  
MARINE LIFE WEST OF THE MARSHALL ISLANDS  
SEPTEMBER 1-20, 1956

Allyn H. Seymour  
Edward E. Held  
Frank G. Lowman  
John R. Donaldson  
Dorothy J. South

Applied Fisheries Laboratory  
University of Washington  
Seattle, Washington

March 15, 1957

Operated by the University of Washington under Contract No.  
AT(45-1)540 with the United States Atomic Energy Commission

DOE ARCHIVES

## ABSTRACT

A survey of the radioactivity in the sea in the region of the North Equatorial Current from the Marshall Islands to the Marianas Islands was made in September 1956. The expedition was sponsored by the United States Atomic Energy Commission, Division of Biology and Medicine, and carried out by the Applied Fisheries Laboratory, University of Washington, with the support and cooperation of the United States Navy.

Plankton samples were taken by oblique tows from 200 meters and water samples were taken from the surface, 25, 50, 100, and 150 meters at 74 stations. The general pattern of distribution of radioactivity shows a sharp decrease east of Bikini and a gradual but irregular decrease west of Eniwetok. A slight degree of contamination is indicated as far to the west as Guam, the western extremity of the survey. Non-fission products account for a large proportion of the radioactivity in plankton and fish samples.

DOE ARCHIVES

## CONTENTS

	Page
Introduction	1
Plans, Equipment and Operations at Sea	6
Collection and Preparation of Samples	9
Methods of Analyses	10
Results of the Survey	11
Plankton	11
Water	13
Continuous Monitoring	24
Fish	26
Discussion of Survey Data	30
Results and Discussion of Radionuclide Analysis	37
Summary	49
Appendix	53

DOE ARCHIVES

SURVEY OF RADIOACTIVITY IN THE SEA AND IN PELAGIC MARINE  
LIFE WEST OF THE MARSHALL ISLANDS SEPTEMBER 1-20, 1956

INTRODUCTION

The amount of radioactivity in the sea and in pelagic marine life during and following weapons tests at the Pacific Proving Ground has been the subject of reports by United States and Japanese laboratories.

Following the 1954 test series, the Japanese survey ship, Shunkotsu-Maru, made a general survey of the amount and distribution of radioactivity in sea water and in some of the marine life in the region west of the Marshall Islands. The report of the Japanese survey<sup>1</sup> indicated that measurable amounts of radioactivity were to be found in the sea even as late as the spring of 1955.

Operation Troll<sup>2</sup> was conducted during the spring of 1955 to measure the level of radioactivity in the sea and the movement of the water mass containing the radioactivity. This was a joint operation of the New York Operations Office, U. S. Atomic Energy Commission, Scripps Institute of Oceanography,

DOE ARCHIVES

University of California, and the Applied Fisheries Laboratory, University of Washington.

The findings of Operation Troll, the 17,419 mile cruise of the Taney from February 25 to May 3, 1955, were summarized as follows:

1. Sea water and plankton samples show the existence of widespread low-level activity in the Pacific Ocean. Water activity ranged from 0-570 d/min/liter and plankton from 3-140 d/min/g wet weight.
2. There is some concentration of the activity in the main current streams, such as the North Equatorial Current. The highest activity was off the coast of Luzon, averaging 190 d/min/liter down to 600 m (April 1, 1955).
3. Analyses of fish indicate no activity approaching the maximum permissible level for foods. The highest activity in tuna fish was 3.5 d/min/g ash, less than 1 percent of the permissible level.
4. Measurements of plankton activity offer a sensitive indication of activity in the ocean.
5. Similar operations would be valuable in assessing the activity from future tests and in gathering valuable data for oceanographic studies.

The Division of Biology and Medicine of the U. S. Atomic Energy Commission requested the Applied Fisheries Laboratory of the University of Washington to conduct surveys of the open sea during 1956 to determine "(a) the levels of introduced radioactivity resulting from the tests in the water, plankton, and fish, and (b) how far the activity extends westward in the North Equatorial current."