DOCUMENT SOURCE
University of California at Berkeley
The Bancroft Library/The University Archives, Berkeley CA
RECORDS SERIES TITLE
JOHN H. LAWRENCE WRITINGS
BANCROFT/UARC ID NO.
87/56 c
CARTON NO.
Carton 1
COPY
FOLDER NAME
OBSERVATIONS OF THE NARCOTIC EFFECT OF XENON
* NOTES KRYPTON INHALATION STUDY @ DONNER
BIBLIOGRAPHY pg. 1/5
FOUND BY/DATE FOUND
K. HOLMES 10/14/94

721934

Copied from originals in The Bancroft Library for reference use only. Copies may not be deposited in other libraries or be sold without express permission. Please contact The Bancroft Library upon completion of research. Permission to reproduce must be obtained in writing from the Bancroft Library, University of California, 94720

R

[*Reprinted from the Journal of Physiology,*
1946, Vol. 105, No. 3, p. 197.]
PRINTED IN GREAT BRITAIN

J. Physiol. (1945) 105, 197–204 615.782:546.29

PRELIMINARY OBSERVATIONS ON THE NARCOTIC EFFECT OF XENON WITH A REVIEW OF VALUES FOR SOLUBILITIES OF GASES IN WATER AND OILS

BY J. H. LAWRENCE, W. F. LOOMIS, C. A. TOBIAS AND F. H. TURPIN

From the Divisions of Medical Physics and Medicine, University of California, Berkeley, California

(*Received* 3 *November* 1945)

The comparative narcotic effects of argon, nitrogen and helium have been previously studied by Behnke (1940) and by Behnke & Yarbrough (1938, 1939). Exposure up to 10 atm. of either air or an 80% argon, 20% oxygen mixture produces progressive narcotic effects in man. These effects are attributed to the dissolved inert gases, nitrogen and argon. Breathed in equivalent concentration, argon has approximately twice the narcotic effect of nitrogen and is twice as soluble as nitrogen in water and in fat. The fat-water solubility ratio of the two gases is nearly the same. Helium, on the other hand, with its lower solubility in water and fat has negligible narcotic effects when compared with argon and nitrogen.

The effect upon the central nervous system of argon, as compared with nitrogen, seems to depend primarily upon the total concentration of the dissolved gas. Reasoning from the Meyer-Overton hypothesis that anaesthetic effects are relative to the fat-water solubility ratio, it is suggested that gases such as krypton and xenon might have striking narcotic properties. These gases have a much greater water solubility and a higher fat-water solubility ratio. It is possible that 80% krypton at sea-level is as narcotic as 6 atm. of air, and that xenon under the same conditions may be equivalent to 25 atm. of air. Aside from the Meyer-Overton hypothesis, calculations of these theoretical effects are subject to considerable speculation in that the dissolved gas distribution ratio in the fat of the critical area in the central nervous system may be unlike that in olive oil. Fig. 1 shows a radio-autograph of a cross-section of a rat previously exposed to an atmosphere containing radiokrypton. After several hours' exposure the animal was killed, frozen and sectioned transversely and a section then placed in the cold, in close apposition to X-ray film. The relatively great uptake of the gas in fat containing tissue is noted by the dark areas in the radio-autograph.

1159789

DOCUMENT SOURCE
University of California at Berkeley
The Bancroft Library/The University Archives, Berkeley CA
RECORDS SERIES TITLE
JOHN H. LAWRENCE WRITINGS
BANCROFT/UARC ID NO.
87/56 c
CARTON NO.
Carton 1
COPY
FOLDER NAME
OBSERVATIONS OF THE NARCOTIC EFFECT OF XENON
NOTES KRYPTON IN HALATION
BIBLIOGRAPHY
Pg. 2/5
FOUND BY/DATE FOUND
K. HOLMES 10/14/04

A single subject in this laboratory, breathing 50% krypton and 50% oxygen in an experimental gas exchange measurement, voluntarily remarked that he had been quite 'dizzy' during the krypton inhalation. This is the report of a single subject. He had had, however, numerous experiences of breathing oxygen and nitrogen mixtures from the gasometric apparatus and, in this particular experiment, he was not aware of the fact that he was breathing some other mixture of gases.

Enough xenon was available to make a preliminary study of the effect of this inert gas upon mice. One of us (J.H.L.) exposed mice individually to a mixture of oxygen and xenon for short periods of time. The mixture was



Fig. 1. The dark areas indicate clouding of the film due to radio krypton. Fat stains of section revealed those areas to have high fat content. (1) Spinal cord. (2) Perirenal fat. (3) Mesenteric fat. (4) Radiokrypton trapped in hair.

continuously circulated through soda lime, and the oxygen content frequently checked so that at least 20% oxygen was always present and CO_2 was removed. Additional oxygen was fed in as needed. The results are presented in Table 1. In the five mice exposed to an atmosphere with high xenon content, but with adequate oxygen, definite central nervous system effects were observed, and these findings might possibly be related to the high fat/water solubility ratio for xenon.

In connexion with certain studies on decompression sickness, it became necessary to know the solubilities of various inert gases in water and fats. By means of the use of radioactive isotopes, one of us (W.F.L.) determined the values for argon, krypton and xenon. An investigation of the literature and tabulation of the solubilities of various gases has given data which are of

Copied from originals in The Bancroft Library
reference use only. Copies may not be deposited
other libraries or institutions without express

Table 1. The effects on mice of breathing mixtures of oxygen and xenon, and of oxygen and krypton

No.	Gas mixture	Exposure	Narcotic effects and remarks
1	Approximately 50% krypton, 50% oxygen	1 hr.	None
2	Approximately 60% xenon, 40% oxygen	20 min.	Within 2 min. animal exhibited slight c vulsive extensor movements of he Developed weakness of hind legs. Co plete recovery 15 min. after removal fr chamber
3	At end of experiment 78% xenon, 22% oxygen	1 hr.	Within 2 min. mouse developed convuls extensor movements of head; at 10 m hind legs seemed partially paralysed; af removal from chamber had ataxia a unsteadiness with complete recovery 15 min.
4	At end of experiment 69% xenon, 30% oxygen	1 hr.	Within 2 min. animal sluggish and c vulsive. Motions of head as above. La convulsive movements of hind legs c veloped
5	At end of experiment 60% xenon, 39% oxygen	45 min.	Within 2 min. developed convulsive mo ments of head and hind legs. On remo from chamber, ataxia for 3 min., th complete recovery
6	At end of experiment 58% xenon, 42% oxygen	30 min.	Within 1½ min. convulsive head moveme as above. This continued througho After removal ataxia for 1 min., th apparent complete recovery

interest in the fields of aviation medicine and respiratory physiology. results are presented in Figs. 2 and 3 and Table 2 and the bibliograph appended.

The values given are correct to two significant figures. Two or more re ences to a given value indicate that two or more investigators agreed (to significant figures) to this value. Great discrepancies in the literature w found, especially in the values for neon and krypton. In cases such as these, various figures were evaluated and those thought to be the most reliable w selected. The values for H_2, O_2, CO and CO_2 were taken from the *Handboo Chemistry and Physics* (1944) and the *Dictionary of Chemical Solubilities* (19 However, the *Handbook* values for the inert gases were in considerable er having been taken from an earlier work of von Antropoff (1910). Antrop earlier values have been listed in the *Handbook of Chemistry and Physics* u the 28th edition (1944), in the *Dictionary of Chemical Solubilities* (1921) in the *International Critical Tables* (1928). Recent authors (Lannung, 19 Valentiner, 1930; Van Liempt & Van Wijk, 1937), including von Antro himself (1919), have shown these values to be in error, especially those neon and krypton. In the case of neon there seem to be only two recent sour of data—that of Lannung (1930) and that of Valentiner (1930). These investigators differed considerably in their results, and it is difficult to de which is the more reliable.

DOCUMENT SOURCE
University of California at Berkeley
The Bancroft Library/The University Archives, Berkeley CA
RECORDS SERIES TITLE
JOHN H. LAWRENCE WRITINGS
BANCROFT/UARC ID NO.
87/86 c
CARTON NO.
Carton 1
COPY
FOLDER NAME
OBSERVATIONS OF THE NARCOTIC EFFECT OF XENON
NOTES KRYPTON INHALATION
BIBLIOGRAPHY pg. 3/5
FOUND BY/DATE FOUND K. HOLMES 10/14/94



Fig. 3. Solubility of inert gases in water. ● von Antropoff (1919). △ *Dictionary of Chem. So* (von Antropoff, 1910) (Argon, Comey). □ Valentiner (1927, 1930). × Lannung (1930 ○ Estreicher (1899). + Cady, Elsey & Berger (1922). ◑ Wiebe & Gaddy (1935). ⊙ Meyr (1913).



Fig. 2. Solubility of gases in water, α = c.c. gas (0° C., 760 mm.) dissolved in 1 c.c. water at temperature of observation.

TABLE 2. Solubilities of gases in water and oils

Gas	Mol. wt.	Half-life of isotope used	Bunsen's absorption coefficient (to 2 significant figures) Solvent, water 22° C.	Solvent, water 37° C.	Solvent, oil 22° C.	Solvent, oil 37° C.	Oil/water solubility ratio 22° C.	Oil/water solubility ratio 37° C.
Chemically inert gases: Helium	4	0·8 sec.	0·0086 (5, 1, 34)	0·0085 (13)	0·015 (3) olive	0·015 (3) olive		1·7 (4) olive
Neon	20		0·0104 (15)	0·0097 (15)				
Nitrogen	28	9·93 min.	0·015 (29)	0·013 (3)	0·052 (30) olive 0·053 (31) corn	0·07 (20) 32° paraffin 0·067 (3, 31) olive 0·063 (31) corn	3·5 olive 4·2 corn	5·2 (4) olive 4·7 (31) corn
Argon	40	110 min.	0·037 (8)	0·026 (4, 15)	0·15* (20) olive	0·14* olive	4·0 olive	5·3 (4) olive
Krypton	83·7	34 hr.	0·059 (28)	0·045 (33, 27)	0·44* olive	0·43* olive	7·5	9·6
Xenon	131·3	34 days	0·13 (33, 27)	0·085 (33)	1·9* olive	1·7* olive	14·5	20·0
Radon	222	3·825 days	0·23 (26, 19)	0·15 (26, 19)	0·25 (16) olive	0·19 (16) olive	110	125
Hydrogen	2	—	0·018 (11)	0·016 (11)	0·042 (31) cottonseed	0·04 (20) 32° paraffin 0·05 (24) 40° cottonseed	2·3	3·1 cottonseed 2·5 paraffin
Oxygen	32	—	0·030 (11)	0·024 (11)	0·11 (31) corn 0·11 (31) cottonseed	0·12 (24) 40° cottonseed	3·7	5·0
CO_2	44	21·5 min.	0·83 (11)	0·56 (11)	1·34 (31) cottonseed	0·876 (24) 40° cottonseed	1·6	1·6
CO	28	21·5 min.	0·022 (11)	0·018 (11)				

The numbers in brackets refer to the references on pp. 203–4.

* Determined in this laboratory by W. F. Loomis, using radioactive isotope.

SUMMARY

Observations on the narcotic effects of xenon are presented. The oil and water solubilities of hydrogen, oxygen, carbon dioxide, nitrogen, helium, neon, argon, krypton, xenon and radon are presented from a review of the literature and from determinations using the radioactive gases.

The work described in this paper was done under a contract recommended by the Committee on Medical Research between the Office of Scientific Research and Development and the University of California.

The xenon and krypton used in these experiments were generously provided by the Linde Air Products Company through the interest of Mr L. A. Bliss.

We wish to thank our colleague, Dr H. B. Jones, for valuable discussions and assistance.

REFERENCES

(1) Akerlof, G. (1935). The solubilities of noble gases in aqueous salt solutions at 25° C. *J. Amer. chem. Soc.* **57**, 1196.

(2) Behnke, A. R. (1940). *Some Physiological Considerations of Inhalation. Anaesthesia and Helium. Anaesthesia and Analgesia.*

(3) Behnke, A. R. & Yarbrough, O. D. (1938). Physiologic studies of helium. *U.S. nav. med. Bull.* **36**, 542.

(4) Behnke, A. R. & Yarbrough, O. D. (1939). Respiratory resistance, oil water solubility and mental effects of argon compared with helium and nitrogen. *Amer. J. Physiol.* **126**, 409.

(5) Cady, H. P., Elsey, H. M. & Berger, E. V. (1922). The solubility of helium in water. *J. Amer. chem. Soc.* **44**, 1456.

(6) Coste, J. H. (1927). Review of nitrogen solubilities. *J. Phys. Chem.* **31**, 81.

(7) *Dictionary of Chemical Solubilities* (1921). 2nd edition. New York: Macmillan Co.

(8) Estreicher, T. (1899). Die Löslichkeitverhältnisse von Argon und Helium im Wasser. *Z. phys. Chem.* **31**, 176.

(9) Giffcken, G. (1904). Beiträge zur Kenntnis der Löslichkeitbeeinflussung. *Z. phys. Chem.* **49**, 257.

(10) Grollman, A. (1929). The solubility of gases in blood and blood fluids. *J. biol. Chem.* **82**, 317.

(11) *Handbook of Chemistry and Physics.* (1941), 25th edition; (1944), 28th edition. Cleveland: Chemical Rubber Publishing Co.

(12) Harned, H. S. & Davis, R. J. (1943). The ionization constant of carbonic acid in water and the solubility of carbon dioxide in water and aqueous salt solutions from 0° to 50° C. *J. Amer. chem. Soc.* **65**, 2030.

(13) Hawkins, J. A. & Shilling, C. W. (1936). Helium solubility in blood at increased pressure. *J. biol. Chem.* **113**, 649.

(14) *International Critical Tables* (1928). 1st edition, 3, 257. McGraw-Hill.

(15) Lannung, A. (1930). The solubilities of helium, neon and argon in water and some organic solvents. *J. Amer. chem. Soc.* **52**, 68.

(16) Lurie (1910). Thesis, Grenoble (from *Int. Crit. Tables* (1928). 1st edition, 3, 257. McGraw-Hill).

(17) Markham, A. E. & Kobe, K. A. (1941). Solubility of carbon dioxide and nitrous oxide in aqueous salt solutions. *J. Amer. chem. Soc.* **63**, 449.

(18) Markham, A. E. & Kobe, K. A. (1941). The solubility of liquids in gases. *Chem. Rev.* **28**, 51 (346 references listed and valuated).

1159192

Copyright 1945 The Bancroft Library for reference use only. Copies may not be deposited in other libraries or institutions without express permission. Please return all copies to The Bancroft Library upon completion of use. Permission to reproduce in full or in part must be obtained in writing from the Director, The Bancroft Library, University of California, Berkeley, California 94720.

DOCUMENT SOURCE University of California at Berkeley
RECORDS SERIES TITLE The Bancroft Library/The University Archives, Berkeley CA
BANCROFTLIBRARY ID NO. JOHN H. LAWRENCE WRITINGS
CARTON NO. B7/Box 6
Carton 1
COPY
FOLDER NAME OBSERVATIONS OF THE NARCOTIC EFFECT OF XENON
NOTES KRYPTON IN HALATION
BIBLIOGRAPHY
Pg. 4/5
FOUND BY/DATE FOUND K. HOLMES 10/14/94

DOCUMENT SOURCE
University of California at Berkeley
The Bancroft Library/The University Archives, Berkeley CA
RECORDS SERIES TITLE
JOHN H. LAWRENCE WRITINGS
BANCROFT/UARC ID NO.
87/86 c
COPY
CARTON NO.
Carton 1
FOLDER NAME
OBSERVATIONS OF THE NARCOTIC EFFECT OF XENON
NOTES KRYPTON INHALATION
BIBLIOGRAPHY pg. 5/5
FOUND BY/DATE FOUND K. HOLMES 10/14/94

Copied from originals in The Bancroft Library for reference use only. Copies may not be deposited in other libraries or institutions without express permission. Please return all copies to the Bancroft Library upon completion of your research. Permission to reproduce or distribute the material contained in this Bancroft Library, University of California, Berkeley, CA 94720.

(19) Meyer, S. (1913). Bemerkung über die Löslichkeit von Radiumemanation und anderen Gasen in Flüssigkeiten. *S.B. Akad. Wiss. Wien.* Abt. IIa, 122, 1281.
(20) Nasini, A. G. & Corinaldi, C. G. (1932). Solubilida della Argo, dell Azoto e dell' idrigeno in Olio de Paraffina. *Atti Soc. Ital. Prog. Sci.* 20, 2264.
(21) Nieloux, M. (1927). Dissolution Aqueuse des Gaz. *C.R. Soc. Biol., Paris*, 97, 1553.
(22) Rakestraw, N. W. & Emmel, V. M. (1938). The solubility of nitrogen and argon in sea water. *J. Phys. Chem.* 42, 1211.
(23) Randall, M. & Failey, C. F. (1927). The activity coefficients of gases in aqueous salt solutions. *Chem. Rev.* 4, 271.
(24) Schaffer, P. S. & Haller, H. S. (1943). The solubility of gases in butter oil, cottonseed oil and lard. *Oil and Soap*, 20, 161.
(25) Seidell, A. (1919). *Solubilities of Inorganic and Organic Substances*, 227 pp. New York: D. Van Nostrand Co.
(26) Valentiner, S. (1927). Über die Löslichkeit der Edelgase in Wasser. *Z. Phys.* 42, 1927.
(27) Valentiner, S. (1930). Bemerkung zu meiner Arbeit: Über die Löslichkeit der Edelgase in Wasser. *Z. Phys.* 61, 563.
(28) Van Liempt, J. A. M. & Van Wijk, W. (1937). Die Löslichkeit von Krypton in verschiedenen Flüssigkeiten. *Rec. Trav. chim. Pays-Bas*, 56, 632.
(29) Van Slyke, D. D., Dillon, R. T. & Margaria, R. (1934). Nitrogen solubility in blood. *J. biol. Chem.* 105, 571.
(30) Vernon, H. M. (1907). Solubility of air in fats and its relation to caisson disease. *Proc. Roy. Soc.* B, 79, 366.
(31) Vibrans, F. C. (1935). Solubility of gases in edible fats and oils. *Oil and Soap*, 12, 14.
(32) von Antropoff, A. (1910). The solubility of xenon, krypton, argon, neon and helium in water. *Proc. Roy. Soc.* A, 83, 480.
(33) von Antropoff, A. (1919). Experimentelle Untersuchung über die Löslichkeit der Edelgase in Flüssigkeiten. *Z. Electro-chem.* 25, 269.
(34) Wiebe, R. & Gaddy, V. L. (1935). The solubility of helium in water at 0°, 25°, 50° and 75° C. and at pressures up to 1000 atmospheres. *J. Amer. chem. Soc.* 57, 847.
(35) Winkler, L. W. (1906). Gesetzmässigkeit bei der Absorbtion der Gase in Flüssigkeiten. *Z. phys. Chem.* 55, 344.

1159793