


407823

**CLASSIFIED**



## Nickel-63 in Marine and Terrestrial Biota, Soil, and Sediment

Thomas M. Beasley  
Edward E. Held

Collection	1320
Date	9
Folder	47

Job	1320
Box	9
Folder	47
HISTORICAL OFFICE INVENTORY CONTROL	

**CLASSIFIED**

## Nickel-63 in Marine and Terrestrial Biota, Soil, and Sediment

**Abstract.** A previously unreported radionuclide, nickel-63 (half-life, 92 years), produced in the testing of nuclear devices, was measured in biological and environmental samples from areas of the Pacific Ocean and the eastern seaboard of the United States. The concentrations of nickel-63 are low (maximum of 163 disintegrations per minute per gram of dry weight), but this radionuclide may be a useful tracer of oceanic processes because of its long half-life and long residence time in the ocean.

A review of the concentrations of radionuclides measured in biota of the Pacific Proving Ground during testing of thermonuclear devices indicated that the production of measurable quantities of the radionuclide  $^{63}\text{Ni}$  was probable. The large amounts of  $^{55}\text{Fe}$ ,  $^{65}\text{Zn}$ ,  $^{57}\text{Co}$ ,  $^{58}\text{Co}$ , and  $^{60}\text{Co}$  reported by Lowman (1) resulted largely from nuclear interactions in the structural material used in and around the device. We concluded

that stable nickel in these materials would yield  $^{63}\text{Ni}$  through the  $(n,\gamma)$  activation process. The cross section for this reaction is large (15 barns).

Detection of  $^{63}\text{Ni}$  by other than specific radiochemical separation and liquid scintillation counting is doubtful. The radionuclide decays by the emission of a beta particle (100 percent) whose maximum energy is only 67 kev (2), an energy intermediate between

those of  $^3\text{H}$  and  $^{14}\text{C}$ . Compared to other radionuclides produced in weapons testing, the radiological hazard from  $^{63}\text{Ni}$  is much less and therefore was of little concern in the evaluation of the effects of radioactivity on biota. The specificity of the required analysis and the lesser hazard probably account for the fact that measurements of the concentrations of  $^{63}\text{Ni}$  in the environment have not been made until now. However, the use of radioisotopes, both artificial and natural, as tracers of oceanic currents and water mixing (3) prompted us to look for  $^{63}\text{Ni}$  since (i) the residence time of nickel in the ocean is  $1.8 \times 10^4$  years; (ii) nickel is a constituent of ferromanganese minerals, comprising some 0.4 percent (by weight) (4); and (iii)  $^{63}\text{Ni}$  has a radioactive half-life of 92 years and is thus a more useful long-term tracer than other radionuclides of the transition elements which have been found in the oceans but which have half-lives of a few years or less.

We separated nickel from biological and environmental samples and other elements by precipitation and solvent extraction. Before dissolution of the samples in concentrated nitric and perchloric acids, 6 mg of stable nickel were added to each sample to serve as both a carrier and a yield determinant. Nickel was concentrated from the digest by precipitation of nickel 1,2-cycloheptanedione dioxime from acid solution (5). The precipitate was collected by filtration, removed from the filter by dissolution in 8M  $\text{HNO}_3$ , and oxidized with perchloric acid to destroy the organic matter; the nickel was concentrated again by a second precipitation with heptoxime. After dissolution and oxidation of the second precipitate,  $\text{Ni}(\text{OH})_2$  was precipitated by the addition of  $\text{NaOH}$ . The precipitate was dissolved in 6N  $\text{HCl}$ , and the resultant solution was extracted with 10 ml of a solution of 10 percent Alamine-336 in xylene. The phases were separated by centrifugation, the organic layer was decanted, and the nickel was again precipitated as the hydroxide. The hydroxide precipitate was prepared for liquid scintillation counting (Packard Instrument liquid scintillation spectrometer model 3375) by dissolution in 0.5

Table 1. Concentration of nickel-63 in environmental samples. Errors given are the 95 percent confidence levels ( $2\sigma$ ) of the count rate measurements.

Location	Sample	Date of collection	Concentration of nickel-63 (disintegrations per minute per gram of dry weight)
<i>Eniwetok Atoll</i>			
Belle Islet	Soil	May 1954	8.2 ± 0.2
Belle Islet	Clam kidney*	August 1964	158.0 ± 2.6
Belle Islet	Clam kidney*	August 1964	67.8 ± 1.6
Elugelab Islet	Crater sediment	August 1964	9.6 ± 0.2
Engebi Islet	Soil	August 1958	7.5 ± .2
Aaraanbiru Islet	Clam kidney†	September 1958	41.2 ± .6
Engebi Islet	Clam kidney‡	August 1964	11.4 ± .6
<i>Bikini Atoll</i>			
Bokonejien Islet	Crater sediment	May 1967	80.0 ± 1.0
Namu Islet	Soil	August 1964	9.8 ± 0.2
Bokororyuru Islet	Clam kidney‡	August 1964	163.0 ± 3.5
<i>Rongelap Atoll</i>			
Kabelle Islet	Soil (0-0.6 cm)	September 1961	3.1 ± 0.2
Kabelle Islet	Soil (0.6-1.3 cm)	September 1961	0.5 ± .06
<i>Christmas Island</i>			
	Clam kidney‡	April 1962	.91 ± .14
<i>Penrhyn Atoll</i>			
	Clam kidney*	April 1962	.42 ± .14
<i>Northeast Pacific Ocean</i>			
	Chaetognaths§	February 1964	.93 ± .80
	Chaetognaths	February 1964	2.3 ± 1.0
	Chaetognaths	April 1964	4.0 ± 3.0
	Chaetognaths	June 1964	4.5 ± 2.4
44°38'N, 125°20'W	Squid	October 1965	0.19 ± 0.08
44°38'N, 125°20'W	Squid	August 1966	.13 ± .08
44°38'N, 125°20'W	Squid	August 1966	.13 ± .08
47°39'N, 173°05'W	Squid	August 1966	.13 ± .08
50°30'N, 167°00'W	Squid	August 1966	.13 ± .08
30°N, 140°W	Squid¶	July 1965	.38 ± .24
<i>Aleutian Islands</i>			
Amchitka Island	Lichen#	October 1965	.18 ± .04
Amchitka Island	Lichen	October 1965	.35 ± .04
<i>Eastern seaboard</i>			
	Composite shellfish	August 1963	.02 ± .002

\* *Tridacna crocea*. † *Tridacna* sp. ‡ *Tridacna gigas*. § *Sagitta elegans*. || *Onychoteuthis* sp. ¶ *Stenoteuthis bartrami*. # Principally *Cladonia pacificia*.

benzene (POPOP) per liter of toluene. We measured the yield by wet-ashing the liquid scintillation solution with  $\text{HNO}_3$  and  $\text{HClO}_4$ , reprecipitating the nickel as the heptoxime complex, and weighing the dried product. Decontamination factors of  $> 10^4$  were observed for the radioisotopes  $^{90}\text{Sr}$ - $^{90}\text{Y}$ ,  $^{55}\text{Fe}$ ,  $^{137}\text{Cs}$ ,  $^{207}\text{Bi}$ ,  $^{106}\text{Ru}$ - $^{106}\text{Rh}$ ,  $^{147}\text{Pm}$ ,  $^{60}\text{Co}$ ,  $^{125}\text{Sb}$ ,  $^{144}\text{Ce}$ ,  $^{95}\text{Zr}$ - $^{95}\text{Nb}$ ,  $^{110\text{m}}\text{Ag}$ ,  $^{226}\text{Ra}$  plus daughters, and  $^{65}\text{Zn}$ . The clean separations obtained suggest that the activity in the low-activity samples is indeed  $^{63}\text{Ni}$ .

Figure 1 shows the relative activity plotted against lower level settings derived from the liquid scintillation counting of a  $^{63}\text{Ni}$  standard and a  $^{59}\text{Ni}$  spike, along with the spectrum obtained from the analysis of a clam kidney that was collected at Bikini Atoll. Clearly, the activity separated from this tissue was  $^{63}\text{Ni}$ . The concentrations of  $^{63}\text{Ni}$  in these samples are listed in Table 1.

Generally, the liquid scintillation counter was set to record 10,000 sample counts or to count for 500 minutes. Background was counted for 500 minutes, and the average rate for the  $^{63}\text{Ni}$  settings was  $15.8 \pm 0.4$  count  $\text{min}^{-1}$ . The detection limit at this background counting rate and at an average overall efficiency of 59 percent for  $^{63}\text{Ni}$  detection was  $1.4 \pm 1.0$  disintegrations per minute per sample at the 95 percent confidence level (6). The values listed in Table 1 are adjusted to unit dry weight and therefore are numerically smaller than the detection limit. However, sample sizes exceeded 10 g, except for the chaetognaths, and the values for all samples were above the calculated detection limit.

The  $^{63}\text{Ni}$  concentrations in marine species ranged from a low of approximately 0.02 disintegration per minute per gram of dry weight for shellfish taken on the eastern seaboard of the

of stable nickel in chaetognaths, which prompted us to analyze this organism for  $^{63}\text{Ni}$ . The concentrations were very low, approaching the limit of detection. This suggests a low concentration of  $^{63}\text{Ni}$  in surface waters in this area and indicates the absence of appreciable amounts of  $^{63}\text{Ni}$  in Columbia River water, which is used upstream as a coolant for the nuclear reactors at the Hanford complex. This observation is consistent with Kirby's (8) radiochemical procedure for the isolation of  $^{65}\text{Ni}$  from Columbia River water (8 to 12 liters) collected near the reactors. This procedure makes possible the measurement of a concentration of  $10^{-8}$   $\mu\text{c}$  of  $^{65}\text{Ni}$  per milliliter. Because of its low specific activity, the activity of  $^{63}\text{Ni}$  is




Fig. 1. Relative activity and discriminator settings for  $^{60}\text{Ni}$  ( $\Delta$ ) and  $^{63}\text{Ni}$  ( $\circ$ ) spike solutions, plus the activity isolated from *Tridacna gigas* clam kidney collected from Bikini Atoll ( $\square$ ). Instrument settings were 20 percent gain and 4 percent window.

1965. The average concentration of  $^{63}\text{Ni}$  for lichens from the Aleutians is lower by only a factor of 2 than the  $^{60}\text{Co}$  concentrations of lichens reported by Hanson *et al.* (10) for Anaktuvak Pass, Alaska; however, the comparison of results from insular species in the Aleutians to those of similar continental species may be invalid.

Some specimens contained sufficient stable nickel to permit comparison of the specific activities of  $^{63}\text{Ni}$ , an indication of the variations that occur between similar organisms. Table 2 shows the results of the measurements of  $^{63}\text{Ni}$  specific activity in clam kidneys collected at test sites and from islands some distance away.

The high specific activity in the clam kidney from Bikini Atoll is due, in part, to the fact that this specimen was taken at Bokororyuru Islet on the western edge of the atoll. Wind-driven surface currents move the lagoon waters toward the southwest and, as a consequence, Bokororyuru Islet receives waterborne radionuclides in addition to local fallout at the time of testing. The presence of  $^{63}\text{Ni}$  in association with short-lived  $^{57}\text{Co}$  (11) in the specimens taken from Christmas Island and Penrhyn Atoll in April 1962, prior to the United States tests of that year, suggests that the source of  $^{63}\text{Ni}$  in these specimens was the Soviet tests of 1961. The evidence is not conclusive, however, since the age of the clams was not determined and it is probable that they were accumulating  $^{63}\text{Ni}$  over a period of time. Bikini and Eniwetok atolls lie in the path of the North Equatorial Current which moves water toward the Asian continent.

Von Arx (12) has described the circulation and refreshment times of both Bikini and Rongelap atolls and has found that the exchange of lagoon water with seawater resulted in winter

the possibility of the production of  $^{59}\text{Ni}$  (half-life, 80,000 years) both by  $(n,\gamma)$  interactions with stable nickel,  $(p,n)$  reactions with stable cobalt, and  $(p,\alpha)$  reactions with  $^{58}\text{Fe}$  (92 percent abundance). Based on abundances, cross sections, and decay constants, we calculated an activity ratio of  $^{63}\text{Ni}$  to  $^{59}\text{Ni}$  of approximately 600 for the  $(n,\gamma)$  production of  $^{59}\text{Ni}$ . Nickel-59 decays by electron capture and is determined by measuring the 6.9-keV x-ray which results from the de-excitation of its daughter,  $^{59}\text{Co}$ . A 3000-minute count

such an inventory and of the rate at which it is injected into the North Equatorial Current would help one to determine its usefulness as a downstream tracer for these waters. However, the giant clam *Tridacna* sp. appears to be an excellent indicator organism, which could be used to delineate the downstream penetration of  $^{63}\text{Ni}$ .

THOMAS M. BEASLEY  
EDWARD E. HELD

Laboratory of Radiation Ecology,  
University of Washington, Seattle

10. W. C. Hanson, D. G. Watson, R. W. Perkins, in *Radioecological Concentration Processes*, B. Aberg and F. B. Hungate, Eds. (Pergamon Press, Oxford, 1967), pp. 233-245.
11. The concentrations of  $^{57}\text{Co}$  were measured at the time of collection and are tabulated in this laboratory.
12. W. S. von Arx, *Trans. Amer. Geophys. Union* **29**, 861 (1948).
13. J. A. Knaus, in *The Sea*, M. N. Hill, Ed. (Interscience, New York, 1963), vol. 2, pp. 235-252.
14. We thank Dr. W. Percy of Oregon State University for supplying samples of chaetognaths; Dr. T. R. Folsom of Scripps Institute of Oceanography and C. H. Fiscus of the Marine Biological Laboratory, Bureau of Commercial Fisheries, for supplying squid specimens. Supported by AEC under contract No. AT(26-1)-269.

3 March 1969

DOE ARCHIVES

UNCLASSIFIED