

one on Nedren (Harry) Island, and one on Amwetak Island. Japtan (David) will be a temporary location from the beginning of the program. The workers from Ujilang will be housed there. They will wait on Japtan, and will live there until the rest of the atoll is ready for occupancy."

BEST COPY AVAILABLE

DOE ARCHIVES

Nervik

-2-

The Magistrate, and Scribe John Abraham also made it clear that while the village sites would be as indicated, all of the islands (and reefs) of the atoll would be used and exploited after the people have returned to Enewetak.

They also stated that the decision of settlement pattern had been reached by the community on Ujilang following a series of lengthy meetings.

I have sent several memoranda to you containing additional pertinent data, mostly regarding dietary patterns. Have you received them?

I ask that you include all additional data in my revised report. Also I note that I have repeated the sentence " It seems unlikely that significant contacts occurred before the nineteenth century." Please remove this sentence from the next to last paragraph on page 1 of the report. It is redundant.

Sincerely,

Jack A. Tobin

cc District Administrator, Marshalls
Community Development Officer, Marshalls
District Director Public Affairs, Marshalls
Dr. Helfrich
Dr. Barr ✓
File

Attach: As stated

DOE ARCHIVES

(3) Freezing, boiling, or processing does not destroy the poison.

(4) The species most likely to be poisonous are:

- RED SNAPPER
- BARRACUDA
- GROUPEERS
- BLACK ULUA (OVER 40 LBS)
- MORAY EELS " " "

Other species may be poisonous but usually not. OPEN OCEAN FISH SUCH AS TUNA, BONITO, WAHOO, MAHI MAHI, AND BILLETISH ARE NEVER AFFECTED BY CIGUATERA.

As a rule of the thumb the silvery-blue reef fish are not affected by ciguatera.

1

.....
Underlining, my own. Note: I obtained the original of this report on Enewetak in October, 1972. J.A.T.

DUB AGENT 124