


Jim Miller

R

407974



UWFL-92

**ATOLL SOIL TYPES IN RELATION TO
THE DISTRIBUTION OF FALLOUT RADIONUCLIDES**

August 1965

Laboratory of Radiation Biology
University of Washington
Seattle, Washington

UNITED STATES ATOMIC ENERGY COMMISSION • DIVISION OF TECHNICAL INFORMATION

DOE ARCHIVES

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

This report has been reproduced directly from the best available copy.

Printed in USA. Price \$3.00. Available from the Clearinghouse for Federal Scientific and Technical Information, National Bureau of Standards, U. S. Department of Commerce, Springfield, Virginia 22151.

ATOLL SOIL TYPES IN RELATION TO THE
DISTRIBUTION OF FALLOUT RADIONUCLIDES

E.E. Held, S.P. Gessel and R.B. Walker

Laboratory of Radiation Biology
University of Washington
Seattle, Washington

Lauren R. Donaldson
Director

August 1965

Operated by the University of Washington under Contract No.
AT(45-1)1385 with the United States Atomic Energy Commission.

cal mixing by animals in old soils and by erosion in young soils

FOOTNOTES

is important in the redistribution of radionuclides near the soil surface. Cs^{137} and Sr^{90} are the principal radionuclides entering a cycle within the soil-plant system. Any loss from this system appears to be small, but a definite conclusion can not be drawn from the data.

TABLE OF CONTENTS

	Page
INTRODUCTION	1
METHODS OF COLLECTION AND MEASUREMENT	4
RESULTS AND DISCUSSION	7
Depth Gradient of Gamma Spectra	12
Old Soil	12
Litter and Guano	14
Young Soil	18
Depth Gradient of Sr ⁹⁰	21
RADIOAUTOGRAPHS	25
Young Soil	25
Undisturbed	25
Eroded	25
Old Soil	28
Horizontal Movement	28
Buried Organic Layers	30
Roots	30
Pumice	31
CONCLUSIONS	33
SUMMARY	35
REFERENCES	37

Nongelap soil, Nongelap gravelly sand, collected
in September 1959.

24

1959


x-ray film approximately one cm below the soil surface <u>in situ</u> .	23
12. Radioautographs and photographs of soil core sections.	26
13. An area of young soil subject to erosion.	27
14. Photograph and radioautograph of old soil showing correspondence of radioactive portions to organic matter.	29

gent land area is about three square miles, consisting of

sixty-one small islets ranging in size from a fraction of an acre to the largest, Rongelap, which is four miles long and one-half mile across at its widest point. There is one small islet on the western reef and the other islets extend along the northern, eastern and southern reefs. The islets on the northern reef are not as well developed as those to the east and south. There are two seasons--a dry season from December to March and a wet season from April to November. Annual rainfall is less than fifty inches, and there is no well-developed fresh-water lens. Some important features of Rongelap Atoll including aerial photographs are given by Wiens (1962).

Classification and mapping of the soil types at Rongelap Atoll were reported by Kenady (1962). The parent material is primarily calcium carbonate, originating from corals, foraminifera, coralline algae and mollusk shells. There is a very small amount of pumice drift in the soils. Since these soils contain no inorganic colloids, exchange capacity and organic content are linearly related. In some areas, particularly along the seaward sides of the islets, buried A_1 horizons are found as deep as eighty inches (Fig. 1). These highly organic horizons presumably result from storm debris covering previously established soil and vegetation.

The pH, determined in the field from a 1:1 soil-water ratio with a Beckman Model N-2 pH meter, is generally between



51

56



Soil pit 35, Ronelap Island. Dark layers are buried horizons.

Fig. 1. Beach Ridge Sand, Rongelap Island. This soil type occurs on the island beach ridges. The pit shown above is seven feet deep and contains six buried horizons.

7 and 8 in the surface horizon but occasional values from 4.6 to 9.5 were found. The pH increases with depth and decreasing organic content.

The amounts of exchangeable cations in the different soil types are given in Table I. The calcium content is so high that more calcium is brought into solution with repeated extractions. Consequently, strontium units have little or no significance relative to atoll soils and, when given, are based on total calcium rather than on exchangeable calcium.

METHODS OF COLLECTION AND MEASUREMENT

The vertical distribution of radioactivity in soil and litter was studied by analyzing samples taken mostly by 1-inch soil increments, in a few cases by 1/4- and 1/8-inch increments and by radioautographs of sections of soil cores prepared by the method of Held et al. (1965). The increment collections were made during both the wet and dry seasons in 1958, 1959, 1961 and 1963. The cores were collected only in 1963. Because there is considerable horizontal variation in the levels of radioactivity (Table II) each set of increments was collected to insure the sampling of a single vertical column.

The large amount of horizontal variability also made it more profitable to compare the relative amounts of radionuclides at different depths from many profiles than to make precise

Table I. Nutrient levels and soil reaction in the surface A₁ horizon of the five most extensive soil series (from Kenady 1962).

Soil Series	Rongelap Gravelly Sand	Gogan Gravelly Sandy Loam	Lomuillal Sand	Beach Ridge Sand	Kabelle Sand
Per Cent Nitrogen	0.57	1.71	0.26	0.09	0.14
Per Cent Organic Matter	16.7	35.6	6.4	4.5	7.7
*Exchange Capacity	22.2	37.7	12.6	3.7	5.7
*Sodium	3.36	4.01	2.68	1.16	1.52
*Magnesium	4.19	11.06	3.21	2.55	1.92
*Potassium	1.95	1.80	0.79	0.37	0.46
Phosphorus (ppm)	81.7	985.2	54.2	32.8	32.1
pH	8.1	7.8	8.4	8.6	8.6

* meq per 100 grams of oven dry soil

Table II. ⁹⁰Sr levels in soil samples from two sets of plots. Samples collected in September 1961. Values in pCi/gm dry. Each plot is 16-1/2 by 33 feet.


Kabelle Islet, Pit 38 Area															
Depth in inches	Plot Sample	A		B		C		D		E		F		G	
		1	2	1	2	1	2	1	2	1	2	1	2	1	2
0-1	Subsample	100	110	94	81	220	280	530	82	40	69	71	100	76	62
	" "	78	100	95	94	240	200	410	130	49	81	49	100	91	73
1-2	" "	33	62	180	60	150	41	45	61	22	54	30	26	13	29
	" "	25	51	180	91	160	43	49	78	24	52	29	22	21	22
2-3	" "	44	50	33	50	65	29	41	49	8	36	19	14	8	17
	" "	36	52	25	69	79	23	41	51	11	32	25	17	7	21

Rongelap Islet, Pit 22 Area												
Depth in inches	Plot Sample	A		B		C		D		Pit 22		
		1	2	1	2	1	2	1	2	1	2	
0-1	Subsample	230	230	130	110	180	170	160	180	130	140	
	" "	230	150	150	170	190	180	150	350	150	200	
1-2	" "	150	190	84	47	100	53	54	140	77	100	
	" "	160	180	89	47	95	61	63	140	79	97	
2-3	" "	63	49	19	6	28	13	25	65	25	43	
	" "	55	50	18	4	30	12	21	65	19	36	

The young soil has little organic material except that in an algal surface crust about 1 cm thick. Hermit crabs, Coenobita perlatus, at the base of a few shrubs, the borrowing ghost crab, Ocypode ceratophthalmia, and ants are the main animals present. Litter accumulation is found only at the bases of the scattered shrubs and is a minor part of this soil system.



Soil pit 6. Black areas are crusts formed by algae.



Top view surface

1 inch

6

17

23

26

RIGHT: Micromonolith of pit 6.



General view of pit 6 area.

Fig. 2. Kabelle Sand. A very young soil composed of fine lagoon sands deposited on the lagoon side of Kabelle Island.



Stand of *Pisonia grandis*, Kabelle Island, pit 38 area.

Fig. 3. Gogan series, Kabelle Island. This is a well-developed, productive soil, usually associated with *Pisonia grandis* and *Cordia subcordata*.



Messerschmidia argenticum at soil pit 12.

Fig. 4. Lomuilaal Sand, Kabelle Island.

Table III. Composition of representative soil types from Rongelap Atoll (after Kenady 1962).

Kabelle Sand, a young soil			
Soil Depth - inches	0-1	1-11	11-38
Per Cent Material > 2 mm	34	8	2
Per Cent Nitrogen	0.22	0.02	0.01
Per Cent Organic Matter	8.1	--	2.6
* Exchange Capacity	6.3	0.3	0.1
* Sodium	1.57	3.01	1.37
* Magnesium	1.37	1.04	1.16
* Calcium	3.01	2.88	2.65
* Potassium	0.57	0.15	0.20
Phosphorus (ppm)	30.0	12.0	12.0
pH	8.9	9.1	9.2

Lomuilal Sand, an old soil					
Soil Depth - inches	0-3	3-10	10-12	12-21	21-48
Per Cent Material > 2 mm	0	5	18	12	2
Per Cent Nitrogen	0.29	0.07	0.08	0.04	0.02
Per Cent Organic Matter	2.8	2.3	2.2	1.9	1.7
* Exchange Capacity	14.2	2.3	2.6	1.1	0.6
* Sodium	2.73	0.73	0.82	0.82	0.84
* Magnesium	4.66	8.36	1.05	0.85	0.83
* Calcium	5.02	1.87	2.50	2.31	2.35
* Potassium	1.09	0.18	0.19	0.16	0.16
Phosphorus (ppm)	105.9	15.1	14.1	5.0	5.0
pH	8.4	8.6	8.3	8.8	9.1

Gogan Gravelly Sandy Loam, an old soil						
Soil Depth - inches	1-0	0-1	1-5	5-12	12-20	20-26
Per Cent Material > 2 mm	10	20	20	27	39	56
Per Cent Nitrogen	1.54	1.96	0.42	0.18	0.07	0.05
Per Cent Organic Matter	21.4	--	5.9	6.8	2.6	2.6
* Exchange Capacity	20.5	43.6	17.9	7.2	2.6	1.7
* Sodium	2.0	3.0	0.8	0.4	0.4	0.4
* Magnesium	7.0	7.4	4.0	2.2	1.2	1.1
* Calcium	10.3	24.2	14.1	6.2	7.7	7.8
* Potassium	--	1.80	--	--	--	--
Phosphorus (ppm)	1330.4	892.8	415.5	216.3	150.6	25.1
pH	7.4	7.1	7.9	8.2	8.6	8.8

* meq per 100 gm dry soil

redistribution of radionuclides in the system.

Sea birds nest in the vegetation and land crabs, predominantly Birgus latro and Coenobita perlatus, burrow in the soil. Earthworms are seldom found and terrestrial isopods, although found in old soils on some islets, are few. There are also few soil insects, which are mainly tenebrionid and carabid beetles as well as ants.

Depth Gradient of Gamma Spectra

Old Soil

The gamma spectra, with background subtracted, of the 0 to 1-inch, 1 to 2-inch, and 9 to 10-inch depth increments from old soil collected in March 1959 are given in Fig. 5. The spectra of the increments between 2 inches and 9 inches have been omitted from the figure for clarity but show a gradual change

from the condition at 1 to 2 inches to that at 9 to 10 inches.

The radionuclides corresponding to the photopeaks are indicated in the figure. The photopeaks of Co^{60} , Zn^{65} , Mn^{54} , Ce^{144} - Pr^{144} and Eu^{155} show significant counts in the surface increments but are not detectable in samples from greater depths. The amounts of Cs^{137} and Sb^{125} decrease and the proportions of Cs^{137} and




Fig. 5. Gamma-ray spectra of equal amounts of old soil from different depths.

Fig. 7 shows the gamma spectra of increments taken from an undisturbed area in 1958. The first spectrum is from the 0 to 1/4-inch depth and the subsequent spectra are from 1/8-inch depth increments to a depth of 1 inch. The highest levels of Sb^{125} , the rare earths, and Co^{60} , which move more slowly than Cs^{137} or Sr^{90} , are in the 1/2 to 5/8-inch increments, whereas the Cs^{137} , which moves most rapidly in this soil type, is in the 3/4 to 7/8-inch increment.

Litter and Guano

The gamma-ray spectrum of litter, consisting of leaves, twigs and floral parts splattered with tern droppings, collected from old soil in 1961 (Fig. 8) shows the Cs^{137} photopeak to be much higher than the Sb^{125} peak. The 1.17 peak of Co^{60} is skewed to the left, indicating the presence of the 1.12-mev




Fig. 6. Gamma-ray spectrum of a 9 to 10-inch-depth increment from old soil compared with the gamma spectra of Sb¹²⁵ and Cs¹³⁷.




Fig. 7. Gamma-ray spectra of depth increments of old soil taken from the surface inch. The origin of each spectrum has been arbitrarily separated from the preceding one by a factor of ten to facilitate inspection of the figure.




Fig. 7. (continued)




Fig. 8. Gamma-ray spectra of litter and guano collected in an area of old soil.

Zn^{65} peak. The presence of Zn^{65} is corroborated by the 0.5-mev peak. Fig. 8 also shows the gamma-ray spectrum of noddy tern guano collected in this area. The 1.12-mev peak predominates over the 1.17 peak of Co^{60} and the 0.51 peak of Zn^{65} is evident. The foliar contribution to the litter contains only Cs^{137} from among the gamma-emitters.

In undisturbed areas Cs^{137} and Sr^{90} are being deposited with the litter and are thus replacing at the surface some of the Cs^{137} and Sr^{90} lost by leaching. There is not sufficient data from the field work to determine whether there eventually will be a loss of these radionuclides from the soil-plant system, or a steady state (excluding physical decay of the radionuclides). Long-term experiments, under simulated field conditions, with monolith lysimeters and controlled and uniform addition of the radionuclides would define this point.

Young Soil

Fig. 9 gives the spectra of the 0 to 1-inch, 1 to 2-inch, and 9 to 10-inch increments of a young soil. Co^{60} , Zn^{65} , Ce^{144} , Pr^{144} , and Eu^{155} were detected only in the surface layers, and with increasing depth the 0.60 to 0.66-mev photopeak region of the spectra shifts toward the 0.60-mev peak of Sb^{125} . The spectrum of the 9 to 10-inch increment is compared with that of an Sb^{125} spike in Fig. 10, showing that the photopeaks of the soil and spike gamma spectra are identical.




Fig. 9. Gamma-ray spectra of equal amounts of young soil from different depths.




Fig. 10. Gamma-ray spectrum of young soil from a depth increment of 9 to 10 inches compared with the gamma spectrum of Sb^{125} .

values from the results of Sr^{90} analyses of subsamples of replicate samples taken from small areas differ by a factor of more than ten (Table II). It is likely that the variability is due largely to the spotty nature of the distribution of the fallout radionuclides, which is evident in the radioautographs discussed below and from X-ray films that were exposed at the soil surface and just below the surface (Fig. 11), and to small differences in the characteristics of the soil within a single soil type.

Sampling by 1/8-inch increments in 1959 of an undisturbed old soil, (Fig. 4 and Table I) on Rongelap Island indicates a gradient of Sr^{90} levels in the top inch of soil (Table V). The levels in the second inch are about one tenth those in the

Table IV. Relative Sr⁹⁰ content of depth increments collected in 1959 from two Rongelap Atoll soil types.

Depth in inches	Percent of total in surface ten inches	
	<u>Old Soil</u> (Pit 4)	<u>Young Soil</u> (Pit 6)
0 - 1	81.0	52.9
1 - 2	4.9	16.6
2 - 3	2.8	16.2
3 - 4	2.0	5.8
4 - 5	1.4	4.0
5 - 6	2.2	2.2
6 - 7	2.0	1.2
7 - 8	0.8	0.7
8 - 9	2.5	0.2
9 - 10	0.6	0.1
Total	100.2	99.9




Fig. 11. Radioautograph obtained by exposing no-screen X-ray film approximately one cm below the soil surface in situ.

Table V. Sr^{90} and Ca in the surface 1-1/2 inches of Rongelap soil, Rongelap Gravelly Sand, collected in September 1959.

Depth in inches	Sr^{90} Percent of total in top 1-1/2 inches	Ca mg/gm
0 - 1/8	27	292
1/8 - 1/4	26	291
1/4 - 3/8	16	315
3/8 - 1/2	10	337
3/4 - 7/8	3.2	346
7/8 - 1	2.7	348
1 - 1-1/8	2.0	306
1-1/8 - 1-1/4	2.6	338
1-1/4 - 1-5/8	1.5	329
1-5/8 - 1-3/4	2.2	328
1-3/4 - 1-7/8	3.6	351
1-7/8 - 1-1/2	2.2	369
Total	99.0	

first inch and remain at approximately the same level from 7/8 inch to 1-1/2 inches in depth.

RADIOAUTOGRAPHS



Young Soil

Undisturbed


Radioautographs of cores taken in 1963 show marked differences in the distribution of radioactivity. Fig. 12 shows photographs and the corresponding radioautographs of sections of cores of young soils from an undisturbed area (Fig. 2) and from an area subject to erosion (Fig. 13) and an old soil (Fig. 3). The radioactivity corresponds closely to the dark area in the photograph of the core from the undisturbed young soil. This dark area is composed almost entirely of a mixture of soil algae, forming a crust which has retained most of the fallout radionuclides.

Eroded

In an eroded area of young soil the radioactivity is associated with large coral fragments which are infiltrated with algae. This area is subject to erosion by both wind and water, which accounts for the coral fragments containing algae and radionuclides occurring below the soil surface. Radioactivity is not associated primarily with smaller particles as



DOT 441031



F19

1940

radioautograph of a section of a soil core collected in a depression on Kabelle Islet. There is a darkened funnel-shaped area at the top center of the radioautograph and a corresponding darker area of high organic content in the core section. The amount of activity in the stem of the "funnel" appears to be too great to be accounted for by leaching from the surface directly above.

Either a depression was filled with radioactive organic matter after fallout, or the depression was filled before fallout and the radionuclides were adsorbed when surface runoff water

interpretation seems more probable since gamma-ray spectra of

PDF ARCHIVES





Fig. 14. Photograph and radioautograph of old soil showing correspondence of radioactive portions to organic matter.

Buried Organic Layers

The organic matter in buried horizons may become important in the vertical distribution of radionuclides in the soils. No detectable gamma-emitters were found in buried horizons during the period 1958-1961. In September 1963 a core was taken through a buried horizon at 14 inches in old soil. Radioautographs of sections of the core show that a small amount of radioactivity has accumulated in the buried horizon. The diffuse distribution of the radioactivity in this case indicates that there is adsorption of the radionuclides from solution as water percolates through the soils, although the possibility of translocation by plants cannot be ruled out. Cs^{137} and naturally occurring K^{40} were the only radionuclides present in the buried horizon.

The accumulation of Cs^{137} in the buried horizons is of particular significance in terms of plant-soil relationships, because a proliferation of roots is often found in the buried horizons and Cs^{137} is the principal fallout radionuclide taken up by atoll plants.

Roots

Roots, which contain Cs^{137} and Sr^{90} , in most instances have higher levels of radioactivity than the surrounding soil except near the soil surface (Fig. 12). The influence of poplar roots on redistribution in a continental soil has been well demonstrated

absorbed by the plants, to be most abundant in the soil adjacent to the roots at depth, if there is appreciable channelization.

Pumice

Radionuclides are also adsorbed by pumice particles. No detailed morphological examinations of the soil sections have been made, but it is obvious in some core sections that a few of the larger "hot spots" several inches below the surface are associated with pumice fragments. The retention of radionuclides by pumice fragments may be of importance in considering soil-plant relationships in a few highly localized areas since proliferation of roots around pumice fragments has been observed (Sachet 1955; Kenady 1962). However, as pumice is rarely found beneath the surface, the effects of this material would not be generally important.

* * *

We can not explain the differences in distribution of radionuclides between soil types but assume that the greater retention

continue to do so for years to come, it is possible that it also retains a variety of mineral nutrients, thus providing a reservoir of nutrients in otherwise barren areas. This reservoir might be tapped upon invasion of the areas by higher plants and the concomitant activity of animals in mixing the upper layers of the soil.

Distribution of the radionuclides in the old soils arises from a combination of processes which are difficult to delineate. No doubt much of the movement is due to leaching and readsorption

10/10/55

of radionuclides moving through the soil. This is certainly true for Cs^{137} and Sr^{90} , which were abundant compared to other radionuclides in the leachates. It also was demonstrated by comparing the gamma-ray spectra of depth increments that there is a more rapid movement of Cs^{137} than of other gamma-emitters present. It should be recalled that the exchange capacity of these soils originates from the organic content, which, as is obvious from the photographs of the core sections and Table III, is far higher in the old than in the young soil. There is thus more opportunity for exchange and retention of radionuclides in old soil. Mechanical mixing, due mainly to the activity of land crabs, plays an important role in redistribution in the surface layers. This effect is obvious in areas where there has been active burrowing, and is probably occurring to a small extent throughout densely vegetated areas as is indicated by the presence of Ce^{144} - Pr^{144} and Eu^{155} in the litter. These radionuclides could only have come directly from the soil by upward mixing since they were not found in the vegetation which contributed significantly to the litter.

CONCLUSIONS

Different plant and soil environments on single islets have a different vertical distribution pattern of radionuclides from the same fallout material. The vertical distribution of

penetration: Cs¹³⁷, Sr⁹⁰, Sb¹²⁵, Co⁶⁰, Zn⁶⁵, Cr¹⁴⁴-Pr¹⁴⁴, Eu¹⁵⁵
and probably other rare earths. In the young soils, consisting almost entirely of parent material, the positions of Cs¹³⁷ and Sb¹²⁵ are reversed and the other radionuclides appear to be more completely retained in the surface algal crust. The maximum concentration of fallout radionuclides remains at the soil surface, a few inches or less in depth, except in areas where there has been erosion. In the eroded areas large particles containing both soil algae and radionuclides are randomly distributed to a depth of a few inches.

There is some horizontal movement of radionuclides but such movement appears to be very localized and thus is of little consequence in the overall picture of distribution.

Cs¹³⁷ and Sr⁹⁰ are the principal radionuclides entering a cycle within the soil-plant system. Any loss from this system

appears to be small (a fraction of one per cent per year), but a definite conclusion can not be drawn from the data.

SUMMARY

1. Rongelap Atoll received a single heavy dose of radioactive fallout in 1954.
2. The atoll soils are calcareous and contain no inorganic colloids; the exchange capacity is related to organic content.
3. Comparison of gamma-ray spectra of depth increments from old and new soils shows that Cs^{137} and Sb^{125} move most readily in the old soil; the principal gamma-emitting radionuclide moving in new soil is Sb^{125} .
4. Sr^{90} moves in both old and young soils.
5. The distribution of radionuclides at the surface is very spotty.
6. There is a vertical gradient in the distribution of radionuclides within the surface one-inch layer.
7. Litter redeposits Cs^{137} and Sr^{90} over the soil surface and bird droppings have added Zn^{65} and Co^{60} .
8. The principal reservoir of radionuclides in young soils is the surface algal crust.
9. In eroded areas radioactivity is associated with large coral fragments, which are infiltrated with algae.
10. There is some localized horizontal movement of radionuclides in old soils.

11. Buried organic horizons contain more Cs^{137} than adjacent soil layers.
12. Roots are generally more radioactive than the surrounding soil except at the soil surface.
13. Pumice particles in the soil adsorb radionuclides.
14. Mechanical mixing by animals in old soils and by erosion in young soils results in a redistribution of radionuclides in the surface layers.
15. Cs^{137} and Sr^{90} are the principal radionuclides entering a cycle within the soil-plant system.

LITERATURE CITED

- Cole, D. W., S. P. Gessel and E. E. Held. 1961. Tension lysimeter studies of ion and moisture movement in glacial till and coral atoll soils. *Soil Sci. Soc. of Am., Proc.* 25, 4.
- Dunning, G. M. (Ed.). 1957. Radioactive contamination of certain areas in the Pacific Ocean from nuclear tests. U.S. Atomic Energy Commission, Washington, D. C.
- Held, E. E., S. P. Gessel, L. J. Matson and R. F. Billings. Autoradiography of sectioned soil cores. *In* Proceedings of [IAEA] Symposium on Radioisotope Sample Measurement Techniques in Medicine and Biology, Vienna, May 24-28, 1965. (In press)
- Kenady, R. M. 1962. The soils of Rongelap Atoll, Marshall Islands. M.S. Thesis, University of Washington Library.
- Miller, C. F. and P. Loeb. 1958. Ionization rate and photon pulse rate of fission products from slow neutron fission of U^{235} . U. S. Naval Radiological Defense Laboratory report USNRDL-TR-247.
- Nugent, L. E., Jr. 1946. Coral reefs in the Gilbert, Marshall, and Caroline Islands. *Geol. Soc. Am., Bull.* No. 57, 8.
- Palmer, H. E., W. C. Hanson, B. I. Griffin, and D. M. Fleming. 1964. Radioactivity measurements in Alaskan Eskimos in 1963. *Science* 144:1620.
- Sachet, M. H. 1955. Pumice and other extraneous volcanic materials on coral atolls. *Atoll Res. Bull.*, No. 37.
- Wiens, H. J. 1962. Atoll environment and ecology. New Haven, Conn., Yale University Press. 532 p.
- Witkamp, M. and M. L. Frank. 1964. First year of movement, distribution and availability of Cs^{137} in the forest floor under tagged tulip poplars. *Rad. Botany* 4(4):485-495.

APR 1965