

Regular Article - Theoretical Physics

Pinning down electroweak dipole operators of the top quark

Markus Schulze^{1,a}, Yotam Soreq^{2,b}

¹ CERN Theory Division, 1211 Geneva 23, Switzerland

Received: 11 April 2016 / Accepted: 13 July 2016 / Published online: 19 August 2016 © The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract We consider hadronic top quark pair production and pair production in association with a photon or a Z boson to probe electroweak dipole couplings in $t\bar{b}W$, $t\bar{t}\gamma$, and $t\bar{t}Z$ interactions. We demonstrate how measurements of these processes at the 13 TeV LHC can be combined to disentangle and constrain anomalous dipole operators. The construction of cross section ratios allows us to significantly reduce various uncertainties and exploit orthogonal sensitivity between the $t\bar{t}\gamma$ and $t\bar{t}Z$ couplings. In addition, we show that angular correlations in $t\bar{t}$ production can be used to constrain the remaining $t\bar{b}W$ dipole operator. Our approach yields excellent sensitivity to the anomalous couplings and can be a further step toward precise and direct measurements of the top quark electroweak interactions.

1 Introduction

The dynamics of top quark production and decay have been extensively studied at hadron colliders. Early measurements at the Tevatron have taught us about the production mechanism of heavy quarks pairs in quantum chromodynamics (QCD) and later evolved into precision measurements of the top quark mass, spin correlations, and the forward-backward asymmetry [1,2]. Similar measurements have been performed during Run-I of the large hadron collider (LHC) which superseded earlier results at an impressive pace [3,4]. A myriad of measurements has profoundly shaped our understanding of top physics in the standard model (SM) and it led to strong exclusion bounds on new physics.

Despite this progress, our understanding of the electroweak interactions of the top quark (i.e. its coupling to electroweak gauge bosons and the Higgs boson) is rather limited. The main reasons are the high production thresholds

Preprint numbers: MIT-CTP/4790, CERN-TH-2016-070.

^a e-mail: markus.schulze@cern.ch

b e-mail: soreqy@mit.edu

of the $t\bar{t}+X$ processes, small branching fractions and large backgrounds. First measurements and searches at 7 and 8 TeV LHC have established some of these SM processes [5–13], but detailed studies of electroweak couplings will only be possible with a larger data set at higher energies during Run-II. These studies will certainly improve our understanding of top interactions with the electroweak sector and potentially probe physics beyond the SM.

The flavor-changing $t\bar{b}W$ interaction is the best-known top quark coupling as it is experimentally accessible through top quark decays in $t\bar{t}$ production and single-top quark processes. This is reflected by precise measurements of W helicity fractions and top quark spin correlations [14-17] which can be translated into bounds on anomalous couplings. All other electroweak interactions of the top quark with the Z, γ , and the Higgs boson are much less explored. For example, the top quark electric charge, which governs the coupling strength of the vector-like $t\bar{t}\gamma$ interaction, is well known to be $Q_t = +2/3$ with a confidence level larger than 5σ [18–21]. However, these determinations were obtained from measuring the electric charges of W boson and b-jet in $t\bar{t}$ production, inferring $Q_t = Q_W + Q_b$. A hard photon was never present in the event sample and the fundamental $t\bar{t}\gamma$ interaction was not probed. Similarly, current LHC data only allows constraints on the vector and axial parts of the $t\bar{t}Z$ vertex with $\mathcal{O}(100\,\%)$ uncertainties [13], while the respective dipole couplings are unconstrained from hadron collider experiments. We note that low-energy observables, such as rare K and B decays [22–24], together with electroweak precision data [25–28] can provide strong constraints on modified $t\bar{t}Z$ interactions. However, these fairly indirect probes are based on either $Z \to b\bar{b}$ decays or highly off-shell top quarks and Z bosons in $b \to s Z^*/\gamma^*$ transitions which rely on additional assumptions on the new physics and are prone to hadronic uncertainties.

This immediately motivates precise studies of the final states $t\bar{t} + Z/\gamma/h$, which yield *direct* sensitivity to the desired couplings. The commencing Run-II of the LHC with a colli-



² Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

466 Page 2 of 10 Eur. Phys. J. C (2016) 76:466

sion energy of 13 TeV will, for the first time, produce sufficiently many events to enable coupling studies. The two final states considered in this paper, $t\bar{t}+\gamma$ and $t\bar{t}+Z(\to\ell\ell)$, with semi-hadronically decaying top quark pairs, have typical fiducial cross sections of $\sim 400\,\mathrm{fb}$ and 5 fb, respectively. Hence, we expect about $40,000\,t\bar{t}\gamma$ and $500\,t\bar{t}Z$ events from $100\,\mathrm{fb}^{-1}$ of integrated luminosity.

Anomalous electroweak top quark coupling studies using these processes have been presented in Refs. [29-49]. For example, the TopFitter collaboration [37] most recently combined various top quark measurements from the Tevatron and the LHC to present a global fit. In Ref. [31] the authors study dedicated differential observables to probe CP-violating top quark decays. Anomalous coupling studies that go beyond the leading order have been presented in Refs. [38,44,45,47,48]. In this work, we go beyond previous studies by combining observables from $t\bar{t}$, $t\bar{t} + \gamma$, and $t\bar{t}+Z$ to investigate sensitivity to the three electroweak dipole operators which enter simultaneously in these processes. We construct cross section ratios to cancel correlated uncertainties and investigate angular asymmetries of the top quark decay products to boost sensitivity to possible effects of new physics.

2 Setup

In the SM, the fundamental interaction vertices of electroweak vector bosons with fermions are given by


$$\Gamma_{q'qV}^{SM} = \bar{q}' \gamma^{\mu} \left(d_{L}^{V} P_{L} + d_{R}^{V} P_{R} \right) q \, \varepsilon_{\mu}^{V}, \tag{1}$$

where $V=\gamma,Z,W^\pm$, and $P_{\rm L,R}$ are the left and right-handed chirality projectors. The respective couplings $d_{\rm L,R}^V$ are fixed by the quantum numbers and gauge symmetries of the SM, see for example Ref. [50]. In this work, we focus on additional contributions from anomalous electroweak dipole moments in the top quark sector. Their coupling structure is given by

$$\delta\Gamma_{q'qV} = \bar{q}' \frac{i\sigma^{\mu\nu}k_{\nu}}{m_t} \left(g_{\rm L}^V P_{\rm L} + g_{\rm R}^V P_{\rm R} \right) q \, \varepsilon_{\mu}^V, \tag{2}$$

where $\sigma^{\mu\nu}=i/2\,[\gamma^\mu,\gamma^\nu]$, $k=p_q-p_{q'}$, and $g^V_{\rm L,R}$ are the dipole couplings. The SM has no such interactions at tree level, but electroweak loop corrections radiatively generate dipole moments. Their size is well below 1 per-mille [51–53] which makes them inaccessible by LHC experiments. Hence, any sizable deviation from zero will indicate an anomalous interaction from physics beyond the SM. In order to understand how deep the new physics scales can be probed, we investigate the effects of Eq. (2) on physical observables.

Various well-motivated models of new physics [53–59] predict sizable top quark dipole moments. In this work, we



$$\mathcal{O}_{uW}^{33} = \left(\bar{q}_{L}\sigma^{\mu\nu}\tau^{I}t_{R}\right)\tilde{H}W_{\mu\nu}^{I},\tag{3}$$

$$\mathcal{O}_{dW}^{33} = \left(\bar{q}_{L}\sigma^{\mu\nu}\tau^{I}b_{R}\right)HW_{\mu\nu}^{I},\tag{4}$$

$$\mathcal{O}_{\mu B\phi}^{33} = (\bar{q}_{L}\sigma^{\mu\nu}t_{R})\,\tilde{H}B_{\mu\nu},\tag{5}$$

where $\tilde{H}=\mathrm{i}\,\tau^2 H^*$, $H=(0,h+v)/\sqrt{2}$ is the Higgs boson doublet with the vacuum expectation value $v=246\,\mathrm{GeV}$. $B_{\mu\nu}(W^I_{\mu\nu})$ is the U(1)_Y(SU(2)_L) gauge field signal strength and I is adjoint SU(2)_L index. $q_{\rm L}=(t_{\rm L},b_{\rm L}),\,b_{\rm R}(t_{\rm R})$ are the quark left-handed doublet and bottom (top) right-handed singlet, respectively. Using the parameterization of Eq. (2) and the operators in Eqs. (3)–(5) we find

$$g_{L}^{W^{-}} = g_{R}^{W^{+}*} = -\frac{e \, m_{t}}{s_{W} M_{W}} \frac{v^{2}}{\Lambda^{2}} C_{dW}^{33*},$$

$$g_{R}^{W^{-}} = g_{L}^{W^{+}*} = -\frac{e \, m_{t}}{s_{W} M_{W}} \frac{v^{2}}{\Lambda^{2}} C_{uW}^{33},$$

$$g_{L}^{\gamma} = g_{R}^{\gamma*} = -\frac{\sqrt{2} \, m_{t} \, v}{\Lambda^{2}} \left(c_{W} C_{uB\phi}^{33*} + s_{W} C_{uW}^{33*} \right),$$

$$g_{L}^{Z} = g_{R}^{Z*} = -\frac{e \, m_{t} \, v^{2}}{\sqrt{2} s_{W} c_{W} M_{Z} \Lambda^{2}} \left(c_{W} C_{uW}^{33*} - s_{W} C_{uB\phi}^{33*} \right),$$
(6)

where e is the electric coupling and $s_W(c_W)$ is sine (cosine) of the weak mixing angle. We emphasize that since the various dipole couplings in $t\bar{b}W$, $t\bar{t}\gamma/Z$ are related via the underlying $SU(2)\times U(1)$ gauge invariance, the coupling degrees of freedom are reduced to only three Wilson coefficients C_{uW}^{33} , C_{dW}^{33} , and $C_{uB\phi}^{33}$. In the expression for $g_{L,R}^{\gamma/Z}$, we see the characteristic weak mixing angle rotation between C_{uW}^{33} and $C_{uB\phi}^{33}$ which will be responsible for the *orthogonal* constraints that we find in $t\bar{t}+\gamma$ and $t\bar{t}+Z$ production below.

At this point we note that there is one more allowed Lorentz structure in addition to Eq. (2), which is relevant for our analysis,

$$\delta\Gamma_{q'qVV'} = \bar{q}' \text{ i}\sigma^{\mu\nu} \left(g_{L}^{VV'} P_{L} + g_{R}^{VV'} P_{R} \right) q \, \varepsilon_{\mu}^{V} \, \varepsilon_{\nu}^{V'}. \tag{7}$$

This four-point vertex enters our calculation only in one specific place: the radiative top quark decay $t \to bW + \gamma$ of the $t\bar{t}\gamma$ final state. In a complete description of $t\bar{t}\gamma$ production, the photon can arise from either the hard production stage (before the top quarks go on-shell), or the top quark decay stage (after one top quark went on-shell). The latter part constitutes more than 50 % of the total $t\bar{t}\gamma$ cross section [60]



Table 1 Contribution of dimension-six Wilson coefficients to various vertices appearing in $t\bar{t}$, $t\bar{t} + \gamma$, and $t\bar{t} + Z$ production

	b	b W W	t γ/Z t
C_{uW}^{33}	\otimes	\otimes	\otimes
$C^{33}_{uW} \\ C^{33}_{dW}$	\otimes	\otimes	
$C^{33}_{uB\phi}$			\otimes

and receives contributions from Eq. (7). It is therefore crucial to account for this contribution in the analysis of $t\bar{t}\gamma$ final states. The two additional dipole couplings in Eq. (7) are given by

$$g_{\mathrm{L,R}}^{W^{\sigma}\gamma} = (-e)\,\sigma\,g_{\mathrm{L,R}}^{W^{\sigma}},\tag{8}$$

where $\sigma=\pm 1$ is the electric charge of the respective W boson. In Table 1 we summarize the anomalous coefficients considered in our analysis and their appearance in various interaction vertices.

The up-to-date hadron collider bounds on the Wilson coefficients of Eqs. (3)–(5) are summarized in Ref. [37] and found to be

$$C_{uW}^{33} \in [-4.0, 3.4] (\Lambda/\text{TeV})^2,$$
 (9)

$$C_{uB\phi}^{33} \in [-7.1, 4.7] (\Lambda/\text{TeV})^2$$
 (10)

at 95% confidence level (CL). Note that in order to infer the bound on C_{uW}^{33} , the contributions of all other top-related dimension-six operators were marginalized, while for $C_{uB\phi}^{33}$, it was assumed that the only new physics contribution is from $C_{uB\phi}^{33}$. Finally, in Ref. [30] we find that

$$C_{dW}^{33} \in [-2.3, +2.3] (\Lambda/\text{TeV})^2,$$
 (11)

at 95% CL. Note that flavor violating contributions from $\mathcal{O}^{33}_{uB\phi}$ can be avoided by alignment to the up sector in flavor space. In that limit, \mathcal{O}^{33}_{uW} induces an irreducible source of flavor violation in the charged current, which is, however, suppressed by off-diagonal CKM matrix elements; for a more detailed discussion see Ref. [61]. Alignment to the down basis will reduce flavor violation originating from \mathcal{O}^{33}_{dW} , see also the discussion in Ref. [62].

In a strict $1/\Lambda^2$ -expansion within an effective field theory even more operators contribute to the processes considered here. For example, the operator $\bar{\ell}\gamma^{\mu}\ell$ $\bar{t}\gamma_{\mu}t$ can enter the $t\bar{t} + Z(\rightarrow \ell\ell)$ process. However, these operators do not exhibit a Breit-Wigner peak structure such as the rest of the amplitude and contribute as a fairly constant function of $m_{\ell\ell}$ around the Z boson mass window of ± 10 GeV, see e.g. Fig. 5 in [41]. Hence, their interference at $\mathcal{O}(\Lambda^{-2})$ integrates to zero and only their squared contribution at $\mathcal{O}(\Lambda^{-4})$ survives. We therefore neglect all operators with fermionic contact interactions. Potential contributions from chromo-magnetic and chromo-electric dipole moments can enter our processes through anomalous top-gluon couplings. However, in this work, we assume that QCD is unaltered by new physics and we do not consider chromo-dipole moments. If QCD is truly non-fundamental, these anomalous interactions are best searched for in jet processes, $gg \to H$ or $t\bar{t}$ production, see e.g. Ref. [63].

Besides dipole interactions, also the strength of the $t\bar{t}Z$ and tbW vector and axial couplings can be altered by new physics. In this work we assume their SM value which is a restrictive assumption on new physics scenarios. Let us therefore comment on possible strategies for a broader analysis in future extensions of our work. One straight-forward solution is to consider the full space of couplings in a six-dimensional analysis (anomalous vector and axial-vector couplings introduce three more operators to the ones considered here [50]). This approach might, however, be challenging given the large number of degrees of freedom and the limited number of events. Another solution may be a careful analysis of kinematics in sequential steps: (1) a first analysis of the process $pp \rightarrow t\bar{t} + \gamma$ can yield information on the dipole moments. Anomalous vector or axial-vector couplings cannot develop thanks to gauge symmetries. (2) Any deviation in $t\bar{t} + \gamma$ immediately predicts anomalies in the $pp \to t\bar{t} + Z$ process, as can be seen from Eq. (6). In Ref. [45] it has been shown that dipole couplings most prominently manifest in energy related contributions such as $p_{T,Z}$. Hence, any remaining discrepancy from anomalous vector and axial-vector couplings can be detected in angular distributions, such as $\Delta \phi_{\ell\ell}$, which



¹ The effects from radiative top quark decays are irrelevant for the $t\bar{t}+Z$ process because the decay $t\to bW+Z$ is suppressed by a phase space factor of 10^{-6} .

466 Page 4 of 10 Eur. Phys. J. C (2016) 76:466

are sufficiently independent of energies. (3) Once anomalous dipole couplings in $t\bar{t}+\gamma/Z$ are established, only one remaining dipole operators in $t\bar{b}W$ interactions remains and can be constrained as outlined in Sect. 4. Additional anomalous vector and axial couplings in the $t\bar{b}W$ interaction can then be constrained by relating analyses of top quark pair and single-top quark production $qb \to W \to q't$ [30]. Finally, we note that scenarios of large CP-violating electric dipole moments have been extensively studied in the literature [54,56,64–66]. These effects arise through CP violation in loop contributions and we expect them to similarly affect the magnetic dipole moments.

Our description of the processes $t\bar{t}, t\bar{t} + \gamma$, and $t\bar{t} + Z(\rightarrow \ell\ell)$ treats top quarks in the narrow-width approximation and includes the full decay chain of the top quarks into a single lepton plus jets final state. All spin correlations are retained. In $t\bar{t} + \gamma$ final states, the photon is allowed to be emitted in the top quark production stage as well as in the decay stage (including the W and W decay products). The Monte Carlo simulation is based on the TOPAZ code [67] developed in Refs. [44,45,68], which we extended to handle all anomalous couplings needed in this analysis. If not otherwise stated, we use the following generic selection cuts for the $t\bar{t}, t\bar{t} + \gamma$, and $t\bar{t} + Z$ processes:


$$p_{\perp}^{\ell} \ge 20 \,\text{GeV}, \quad |y_{\ell}| \le 2.5, \quad E_{\perp}^{\text{miss}} \ge 20 \,\text{GeV},$$

 $p_{\perp}^{j,b} \ge 20 \,\text{GeV}, \quad |y_{j}| \le 2.5, \quad |y_{b}| \le 2.0.$ (12)

Jets are defined by the anti- $k_{\rm T}$ jet algorithm [69] with R=0.4. In addition, for $t\bar{t}+Z$ production we require an invariant mass cut of $|m_{\ell\ell}-M_Z|\leq 10\,{\rm GeV}$. For $t\bar{t}+\gamma$ we require the isolation cuts $R_{\gamma j}=R_{\gamma\ell}=0.4$ for photons with $p_\perp^\gamma\geq p_{\perp,{\rm cut}}^\gamma=20\,{\rm GeV}$ and $|y_\gamma|\leq 2.5$. In accordance with the findings at NLO QCD [45,60], we set the central renormalization and factorization scales to $\mu=m_t$ for $t\bar{t}$ and $t\bar{t}+\gamma$ production, and $\mu=m_t+M_Z/2$ for $t\bar{t}+Z$ production. We use NNPDF3.0 [70] parton distribution functions, $m_t=173.2\,{\rm GeV},\,M_Z=91.1876\,{\rm GeV},\,M_W=80.399\,{\rm GeV},\,G_{\rm F}=1.16639\times 10^{-5}\,{\rm GeV}^{-2},$ and use $\alpha=1/137$ in the $t\bar{t}\gamma$ process.

3 Cross section ratios

In the following we study the sensitivity of different cross section ratios to the anomalous dipole couplings. Ratios of observables have the advantage that leading uncertainties on e.g. α_s and parton distribution functions (PDFs) largely cancel. Even higher-order corrections are expected to cancel to some extent, provided that the cross sections are probed in similar regions of phase space. Experimental uncertainties



A similar idea of employing ratios has been presented in Ref. [71] for measuring the top quark Yukawa coupling at the 100 TeV future circular collider. The authors consider the cross section ratio of $t\bar{t}+H$ over $t\bar{t}+Z$ and demonstrate that a precision of 1 % can be reached. This certainly extreme precision is obtained thanks to the kinematic similarities of the two processes and the enormous event rate at a 100 TeV collider. Hence, in the context of this work it seems suggestive to study the ratio $\sigma_{t\bar{t}Z}/\sigma_{t\bar{t}\gamma}$. Yet, we refrain from doing so and instead construct two ratios

$$\mathcal{R}_{\gamma} = \sigma_{t\bar{t}\gamma} / \sigma_{t\bar{t}}, \quad \mathcal{R}_{Z} = \sigma_{t\bar{t}Z} / \sigma_{t\bar{t}}, \tag{13}$$

with respect to the $t\bar{t}$ cross section. This allows one to benefit from the large $\sigma_{t\bar{t}}$ cross section which has almost no statistical uncertainties. Moreover, we will find that $\sigma_{t\bar{t}Z}$ and $\sigma_{t\bar{t}\nu}$ have orthogonal dependence on the anomalous couplings which is distinctly exposed in the ratios of Eq. (13). Of course, it is no longer given that uncertainties cancel in these ratios because the $t\bar{t} + \gamma/Z$ and $t\bar{t}$ processes probe very different energies and phase spaces. For example, using the cuts in Eq. (12) we find at leading order an average center-of-mass energy of $\langle \hat{E} \rangle \approx 525$ GeV for the $t\bar{t}$ process, but $\langle \hat{E} \rangle \approx 630$ GeV and 860 GeV for $t\bar{t} + \gamma$ and $t\bar{t} + Z$, respectively. Hence, the parton distribution functions are evaluated at significantly different values of Q^2 and a cancellation of uncertainties is not guaranteed. We circumvent this issue by applying additional (mild) invariant mass cuts on the two $t\bar{t}$ cross sections in Eq. (13) to increase the average center-of-mass energy. In particular, we request $m_{t\bar{t}} \geq 470$ GeV for $\sigma_{t\bar{t}}$ in \mathcal{R}_{γ} , and $m_{t\bar{t}} \geq 700 \text{ GeV for } \sigma_{t\bar{t}} \text{ in } \mathcal{R}_Z.$ We verified that the average center-of-mass energy in these two $t\bar{t}$ processes matches the one of the respective $t\bar{t} + \gamma/Z$ process. We note that the experimental reconstruction of the $t\bar{t}$ invariant mass is well known to involve large systematic errors. One could therefore worry how this affects the above mentioned cut on $m_{t\bar{t}}$ and how it may spoil the cancellation of uncertainties in the cross section ratios. We therefore investigated variations of this cut and its impact on the ratios \mathcal{R}_{γ} and \mathcal{R}_{Z} . While the absolute values obviously change with different values of the cut, we find that our uncertainty estimates presented below are completely unchanged. Even in the extreme case where we remove the $m_{t\bar{t}}$ cuts entirely, our conclusions remain the same as presented below.

To explicitly quantify cancellations of uncertainties in the ratios, we evaluate the cross sections at NLO QCD and study variations of parton distribution functions. This also allows us to obtain error estimates that will be important when estimating sensitivity to the dipole couplings. We partially use results from the NLO QCD computations of $t\bar{t} + \gamma$ and $t\bar{t} + Z$ in Refs. [44,60] and recompute the NLO $t\bar{t}$ cross sections with



Eur. Phys. J. C (2016) 76:466 Page 5 of 10 466

the same input parameters and the above mentioned cut on $m_{t\bar{t}}$. The higher average center-of-mass energy of the $t\bar{t}$ cross sections also calls for adapting the renormalization and factorization scales. We find it natural to choose $\mu = m_t + p_{\perp, \text{cut}}^{\gamma}$ and $\mu = m_t + M_Z/2$ for the two $t\bar{t}$ cross sections in Eq. (13), respectively. Using this setup, we find

$$\mathcal{R}_{\gamma}^{\text{SM}} \times 10^{-3} = \begin{cases} 11.4_{+0.7\%}^{-0.7\%} & \text{at LO,} \\ 12.6_{-1.8\%}^{+3.1\%} & \text{at NLO QCD,} \end{cases}$$
 (14)

$$\mathcal{R}_{\gamma}^{\text{SM}} \times 10^{-3} = \begin{cases} 11.4_{+0.7\%}^{-0.7\%} & \text{at LO,} \\ 12.6_{-1.8\%}^{+3.1\%} & \text{at NLO QCD,} \end{cases}$$

$$\mathcal{R}_{Z}^{\text{SM}} \times 10^{-4} = \begin{cases} 2.27_{+2.0\%}^{-1.7\%} & \text{at LO,} \\ 1.99_{+2.8\%}^{-1.9\%} & \text{at NLO QCD.} \end{cases}$$
(15)

The upper (lower) values correspond to the lower (upper) scale variation by a factor of two around the respective central scale. We observe a scale dependence of ± 1 % and ± 2 % at LO for \mathcal{R}_{ν} and \mathcal{R}_{Z} , respectively. These values are slightly increased to $\pm (2-3)$ % at NLO QCD. This constitutes a remarkable stability with respect to scale variation when compared to the cross sections themselves which exhibit a dependence of about ± 20 % at NLO. It should be noted that the LO scale variation is far outside the NLO result. This is to be expected as $\alpha_s(\mu_R)$ cancels exactly and the only remaining source of scale dependence arises from unmatched q^2 dependence in the parton distribution functions. Only our NLO results develop, for the first time, logarithms of the scales with process-dependent coefficients that yield a good error estimate. Hence only the NLO result should be considered a physical meaningful prediction. For our coupling constraints below we will choose the largest value of the NLO scale variation, $\Delta \mathcal{R}_{Z/\gamma} = 3\%$, as our uncertainty for both ratios.

Given the fiducial cross sections of about $\sim 5 (400)$ fb for $t\bar{t}Z(t\bar{t}\gamma)$ production, the statistical error is expected to be sub-dominant after an integrated luminosity of about 250 fb⁻¹. This argument is supported by a first measurement of \mathcal{R}_{γ} by the CMS collaboration [10] at 8 TeV. They find the value $\mathcal{R}_{\gamma}(8\,\text{TeV}) = 10.7 \times 10^{-3} \pm 6.5\,\%$ (stat.) \pm 25% (syst.) from an integrated luminosity of $19.7 \, \text{fb}^{-1}$. The dominant systematics arise from background modeling $(\pm 23 \%)$ which have the potential to be improved in future analyses.

We also investigate uncertainties from parton distribution functions and find the following results from using three different sets of parton distribution functions:

$$\mathcal{R}_{\gamma}^{\text{LO}} \times 10^{-3} = \begin{cases} 11.5 & \text{with NNPDF3.0 [70],} \\ 11.4 & \text{with CTEQ6L1 [72],} \\ 11.5 & \text{with MSTW08 [73],} \end{cases}$$
 (16)

$$\mathcal{R}_{Z}^{LO} \times 10^{-4} = \begin{cases} 2.29 & \text{with NNPDF3.0,} \\ 2.27 & \text{with CTEQ6L1,} \\ 2.27 & \text{with MSTW08.} \end{cases}$$
 (17)

We observe very stable results with variations at the level of 1 %, which have to be compared to ± 10 % variations on the cross sections themselves. Again, we find confirmation that the ratios are true precision observables.

Finally, let us briefly comment on the impact of electroweak corrections. Compared to the QCD corrections, they are expected to be much less universal for the $t\bar{t}$ and $t\bar{t} + \gamma/Z$ processes. Hence, on the one hand, a cancellation is most likely incomplete. On the other hand, the electroweak corrections on the cross sections are known to be in the few percent range $(\mathcal{O}(\alpha))$ [74,75]. This leads to a (minor) shift in the absolute values of the ratios but it does not at all affect our estimate of uncertainties.

Let us now turn to studying the effects of anomalous electroweak dipole moments on the cross section ratios. Given two (pseudo-)observables, \mathcal{R}_{γ} and \mathcal{R}_{Z} , we investigate their dependence on the Wilson coefficients C_{uW}^{33} and $C^{33}_{uB\phi}$ (neglecting operator mixing and running from beyond the LO). The remaining coefficient C_{dW}^{33} we examine through angular asymmetries in $t\bar{t}$ production in the following section. Since only total cross sections enter this analysis, we are not sensitive to the tail of energy-related distributions where possible issues with unitarity violating EFT operators could appear. We vary the numerical values of C_{uW}^{33} and $C_{uB\phi}^{33}$ between [-4, 4] in steps of 1 and compute 81 LO cross sections for each of the $t\bar{t}$, $t\bar{t} + \gamma$, and $t\bar{t} + Z$ processes. We then perform a two-dimensional analytic fit of the ratios to present our results. Figure 1 shows the deviation of the anomalous ratios from the SM value for $t\bar{t}\gamma$ on the left and $t\bar{t}Z$ in the middle. These two plots strikingly show the *orthogonal* dependence of the two ratios on the dipole couplings. This is a consequence of the already mentioned $s_{\rm W}$ rotation pattern of $g_{\rm L,R}^{\gamma,Z}$ in Eq. (6). The white (dark green) bands in left and middle plot of Fig. 1 indicate the parameter spaces where the anomalous cross section ratios deviate from the SM by less than the assumed 1 (2) standard deviation of the assumed uncertainty. Hence, all anomalous couplings outside the dark green bands can be excluded if the measurement is in agreement with the SM at the 2σ level. In the third plot, on the right of Fig. 1, we show the combination of the two constraints using a naive χ^2 combination. This noticeably leads to a striking improvement of the constraints as the orthogonal dependence on the Wilson coefficients allows us to completely bound the parameter space without a blind direction. We note that these projected bounds are stronger by factors of 2–3 compared to the current bounds from 7 and 8 TeV cross section measurements. Moreover, a better understanding of the anomalous $pp \rightarrow b\bar{b}\ell\nu jj + \gamma$ process beyond



466 Page 6 of 10 Eur. Phys. J. C (2016) 76:466




Fig. 1 Cross section ratios \mathcal{R}_{γ} (*left*) and \mathcal{R}_{Z} (*middle*) normalized to their SM values ($\mathcal{R}_{\gamma/Z}^{\mathrm{SM}}$) as a function of the anomalous dipole operator couplings. The *contours* show the deviation from the SM value in

steps of 3, 6 and 9 %. On the *right* we show the 1, 2, 3σ contours from combining \mathcal{R}_{γ} and \mathcal{R}_{Z} with an assumed uncertainty of $\Delta \mathcal{R}_{Z/\gamma} = 3$ %




Fig. 2 Angular distributions in $pp \to t\bar{t} \to b\bar{b} \,\ell\nu \,jj$ at 13 TeV used to construct the asymmetries $A_{\theta^*_i}$ and A_{α}

the LO will further improve these results. Such a calculation is currently not available and we refer to future work.

4 Angular asymmetries in $t\bar{t}$ production

In this section we expand and complement the constraints obtained from $t\bar{t}+\gamma$ and $t\bar{t}+Z$ in the previous section. We make use of large $t\bar{t}$ cross section at the LHC to define asymmetries of angular distributions from the top quark decay products in order to constrain the remaining Wilson coefficient C_{dW}^{33} , and to over-constrain C_{uW}^{33} . As before, we use the lepton+jet final state of the $t\bar{t}$ system. Similar ideas have been presented in Ref. [29], which puts more emphasis on probing the complex phases of the anomalous couplings. Here, we consider the two angles, defined by

$$\cos \theta_{\ell}^* = \frac{\hat{\mathbf{p}}_{\ell} \cdot \tilde{\mathbf{p}}_{W}}{|\hat{\mathbf{p}}_{\ell}| |\tilde{\mathbf{p}}_{W}|}, \quad \cos \alpha = \frac{\mathbf{p}_{t} \cdot \tilde{\mathbf{p}}_{W}}{|\mathbf{p}_{t}| |\tilde{\mathbf{p}}_{W}|}, \tag{18}$$

where $\hat{\mathbf{p}}_{\ell}$ is the lepton momentum in the corresponding W rest frame, $\tilde{\mathbf{p}}_{W}$ is the W momentum in the corresponding top quark rest frame, and \mathbf{p}_{t} is the top quark momentum in the laboratory frame. Their kinematic distributions are shown in Fig. 2 for the SM and two anomalous coupling choices. From these distributions we construct asymmetries

$$A_{\phi}(c_0) = \frac{\sigma(\cos\phi < c_0) - \sigma(\cos\phi > c_0)}{\sigma(\cos\phi < c_0) + \sigma(\cos\phi > c_0)},\tag{19}$$

and use $A_{\theta_\ell^*}(-0.1)$ and $A_\alpha(0.0)$ to maximize their value in our analysis. We evaluated the SM asymmetries at NLO QCD and find perturbative shifts of only $\mathcal{O}(+0.5\,\%)$. In our coupling analysis below we will assume a slightly inflated uncertainty of $\pm 4\,\%$. On the experimental side, we expect a similar precision because of the large statistical sample at 13 TeV and the fact that existing measurements at 8 TeV already reach 10 % precision for spin asymmetries [76,77].



Eur. Phys. J. C (2016) 76:466 Page 7 of 10 466




Fig. 3 Angular asymmetries $A_{\theta_{\ell}^*}(-0.1)$ (*left*) and $A_{\alpha}(0.0)$ (*middle*) as a function of the two Wilson coefficients in $t\bar{t}$ production relative to the their SM values. *Right* χ^2 combination of the two asymmetries assuming an uncertainty of $\pm 4\%$

To probe the sensitivity of the two asymmetries to the dipole couplings we vary C_{dW}^{33} and C_{uW}^{33} between [-4, 4] in steps of 0.5. Hence, we perform 324 computations at leading order and sub-sequentially fit the results to an analytic parameterization that is shown in Fig. 3. Interestingly, the functional dependence of the asymmetries on the dipole operator couplings is opposite: $A_{\theta_{\ell}^*}$ has a concave shape (Fig. 3, left) as a function of C_{uW}^{33} and C_{dW}^{33} , whereas A_{α} (middle) is convexly shaped. This feature allows us to combine the two constraints, shown on the right of Fig. 3, in order to break the (white) invariance bands of the two separate observables. As a result, both anomalous operators are clearly bounded in the combination plot. Moreover, the constraints on C_{nW}^{33} can be further utilized in conjunction with the bounds obtained from cross section ratios (Fig. 1, right) as they exclude almost the entire region of positive C_{uW}^{33} values. Turning the argument around, two independent measurements of (1) cross section ratios and (2) angular asymmetries can also be used to overconstrain the Wilson coefficient C_{uW}^{33} .

5 Summary

In this paper we investigated the prospects of constraining electroweak dipole moments of the top quark at the 13 TeV LHC. The SM radiatively generates these couplings through electroweak loop corrections, which turn out to be too small to be observed at the LHC. This opens up the possibility to search for sizable deviations from zero as a way to probe new physics in the top quark sector.

We considered all anomalous dipole interactions between the top quark and the electroweak gauge bosons in $\bar{t}bW$, $t\bar{t}\gamma$, and $t\bar{t}Z$ interactions, and we showed that the processes $pp \to t\bar{t}$ and $pp \to t\bar{t} + \gamma/Z$ are ideal probes for studying them. While these couplings enter simultaneously in various places, electroweak gauge symmetries relate them and lead to only a small number of relevant operators. In

order to constrain and disentangle them, we propose the study of cross section ratios to make use of *orthogonal* sensitivity to anomalous operators entering $t\bar{t}\gamma$ and $t\bar{t}Z$. We carefully investigated the ratios at NLO QCD and verify a strong reduction of uncertainties related to parton distribution functions and higher-order corrections. Experimental systematics such as luminosity or jet energy scale uncertainties are expected to drop out as well, yielding true precision observables.

The final results of our study confirm this picture as we find that 13 TeV data can place firm bounds on the contributing operators. Marginalizing over other operators, we find sensitivity of $C_{uW}^{33} = [-1.2, +1.4] (\Lambda/\text{TeV})^2$ and $C_{uB\phi}^{33} = [-1.9, +1.2] (\Lambda/\text{TeV})^2$ at the 95 % CL from combining the cross section ratios, assuming that the theoretical accuracy of 3% is matched by the experimental one.

We corroborate our results by studying angular asymmetries in $t\bar{t}$ production to bound the last operator in our analysis and find the sensitivity $C_{dW}^{33} = [-2.0, +2.0] \, (\Lambda/\text{TeV})^2$. Since the angular asymmetries are also sensitive to C_{uW}^{33} , they can yield the independent constraint $C_{uW}^{33} = [-0.8, +0.1] \, (\Lambda/\text{TeV})^2$, which can be further used to boost our results from cross section ratios. Altogether, our proposal to construct four precision observables allows one to pin down all of the three anomalous electroweak dipole operators without a blind direction.

We note that for a 100 TeV pp collider the $t\bar{t}Z$ and $t\bar{t}\gamma$ cross sections are about a factor of 30 larger than at the 13 TeV LHC. Thus, the statistical error will be sub-dominant only after a few tens fb⁻¹ of integrated luminosity. In order to fully exploit the potential of a 1 ab⁻¹ data set, the theoretical predictions should be improved by one order of magnitude. Theoretical control of cross section ratios at the per-mille level may be possible once predictions at next-to-next-to-leading order QCD are available for $t\bar{t}+\gamma/Z$. This does not seem unrealistic on the relevant time scale and would boost the constraints by more than 100%.



466 Page 8 of 10 Eur. Phys. J. C (2016) 76:466


Future work on improving our results could be a more precise understanding of uncertainties of the cross section ratios. For example, it would be desirable to have the complete NLO predictions for the $t\bar{t}+\gamma$ process with anomalous couplings. A fully realistic analysis also needs to consider backgrounds which we neglected in this work. The incorporation of single-top quark processes into our analysis or more differential observables will certainly further strengthen sensitivity to new physics.

Acknowledgments We thank Michelangelo Mangano, Gilad Perez, Raoul Röntsch for comments on the manuscript and Jesse Thaler for helpful discussions. M.S. is grateful for the hospitality at MIT-CTP where this project was initiated. Y.S. is supported by the U.S. Department of Energy (DOE) under cooperative research agreement DE-SC-00012567.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Funded by SCOAP³.

References

- F. Deliot, D.A. Glenzinski, Top quark physics at the Tevatron. Rev. Mod. Phys. 84, 211 (2012). doi:10.1103/RevModPhys.84.211
- D. Wicke, Properties of the top quark. Eur. Phys. J. C 71, 1627 (2011). doi:10.1140/epjc/s10052-011-1627-0
- V. del Duca, E. Laenen, Top physics at the LHC. Int. J. Mod. Phys. A 30(35), 1530,063 (2015). doi:10.1142/S0217751X1530063X
- F.P. Schilling, Top quark physics at the LHC: a review of the first two years. Int. J. Mod. Phys. A 27, 1230,016 (2012). doi:10.1142/ S0217751X12300165
- 5. G. Aad et al., Measurement of the $t\bar{t}W$ and $t\bar{t}Z$ production cross sections in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector. JHEP 11, 172 (2015). doi:10.1007/JHEP11(2015)172
- 6. G. Aad et al., Observation of top-quark pair production in association with a photon and measurement of the $t\bar{t}\gamma$ production cross section in pp collisions at $\sqrt{s}=7$ TeV using the ATLAS detector. Phys. Rev. D **91**(7), 072,007 (2015). doi:10.1103/PhysRevD.91. 072007
- 7. G. Aad et al., Search for $H \to \gamma \gamma$ produced in association with top quarks and constraints on the Yukawa coupling between the top quark and the Higgs boson using data taken at 7 TeV and 8 TeV with the ATLAS detector. Phys. Lett. B **740**, 222–242 (2015). doi:10.1016/j.physletb.2014.11.049
- G. Aad et al., Search for the associated production of the Higgs boson with a top quark pair in multilepton final states with the ATLAS detector. Phys. Lett. B 749, 519–541 (2015). doi:10.1016/ j.physletb.2015.07.079
- 9. G. Aad et al., Search for the standard model Higgs boson produced in association with top quarks and decaying into $b\bar{b}$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector. Eur. Phys. J. C **75**(7), 349 (2015). doi:10.1140/epjc/s10052-015-3543-1
- V. Khachatryan et al., Measurement of the inclusive top-quark pair + photon production cross section in the muon + jets channel in pp collisions at 8 TeV. CMS-PAS-TOP-13-011 (2014)



- V. Khachatryan et al., Search for a standard model Higgs boson produced in association with a top-quark pair and decaying to bottom quarks using a matrix element method. Eur. Phys. J. C 75(6), 251 (2015). doi:10.1140/epjc/s10052-015-3454-1
- 13. V. Khachatryan et al., Observation of top quark pairs produced in association with a vector boson in pp collisions at $\sqrt{s} = 8$ TeV. JHEP **01**, 096 (2016). doi:10.1007/JHEP01(2016)096
- 14. G. Aad et al., Search for anomalous couplings in the Wtb vertex from the measurement of double differential angular decay rates of single top quarks produced in the t-channel with the ATLAS detector (2015). arXiv:1510.03764
- 15. G. Aad et al., Measurement of the correlations between the polar angles of leptons from top quark decays in the helicity basis at $\sqrt{s} = 7\text{TeV}$ using the ATLAS detector. Phys. Rev. D **93**(1), 012,002 (2016). doi:10.1103/PhysRevD.93.012002
- T. Aaltonen et al., Combination of CDF and D0 measurements of the W boson helicity in top quark decays. Phys. Rev. D 85, 071,106 (2012). doi:10.1103/PhysRevD.85.071106
- 17. V. Khachatryan et al., Measurement of the W boson helicity in events with a single reconstructed top quark in pp collisions at $\sqrt{s}=8$ TeV. JHEP **01**, 053 (2015). doi:10.1007/JHEP01(2015)053
- 18. G. Aad et al., Measurement of the top quark charge in pp collisions at *sqrts* = 7 TeV in the ATLAS experiment. ATLAS-CONF-2011-141 (2011)
- T. Aaltonen, et al., Exclusion of exotic top-like quarks with -4/3 electric charge using jet-charge tagging in single-lepton ttbar events at CDF. Phys. Rev. D 88(3), 032,003 (2013). doi:10.1103/PhysRevD.88.032003
- V.M. Abazov et al., Measurement of the electric charge of the top quark in *tt* events. Phys. Rev. D **90**(5), 051,101 (2014). doi:10.1103/PhysRevD.90.051101, doi:10.1103/PhysRevD.90.079904 [Erratum: Phys. Rev. D 90, no. 7, 079904(2014)]
- V. Khachatryan, et al., Constraints on the top-quark charge from top-pair events. CMS-PAS-TOP-11-031 (2012)
- 22. J. Brod, A. Greljo, E. Stamou, P. Uttayarat, Probing anomalous $t\bar{t}Z$ interactions with rare meson decays. JHEP **02**, 141 (2015). doi:10. 1007/JHEP02(2015)141
- B. Grzadkowski, M. Misiak, Anomalous Wtb coupling effects in the weak radiative B-meson decay. Phys. Rev. D 78, 077,501 (2008). doi:10.1103/PhysRevD.84.059903, doi:10.1103/ PhysRevD.78.077501. [Erratum: Phys. Rev.D84,059903(2011)]
- J.F. Kamenik, M. Papucci, A. Weiler, Constraining the dipole moments of the top quark. Phys. Rev. D 85, 071,501 (2012). doi:10. 1103/PhysRevD.88.039903, doi:10.1103/PhysRevD.85.071501. [Erratum: Phys. Rev. D 88, no. 3, 039903 (2013)]
- 25. J. Abdallah et al., A study of b anti-b production in e+e collisions at s**(1/2) = 130-GeV 207-GeV. Eur. Phys. J. C **60**, 1–15 (2009). doi:10.1140/epjc/s10052-009-0917-2
- J. de Blas, M. Chala, J. Santiago, Renormalization group constraints on new top interactions from electroweak precision data. JHEP 09, 189 (2015). doi:10.1007/JHEP09(2015)189
- F. Larios, M.A. Perez, C.P. Yuan, Analysis of tbW and ttZ couplings from CLEO and LEP/SLC data. Phys. Lett. B 457, 334–340 (1999). doi:10.1016/S0370-2693(99)00541-9
- S. Schael et al., Precision electroweak measurements on the Z resonance. Phys. Rep. 427, 257–454 (2006). doi:10.1016/j.physrep. 2005.12.006
- J.A. Aguilar-Saavedra, J. Bernabeu, W polarisation beyond helicity fractions in top quark decays. Nucl. Phys. B 840, 349–378 (2010). doi:10.1016/j.nuclphysb.2010.07.012



Eur. Phys. J. C (2016) 76:466 Page 9 of 10 466

- J.A. Aguilar-Saavedra, N.F. Castro, A. Onofre, Constraints on the Wtb vertex from early LHC data. Phys. Rev. D 83, 117,301 (2011). doi:10.1103/PhysRevD.84.019901, doi:10.1103/ PhysRevD.83.117301
- J.A. Aguilar-Saavedra, S.A. dos Santos, New directions for top quark polarization in the *t*-channel process. Phys. Rev. D 89(11), 114,009 (2014). doi:10.1103/PhysRevD.89.114009
- 32. J.A. Aguilar-Saavedra, E. Ivarez, A. Juste, F. Rubbo, Shedding light on the $t\bar{t}$ asymmetry: the photon handle. JHEP **04**, 188 (2014). doi:10.1007/JHEP04(2014)188
- F. Bach, T. Ohl, Anomalous top couplings at hadron colliders revisited. Phys. Rev. D86, 114,026 (2012). doi:10.1103/PhysRevD.86. 114026
- U. Baur, A. Juste, L.H. Orr, D. Rainwater, Probing electroweak top quark couplings at hadron colliders. Phys. Rev. D 71, 054,013 (2005). doi:10.1103/PhysRevD.71.054013
- E.L Berger, Q.H. Cao, I. Low, Model independent constraints among the Wtb, Zb anti-b, and Zt anti-t couplings. Phys. Rev. D 80, 074,020 (2009). doi:10.1103/PhysRevD.80.074020
- C. Bernardo, N.F. Castro, M.C.N. Fiolhais, H. Gonalves, A.G.C. Guerra, M. Oliveira, A. Onofre, Studying the *Wtb* vertex structure using recent LHC results. Phys. Rev. D 90(11), 113,007 (2014). doi:10.1103/PhysRevD.90.113007
- A. Buckley, C. Englert, J. Ferrando, D.J. Miller, L. Moore, M. Russell, C.D. White, Constraining top quark effective theory in the LHC Run II era (2015). arXiv:1512.03360
- O.B. Bylund, F. Maltoni, I. Tsinikos, E. Vryonidou, C. Zhang, Probing top quark neutral couplings in the standard model effective field theory at NLO QCD (2016). arXiv:1601.08193
- 39. Q.H. Cao, B. Yan, J.H. Yu, C. Zhang, A general analysis of *Wtb* anomalous couplings (2015). arXiv:1504.03785
- 40. V. Cirigliano, W. Dekens, J. de Vries, E. Mereghetti, Is there room for CP violation in the top-Higgs sector? (2016). arXiv:1603.03049
- G. Durieux, F. Maltoni, C. Zhang, Global approach to top-quark flavor-changing interactions. Phys. Rev. D 91(7), 074,017 (2015). doi:10.1103/PhysRevD.91.074017
- V.A. Prasath, R.M. Godbole, S.D. Rindani, Longitudinal top polarisation measurement and anomalous *Wtb* coupling. Eur. Phys. J. C 75(9), 402 (2015). doi:10.1140/epjc/s10052-015-3601-8
- S.D. Rindani, P. Sharma, Probing anomalous tbW couplings in single-top production using top polarization at the large hadron collider. JHEP 11, 082 (2011). doi:10.1007/JHEP11(2011)082
- 44. R. Rontsch, M. Schulze, Constraining couplings of top quarks to the Z boson in $t\bar{t}$ + Z production at the LHC. JHEP **07**, 091 (2014). doi:10.1007/JHEP09(2015)132, doi:10.1007/JHEP07(2014)091. [Erratum: JHEP09,132(2015)]
- R. Rontsch, M. Schulze, Probing top-Z dipole moments at the LHC and ILC. JHEP 08, 044 (2015). doi:10.1007/JHEP08(2015)044
- A. Tonero, R. Rosenfeld, Dipole-induced anomalous top quark couplings at the LHC. Phys. Rev. D 90(1), 017,701 (2014). doi:10.1103/PhysRevD.90.017701
- C. Zhang, Effective field theory approach to top-quark decay at next-to-leading order in QCD. Phys. Rev. D 90(1), 014,008 (2014). doi:10.1103/PhysRevD.90.014008
- C. Zhang, Single top production at next-to-leading order in the standard model effective field theory (2016). arXiv:1601.06163
- C. Zhang, N. Greiner, S. Willenbrock, Constraints on non-standard top quark couplings. Phys. Rev. D 86, 014,024 (2012). doi:10.1103/ PhysRevD.86.014024
- J.A. Aguilar-Saavedra, A minimal set of top anomalous couplings. Nucl. Phys. B 812, 181–204 (2009). doi:10.1016/j.nuclphysb.2008. 12.012
- J. Bernabeu, D. Comelli, L. Lavoura, J.P. Silva, Weak magnetic dipole moments in two Higgs doublet models. Phys. Rev. D 53, 5222–5232 (1996). doi:10.1103/PhysRevD.53.5222

 A. Czarnecki, B. Krause, On the dipole moments of fermions at two loops. Acta Phys. Polon. B 28, 829–834 (1997)

- W. Hollik, J.I. Illana, S. Rigolin, C. Schappacher, D. Stockinger, Top dipole form-factors and loop induced CP violation in supersymmetry. Nucl. Phys. B 551, 3–40 (1999). doi:10.1016/ S0550-3213(99)00396-X, doi:10.1016/S0550-3213(99)00201-1. [Erratum: Nucl. Phys. B557,407(1999)]
- K. Agashe, G. Perez, A. Soni, Collider signals of top quark flavor violation from a warped extra dimension. Phys. Rev. D 75, 015,002 (2007). doi:10.1103/PhysRevD.75.015002
- C. Grojean, O. Matsedonskyi, G. Panico, Light top partners and precision physics. JHEP 10, 160 (2013). doi:10.1007/ JHEP10(2013)160
- T. Ibrahim, P. Nath, The top quark electric dipole moment in an MSSM extension with vector like multiplets. Phys. Rev. D 82, 055,001 (2010). doi:10.1103/PhysRevD.82.055001
- T. Ibrahim, P. Nath, The chromoelectric dipole moment of the top quark in models with vector like multiplets. Phys. Rev. D 84, 015,003 (2011). doi:10.1103/PhysRevD.84.015003
- A.L. Kagan, G. Perez, T. Volansky, J. Zupan, General minimal flavor violation. Phys. Rev. D 80, 076,002 (2009). doi:10.1103/ PhysRevD.80.076002
- 59. F. Richard, Can LHC observe an anomaly in $t\bar{t}Z$ production? (2013). arXiv:1304.3594
- K. Melnikov, M. Schulze, A. Scharf, QCD corrections to top quark pair production in association with a photon at hadron colliders. Phys. Rev. D 83, 074,013 (2011). doi:10.1103/PhysRevD.83. 074013
- P.J. Fox, Z. Ligeti, M. Papucci, G. Perez, M.D. Schwartz, Deciphering top flavor violation at the LHC with *B* factories. Phys. Rev. D 78, 054,008 (2008). doi:10.1103/PhysRevD.78.054008
- J. Drobnak, S. Fajfer, J.F. Kamenik, Probing anomalous tWb interactions with rare B decays. Nucl. Phys. B 855, 82–99 (2012). doi:10.1016/j.nuclphysb.2011.10.004
- J.A. Aguilar-Saavedra, B. Fuks, M.L. Mangano, Pinning down top dipole moments with ultra-boosted tops. Phys. Rev. D 91, 094,021 (2015). doi:10.1103/PhysRevD.91.094021
- W. Bernreuther, T. Schroder, T.N. Pham, CP violating dipole form-factors in e+ e- -> anti-t t. Phys. Lett. B 279, 389–396 (1992). doi:10.1016/0370-2693(92)90410-6
- W. Dekens, J. de Vries, J. Bsaisou, W. Bernreuther, C. Hanhart, U.G. Meiner, A. Nogga, A. Wirzba, Unraveling models of CP violation through electric dipole moments of light nuclei. JHEP 07, 069 (2014). doi:10.1007/JHEP07(2014)069
- 66. J. Hernandez-Sanchez, F. Procopio, G. Tavares-Velasco, J.J. Toscano, CP-odd static electromagnetic properties of the W gauge boson and the t quark via the anomalous tbW coupling. Phys. Rev. D 75, 073,017 (2007). doi:10.1103/PhysRevD.75.073017
- M. Schulze, R. Röntsch, Topaz. https://github.com/ TOPAZdevelop/TOPAZ (2014)
- K. Melnikov, M. Schulze, NLO QCD corrections to top quark pair production and decay at hadron colliders. JHEP 08, 049 (2009). doi:10.1088/1126-6708/2009/08/049
- M. Cacciari, G.P. Salam, G. Soyez, The anti-k(t) jet clustering algorithm. JHEP 04, 063 (2008). doi:10.1088/1126-6708/2008/04/063
- R.D. Ball et al., Parton distributions for the LHC Run II. JHEP 04, 040 (2015). doi:10.1007/JHEP04(2015)040
- M.L. Mangano, T. Plehn, P. Reimitz, T. Schell, H.S. Shao, Measuring the top Yukawa coupling at 100 TeV. J. Phys. G 43(3), 035,001 (2016). doi:10.1088/0954-3899/43/3/035001
- J. Pumplin, D.R. Stump, J. Huston, H.L. Lai, P.M. Nadolsky, W.K. Tung, New generation of parton distributions with uncertainties from global QCD analysis. JHEP 07, 012 (2002). doi:10.1088/ 1126-6708/2002/07/012



466 Page 10 of 10 Eur. Phys. J. C (2016) 76:466

A.D. Martin, W.J. Stirling, R.S. Thorne, G. Watt, Parton distributions for the LHC. Eur. Phys. J. C 63, 189–285 (2009). doi:10. 1140/epjc/s10052-009-1072-5

- S. Frixione, V. Hirschi, D. Pagani, H.S. Shao, M. Zaro, Electroweak and QCD corrections to top-pair hadroproduction in association with heavy bosons. JHEP 06, 184 (2015). doi:10.1007/JHEP06(2015)184
- J.H. Kuhn, A. Scharf, P. Uwer, Weak interactions in top-quark pair production at hadron colliders: an update. Phys. Rev. D 91(1), 014,020 (2015). doi:10.1103/PhysRevD.91.014020
- 76. G. Aad et al., Measurement of spin correlation in top–antitop quark events and search for top squark pair production in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector. Phys. Rev. Lett. **114**(14), 142,001 (2015). doi:10.1103/PhysRevLett.114.142001
- 77. V. Khachatryan, et al., Measurements of $t\bar{t}$ spin correlations and top quark polarization using dilepton final states in pp collisions at $\sqrt{s} = 8 \text{ TeV } (2016)$. arXiv:1601.01107

