



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

LLNL-JRNL-741164

A Convenient and Safer Synthesis of Diaminoglyoxime

E. C. Johnson, J. J. Sabatini, N. B. Zuckerman

October 13, 2017

Organic Process and Research Development


A Convenient and Safer Synthesis of Diaminoglyoxime

Eric C. Johnson^a, Jesse J. Sabatini^{*a}, Nathaniel B. Zuckerman^{b*}

^aUS Army Research Laboratory, Energetics Technology Branch, Aberdeen Proving Ground, MD
21005, United States

^bLawrence Livermore National Laboratory, Material Science Division, Livermore, CA
94550, United States

TOC Graphic



KEYWORDS: Energetic Materials, Synthesis, Process Chemistry, Heterocycles

ABSTRACT

A new procedure for the synthesis and isolation of diaminoglyoxime (DAG) is described. A previous procedure involved treating glyoxal with two equivalents each of hydroxylammonium chloride and sodium hydroxide to form glyoxime, followed by further treatment of this intermediate with two additional equivalents of hydroxylammonium chloride and sodium hydroxide at 95 °C to form DAG. Two recrystallizations were needed to obtain the desired product in pure form. Another previous procedure employed glyoxal in the presence of four equivalents each of hydroxylammonium chloride and sodium hydroxide at 95 °C to form DAG. Though this latter procedure gives product after a few hours, yields do not exceed 40%, and the reaction is prone to thermal runaway. Furthermore, the use of decolorizing carbon, and recrystallization of the crude solid is necessary to obtain a pure product.

The new disclosed procedure involves treating a preheated aqueous hydroxylamine solution (50 wt. %, ten equivalents) with aqueous glyoxal (40 wt. %), followed by heating at 95 °C for 72-96 h. The reaction is cooled to room temperature, and then to 0-5 °C to obtain DAG in pure form, without recrystallization or decolorizing carbon in 77-80% yield. The exothermic nature of the reaction is also minimized by this updated process.

INTRODUCTION

Diaminoglyoxime (DAG) is a popular intermediate for the synthesis of a multitude of energetic materials. This not only includes materials stemming from the *bis*-1,2,4-oxadiazole ring system¹, but also the many compounds derived from diaminofurazan (DAF).^{2, 3} Some representative molecules that fall into these two classes of materials are summarized in Figure 1.





Figure 1. Energetic materials obtained from the DAG intermediate.

Though DAG has been synthesized by numerous methods, the most popular methods to-date are via a two-step synthesis⁴ or a one-step synthesis.⁵ In the two-step procedure, as summarized in Scheme 1, glyoxal is first converted to glyoxime (**1**) upon exposure to two equivalents of hydroxylammonium chloride and two equivalents of aqueous sodium hydroxide. Following recrystallization, **1** is then treated with these same reaction conditions, plus exposure to heating for several hours to provide crude DAG (**2**). Crude DAG is recrystallized from boiling water to afford DAG as a yellow crystalline solid. The overall yield for the two-step procedure is 44%.


Scheme 1. Previous two-step synthesis of diaminoglyoxime.



In the one-step procedure (Scheme 2), DAG is obtained as a crude material by heating an aqueous solution of glyoxal, four equivalents of hydroxylammonium chloride and four equivalents of sodium hydroxide for several hours. After isolation of the yellow, crystalline material, the crude product is redissolved in hot water, treated with decolorizing carbon and filtered hot to obtain pure


DAG as a white, crystalline solid. While convenient as a one-step procedure that provides pure material in a matter of a few hours, this method is prone to thermal runaway, and the yields are typically a maximum of ca. 40%.

Scheme 2. Previous one-step synthesis of diaminoglyoxime.



Given the aforementioned yield and lengthy processing issues, the development of a simpler process that yields DAG in a higher yield would provide a significant benefit to the energetic materials community. Such a high yielding and simple process to obtain DAG from glyoxal is given in Scheme 3. In this procedure, an excess of 50% aqueous hydroxylamine is heated to 95 °C with stirring, and a 40% aqueous solution of glyoxal is then added dropwise over an hour. Following addition, the flask is fitted with a reflux condenser, and the reaction mixture is stirred for 72-96 hours. Following this time, the reaction mixture is slowly cooled to room temperature with stirring, and is then further cooled to 0-5 °C with stirring. Filtration of the resulting white crystalline solid affords the product in a greatly improved yield of 77-80%. Treatment with decolorizing carbon and further recrystallizations are not necessary to obtain a pure product.

Scheme 3. New one-step procedure for the synthesis of diaminoglyoxime.



Optimization of the new, all aqueous procedure focused initially on the protocols of previous reports using hydroxylammonium salts in the one-pot formation of DAG from glyoxal.⁶ The glyoxal solution was added to aqueous hydroxylamine (five molar equivalents) in such a manner to keep the reaction below 5 °C. Halfway through the addition, glyoxime precipitated, and following the addition, the mixture was warmed to 20 °C. The thick white precipitate was then heated in one of two manners: heating rate controlled by the temperature of the reaction (T_r) over ten minutes, or ramping the temperature of the heating jacket (T_j) immediately to 95-100 °C. During the initial temperature ramp, the reaction was heterogeneous and the slope of T_r vs time steadily climbed at ~5 °C/min, but at approximately 75 °C, the solids went into solution and the rate of T_r increase doubled. In trials where T_r was used to ramp temperature, heating rate of the jacket was immediately reduced by the control software to limit the reaction temperature rate during this event. The reaction temperature would plateau just around the input temperature and no thermal runaway would occur.⁷ When T_j was used to control the reaction heating, the same pattern of dissolution and heating rate increase were observed while the reaction temperature exceeded the T_r input and gradually reduced to the input temperature. There were no runaway reactions observed in any case where only aqueous reagents were used. At this point, it was determined that more than five equivalents of hydroxylamine was necessary to drive the reaction to completion due to significant discoloration in the final material, and the presence of impurities inseparable by recrystallization. In some cases, where only five equivalents of hydroxylamine was used, discoloration was not observed, but the recovered material had a significantly depressed melting/decomposition point, much closer to that of diaminoglyoxime. Resubmission of this material to heating in additional aqueous hydroxylamine led to a pure, colorless DAG material.

By adding the glyoxal over a period of 30 minutes to the preheated aqueous hydroxylamine ($T_r = 90\text{ }^\circ\text{C}$), there is no precipitation/dissolution of intermediates, and the bulk of the reaction heat is evolved instantaneously as each drop is consumed, minimizing the number of molecules reacting at any given moment. Thus, the likelihood of thermal runaway is mitigated in this procedure by minimizing the presence of reactive species during dissolution and subsequent conversion of glyoxime to DAG. Heat flow calorimetry⁸ (HFC) was used to quantitate heat evolution during the glyoxal addition step. The HFC experiment revealed a heat release of -93.5 kJ/mol over the glyoxal addition period (30 minutes), and immediately following the addition, the reaction temperature quickly returned to the input temperature. Heating was controlled by maintaining the jacket temperature between $95\text{-}100\text{ }^\circ\text{C}$, which translated to approximately $90\text{-}95\text{ }^\circ\text{C}$ in reaction temperature. This new method has been performed reproducibly multiple times on the 50 g scale, and was then subsequently scaled-up successfully at the 250 g scale.

Experimental

Chemicals and solvents were used as received from Sigma-Aldrich. ^1H NMR spectra were recorded using an Anastazi 90 MHz instrument. The chemical shifts quoted in ppm in the text refer to typical standard tetramethylsilane (^1H) in CDCl_3 as the solvent. Infrared spectra were measured with a Bruker Alpha-P FTIR instrument. Decomposition temperatures were measured at a heating rate of $5\text{ }^\circ\text{C/min}$ using a TA Instruments Q10 DSC instrument.

81 g scale: A 1 L round-bottom flask equipped with a stirbar was charged with aqueous hydroxylamine (50 wt %, 569 mL, 8.62 mol) and was immersed in an oil bath. The oil bath was heated to $95\text{ }^\circ\text{C}$, the aqueous hydroxylamine was stirred for 30 minutes, and glyoxal (40 wt. %, 125 mL, 0.862 mol) was added via a pressure-equalizing addition funnel over 30 min. Following

the addition, the reaction was allowed to stir for 72-96 hours. The reaction mixture was removed from the oil bath, and was slowly cooled to 20 °C with stirring, during which time, a white solid appeared. The reaction mixture was immersed into an ice bath and was cooled to 0-5 °C, followed by stirring for one hour. The white solid was collected by Buchner filtration, rinsed with minimal cold water, and vacuum dried overnight to afford 81.3 g (80%) of **2** as a white, crystalline solid; mp = 210.75 °C; ¹H NMR (90 MHz, DMSO-*d*₆) δ 9.73 (s, 2H), 5.17 (s, 4H); IR (neat) cm⁻¹ 3646.82 (m), 3361.46 (s), 1641.62 (m), 1569.84 (s); T_m = 210.75 °C; T_{peak} = 223.71 °C.

250 g: A 2 L glass jacketed reactor with an overhead stirrer was charged with aqueous hydroxylamine (50 wt %, 1.69 L, 27.5 mol) and the jacket was heated to 100 °C. Once the process solution temperature stabilized (~ 85 °C), glyoxal (40 wt. %, 315 ml, 2.74 mol) was added over 1.5 h. Following the addition, the reaction was allowed to stir for 4 d and was then cooled to 20 °C over 20 h with stirring at 40 rpm. Prior to filtering, the reactor was cooled to 0-5 °C over three hours, followed by stirring for three additional hours. Following filtration, the isolated crystals, identified as DAG (**2**), were rinsed with minimal cold water and vacuum dried overnight (240 g, 74 % yield). An additional 9.5 g (3%) of **2** was recovered from the mother liquor upon sitting.

The thermostat, dosing pump, and stirrer were controlled with a Mettler Toledo RX-10, which allowed for a pre-programmed reaction sequence and over-temperature control (automatic cooling if reaction temperature reached a pre-determined set point).

Supporting Information. The synthetic procedure of **2**, its ¹H NMR, IR, and DSC trace are provided in the Supporting Information. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Corresponding Author

*E-mail: jesse.j.sabatini.civ@mail.mil; Phone: 410-278-0235; zuckerman2@llnl.gov; Phone: 925-423-8457

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

ACKNOWLEDGMENT

The authors are indebted to the US Army and to Lawrence Livermore National Laboratory for financial support in carrying out this work. This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. Zuckerman is grateful for financial and support from the DOE/DoD Joint Munitions Program and the DOE Campaign 2 Program.

ABBREVIATIONS

NaOH = sodium hydroxide, NH_2OH = hydroxylamine, $\text{NH}_2\text{OH}\cdot\text{HCl}$ = hydroxylamine hydrochloride, DAG = diaminoglyoxime

REFERENCES

1. Kettner, M. A., Karaghiosoff, K., Klapötke, T. M., Sućeska, M., Wunder, S., *Chem. Eur. J.*, **2014**, *20*, 7622-7631.
2. Boyer, J., Gunasekaran, A. & Trudell, M. *Heteroatom Chemistry*, **1994**, *5*, 441-446.
3. Rakitin, O., Zalesova, A. S., Kulikov, A. S., Makhova, N. N. ; Godovikova, T. I., Khmel'nitskii, L. I., *Rus. Chem. Bull.*, **1993**, *42*, 1865-1870.

4. Gunasekaran, A., Jayachandran, T., Boyer, J., Trudell M. *J. Het. Chem.*, **1995**, *32*, 1405-1407.
5. Trudell, M., Zelenin, A. *J. Het. Chem.*, **1997**, *34*, 1057-1060.
6. An EasyMax 102 or 402 synthesis workstation with iControl 5.4 software (Mettler Toledo) was used to follow and control the reaction conditions.
7. The authors experienced thermal runaway with the hydroxylammonium salt procedures when heat was not actively controlled/removed during the initial heating ramp following dissolution of the glyoxime.
8. A Mettler-Toledo HFCal accessory was used for calorimeter measurements of the glyoxal addition at elevated temperature in the EasyMax 102.