

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

Discovery and functional characterization of a yeast sugar alcohol phosphatase

Journal:	<i>ACS Chemical Biology</i>
Manuscript ID	Draft
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Xu, Yifan; Princeton University, Lu, Wenyun; Princeton University Chen, Jonathan; Princeton University Johnson, Sarah; Princeton University Gibney, Patrick; Cornell University Thomas, David; Princeton University Brown, Greg; University of Toronto, BBDMR May, Amanda; University of Tennessee Campagna, Shawn; University of Tennessee Yakunin, Alexander; University of Toronto, Department of Chemical Engineering and Applied Chemistry Botstein, David; Princeton University Rabinowitz, Joshua; Princeton University

SCHOLARONE™
Manuscripts

Discovery and functional characterization of a yeast sugar alcohol phosphatase

Yi-Fan Xu^{1,2}, Wenyun Lu¹, Jonathan C. Chen^{1,2}, Sarah A. Johnson^{1,2}, Patrick A. Gibney¹, David G. Thomas¹, Greg Brown⁴, Amanda L. May⁵, Shawn R. Campagna⁵, Alexander F. Yakunin⁴, David Botstein^{1,3}, Joshua D. Rabinowitz^{1,2*}

¹Lewis Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ 08544, USA.

²Department of Chemistry, Princeton University, Princeton, NJ 08544, USA. ³Department of Molecular Biology, Princeton University, Princeton, NJ 08544, USA

⁴Department of Chemical Engineering and Applied Chemistry, Banting and Best Department of Medical Research, University of Toronto, Toronto, ON M5S 3E5, Canada

⁵Department of Chemistry, University of Tennessee, Knoxville, TN 37996, USA

*To whom correspondence should be addressed. E-mail: joshr@princeton.edu

- Yi-Fan Xu and Wenyun Lu contributed equally to this work

Abstract

Sugar alcohols (polyols) exist widely in nature. While some specific sugar alcohol phosphatases are known, there is no known phosphatase for some important sugar alcohols (e.g., sorbitol-6-phosphate). Using liquid chromatography-mass spectrometry-based metabolomics, we screened yeast strains with putative phosphatases of unknown function deleted. We show that the yeast gene *YNL010W*, which has close homologues in all fungi species and some plants, encodes a sugar alcohol phosphatase. We term this enzyme, which hydrolyzes sorbitol-6-phosphate, ribitol-5-phosphate, and (D)-glycerol-3-phosphate, polyol phosphatase 1 or *PYP1*. Polyol phosphates are structural analogs of the enediol intermediate of phosphoglucose isomerase (Pgi). We find that sorbitol-6-phosphate and ribitol-5-phosphate inhibit Pgi and that Pyp1 activity is important for yeast to maintain Pgi activity in the presence of environmental sugar alcohols. Pyp1 expression is strongly positively correlated with yeast growth rate, presumably because faster growth requires greater glycolytic and accordingly Pgi flux. Thus, yeast express the previously uncharacterized enzyme Pyp1 to prevent inhibition of glycolysis by sugar alcohol phosphates. Pyp1 may be useful for engineering sugar alcohol production.


Introduction

Polyols, the reduced forms of sugars, are an important natural family of carbohydrates¹⁻². Glycerol, the simplest polyol, is the backbone of phospholipids and is excreted by many microbes in response to stress³⁻⁴. Longer chain polyols include erythritol, ribitol, xylitol, arabitol, sorbitol, and mannitol, all of which exist only in the (D)-form in nature and are usually found in plants. Due to the inability of most organisms to assimilate long chain polyols into glycolysis, they are often regarded as inert solutes with unclear physiological functions. While the biological function of polyols has remained obscure, they have gained commercial interest in the last decade due to their increased usage as a sugar substitute by the food industry⁵.

Many fungi species are able to produce polyols via reduction of the corresponding sugar⁶. To engineer polyol production, the standard approach is to first dephosphorylate the sugar phosphate and then reduce the resulting sugar to the polyol⁷⁻¹⁰. This engineering approach differs from the natural glycerol production pathway, where (L)-glycerol-3-phosphate (a.k.a. *sn*-glycerol-3-phosphate) is first made from the reduction of dihydroxyacetone phosphate, followed by the dephosphorylation of the phosphorylated polyol¹¹⁻¹². This natural approach avoids accumulation of the potentially toxic metabolite dihydroxyacetone (DHA). Moreover, it avoids making a dephosphorylated sugar that might escape from the cell. However, this series of reactions has not been feasible as an engineering strategy due to lack of a suitable polyol phosphatase.

Because many enzymes remain unannotated even in the best studied organisms such as Baker's yeast¹³ it is possible that various polyol phosphate phosphatases exist but have yet to be characterized. One approach to metabolic enzyme annotation is to knock out the corresponding gene and assess the metabolome of the resulting strain. In general, substrates of the enzyme are likely to increase and products to decrease. This approach has now been successfully used to assign function to a variety of enzymes¹⁴⁻²².

Using this approach and subsequent biochemical studies, here we show that *YNL010W*, a gene conserved across all fungi species and some plants, encodes a polyol phosphatase (Pyp1). We further show that the enzyme prevents polyol phosphate accumulation in yeast, and that this is physiologically important due to polyols being Pgi transition state analogues (**Scheme 1**). Thus, through assigning function to this previously unannotated yeast gene, we identify both a new glycolytic regulatory mechanism and a promising new enzyme for polyol metabolic engineering.

**Scheme 1**

Results

A yeast *Pyp1* deletion mutant accumulates octulose-8-phosphate and polyol phosphates

We screened yeast strains, with putative phosphatases of unknown function deleted, for changes in metabolite concentrations. Yeast were grown in glucose minimal media, and metabolites were extracted into 40:40:20 methanol: acetonitrile: water, followed by metabolome analysis by reversed-phase ion-pairing liquid chromatography – high resolution mass spectrometry (LC-MS). We found that the deletion of *PYP1*, formerly known as the uncharacterized gene *YNL010W*, while not significantly altering the concentrations of most metabolites, led to the statistically significant (false discovery rate < 0.05) accumulation of three compounds in negative ion mode (**Figure 1A**). The metabolites' formulae were obtained by labeling cells with ^{13}C and ^{15}N and observing no shift from nitrogen labeling and a shift of +8, +5, or +4 daltons from carbon labeling (**Figure 1B**)²³. Exact masses of these compounds matched putative formulae of $\text{C}_8\text{H}_{17}\text{O}_{11}\text{P}_1$, $\text{C}_5\text{H}_{13}\text{O}_8\text{P}_1$ and $\text{C}_4\text{H}_{11}\text{O}_7\text{P}_1$. Searching for these formulae in the KEGG database returned the metabolites octulose-8-phosphate ($\text{C}_8\text{H}_{17}\text{O}_{11}\text{P}_1$); ribitol-5-phosphate, arabitol-5-phosphate, and xylitol-5-phosphate ($\text{C}_5\text{H}_{13}\text{O}_8\text{P}_1$); and erythritol-4-phosphate ($\text{C}_4\text{H}_{11}\text{O}_7\text{P}_1$) (**Figure 1C**).

To identify the five- and four-carbon polyol phosphates, we performed an isotope labeling experiment involving switching *pyp1Δ* cells from U- ^{13}C -glucose to unlabeled ribitol, arabitol, xylitol or erythritol. Although baker's yeast cannot effectively utilize these polyols as carbon sources, in the absence of glucose, they slowly transport and phosphorylate them. We found that feeding ribitol and erythritol resulted in the build-up of intracellular metabolites with exact mass and LC retention time matching the $\text{C}_5\text{H}_{13}\text{O}_8\text{P}_1$ and $\text{C}_4\text{H}_{11}\text{O}_7\text{P}_1$ that accumulated with *Pyp1* deletion (**Figure 1D**). Feeding of arabitol and xylitol resulted in the accumulation of polyol phosphates with different retention times (**Figure 1D**). Based on these results, we synthesized ribitol-5P and confirmed that exact mass and retention time matched to the endogenous 5-carbon sugar alcohol (**Figure 1E**). Octulose-8P has been reported in yeast previously¹⁸, and the chromatographic retention time of the accumulated $\text{C}_8\text{H}_{17}\text{O}_{11}\text{P}_1$ exactly matched the synthetic octulose-8P standard (**Figure 1F**). Thus, the compounds that accumulate in the *pyp1Δ* strain are octulose-8P, ribitol-5P and erythritol-4P.

Ribitol-5P and erythritol-4P are polyol phosphates, but octulose-8P is a sugar phosphate. We were curious why *Pyp1* deletion resulted in accumulation of metabolites from these different structural families. The most common conformer of the seven carbon sugar sedoheptulose and its derivatives in solution is β -furanose²⁴. Octulose-8P likely has a similar β -furanose ring structure (**Figure 1C**), which

1 has three carbons (6'-8') of octulose out of the ring, forming a tail that resembles (D)-glycerol-3P, a
2 polyol phosphate. Note that (D)-glycerol-3P is the same compound as (L)-glycerol-1P, which is the
3 enantiomer of the common central carbon metabolite (L)-glycerol-3P (a.k.a. sn-glycerol-3P). We use
4 the (D)-glycerol-3P nomenclature to emphasize the structural similarity to longer (D)-polyol phosphates,
5 including ribitol-5P and sorbitol-6P (**Figure 1C**).

12 **Recombinant Pyp1 dephosphorylates ribitol-5-phosphate, sorbitol-6-phosphate, and (D)-** 13 **glycerol-3-phosphate**

14 Although polyols are excretion products of fungi and plants, polyol phosphates were not previously
15 thought to be an intermediate in this pathway (sugar phosphate → sugar → polyol). To determine if the
16 accumulated compounds were indeed the substrates of Pyp1, we analyzed the biochemical activity of
17 the purified recombinant Pyp1 on these compounds. Incubation with Pyp1 led to the depletion of ribitol-
18 5P and the accumulation of ribitol (**Figure 2A**), whereas no detectable phosphatase activity was found
19 against octulose-8P or erythritol-4P (**Figure 2B**). Because flux through octulose-8P and erythritol-4P is
20 very low in cells, minimal Pyp1 activity may nevertheless be sufficient to alter the cellular concentration
21 ¹⁸. Alternatively, Pyp1 may need to bind an activator or other substrate in order to hydrolyze these
22 species, or the accumulation of octulose-8P and erythritol-4P may be an indirect consequence of Pyp1
23 loss (e.g. occurring secondary to build-up of a more preferred substrate).

24 To better define the range of substrates of Pyp1, we also measured its activity against erythrose-4P,
25 sorbitol-6P, racemic glycerol-3P, and (L)-glycerol-3P. Incubation with Pyp1 led to the hydrolysis of
26 erythrose-4P, sorbitol-6P and racemic glycerol-3P, but not (L)-glycerol-3P (**Figure 2B**). Thus, unlike the
27 known glycerol-3-phosphatases (Hor2 and Rhr2) in the glycerol biosynthetic pathway, which act on
28 both (L) and (D)-glycerol-3P ¹², Pyp1 is specific to (D)-glycerol-3P. No other phosphatase activity was
29 found for other common phosphorylated metabolites (see **Supplementary Table 1**). The best
30 biochemical substrates for Pyp1 do not perfectly match those that accumulated in cells, likely because
31 cellular accumulation depends on the absence of other routes of metabolizing the compounds.

32 **Source of ribitol-5P in glucose-grown yeast**

1 Although polyol phosphates have not been previously described in Baker's yeast, they have been noted
2 in other fungi²⁵. There are two likely routes for their cellular biosynthesis. The first involves the
3 reduction of the corresponding sugar phosphate and the second involves phosphorylation of polyols.
4 Both activities have been described in fungi but the responsible genes remain unknown²⁵. In Baker's
5 yeast grown on glucose, the only known polyol is glycerol, and thus polyol phosphates are likely made
6 from the reduction of sugar phosphates (as in *Haemophilus influenza*, *Staphylococcus aureus*²⁶, and
7 *Escherichia coli*²⁷). Consistent with this, deletion of *Zwf1*, the first step in the oxidative pentose
8 phosphate pathway (PPP), eliminated the endogenous peak for ribitol-5P, presumably due to
9 decreased levels of ribose-5P and ribulose-5P (**Figure 3**). This is consistent with ribitol-5P being made
10 via the reduction of ribose-5P or ribulose-5P in yeast grown on glucose.
11
12
13
14
15
16
17
18
19
20

21 **Growth on sorbitol requires Pyp1**

22
23 When polyols are present in the growth environment, polyol phosphates can be made by their
24 phosphorylation. Of the natural long-chain polyols, the only one known to support yeast growth as the
25 sole carbon source is sorbitol²⁸. Catabolism of sorbitol begins with its dehydrogenation to fructose,
26 which then enters central metabolism. Due to the structural similarity of sorbitol to glucose, we
27 hypothesized that yeast might sometimes erroneously phosphorylate sorbitol into sorbitol-6P, which
28 could be toxic in excess. To explore this possibility, we fed both wild type and *pyp1Δ* strains minimal
29 media containing sorbitol as the sole carbon source. While wild type cells grew to saturation after a long
30 lag phase, *pyp1Δ* cells never grew (**Figure 4A**). Metabolome profiling revealed that sorbitol-6P levels in
31 the *pyp1Δ* strain were much higher than the wild type strain (**Figure 4B**). These results are consistent
32 with Pyp1 being required to dephosphorylate sorbitol-6P, which otherwise accumulates to toxic levels,
33 thereby precluding growth on sorbitol.
34
35
36
37
38
39
40

41 To investigate whether Pyp1 is important for growth on other polyol substrates, we fed *pyp1Δ* yeast
42 glycerol or mannitol. The *pyp1Δ* cells grew poorly on glycerol. While wild type cells managed to reach
43 saturation after a long lag phase on mannitol, *pyp1Δ* cells were unable to grow on mannitol (**Figure 4C**).
44 Thus, Pyp1 contributes to polyol phosphate detoxification, and such detoxification is required for growth
45 on diverse polyol substrates.
46
47
48
49

50 To more directly ascertain whether the impaired growth was a result of polyol phosphate toxicity, we fed
51 yeast trehalose (a dimer of glucose that can be slowly utilized by yeast resulting in carbon-limited
52
53

1 growth²⁹⁻³⁰) with or without addition of ribitol. In wild type yeast, addition of ribitol to the medium resulted
2 in accumulation of intracellular ribitol-5P, without impacting growth rate (**Figure 4D**). In *pyp1* yeast, the
3 ribitol-5P levels rose yet higher, resulting in a ~20% decrease in the growth rate (**Figure 4D**). Thus,
4 buildup of polyol phosphate compounds impairs yeast growth, including on substrates other than
5 polyols.
6
7
8
9

10 11 12 **Polyol phosphates are inhibitors of phosphoglucose isomerase (Pgi)**

13
14
15 One potential mechanism by which polyol phosphates could impair cell growth is through metabolic
16 enzyme inhibition. Phosphoglucose isomerase (Pgi1) catalyzes the interconversion between glucose-
17 6P and fructose-6P in glycolysis and is essential for yeast to grow on either glucose or fructose as the
18 sole carbon source³¹. Sorbitol-6P is structurally similar to the enediol transition state of the Pgi reaction
19 (**Scheme 1**) and sorbitol-6P and other structural mimics of the enediol intermediate are known Pgi
20 inhibitors³². We confirmed that sorbitol-6P and less potently ribitol-5P inhibit Pgi (**Figure 5A**). Motivated
21 by these observations, we sought to determine whether polyol phosphates significantly and selectively
22 inhibit Pgi in yeast cells. We hypothesized that two metabolic hallmarks of Pgi inhibition would be
23 increased glucose-6P and decreased glycolytic intermediates downstream of fructose-6P, with
24 fructose-1, 6-bisphosphate (FBP) a convenient marker compound. To evaluate whether decreased Pgi
25 activity indeed induces these metabolic changes, we constructed a yeast strain with PGI under control
26 of an estradiol-inducible promoter. In the low induction condition (1 nM estradiol, Pgi protein level ~1/7
27 of WT cells), we observed both increased glucose-6P and decreased FBP (**Figure 5B**).
28
29
30
31
32
33
34
35

36 We then tested whether ribitol-5P accumulation results in these metabolic hallmarks of Pgi deficiency.
37 Cells were initially grown in trehalose + ribitol and then switched to glucose + ribitol to enhance
38 glycolytic flux. Relative to wild-type yeast, the *pyp1Δ* strain accumulated dramatically more ribitol-5P
39 (**Figure 5C**). Critically, the yeast lacking Pyp1 also manifested both hallmarks of physiological Pgi
40 inhibition, increased glucose-6P and decreased FBP (**Figure 5C**).
41
42
43
44
45
46
47
48
49
50
51
52
53

Discussion

Sorbitol-6-phosphatase activity has been found in apple leaves³³ and silk worms³⁴. The gene encoding this enzyme, however, has not been discovered. In engineered fungi, sorbitol production has been achieved by expressing bacterial sorbitol-6P dehydrogenase, but again the gene encoding the required phosphatase activity remained missing³⁵. Here we identify the previously unannotated yeast gene *YNL010W*, which we now term *PYP1*, as a polyol phosphate phosphatase. Pyp1 dephosphorylates a variety of compounds with the common structural motif of a (D)-glycerol-3-phosphate tail, including (D)-glycerol-3P, erythrose-4P, ribitol-5P, and sorbitol-6P (**Figure 2B**). It is likely that Pyp1 would also dephosphorylate erythritol-4P, arabitol-5P, and xylitol-5P. The ability of Pyp1 to hydrolyze a variety of sugar alcohol phosphates gives rise to an intriguing opportunity to employ Pyp1 in the production of diverse sugar alcohol consumables.

One functional role of Pyp1 is to limit the polyol phosphate concentrations in cells. High levels of polyol phosphates impair yeast growth, at least in part by inhibition of the upper glycolytic enzyme phosphoglucose isomerase (Pgi), for which sorbitol-6-phosphate is a transition state analogue. Other sugar phosphate isomerases, such as triose-phosphate isomerase and ribose-5-phosphate isomerase, have similar enediol reaction intermediates³⁶⁻³⁷, and thus may also be inhibited by polyol phosphates. Similar to strong inhibition of Pgi by sorbitol-6 phosphate, triose-phosphate isomerase may be inhibited by (D)-glycerol-3P, and ribose-5-phosphate isomerase by ribitol-5P or xylitol-5P. Flux through each of these enzymes tends to increase with faster yeast growth rate. For example, ribose-5-phosphate isomerase is required to feed ribosome biogenesis. Thus rapidly growing yeast cells may be particularly sensitive to enzyme inhibition by polyol phosphates. To determine whether Pyp1 function is associated with growth rate, we analyzed its expression as a function of growth rate across 25 different chemostat conditions³⁸. Interestingly, Pyp1's expression was strongly positively correlated with growth rate, regardless of the limiting nutrient used (top 6% of all transcripts in genome) (**Figure 6**). Thus, unlike most protective or detoxification genes (e.g. against heat, osmolarity or oxidative stress), which are highly expressed under slower growth conditions³⁸⁻³⁹, *PYP1* is a fast-growth gene that maintains high Pgi flux by dephosphorylating polyol phosphates.

The transition state inhibition by polyol phosphates and their derivatives is not limited to sugar phosphate isomerase. In plants, ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO) also has an enediol intermediate and is known to be inhibited by 2-carboxy-D-arabitol-1-phosphate (CA1P)⁴⁰.

1 The corresponding 2-carboxy-D-arabitol-1-phosphatase is activated by light and controls cellular level
2 of CA1P. In darkness, RuBisCO is inhibited and also protected by CA1P from proteolysis⁴¹.
3
4

5 Pyp1 is conserved across fungi species and some plants and bacteria, including *Ricinus communis*,
6 *Dehalococcoides ethenogenes*, and *Bacillus anthracis*⁴². The highly conserved nature of Pyp1 in fungi
7 indicates the importance of clearing polyol phosphates. It is likely that the homologous enzymes play
8 the same role in bacteria and plants. Although we have not identified a Pyp1 homolog in mammals, nor
9 are linear polyol phosphates longer than three carbons known mammalian metabolites, polyols are also
10 produced in humans and can accumulate in disease⁴³. For example, during hyperglycemia (e.g., due to
11 diabetes), sorbitol accumulates in a number of organs due to the action of aldose reductase on glucose,
12 which may contribute to diabetic complications including diabetic retinopathy, peripheral neuropathy
13 and diabetic kidney disease⁴⁴⁻⁴⁶. It is unclear whether such sorbitol sometimes becomes
14 phosphorylated to sorbitol-6P and, if so, whether sorbitol-6P contributes to disease pathology.
15
16
17
18
19
20
21

22 Beyond their potential cellular toxicity, polyol phosphates may also have a regulatory role in central
23 carbon metabolism. Pgi sits at the branch point between glycolysis and the oxidative pentose
24 phosphate pathway. Despite being ideally situated to regulate the branching ratio between these
25 pathways (**Figure 5**), Pgi is not an allosteric enzyme, with no physiological regulators known^{32, 47}. It is
26 tempting to speculate that polyol phosphates might serve as endogenous Pgi regulators, with Pgi an
27 enzyme controlled by active site competition⁴⁸⁻⁵³. Further investigation is merited to see whether polyol
28 phosphates contribute to physiological metabolic flux control.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

Experimental Procedures

Chemicals and reagents

Most chemicals, reagents and media components in the study, including (D)-sorbitol-6-phosphate (S1753), (D)-ribose-5-phosphate (83875), (D)-erythritol-4-phosphate (08897), (D)-erythrose-4-phosphate (E0377), (DL)-glycerol-3-phosphate (61668), (L)-glycerol-3P (94124), (D)-glycerol-3P (92034), NaBH₄ (452882), (D)-sorbitol (S1876), ribitol (adonitol, A5502), (D)-arabitol (A3381), xylitol (X3375), erythritol (PHR1479), tributylamine (90780), and acetic acid (695092), were obtained from Sigma-Aldrich (St. Louis, MO). LCMS-grade acetonitrile (A955), methanol (A456), water (W6) were obtained from Fisher Scientific (Waltham, MA). ¹³C₆-glucose (CLM-1396) was from Cambridge Isotope Laboratories (Tewksbury, MA).

Yeast strains and media

Yeast strains were derived from prototrophic S288C. Prototrophic deletions were created by homologous recombination using the allele amplified by PCR from the synthetic genetic analysis (SGA) deletion set⁵⁴. The Pgi1 inducible strain was generated as described:⁵⁵ human estradiol was used to induce *PGI1* via insertion of a synthetic promoter involving chimeric transcription factor-estrogen receptor in front of *PGI1*. 1 nM and 5 nM estradiol were added to achieve low and high expression levels of yeast Pgi1, resulting in a 2.5-fold difference in Pgi1 protein level⁵⁵.

Cells were grown in minimal media comprising 6.7g/L Difco Yeast Nitrogen Base without amino acids plus 2% (w/v) glucose, sorbitol or mannitol. Glycerol medium was composed of 6.7g/L Difco Yeast Nitrogen Base without amino acids, 0.79 g/L Complete Supplement Mixture (Sunrise Science, San Diego, CA) and 3% (v/v) glycerol. Trehalose minimal medium was prepared by mixing 6.7g/L Difco Yeast Nitrogen Base without amino acids and 1% (w/v) trehalose, and adjusting pH to 4.8 by adding succinic acid.

Yeast culture conditions and extraction

The metabolome of batch culture *Saccharomyces cerevisiae* was characterized as described previously⁵⁶: Saturated overnight cultures were diluted 1:30 and grown in liquid media in a shaking flask to A₆₀₀ of ~ 0.6. A portion of the cells (3 mL) were filtered onto a 50 mm nylon membrane filter (Millipore, Billerica, MA), which was immediately transferred into -20 °C extraction solvent (40:40:20 acetonitrile/methanol/water). For carbon upshift, 100 mL of cell culture grown on trehalose at A₆₀₀ of

1 ~0.6 were poured onto a 100 mm cellulose acetate membrane filter (Sterlitech, Kent, WA) resting on a
2 vacuum filter holder with a 1000 mL funnel (Kimble Chase, Vineland, NJ) and were washed with 100
3 mL pre-warmed (30°C) glucose minimal medium. Immediately after the wash media went through, the
4 filter was taken from the holder and the cells were washed into a new flask containing 100 mL pre-
5 warmed (30 °C) glucose minimal medium. Samples were then taken at the indicated time points after
6 the switch, and filtered and quenched as described above.
7
8
9
10

11 12 13 **LC-MS metabolite measurement**

14
15 Cell extracts were analyzed by reversed phase ion-pairing liquid chromatography (LC) coupled by
16 electrospray ionization (ESI) (negative mode) to a high-resolution, high-accuracy mass spectrometer
17 (Exactive; Thermo Fisher Scientific, Waltham, Massachusetts) operating in negative ion mode scanning
18 m/z 70-1000 with 100,000 resolution at m/z 200. Liquid chromatography separation was achieved on a
19 Synergy Hydro-RP column (100 mm × 2 mm, 2.5 μm particle size, Phenomenex, Torrance, CA) with
20 the following gradient: 0 min, 0% B; 2.5 min, 0% B; 5 min, 20% B; 7.5 min, 20% B; 13 min, 55% B; 15.5
21 min, 95% B; 18.5 min, 95% B; 19 min, 0% B; 25 min, 0% B. Solvent A is 97:3 water/methanol with 10
22 mM tributylamine and 15 mM acetic acid; solvent B is methanol. Other LC parameters are autosampler
23 temperature 4 °C, injection volume 10 μL, and column temperature 25 °C. Peaks differing between
24 wild-type and *pyp1Δ* strain were determined using the in-house developed, open-source software
25 MAVEN⁵⁷⁻⁵⁸. Compounds' identities were verified by mass and retention time match to authenticated
26 standards. Differences in metabolome between wild type and *pyp1Δ* strains were tested for significance
27 using Student's T-test. The resulting P-values were then corrected using the Benjamini-Yekutieli False
28 Discovery Rate (FDR) model⁵⁹.
29
30
31
32
33
34
35
36
37

38 The number of C and N atoms in each accumulated compound was determined by the method
39 described²³. Yeast batch cultures were grown with uniformly labeled glucose or ammonium sulfate
40 (Cambridge Isotopes, Andover, MA) for > 20 generations to ensure complete labeling of the
41 metabolome.
42
43
44

45 Absolute intracellular metabolite concentrations in steadily growing wild type and *pyp1Δ* cells were
46 determined as described previously⁶⁰. Metabolite concentrations after perturbations were computed
47 based on fold-change in ion counts relative to steadily-growing cells (grown and analyzed in parallel)
48 multiplied by the known absolute concentration in the steadily growing cells, as determined using an
49 isotope ratio-based approach⁶¹.
50
51
52
53

Synthesis of ribitol-5-phosphate and octulose-8-phosphate

Synthesis of ribitol-5-phosphate was performed as described previously through the reduction of ribose-5-phosphate using NaBH_4 ⁶². Synthesis of (D)-glycero-D-altro-octulose-8-phosphate was performed enzymatically as described⁶³.

Protein purification and enzymatic assays

For Pyp1's polyol phosphatase activity assay, a yeast Open Reading Frame (ORF) strain with an expression vector containing C-terminal His-tagged *PYP1* (Open Biosystems, Thermo Fisher Scientific, San Jose, CA) was grown on galactose to induce Pyp1 expression. The resulting cells were lysed using glass beads and His-tagged Pyp1 was purified using Qiagen Ni-NTA spin columns according to the manufacturer's instructions. Phosphatase activity against ribitol-5-phosphate, sorbitol-6-phosphate, erythritol-4-phosphate and octulose-8-phosphate was determined by monitoring the increase of ribitol, sorbitol, erythritol or octulose using LC-MS. The reaction mixture contained 100 mM Tris-HCl at the pH 8.0, 10 mM MgCl_2 and a range of substrate concentrations (0.01 to 5 mM).

Purified Pyp1 was also assayed against other compounds with similar structures and 90 common phosphorylated metabolites (see **Supplementary Table** for the list of compounds). Briefly, the gene encoding Pyp1 was amplified by PCR using *S. cerevisiae* genomic DNA. The amplified fragments were cloned into a modified pET15b vector (Novagen, Darmstadt, Germany) and overexpressed in the *E. coli* BL21 (DE3) Gold strain (Stratagene, La Jolla, California) as previously described⁶⁴. The recombinant protein was purified using metal ion affinity chromatography on nickel chelate resin (Qiagen, Hilden, Germany) to high homogeneity and stored at $-80\text{ }^\circ\text{C}$. Purified Pyp1 was then screened for phosphatase activity as described previously⁶⁵. Due to the difficulty of obtaining (D)-glycerol-3P, the activity of Pyp1 on (D)-glycerol-3P was determined using racemic glycerol-3P as the substrate. Because Pyp1 has minimal activity on (L)-glycerol-3P, its activity on racemic glycerol-3P was taken as the activity on (D)-glycerol-3P. Compounds with specific activity higher than $0.1\text{ }\mu\text{mol}/\text{mg}/\text{min}$, as well as erythritol-4P, are shown in **Figure 2B**.

To measure the inhibition of Pgi by sorbitol-6-phosphate and ribitol-5-phosphate, yeast Pgi1 was purchased from Sigma Aldrich (St. Louis, MO). Phosphoglucose isomerase activity was determined by adding fructose-6-phosphate and monitoring the appearance of glucose-6-phosphate using LC/MS. We

1 found such LC/MS based assay is consistently more sensitive and accurate than the more typical
2 colorimetric-based assay, which involves coupling the Pgi1 activity with glucose-6-phosphate
3 dehydrogenase activity and monitoring the appearance of NADH. The reaction mixture contained 100
4 mM Tris-HCl at pH 8.0, 10 mM MgCl₂, 0.6 mM fructose-6-phosphate (the physiological concentration in
5 cells grown exponentially on glucose), and a range of sorbitol-6-phosphate and ribitol-5-phosphate
6 concentrations (0.05 to 5 mM). The resulting data were fitted to the Hill equation using the GraphPad
7 Prism Software.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53


Acknowledgments

We thank Scott McIsaac, Sanford Silverman, Sean Hackett and Xiaoyang Su for helpful discussions. This research was funded by NSF CAREER award MCB-0643859, Joint DOE-AFOSR Award DOE DE-SC0002077 - AFOSR FA9550-09-1-0580, NSF grant CBET-0941143, NIH R01 grant CA163591, and DOE Center for Advanced Bioenergy and Bioproducts Innovation (CABBI) DE-SC0018420 to J.D. R. and CA211437 to W.L., with additional support from the Princeton University Center for Quantitative Biology (P50 GM071508) and from the Government of Canada through Genome Canada, Ontario Genomics Institute (2009-OGI-ABC-1405), and Ontario Research Fund (ORF-GL2-01-004), and from NSF grant OCE-1233964.

Author Contributions:

Y.F.X., J.C.C., W.Y.L., D.B., and J.D.R. designed experiments and analyses. Y.F.X. and J.C.C. performed metabolomics, strain construction, yeast physiology and biochemistry. W.Y.L. contributed to the identification of polyol phosphates and biochemistry. S.A.J., A.L.M. and S.R.C. contributed to organic synthesis and biochemistry. G.B. and A.F.Y. contributed to biochemistry. D.G.T. and P.A.G. contributed to strain constructions. Y.F.X., W.Y.L. and J.D.R. wrote the paper with input from all authors.

Figure Legends:

Figure 1. Metabolomic phenotype of *pyp1*Δ cells.

(A). Metabolite profiles from glucose-grown wild type and *pyp1*Δ yeast cells detected by untargeted negative ion mode LC-high resolution MS. **1**, **2**, and **3** represent compounds with significantly different signal intensity between wild type and *pyp1*Δ cells. The x-axis represents the average of signal intensity in WT cells ($N = 3$). The y-axis represents the average of signal intensity in *pyp1*Δ cells ($N = 3$). Adduct

1 and isotopic peaks were excluded. **(B)**. Impact of ^{13}C - and ^{15}N -labeling on peaks of interest.
2
3 Compounds **1**, **2** and **3** were assigned formula as $\text{C}_8\text{H}_{17}\text{O}_{11}\text{P}_1$, $\text{C}_5\text{H}_{13}\text{O}_8\text{P}_1$ and $\text{C}_4\text{H}_{11}\text{O}_7\text{P}_1$ respectively.
4 **(C)**. Chemical structures of octulose-8P, ribitol-5P and erythritol-4P. **(D)**. Extracted ion chromatograms
5 of polyol phosphates produced from ^{13}C -glucose (above) or by phosphorylation of ^{12}C -polyols fed to the
6 *pyp1* Δ cells (below). *pyp1* Δ cells growing exponentially on 2% ^{13}C -glucose was switched to 2% ^{12}C -
7 polyols. Yeast metabolome right before the switch and four hours after the switch were analyzed by LC-
8 MS. Retention time identifies the glucose-derived five carbon polyol phosphate as ribitol-5-phosphate
9 and the four carbon polyol phosphate as erythritol-4P. **(E and F)**. Extracted ion chromatogram for
10 endogenous ribitol-5P **(E)** and octulose-8P **(F)** compared to synthetic standards.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53




Figure 2. Pyp1 hydrolyzes (D)-polyol phosphates.

(A). Purified Pyp1's phosphatase activity was assayed against 0.5 mM ribitol-5P by LC-MS. Incubation with Pyp1 led to the depletion of ribitol-5P (left panel) and the accumulation of ribitol (right panel). (B). Pyp1's (D)-polyol phosphatase activity was measured in the presence of 0.5 mM substrate and 5 mM Mg^{2+} (pH = 7.0, 30°C) by monitoring the appearance the polyol. See **Experimental Procedures** for details. The y-axis represents specific phosphatase activity ($\mu\text{ moles product per min per mg of protein}$) (mean \pm range, N = 2).




Figure 3. Intracellular levels of ribitol-5-phosphate and ribulose-5-phosphate/xylulose-5-phosphate in wild type and *zwf1*Δ strain.

The metabolomes of wild type and *zwf1*Δ strains growing on 2% glucose were measured by LC-MS. The y-axis represents the relative level for select metabolites (mean ± range, N = 2).




Figure 4. Pyp1 accelerates yeast growth on or in the presence of polyols.

(A). Growth of wild type and *pyp1*Δ yeast on sorbitol. Yeast cells grown on YPD medium were switched to minimal media containing 2% sorbitol as the carbon source. The x-axis represents days after the switch and the y-axis represents optical density (A_{600}). (B). Absolute concentration of sorbitol-6-phosphate in wild type and *pyp1*Δ cells in the experiment shown in (A). The x-axis represents days after the switch and the y-axis represents absolute intracellular concentration (mean \pm range, N = 2). (C). Growth of wild type and *pyp1*Δ cells on glycerol and mannitol. Yeast cells grown on YPD medium were switched to medium containing complete supplement mixture (CSM) and 3% glycerol as the carbon source or minimal medium containing 2% mannitol as the carbon source. The x-axis represents time after the switch and the y-axis represents optical density (A_{600}). (D). Impact of ribitol in the presence of trehalose as the carbon source. Absolute concentration of ribitol-5P in wild type and *pyp1*Δ cells (left panel) and their growth rates (right panel) on minimal media containing 1% trehalose +/- 1% ribitol.




Figure 5. Polyol phosphates inhibit phosphoglucose isomerase *in vitro* and in growing yeast.

(A). Activity of the purified Pgi1 was assayed in the presence of 0.6 mM fructose-6-phosphate (substrate), 5 mM Mg^{2+} (pH = 7.0, 30°C) and a range of 0.05–5 mM of sorbitol-6P or ribitol-5P by monitoring the appearance of glucose-6-phosphate. The x-axis represents polyol phosphate concentration and the y-axis represents relative Pgi1 activity (mean \pm range of N = 2 replicates). (B). Metabolome profiling of cells with high (100 nM estradiol, comparable to WT protein level, shown in black) and low (1 nM estradiol, \sim 1/7 of WT protein level, shown in grey) Pgi1 induction. This experiment is used to define a characteristic low-Pgi metabolome. (C). Metabolome profiling of *pyp1Δ* (dark red) and wild type (blue) cells at t = 1 h after switching from trehalose + ribitol to glucose + ribitol. This experiment shows that PYP1 deletion results, in the presence of ribitol, in a characteristic low-Pgi metabolome.




Figure 6. Pyp1 is highly expressed and functionally important in rapid growth.

Relative expression level of *PYP1* at different growth rates in C-, N-, P- and S-limited chemostats ³⁸. The x-axis represents different growth rates and the y-axis represents relative expression level of *PYP1*.


References:

1. Teo, G.; Suzuki, Y.; Uratsu, S. L.; Lampinen, B.; Ormonde, N.; Hu, W. K.; DeJong, T. M.; Dandekar, A. M., Silencing leaf sorbitol synthesis alters long-distance partitioning and apple fruit quality. *Proc Natl Acad Sci U S A* **2006**, *103* (49), 18842-7.
2. Shindou, T.; Sasaki, Y.; Miki, H.; Eguchi, T.; Hagiwara, K.; Ichikawa, T., Determination of Erythritol in Fermented Foods by High-Performance Liquid-Chromatography. *J Food Hyg Soc Jpn* **1988**, *29* (6), 419-422.
3. Pahlman, A. K.; Granath, K.; Ansell, R.; Hohmann, S.; Adler, L., The yeast glycerol 3-phosphatases Gpp1p and Gpp2p are required for glycerol biosynthesis and differentially involved in the cellular responses to osmotic, anaerobic, and oxidative stress. *J Biol Chem* **2001**, *276* (5), 3555-63.
4. Nevoigt, E.; Stahl, U., Osmoregulation and glycerol metabolism in the yeast *Saccharomyces cerevisiae*. *FEMS Microbiol Rev* **1997**, *21* (3), 231-41.
5. Bradshaw, D. J.; Marsh, P. D., Effect of Sugar Alcohols on the Composition and Metabolism of a Mixed Culture of Oral Bacteria Grown in a Chemostat. *Caries Res* **1994**, *28* (4), 251-256.
6. Chang, Q.; Griest, T. A.; Harter, T. M.; Petrash, J. M., Functional studies of aldo-keto reductases in *Saccharomyces cerevisiae*. *Bba-Mol Cell Res* **2007**, *1773* (3), 321-329.
7. Moon, H. J.; Jeya, M.; Kim, I. W.; Lee, J. K., Biotechnological production of erythritol and its applications. *Applied Microbiology and Biotechnology* **2010**, *86* (4), 1017-1025.
8. Povelainen, M.; Miasnikov, A. N., Production of D-arabitol by a metabolic engineered strain of *Bacillus subtilis*. *Biotechnology journal* **2006**, *1* (2), 214-9.
9. Povelainen, M.; Miasnikov, A. N., Production of xylitol by metabolically engineered strains of *Bacillus subtilis*. *Journal of biotechnology* **2007**, *128* (1), 24-31.
10. Toivari, M. H.; Maaheimo, H.; Penttila, M.; Ruohonen, L., Enhancing the flux of D-glucose to the pentose phosphate pathway in *Saccharomyces cerevisiae* for the production of D-ribose and ribitol. *Appl Microbiol Biotechnol* **2010**, *85* (3), 731-9.
11. Albertyn, J.; Hohmann, S.; Thevelein, J. M.; Prior, B. A., GPD1, which encodes glycerol-3-phosphate dehydrogenase, is essential for growth under osmotic stress in *Saccharomyces cerevisiae*, and its expression is regulated by the high-osmolarity glycerol response pathway. *Mol Cell Biol* **1994**, *14* (6), 4135-44.
12. Norbeck, J.; Pahlman, A. K.; Akhtar, N.; Blomberg, A.; Adler, L., Purification and characterization of two isoenzymes of DL-glycerol-3-phosphatase from *Saccharomyces cerevisiae* - Identification of the corresponding GPP1 and GPP2 genes and evidence for osmotic regulation of Gpp2p expression by the osmosensing mitogen-activated protein kinase signal transduction pathway. *Journal of Biological Chemistry* **1996**, *271* (23), 13875-13881.
13. Ellens, K. W.; Christian, N.; Singh, C.; Satagopam, V. P.; May, P.; Linster, C. L., Confronting the catalytic dark matter encoded by sequenced genomes. *Nucleic Acids Res* **2017**, *45* (20), 11495-11514.
14. Fiehn, O.; Kopka, J.; Dormann, P.; Altmann, T.; Trethewey, R. N.; Willmitzer, L., Metabolite profiling for plant functional genomics. *Nat Biotechnol* **2000**, *18* (11), 1157-61.
15. Raamsdonk, L. M.; Teusink, B.; Broadhurst, D.; Zhang, N.; Hayes, A.; Walsh, M. C.; Berden, J. A.; Brindle, K. M.; Kell, D. B.; Rowland, J. J.; Westerhoff, H. V.; van Dam, K.; Oliver, S. G., A functional genomics strategy that uses metabolome data to reveal the phenotype of silent mutations. *Nat Biotechnol* **2001**, *19* (1), 45-50.
16. Saghatelian, A.; Cravatt, B. F., Discovery metabolite profiling--forging functional connections between the proteome and metabolome. *Life sciences* **2005**, *77* (14), 1759-66.
17. Yonekura-Sakakibara, K.; Tohge, T.; Matsuda, F.; Nakabayashi, R.; Takayama, H.; Niida, R.; Watanabe-Takahashi, A.; Inoue, E.; Saito, K., Comprehensive flavonol profiling and transcriptome coexpression analysis leading to decoding gene-metabolite correlations in *Arabidopsis*. *The Plant cell* **2008**, *20* (8), 2160-76.

18. Clasquin, M. F.; Melamud, E.; Singer, A.; Gooding, J. R.; Xu, X.; Dong, A.; Cui, H.; Campagna, S. R.; Savchenko, A.; Yakunin, A. F.; Rabinowitz, J. D.; Caudy, A. A., Riboneogenesis in yeast. *Cell* **2011**, *145* (6), 969-80.
19. Clasquin, M. F.; Melamud, E.; Rabinowitz, J. D., LC-MS data processing with MAVEN: a metabolomic analysis and visualization engine. *Current protocols in bioinformatics / editorial board, Andreas D. Baxevanis ... [et al.]* **2012**, Chapter 14, Unit14 11.
20. Quanbeck, S. M.; Brachova, L.; Campbell, A. A.; Guan, X.; Perera, A.; He, K.; Rhee, S. Y.; Bais, P.; Dickerson, J. A.; Dixon, P.; Wohlgemuth, G.; Fiehn, O.; Barkan, L.; Lange, I.; Lange, B. M.; Lee, I.; Cortes, D.; Salazar, C.; Shuman, J.; Shulaev, V.; Huhman, D. V.; Sumner, L. W.; Roth, M. R.; Welti, R.; Illarlan, H.; Wurtele, E. S.; Nikolau, B. J., Metabolomics as a Hypothesis-Generating Functional Genomics Tool for the Annotation of Arabidopsis thaliana Genes of "Unknown Function". *Frontiers in plant science* **2012**, *3*, 15.
21. Su, X.; Chen, W.; Lee, W.; Jiang, H.; Zhang, S.; Lin, H., YBR246W is required for the third step of diphthamide biosynthesis. *J Am Chem Soc* **2012**, *134* (2), 773-6.
22. Xu, Y. F.; Letisse, F.; Absalan, F.; Lu, W.; Kuznetsova, E.; Brown, G.; Caudy, A. A.; Yakunin, A. F.; Broach, J. R.; Rabinowitz, J. D., Nucleotide degradation and ribose salvage in yeast. *Mol Syst Biol* **2013**, *9*, 665.
23. Hegeman, A. D.; Schulte, C. F.; Cui, Q.; Lewis, I. A.; Huttlin, E. L.; Eghbalnia, H.; Harms, A. C.; Ulrich, E. L.; Markley, J. L.; Sussman, M. R., Stable isotope assisted assignment of elemental compositions for metabolomics. *Anal Chem* **2007**, *79* (18), 6912-21.
24. Kuchel, P. W.; Berthon, H. A.; Bubb, W. A.; McIntyre, L. M.; Nygh, N. K.; Thorburn, D. R., C-13 and P-31 Nmr-Studies of the Pentose-Phosphate Pathway in Human Erythrocytes. *Biomed Biochim Acta* **1990**, *49* (2-3), S105-S110.
25. Jennings, D. H., Polyol Metabolism in Fungi. *Adv Microb Physiol* **1984**, *25*, 149-193.
26. Pereira, M. P.; Brown, E. D., Bifunctional catalysis by CDP-ribitol synthase: Convergent recruitment of reductase and cytidylyltransferase activities in Haemophilus influenzae and Staphylococcus aureus. *Biochemistry* **2004**, *43* (37), 11802-11812.
27. Novotny, M. J.; Reizer, J.; Esch, F.; Saier, M. H., Purification and Properties of D-Mannitol-1-Phosphate Dehydrogenase and D-Glucitol-6-Phosphate Dehydrogenase from Escherichia-Coli. *Journal of Bacteriology* **1984**, *159* (3), 986-990.
28. Sarthy, A. V.; Schopp, C.; Idler, K. B., Cloning and Sequence Determination of the Gene Encoding Sorbitol Dehydrogenase from Saccharomyces-Cerevisiae. *Gene* **1994**, *140* (1), 121-126.
29. Jules, M.; Guillou, V.; Francois, J.; Parrou, J. L., Two distinct pathways for trehalose assimilation in the yeast Saccharomyces cerevisiae. *Appl Environ Microbiol* **2004**, *70* (5), 2771-8.
30. Walther, T.; Novo, M.; Rossger, K.; Letisse, F.; Loret, M. O.; Portais, J. C.; Francois, J. M., Control of ATP homeostasis during the respiro-fermentative transition in yeast. *Mol Syst Biol* **2010**, *6*, 344.
31. Aguilera, A., Deletion of the phosphoglucose isomerase structural gene makes growth and sporulation glucose dependent in Saccharomyces cerevisiae. *Mol Gen Genet* **1986**, *204* (2), 310-6.
32. Milewski, S.; Janiak, A.; Wojciechowski, M., Structural analogues of reactive intermediates as inhibitors of glucosamine-6-phosphate synthase and phosphoglucose isomerase. *Archives of Biochemistry and Biophysics* **2006**, *450* (1), 39-49.
33. Zhou, R.; Cheng, L. L.; Wayne, R., Purification and characterization of sorbitol-6-phosphate phosphatase from apple leaves. *Plant Sci* **2003**, *165* (1), 227-232.
34. Oda, Y.; Iwami, M.; Sakurai, S., Membrane-bound sorbitol 6-phosphatase in fat body cells controls the dynamics of sorbitol 6-phosphate, a major hemolymph sugar in the silkworm. *Insect Biochem Molec* **2005**, *35* (11), 1284-1292.
35. Ladero, V.; Ramos, A.; Wiersma, A.; Goffin, P.; Schanck, A.; Kleerebezem, M.; Hugenholtz, J.; Smid, E. J.; Hols, P., High-level production of the low-calorie sugar sorbitol by Lactobacillus plantarum






- 1 through metabolic engineering. *APPLIED AND ENVIRONMENTAL MICROBIOLOGY* **2007**, *73* (6),
2 1864-1872.
- 3
- 4 36. Zhang, R.; Andersson, C. E.; Savchenko, A.; Skarina, T.; Evdokimova, E.; Beasley, S.;
5 Arrowsmith, C. H.; Edwards, A. M.; Joachimiak, A.; Mowbray, S. L., Structure of Escherichia coli ribose-
6 5-phosphate isomerase: a ubiquitous enzyme of the pentose phosphate pathway and the Calvin cycle.
7 *Structure* **2003**, *11* (1), 31-42.
- 8 37. Komives, E. A.; Chang, L. C.; Lolis, E.; Tilton, R. F.; Petsko, G. A.; Knowles, J. R., Electrophilic
9 catalysis in triosephosphate isomerase: the role of histidine-95. *Biochemistry* **1991**, *30* (12), 3011-9.
- 10 38. Brauer, M. J.; Huttenhower, C.; Airoidi, E. M.; Rosenstein, R.; Matese, J. C.; Gresham, D.; Boer,
11 V. M.; Troyanskaya, O. G.; Botstein, D., Coordination of growth rate, cell cycle, stress response, and
12 metabolic activity in yeast. *Molecular biology of the cell* **2008**, *19* (1), 352-367.
- 13 39. Gasch, A. P.; Spellman, P. T.; Kao, C. M.; Carmel-Harel, O.; Eisen, M. B.; Storz, G.; Botstein,
14 D.; Brown, P. O., Genomic expression programs in the response of yeast cells to environmental
15 changes. *Molecular biology of the cell* **2000**, *11* (12), 4241-4257.
- 16 40. Andralojc, P. J.; Dawson, G. W.; Parry, M. A.; Keys, A. J., Incorporation of carbon from
17 photosynthetic products into 2-carboxyarabinitol-1-phosphate and 2-carboxyarabinitol. *The Biochemical*
18 *journal* **1994**, *304* (Pt 3), 781-6.
- 19 41. Khan, S.; Andralojc, P. J.; Lea, P. J.; Parry, M. A., 2'-carboxy-D-arabinitol 1-phosphate protects
20 ribulose 1, 5-bisphosphate carboxylase/oxygenase against proteolytic breakdown. *European journal of*
21 *biochemistry / FEBS* **1999**, *266* (3), 840-7.
- 22 42. Gibney, P. A.; Hickman, M. J.; Bradley, P. H.; Matese, J. C.; Botstein, D., Phylogenetic Portrait
23 of the Saccharomyces cerevisiae Functional Genome. *G3 (Bethesda)* **2013**, *3* (8), 1335-40.
- 24 43. Lee, A. Y. W.; Chung, S. K.; Chung, S. S. M., Demonstration That Polyol Accumulation Is
25 Responsible for Diabetic Cataract by the Use of Transgenic Mice Expressing the Aldose Reductase
26 Gene in the Lens. *Proceedings of the National Academy of Sciences of the United States of America*
27 **1995**, *92* (7), 2780-2784.
- 28 44. Nishikawa, T.; Edelstein, D.; Du, X. L.; Yamagishi, S.; Matsumura, T.; Kaneda, Y.; Yorek, M. A.;
29 Beebe, D.; Oates, P. J.; Hammes, H. P.; Giardino, I.; Brownlee, M., Normalizing mitochondrial
30 superoxide production blocks three pathways of hyperglycaemic damage. *Nature* **2000**, *404* (6779),
31 787-90.
- 32 45. Brownlee, M., Biochemistry and molecular cell biology of diabetic complications. *Nature* **2001**,
33 *414* (6865), 813-20.
- 34 46. Schrijvers, B. F.; De Vriese, A. S.; Flyvbjerg, A., From hyperglycemia to diabetic kidney disease:
35 the role of metabolic, hemodynamic, intracellular factors and growth factors/cytokines. *Endocrine*
36 *reviews* **2004**, *25* (6), 971-1010.
- 37 47. Noltmann, E. A., *Aldose-ketose isomerases*. Academic Press: 1972; Vol. 6.
- 38 48. Goyal, S.; Yuan, J.; Chen, T.; Rabinowitz, J. D.; Wingreen, N. S., Achieving optimal growth
39 through product feedback inhibition in metabolism. *PLoS computational biology* **2010**, *6* (6), e1000802.
- 40 49. Heinrich, R.; Rapoport, T. A., Linear Steady-State Treatment of Enzymatic Chains - General
41 Properties, Control and Effector Strength. *European Journal of Biochemistry* **1974**, *42* (1), 89-95.
- 42 50. Fell, D., *Understanding the Control of Metabolism*. Portland Press: 1997.
- 43 51. Kacser, H.; Burns, J. A.; Fell, D. A., The Control of Flux. *Biochemical Society Transactions* **1995**,
44 *23* (2), 341-366.
- 45 52. Kell, D. B.; Westerhoff, H. V., Metabolic Control-Theory - Its Role in Microbiology and
46 Biotechnology. *Fems Microbiology Reviews* **1986**, *39* (4), 305-320.
- 47 53. Hofmeyr, J. H. S.; Cornishbowden, A., Quantitative Assessment of Regulation in Metabolic
48 Systems. *European Journal of Biochemistry* **1991**, *200* (1), 223-236.
- 49
50
51
52
53
54
55
56
57
58
59
60



- 1
2 54. Tong, A. H.; Evangelista, M.; Parsons, A. B.; Xu, H.; Bader, G. D.; Page, N.; Robinson, M.;
3 Raghbizadeh, S.; Hogue, C. W.; Bussey, H.; Andrews, B.; Tyers, M.; Boone, C., Systematic genetic
4 analysis with ordered arrays of yeast deletion mutants. *Science* **2001**, *294* (5550), 2364-8.
- 5 55. Mclsaac, R. S.; Oakes, B. L.; Wang, X.; Dummit, K. A.; Botstein, D.; Noyes, M. B., Synthetic
6 gene expression perturbation systems with rapid, tunable, single-gene specificity in yeast. *Nucleic*
7 *Acids Res* **2013**, *41* (4), e57.
- 8 56. Xu, Y. F.; Zhao, X.; Glass, D. S.; Absalan, F.; Perlman, D. H.; Broach, J. R.; Rabinowitz, J. D.,
9 Regulation of yeast pyruvate kinase by ultrasensitive allostery independent of phosphorylation.
10 *Molecular cell* **2012**, *48* (1), 52-62.
- 11 57. Lu, W. Y.; Clasquin, M. F.; Melamud, E.; Amador-Noguez, D.; Caudy, A. A.; Rabinowitz, J. D.,
12 Metabolomic Analysis via Reversed-Phase Ion-Pairing Liquid Chromatography Coupled to a Stand
13 Alone Orbitrap Mass Spectrometer. *Analytical Chemistry* **2010**, *82* (8), 3212-3221.
- 14 58. Melamud, E.; Vastag, L.; Rabinowitz, J. D., Metabolomic Analysis and Visualization Engine for
15 LC-MS Data. *Analytical Chemistry* **2010**, *82* (23), 9818-9826.
- 16 59. Benjamini, Y. Y., Daniel, The control of the false discovery rate in multiple testing under
17 dependency. *Ann Stat* **2001**, *20* (4), 1165-1188.
- 18 60. Bennett, B. D.; Yuan, J.; Kimball, E. H.; Rabinowitz, J. D., Absolute quantitation of intracellular
19 metabolite concentrations by an isotope ratio-based approach. *Nature Protocols* **2008**, *3* (8), 1299-1311.
- 20 61. Bennett, B. D.; Kimball, E. H.; Gao, M.; Osterhout, R.; Van Dien, S. J.; Rabinowitz, J. D.,
21 Absolute metabolite concentrations and implied enzyme active site occupancy in Escherichia coli.
22 *Nature Chemical Biology* **2009**, *5* (8), 593-599.
- 23 62. Egan, W.; Schneerson, R.; Werner, K. E.; Zon, G., Structural Studies and Chemistry of Bacterial
24 Capsular Polysaccharides - Investigations of Phosphodiester-Linked Capsular Polysaccharides Isolated
25 from Hemophilus-Influenzae Types a, B, C, and F - Nmr Spectroscopic Identification and Chemical
26 Modification of End Groups and the Nature of Base-Catalyzed Hydrolytic Depolymerization. *Journal of*
27 *the American Chemical Society* **1982**, *104* (10), 2898-2910.
- 28 63. Kapuscinski, M.; Franke, F. P.; Flanigan, I.; Macleod, J. K.; Williams, J. F., Improved Methods
29 for the Enzymic Preparation and Chromatography of Octulose Phosphates. *Carbohydrate research*
30 **1985**, *140* (1), 69-79.
- 31 64. Kuznetsova, E.; Xu, L.; Singer, A.; Brown, G.; Dong, A.; Flick, R.; Cui, H.; Cuff, M.; Joachimiak,
32 A.; Savchenko, A.; Yakunin, A. F., Structure and activity of the metal-independent fructose-1,6-
33 bisphosphatase YK23 from Saccharomyces cerevisiae. *J Biol Chem* **2010**, *285* (27), 21049-59.
- 34 65. Kuznetsova, E.; Proudfoot, M.; Gonzalez, C. F.; Brown, G.; Omelchenko, M. V.; Borozan, I.;
35 Carmel, L.; Wolf, Y. I.; Mori, H.; Savchenko, A. V.; Arrowsmith, C. H.; Koonin, E. V.; Edwards, A. M.;
36 Yakunin, A. F., Genome-wide analysis of substrate specificities of the Escherichia coli haloacid
37 dehalogenase-like phosphatase family. *J Biol Chem* **2006**, *281* (47), 36149-61.
- 38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53







B



Compound	$^{12}\text{C} + ^{14}\text{N}$		$^{13}\text{C} + ^{14}\text{N}$		$^{12}\text{C} + ^{15}\text{N}$		Neutral Formula
	$[\text{M}-\text{H}]_{\text{obs}}$	ppm error	$[\text{M}-\text{H}]_{\text{obs}}$	ppm error	$[\text{M}-\text{H}]_{\text{obs}}$	ppm error	
1	319.04373	+0.5	327.07004	-1.1	319.04343	-0.5	$\text{C}_8\text{H}_{17}\text{O}_{11}\text{P}_1$
2	231.02762	+0.4	236.04439	+0.4	231.02731	-1.0	$\text{C}_5\text{H}_{13}\text{O}_8\text{P}_1$
3	201.01691	-0.3	205.03038	-0.9	201.01682	-0.7	$\text{C}_4\text{H}_{11}\text{O}_7\text{P}_1$



A**B**

ribitol-5P**ribose-5P****Ru5P/Xu5P**1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32



A**B****C**