

Rotation curves and scaling relations of extremely massive spiral galaxies

Enrico M. Di Teodoro¹,^{1,2}★ Lorenzo Posti³,³ Patrick M. Ogle,² S. Michael Fall² and Thomas Jarrett⁴⁴

¹Department of Physics & Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA

²Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

³Université de Strasbourg, CNRS UMR 7550, Observatoire astronomique de Strasbourg, 11 rue de l'Université, F-67000 Strasbourg, France

⁴Department of Astronomy, University of Cape Town, Private Bag X3, Rondebosch 7701, Cape Town, South Africa

Accepted 2021 September 6. Received 2021 September 3; in original form 2021 May 3

ABSTRACT

We study the kinematics and scaling relations of a sample of 43 giant spiral galaxies that have stellar masses exceeding $10^{11} M_{\odot}$ and optical discs up to 80 kpc in radius. We use a hybrid 3D–1D approach to fit 3D kinematic models to long-slit observations of the H α –[N II] emission lines and we obtain robust rotation curves of these massive systems. We find that all galaxies in our sample seem to reach a flat part of the rotation curve within the outermost optical radius. We use the derived kinematics to study the high-mass end of the two most important scaling relations for spiral galaxies: the stellar/baryonic mass Tully–Fisher relation and the Fall (mass–angular momentum) relation. All galaxies in our sample, with the possible exception of the two fastest rotators, lie comfortably on both these scaling relations determined at lower masses, without any evident break or bend at the high-mass regime. When we combine our high-mass sample with low-mass data from the Spitzer Photometry & Accurate Rotation Curves catalogue, we find a slope of $\alpha = 4.25 \pm 0.19$ for the stellar Tully–Fisher relation and a slope of $\gamma = 0.64 \pm 0.11$ for the Fall relation. Our results indicate that most, if not all, of these rare, giant spiral galaxies are scaled up versions of less massive discs and that spiral galaxies are a self-similar population of objects up to the very high-mass end.

Key words: galaxies: evolution – galaxies: haloes – galaxies: kinematics and dynamics – galaxies: spiral.

1 INTRODUCTION

Disc galaxies follow tight, and approximately featureless, scaling relations between some of their basic global properties, such as mass, velocity, and angular momentum. The most important scaling laws are arguably the Tully & Fisher (1977) relation, between the stellar mass or luminosity of a galaxy and its rotation velocity V , and the Fall (1983) relation, between stellar mass M_{\star} and specific angular momentum $j_{\star} \equiv J_{\star}/M_{\star}$. The stellar Tully–Fisher is a power law $M_{\star} \propto V^{\alpha}$ with $\alpha \sim 4$ –5 depending on the prescriptions used to estimate stellar masses (McGaugh & Schombert 2015; Ponomareva et al. 2018) and depending on how the rotation velocities are defined (Verheijen 2001; Lelli et al. 2019). When the baryonic mass M_{bar} (stars + cold gas) is used instead of the stellar mass, the baryonic Tully–Fisher relation (McGaugh et al. 2000) becomes an extremely tight power law $M_{\text{bar}} \propto V^{\alpha'}$, with $\alpha' \sim 4$ and intrinsic scatter ~ 0.1 dex, extending down to the lowest mass dwarf galaxies (e.g. McGaugh 2012; Lelli, McGaugh & Schombert 2016b; Lelli et al. 2019). The Fall relation is also a power law in the form $j_{\star} \propto M_{\star}^{\gamma}$, with slope $\gamma \sim 0.6$ and a small intrinsic scatter of ~ 0.15 dex, ranging from dwarf galaxies to massive discs ($7 \lesssim \log M_{\star}/M_{\odot} \lesssim 11$, Romanowsky & Fall 2012; Posti et al. 2018b; Mancera Piña et al. 2021a, b).

In the standard Λ cold dark matter (Λ CDM) paradigm, galaxies form within the gravitational potential of dark matter haloes, whose assembly by hierarchical clustering is relatively well understood (e.g. Mo, van den Bosch & White 2010). Because the Λ CDM perturbation

spectrum is featureless and because gravity is a scale-free force, the global structure of haloes is characterized by simple power-law scaling relations. Some of these basic relations for dark matter haloes are $M_{\text{h}} \propto V_{\text{h}}^3$, between halo mass M_{h} and virial velocity V_{h} , and $j_{\text{h}} \propto M_{\text{h}}^{2/3}$, between halo-specific angular momentum j_{h} and halo mass. While the $M_{\text{h}} - V_{\text{h}}$ relation follows from the definition of the virial radius, the $j_{\text{h}} - M_{\text{h}}$ relation is a consequence of the tidal torques exerted by surrounding haloes and the resulting mass independence of the spin parameter λ (Peebles 1969; Bullock et al. 2001).

The scaling laws of dark matter haloes are analogous to those of disc galaxies, with also similar slopes, suggesting a co-evolution of dark and baryonic matter. A convenient formalism that we can use to study galaxy scaling laws is to refer to the underlying relations for dark matter and to focus on the ratios $f_X \equiv X_{\star}/X_{\text{halo}}$ for a given quantity X , such as mass M , velocity V , or specific angular momentum j . With these ratios, we can rewrite the Tully–Fisher and Fall relations referencing to the analogous relations for dark matter, as $M_{\star} \propto f_M (V_{\text{flat}}/f_V)^3$ and $j_{\star} \propto f_j (M_{\star}/f_M)^{2/3}$ (e.g. Posti et al. 2019b). If f_M , f_j , and f_V were constants and did not scale with mass, then galaxies would be fully homologous to dark haloes and the shape of their scaling relations would be identical. Conversely, any significant differences between the observed slopes of the Tully–Fisher and Fall relations and the dark halo values (3 and 2/3) would indicate that some combination of f_M , f_j , and f_V must be mass dependent (e.g. Navarro & Steinmetz 2000; Dutton & van den Bosch 2012; Ferrero et al. 2017; Posti et al. 2018a).

In this framework, the remarkable similarity between galaxy and dark halo scaling laws is quite puzzling. This is because the relation between stellar mass and halo mass of the general population of galaxies, which can be expressed as the $f_M - M_{\star}$ or the $f_M - M_{\text{h}}$

★ E-mail: editeodoro@jhu.edu

relation, is not a simple, featureless power-law relation. The current consensus is that f_M increases up to a peak, at around $M_\star \sim 5 \times 10^{10} M_\odot$ ($M_h \sim 10^{12} M_\odot$), and then decreases with mass (see e.g. the review by Wechsler & Tinker 2018). Such a highly non-linear relation should induce a prominent feature, similar to a break, on the scaling laws of disc galaxies at the mass scale where f_M peaks. However, this is not observed, which implies that either the other ratios f_j and f_V are not constants (Cattaneo, Salucci & Papastergis 2014; Posti et al. 2018a; Lapi, Salucci & Danese 2018) or that the $f_M - M_\star$ relation of disc galaxies is actually a power law, contrary to estimates of the stellar-to-halo mass relation based on abundance matching (Posti, Fraternali & Marasco 2019a; Posti et al. 2019b; Posti & Fall 2021).

This issue has been studied by Lapi et al. (2018), using a compilation of stacked $H\alpha$ rotation curves of 550 nearby disc galaxies. They find indications that the scaling laws have some deviations from power laws and that the ratios f_M , f_V , and f_j are not constant, but they all increase with stellar mass up to a peak at $M_\star \sim 5 \times 10^{10} M_\odot$ and then decrease with mass. Their $H\alpha$ rotation curves, however, extend only up to one to two optical radii of the galaxies, which is typically not enough to reach the flat and dark matter-dominated parts of the rotation curves (van Albada et al. 1985; Kent 1987, 1988), and are the result of stacking, which potentially biases the results as it attenuates individual differences (e.g. Noordermeer et al. 2007). Posti et al. (2019b) re-addressed this issue with a detailed analysis of a smaller sample of nearby discs with both extended atomic hydrogen ($H I$) rotation curves and photometry at $3.6 \mu\text{m}$. Analyzing each individual curve and surface brightness profile, they determined the disc scaling laws over a large mass range ($6.5 \lesssim \log M_\star/M_\odot \lesssim 11.2$), finding no deviations from power laws, and showed that f_j and f_V are constant and close to unity across the entire mass range, while f_M varies as a simple power law of M_\star . From these results, they concluded that spirals are a self-similar population of objects, with the stellar-to-halo mass ratio f_M scaling as a power law of M_\star because of feedback from young stars (Dekel & Silk 1986).

A break in the scaling relations might still be present at higher masses than those explored by Posti et al. (2019b), i.e. at $\log M_\star/M_\odot \gtrsim 11.2$. Finding such a feature would imply that the self-similarity of discs breaks at a larger mass scale than expected, suggesting that some physical process, such as inefficient cooling of proto-galactic gas, feedback from active galactic nuclei (AGNs), or galaxy merging, are becoming important at that mass scale. Ogle et al. (2019b) recently found indications for such a break in the Tully–Fisher and Fall relations of super-luminous spiral galaxies, a rare population of giant, star-forming discs with stellar masses $M_\star \simeq 2 - 7 \times 10^{11} M_\odot$ (Ogle et al. 2016, 2019a). This study relied on long-slit $H\alpha$ spectra for 23 massive spirals, from which they derived rotation velocities typically out to the optical radii. Since this result is potentially transformational for our understanding of disc galaxies, in this paper we set out to scrutinize in detail these findings through a more accurate characterization of the kinematics and the dynamics of a larger sample of extremely massive spiral galaxies. We develop a new technique to derive robust rotation curves through modelling of long-slit $H\alpha$ -[N II] emission-line observations. We use this technique to investigate the very high-mass ends of the Tully–Fisher and Fall relations and their connections to the relations at lower masses from previous works.

The remainder of this paper is structured as follows. Section 2 introduces our galaxy sample and the long-slit data analyzed in this work. Sections 3 and 4 describe the techniques used to derive stellar surface density profiles and galaxy kinematics, respectively. We present $H\alpha$ rotation curves in Section 5, while in Section 6 we

build the Tully–Fisher and Fall scaling relations for massive spiral galaxies. We discuss our findings in Section 7 and we conclude in Section 8. Throughout this paper, we use a flat Λ CDM cosmology with $\Omega_{m,0} = 0.307$, $\Omega_{b,0} = 0.048$, and $H_0 = 67.7 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (Planck Collaboration XIII 2016). In this cosmology framework, 1 arcsec corresponds to 1.9 kpc and the lookback time is 1.3 Gyr at $z \simeq 0.1$.

2 GALAXY SAMPLE AND KINEMATIC DATA

Our work is based on a sample of 51 massive star-forming galaxies for which we have collected long-slit spectroscopy observations. Twenty-three galaxies are in common with Ogle et al. (2019b). The parent sample of massive spiral galaxies was selected from a combination of the most optically luminous Sloan Digital Sky Survey (SDSS, Eisenstein et al. 2011) galaxies (OGC catalogue, Ogle et al. 2019a) and a lower-luminosity sample of infrared-selected galaxies (Ogle et al. 2019b) from the 2-Micron All-Sky Survey (2MASS) Extended Source Catalog (2MASX, Jarrett et al. 2000; Skrutskie et al. 2006) consisting of about 54 times as many galaxies. The SDSS sub-sample was selected from the ~ 1600 most optically luminous galaxies of all types at redshift $z < 0.3$, based on SDSS redshift and r -band photometry (see Ogle et al. 2019a, for details). The 2MASX sub-sample was selected from the $\sim 86\,000$ galaxies of all types that have SDSS redshifts and K_s -band luminosity $L_{K_s} > 2 \times 10^{11} L_\odot$. The 2MASX sample was further restricted to the 19 542 massive, equatorial galaxies within the SDSS footprint with declination $\delta_{J2000} < 9^\circ$ that are observable by the Southern African Large Telescope (SALT, Buckley, Swart & Meiring 2006). To ensure reliable de-projected rotation velocities, we made a cut on galaxy ellipticity ϵ , using bulge-disc decomposition from Simard et al. (2011), which yielded 4830 high- ϵ ellipticals, lenticulars, and spiral galaxies with inclination angle $i > 40^\circ$. We visually identified 237 spiral galaxies in this subset by inspecting three-colour SDSS DR13 images using the Sky Server Chart service. Our final sample was selected from this final subset with preference for the largest actively star-forming spirals. All 51 galaxies in our sample have SDSS images in $u-g-r-z-i$ filters, deeper $g-r-z$ images from the Dark Energy Spectroscopic Instrument (DESI) Legacy Surveys (Legacy, hereinafter, Dey et al. 2019), and J -, H -, and K_s -band images from 2MASS.

To derive galaxy kinematics, we use long-slit observations of the $H\alpha$ emission line at 6562.80 \AA and of the [N II] emission-line doublet at $6548.05 - 6583.45 \text{ \AA}$. Long-slit data for 48 galaxies were obtained with the Robert Stobie Spectrograph (RSS, Burgh et al. 2003) on SALT, using the 1800 line mm^{-1} volume-phase holographic grating, which gives a resolving power of 4200–5300 at the wavelengths corresponding to the redshifted $H\alpha$ -[N II] lines. Three galaxies were observed with the Double Spectrograph on the Palomar Hale Telescope, using the 1200 line mm^{-1} grating with a constant resolving power of 6000. The width of the long-slit is $1''$, while the typical seeing of observations is $1.5'' - 2''$. We refer to Ogle et al. (2019b) for more details about the instrumental setup and data reduction.

Because only emission lines are required for our kinematic analysis, the galaxy continuum was removed from the long-slit observations: at each spatial position, a first-degree polynomial was fitted to the continuum emission in the spectral range free from the emission lines. The best-fitting polynomials were then subtracted from the data, leaving only the emission from the $H\alpha$ -[N II] lines. The $H\alpha$ -[N II] emission lines in these galaxies typically extend out to $2-4R_d$, where R_d is the scale radius of an exponential disc. From our initial sample of 51 galaxies, we excluded eight galaxies for which

we could not derive robust rotation curves because of either strong asymmetries in their emission-line profiles or insufficient signal-to-noise (S/N) ratio in the data. This leaves us with a final sample of 43 massive spiral galaxies, having redshift in the range of $0.04 \lesssim z \lesssim 0.28$ and morphology spanning from early-type discs (e.g. Sa, Sab) to late-type discs (e.g. Sb, Sc). The main properties of the sample are listed in Table 1. Galaxies excluded from our kinematic analysis are also listed at the bottom of Table 1 and their $H\alpha$ data can be found in Fig. A2 of Appendix A. In the remainder of the paper, we will refer to individual galaxies in our sample either by using the alternative names listed in the second column of Table 1 or by cutting down the 2MASS name to the first four digits of the sky coordinates (e.g. 2MASXJ1022).

3 STELLAR MASSES AND SURFACE DENSITY PROFILES

W1-band photometry at $3.4 \mu\text{m}$ from the Wide-field Infrared Survey Explorer (WISE, Wright et al. 2010) was used to calculate the total stellar content of our galaxies, applying the data pipeline and the methods described in Jarrett et al. (2019) (see their Section 2.4). Foreground stars and other contaminating sources were identified and removed from the images and the background was measured in an elliptical annulus and subtracted. Total flux was measured by fitting the radial surface brightness profile with a double Sérsic function and integrating the extrapolated profile out to $3R_d$, where R_d is the disc scale length. This is never more than 10 per cent more than the isophotal flux measured at the 1σ outer isophote. The W1-band flux was K -corrected using the 2MASS + WISE spectral energy distribution and used to compute the W1 luminosity L_{W1} relative to the Sun (Cluver et al. 2014). Finally, the total mass in stars M_* was estimated assuming a constant mass-to-light ratio $M_*/L_{W1} = 0.6$ with a typical uncertainty of ~ 40 per cent (Meidt et al. 2014; Norris et al. 2014; Röck et al. 2015). Errors on the final stellar masses are of the order 0.2 dex and are dominated by the uncertainty on the mass-to-light ratio. Dust obscuration is expected to be negligible or small at $3.4 \mu\text{m}$. The fourth column of Table 1 lists our derived stellar masses M_* . All galaxies in our sample have stellar masses that exceed $10^{11} M_\odot$, with the most massive one reaching a $M_* \simeq 6 \times 10^{11} M_\odot$. Star formation rates (SFRs) for our galaxies were estimated from the WISE W3-band at $12 \mu\text{m}$, following the prescriptions of Cluver et al. (2017) (see their equation 4). Our galaxies have dust-obscured SFRs ranging from a few $M_\odot \text{yr}^{-1}$ up to a few tens $M_\odot \text{yr}^{-1}$ (column 5 in Table 1).

We use z -band images from the Legacy surveys to derive stellar surface density profiles $\Sigma_*(R)$, under the assumption that light traces stellar mass. Surface photometry analysis is performed using the PHOTUTILS package (Bradley et al. 2020) implemented within the ASTROPY PYTHON library (Astropy Collaboration 2013, 2018). A median sky value is determined after masking contaminating sources and subtracted from each image. We then measure azimuthally averaged surface brightness profiles by fitting a set of elliptical isophotes to galaxy images, using the iterative method described in Jedrzejewski (1987). Each ellipse is defined by four parameters: the galaxy centre (x_0, y_0) , the position angle ϕ of the ellipse's major axis with respect to the North direction, and the ellipticity $\epsilon \equiv 1 - b/a$, where b/a is the ratio of the minor axis b to the major axis a . We provide an initial guess ellipse and we let all parameters free during the ellipse fitting. During the procedure, pixels deviating more than 3σ from the average value within an ellipse, which may represent unresolved background/foreground objects, were automatically masked. Total errors for each ellipse

were calculated as $\sigma_\Sigma^2(R) = \sigma_{\text{rms}}^2(R) + \sigma_{\text{sky}}^2$, where $\sigma_{\text{rms}}(R)$ is the root-mean-square (rms) variation around each ellipse and σ_{sky} is the error on the sky determination. Typically, the first term dominates the error budget in the inner regions of a galaxy, while the latter dominates in the outermost regions. Best-fitting position angles of the outermost isophotes generally agree with the long-slit position angles within a few degrees. Fig. 1 illustrates examples of surface density photometry for three galaxies in our sample. Bottom panels show their stellar surface brightness profiles derived from the z -band images along the best-fitting ellipses (red lines) displayed in the top panels.

4 KINEMATIC MODELLING

We exploited a hybrid three-dimensional–one-dimensional (3D)–(1D) approach to derive accurate rotation curves from our long-slit spectroscopic data. Our procedure starts with building 3D emission-line models through the kinematic code ^{3D}BAROLO (Di Teodoro & Fraternali 2015). ^{3D}BAROLO uses a 3D tilted-ring model to simulate 3D emission-line spectral data cubes (two spatial and one spectral dimension) for a galaxy with given geometrical and kinematic parameters. The geometry is characterized through five parameters: the central positions of the galaxy (x_0, y_0) , the inclination angle i with respect to the line of sight (90° for an edge-on disc), the position angle ϕ of the disc's major axis with respect to the North direction, and the scale height Z_0 of the disc. The kinematics is defined by three main parameters: the redshift z , the rotation velocity V_{rot} , and the intrinsic gas velocity dispersion σ_{gas} . A 3D modelling allows to account easily for the instrumental biases introduced by the point spread function (PSF) of the telescope, which can be important for galaxies resolved with just a few resolution elements (beam smearing, e.g. Begeman 1987), and by the spectral resolution of the spectrograph. In addition, many of our galaxies have $i > 70^\circ$ and a 3D model allows us to deal with highly inclined systems, overcoming the well-known issues of deriving rotation curve in nearly edge-on galaxies (e.g. Fraternali, Sancisi & Kamphuis 2011).

Throughout the fitting procedure, we constructed 3D models using ring widths of $1.5''$ and fixing the galaxy geometry. The galaxy centre was set to the position corresponding to the peak flux of the continuum emission in the long-slit data. The position angle was arbitrarily fixed to $\phi = 90^\circ$ and the disc thickness to $Z_0 = 500 \text{ pc}$. We stress that the discs of these galaxies seem to be extremely thin, as demonstrated by our edge-on systems' optical images, implying that the effect of the disc scale height on the derived rotation curve is most likely negligible. The inclination angle was calculated from the axis ratios from the isophotal fit of the optical images (Section 3), i.e. $\cos^2 i = [(b/a)^2 - q_0^2]/(1 - q_0^2)$, where a and b are the major and minor semi-axes, respectively, and we assume $q_0 = 0.16$ for these thin discs (e.g. Fouque et al. 1990). To avoid issues with the possible presence of a spheroidal bulge in the inner regions of our galaxies, we used a b/a averaged over the outermost isophotes only. Keeping fixed the galaxy geometry to the above parameters, we fitted for the galaxy redshift z , the rotation velocity at each radius $V_{\text{rot}}(R)$, and a constant value for the gas velocity dispersion σ_{gas} .

Model data cubes were simulated using a spatial pixel size of $0.254''$ and a spectral channel width of $0.3 - 0.4 \text{ \AA}$, corresponding to the spatial and spectral pixels of the long-slit data, respectively. An instrumental spectral broadening of $\sigma_{\text{instr}} \simeq 1 \text{ \AA}$ was added in quadrature to the gas intrinsic velocity dispersion σ_{gas} to take into account the finite resolving power of spectrographs. Three-dimensional models were convolved with a circular two-dimensional

Table 1. Massive spiral galaxy sample analyzed in this work. Columns: (1) primary name from either 2MASS or SDSS surveys; (2) alternative name used in this paper; (3) redshift from kinematic modelling; (4) stellar mass from W1 WISE aperture photometry (typical error is 0.2 dex); (5) star formation rates from W3 WISE luminosity, with typical uncertainty of 30%–40%; (6) inclination angle from z -band axial ratio (typical error is 5°); (7) Position angle of the major axis (counterclockwise from North direction); (8) velocity of the flat part of the rotation curve; (9) warm-ionized gas velocity dispersion; and (10) total specific angular momentum. We also list galaxies excluded from our final sample, specifying the reason why we were not able to determine rotation curves for them.

Name	Alt. name	z	$\log \frac{M_*}{M_\odot}$	SFR $M_\odot \text{ yr}^{-1}$	i deg	P.A. deg	V_{flat} km/s	σ_{gas} km/s	$\log \frac{j_*}{\text{kpc km s}^{-1}}$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
2MASXJ00083819 – 0044064	–	0.03952	11.24	9	57	16	275 ± 20	41 ± 6	3.45 ± 0.11
2MASXJ01431323 + 0034405	–	0.17771	11.60	19	71	128	361 ± 30	28 ± 5	3.78 ± 0.13
2MASXJ02052519 + 0015129	–	0.17186	11.57	19	52	148	348 ± 29	25 ± 4	3.77 ± 0.13
2MASXJ02152347 – 0937283	–	0.21625	11.51	9	68	22	364 ± 28	19 ± 4	3.91 ± 0.12
2MASXJ02201242 – 0832228	–	0.11015	11.23	9	87	1	252 ± 16	23 ± 5	3.69 ± 0.10
2MASXJ02205358 + 0020002	–	0.13957	11.26	14	63	52	286 ± 17	29 ± 5	3.75 ± 0.10
2MASXJ09034365 + 0139334	–	0.10351	11.31	8	72	126	265 ± 15	28 ± 3	3.63 ± 0.15
2MASXJ09484652 + 0635428	–	0.08824	11.14	4	67	155	311 ± 19	29 ± 5	3.77 ± 0.13
SDSSJ095727.02 + 083501.7	OGC 0441	0.25611	11.60	11	44	92	384 ± 41	33 ± 6	3.96 ± 0.12
2MASXJ09590646 + 0559293	–	0.09290	11.30	12	90	34	284 ± 18	23 ± 4	3.73 ± 0.13
2MASXJ10082752 + 0304597	–	0.16839	11.38	16	76	152	303 ± 18	30 ± 4	3.89 ± 0.10
2MASXJ10102901 + 0106588	–	0.10219	11.44	10	50	30	359 ± 27	27 ± 4	3.81 ± 0.11
2MASXJ10222648 + 0911396	–	0.09143	11.42	17	71	75	307 ± 21	25 ± 7	3.70 ± 0.10
2MASXJ10304263 + 0418219	OGC 0926	0.16105	11.66	34	50	64	325 ± 28 ^b	28 ± 5	3.70 ± 0.11
2MASXJ10494004 + 0026155	–	0.09103	11.14	6	83	102	270 ± 17	27 ± 5	3.61 ± 0.10
2MASXJ11052843 + 0736413	2MFGC 08638	0.15244	11.59	24	89	80	458 ± 21	26 ± 6	4.16 ± 0.09
2MASXJ11193844 + 0047233	–	0.17419	11.44	15	60	129	413 ± 24	25 ± 3	3.81 ± 0.10
2MASXJ11232039 + 0018029	–	0.14461	11.43	18	79	74	386 ± 22	20 ± 6	3.94 ± 0.08
2MASXJ11483552 + 0325268	–	0.12007	11.42	13	76	29	293 ± 18	25 ± 5	3.84 ± 0.10
2MASXJ11495671 + 0457570	–	0.09303	11.25	13	80	14	260 ± 21	28 ± 4	3.61 ± 0.10
2MASXJ11535621 + 4923562	OGC 0586	0.16673	11.64	38	66	128	338 ± 27 ^b	20 ± 5	3.89 ± 0.16
2MASXJ12003406 + 0510290	–	0.13060	11.29	10	80	123	273 ± 19	29 ± 5	3.72 ± 0.10
2MASXJ12422564 + 0056492	–	0.07964	11.24	10	55	46	270 ± 21	32 ± 5	3.40 ± 0.11
2MASXJ12592630 – 0146580	–	0.08338	11.23	8	58	80	317 ± 23	31 ± 5	3.64 ± 0.09
2MASXJ13033075 – 0214004	2MFGC 10372	0.08425	11.37	14	81	42	301 ± 19	30 ± 6	3.52 ± 0.11
2MASXJ13270366 – 0002093	–	0.14062	11.18	10	80	71	283 ± 20	25 ± 4	3.70 ± 0.10
2MASXJ13395130 + 0302174	–	0.11562	11.42	9	51	172	364 ± 29	28 ± 5	3.78 ± 0.13
2MASXJ13450793 – 0247470	–	0.07520	11.24	7	90	138	288 ± 18	27 ± 5	3.75 ± 0.10
SDSSJ143447.86 + 020228.6	OGC 1312	0.27967	11.60	35	63	88	364 ± 23	25 ± 4	3.81 ± 0.11
2MASXJ15154614 + 0235564	2MFGC 12344	0.14049	11.74	22	78	150	530 ± 24	21 ± 4	4.11 ± 0.10
2MASXJ15172226 – 0216475	–	0.11725	11.06	10	79	143	274 ± 19	22 ± 3	3.60 ± 0.11
2MASXJ15404057 – 0009331	–	0.07849	11.39	13	66	44	310 ± 21	22 ± 6	3.65 ± 0.09
2MASXJ16184003 + 0034367	–	0.16734	11.67	19	69	78	383 ± 25 ^b	23 ± 4	3.93 ± 0.11
2MASXJ16245634 + 0714325	–	0.15564	11.57	22	55	155	298 ± 21	46 ± 6	3.72 ± 0.10
2MASXJ20541957 – 0055204	–	0.21014	11.41	24	63	2	316 ± 22	19 ± 4	3.77 ± 0.10
2MASXJ21175224 – 0057041	–	0.18022	11.38	19	78	174	303 ± 22	20 ± 4	3.82 ± 0.10
2MASXJ21362206 + 0056519	–	0.10378	11.47	12	62	113	348 ± 24	26 ± 4	3.74 ± 0.10
2MASXJ21384311 – 0052162	–	0.08306	11.20	3	58	78	291 ± 23	28 ± 5	3.56 ± 0.11
2MASXJ21431882 – 0820164	–	0.06249	11.13	4	75	82	310 ± 20	22 ± 5	3.57 ± 0.09
2MASXJ21445295 – 0804037	–	0.12781	11.35	7	38	187	288 ± 26	25 ± 4	3.62 ± 0.18
2MASXJ22073122 – 0729223	–	0.06332	11.20	10	71	84	223 ± 19	33 ± 6	3.45 ± 0.09
2MASXJ22520878 + 0015154	–	0.15176	11.25	12	80	45	292 ± 19	24 ± 5	3.62 ± 0.10
2MASXJ23130513 – 0033477	–	0.11116	11.20	11	56	56	279 ± 27	29 ± 5	3.48 ± 0.11
Discarded galaxies									
2MASXJ09394584 + 0845033 ^a	–	0.13784	11.45	28	55	120	Highly asymmetric kinematics		
2MASXJ13180708 + 0502353	–	0.09626	11.26	5	87	39	Highly asymmetric kinematics		
2MASXJ13451949 + 0058117	–	0.16447	11.31	7	79	146	Low S/N (marginal detection)		
2MASXJ15592695 + 0842570	–	0.19962	11.52	14	46	59	Low S/N (marginal detection)		
2MASXJ16014061 + 2718161 ^a	OGC 1304	0.16466	11.63	23	60	116	Asymmetric: approaching side not well defined		
2MASXJ16262411 + 0841036	–	0.21553	11.43	9	72	28	Asymmetric: receding side not well defined		
2MASXJ16394598 + 4609058 ^a	OGC 0139	0.24714	11.74	45	76	115	Low S/N + Anomalous emission		
2MASXJ21020534 – 0647558	–	0.12730	11.84	59	52	153	Low S/N + AGN contamination		

^aGalaxy in common with Ogle et al. (2019b).

^bDetermination of V_{flat} may be affected by atmospheric absorption.




Figure 1. Surface photometry for three massive spiral galaxies in our sample. *Top panels:* z -band images from the Legacy survey (Dey et al. 2019) with overlaid some of the best-fitting ellipses in red. *Bottom panels:* average surface brightness over the best-fitting ellipses. The dashed grey line denotes an exponential function fit to the outer parts of the surface brightness profile.

Gaussian kernel with full width at half-maximum of $1.5''$, representing the typical seeing of long-slit observations (Ogle et al. 2019b). From these 3D model data cubes, we extracted a $1''$ -wide position-wavelength slice along the kinematic major axis, representing a simulated long-slit observation, and we fitted the model parameters $\mathbf{p} = \{z, V_{\text{rot}}(R), \sigma_{\text{gas}}\}$ by minimizing the absolute residuals between the simulated and the real data:

$$\mathcal{F}(\mathbf{p}) = \frac{1}{N} \sum_{x,\lambda} |f_{\text{data}}(x, \lambda) - f_{\text{model}}(x, \lambda, \mathbf{p})|, \quad (1)$$

where $f_{\text{data}}(x, \lambda)$ and $f_{\text{model}}(x, \lambda)$ are the observed and predicted flux densities at a given position x and wavelength λ , respectively, and the summation extends over all the N pixels with S/N ratio >2.5 in the continuum-subtracted slit data. Absolute residuals give more weight to regions where the emission is faint and diffuse with respect to the square of residuals (Di Teodoro & Fraternali 2015). The optimization of the objective function $\mathcal{F}(\mathbf{p})$ made use of a Multilevel Coordinate Search algorithm (Huyer & Neumaier 1998). The above procedure was applied preferentially to the $H\alpha$ emission line only, which is usually stronger than the $[\text{N II}]$ -doublet. For two galaxies (OGC0586 and 2MASXJ2138), for which part of the $H\alpha$ line was affected by the subtraction of atmospheric sky lines, we fitted our kinetic model to three emission lines simultaneously (e.g. Di Teodoro et al. 2018).

Uncertainties on the derived parameters, in particular on the rotation curve $V_{\text{rot}}(R)$, are dominated by (1) the relatively low S/N ratio of the data, (2) systematic errors on the assumed inclination angle i , and (3) possible misalignment of the long-slit with respect to a galaxy's major axis (i.e. an error on the position angle ϕ). To take into account these three important effects, we estimated parameter

uncertainties via bootstrapping. We produced 10 k realizations of our best-fitting parameters: in each realization, a $1\sigma_{\text{RMS}}$ Gaussian noise was added to the long-slit data, while i and ϕ values were drawn from normal distributions with mean given by our fiducial values and standard deviation of 5° . We checked that bootstrap distributions of our free parameters were nearly Gaussian after 10 k iterations and we assumed the values at 15.87th and the 84.13th percentiles as 1σ deviation errors.

5 ROTATION CURVES AND GAS VELOCITY DISPERSION

Fig. 2 illustrates our best-fitting models and the corresponding $H\alpha$ rotation curves for three fast rotators in our sample. Models for all 43 galaxies can be found in Fig. A1 of Appendix A. We show composite g, r, z images from the Legacy surveys (*left-hand panels*), the $H\alpha$ emission along the major axes from our long-slit observations (*middle panels*), and our best-fitting rotation curves (*right-hand panels*). Grey contours in middle panels denote the data and red contours our best-fitting models, with yellow points representing the rotation velocities projected along the line of sight. Empty points in the rightmost panels highlight regions where the derived rotation velocities are less reliable due to possible extinction and projection effects that are not taken into account during the kinematic modelling. Unlike the atomic hydrogen line at 21 cm (H I), the $H\alpha$ line is not optically thin and extinction may occur, especially in the innermost regions of highly inclined systems. For example, a hole in the ionized gas distribution or an extinguished region with a high rotation velocity in the inner parts of a nearly edge-on galaxy may be hardly visible




Figure 2. Rotation curves for our three fastest rotation galaxies. *Left:* RGB images in the g -, r -, and z -bands from the Legacy surveys (Dey et al. 2019). *Middle:* Continuum-subtracted $H\alpha$ long-slit data from SALT, shown in grey scale and contours, and our best-fitting kinematic models (red contours). Some of the $[\text{N II}]$ -doublet emission is also visible at the top/bottom edges of the plots. Contours are drawn at $2^n \times 2.5\sigma_{\text{rms}}$, where σ_{rms} is the rms noise of the data and $n = 0, 1, 2, 3$, and 4. Yellow dots denote the derived rotation velocities projected along the line of sight (i.e. $V_{\text{rot}}\sin i$). *Right:* Best-fitting rotation curves (red circles). Errors in the rotation curves include systematic uncertainties on the inclination angle and long-slit positioning. White empty dots highlight radii where the rotation velocity cannot be robustly determined because of possible projection/absorption effects. The grey horizontal line and band represent the velocity of the flat part of the rotation curve and its error, respectively.

in our $H\alpha$ data. To better understand these effects, we build a set of kinematic models starting from our best-fitting rotation curve and by progressively removing emission from the inner rings of each galaxy until the model starts to deviate significantly from the data. This procedure allows us to identify the minimum radius where the rotation curve is reliable. Finally, for galaxies where the line emission is considerably non-axisymmetric with respect to the galaxy centre, we preferentially fit only the side that is either more regular or more extended in radius. An example of a galaxy with asymmetric emission is 2MASXJ1123, visible in the second row of Fig. 2.

All galaxies in our sample have rotation curves that reach a flat part in the outer regions of the optical disc. The shape of rotation curves of spiral galaxies has been observed to have a clear dependence on galaxy luminosity (e.g. Persic, Salucci & Stel 1996). In particular, massive spiral galaxies typically have rotation curves that sharply rise in the inner regions and then quickly approach an asymptotic value. Some earlier-type massive disc galaxies are known to exhibit slowly declining rotation curves that flatten out at large radii (e.g. Kent 1988; Noordermeer et al. 2007). As expected, most galaxies in our sample show steeply rising rotation curves that flatten well within 2–

$3R_e$, where R_e is the effective radius. Notable exceptions are our two fastest rotating discs, 2MFGC08638 and 2MFGC12344 (see first and third rows in Fig. 2). Although the inner rotation velocities of these galaxies could be very uncertain because of their high inclination angles, the rotation curves seem to rise slowly and reach a flat region only in the outermost regions of the stellar disc. This evidence might indicate that the dynamics of these galaxies is dominated by the dark matter halo even in the inner regions. These two galaxies also reach extremely large rotation velocities, with 2MFGC12344 arguably setting the record for the most rapidly rotating spiral galaxy currently known (see also UGC12591, Giovanelli et al. 1986). Some galaxies in our sample have rotation curves that peak very near their centres (e.g. 2MFGC10372, 2MASXJ1200), highlighting the presence of massive bulges that are also visible in the optical images. Although a few galaxies show a slight decline in some part of the optical rotation curve, none seems to have a rotation velocity that is still declining significantly at the outermost measured radius, with the possible exceptions of 2MASXJ0205 and 2MASXJ0215.

Together with the rotation velocity, our kinematic modelling returns the velocity dispersion of the warm ionized gas that originates the $H\alpha$ -[N II] emission lines. Our best-fitting value is corrected for observational biases, in particular, for the PSF and for the instrumental broadening, and therefore represents the intrinsic velocity dispersion of gas, accounting for thermal and turbulent broadening. We find that the 1D velocity dispersion in our sample ranges between 22 km/s and 33 km/s (see Table 1), values fully consistent with previous measurements of gas velocity dispersion of the warm ionized medium in less massive spiral galaxies (e.g. Andersen et al. 2006; Martinsson et al. 2013). This implies that the kinematics of these systems is strongly dominated by rotational motions, with $V_{\text{rot}}/\sigma_{\text{gas}} \gg 10$ for all galaxies. Overall, from the shape of the rotation curves and from the gas velocity dispersions, we can conclude that these spiral galaxies seem akin to less massive discs from a kinematical point of view.

6 SCALING RELATIONS

6.1 Tully–Fisher relations

The Tully–Fisher relation comes in different flavours, depending on the way mass, luminosity, or rotation velocity is measured. Historically, luminous matter has been expressed through galaxy luminosity in different bands (e.g. Karachentsev et al. 2002), stellar mass (e.g. Torres-Flores et al. 2011), or baryonic mass (stars + cold gas, e.g. McGaugh 2005). Widely used velocity definitions (see e.g. Lelli et al. 2019, for a detailed discussion) include the width of the global H I profile W_{HI} , the circular velocity $V_{2.2}$ at $2.2R_d$, the maximum rotation velocity V_{max} , and the mean velocity along the flat part of the rotation curve V_{flat} . Several studies have shown that the latter velocity definition, V_{flat} , minimizes the scatter of the relation (Verheijen 2001; Lelli et al. 2019).

We use the stellar masses derived in Section 3 and the flat part of the rotation curves modelled in Section 4 to build the stellar Tully–Fisher relation $M_\star - V_{\text{flat}}$ for our massive spiral galaxies. Following Lelli et al. (2016b), we calculate V_{flat} using a simple iterative procedure that quantifies the level of flatness of a curve. This algorithm starts from the last points of a rotation curve and proceeds inwards, evaluating the variation in velocity with respect to the average flat velocity V_a until a deviation larger than 5 per cent is found (for details, see section 2.2 of Lelli et al. 2016b). All galaxies in our sample happen to have rotation curves that are flat within 5 per cent over at least three velocity points. Under the assumption that rotation curves remain

flat at larger radii, our measured V_{flat} corresponds to the asymptotic rotation velocity. The uncertainty on V_{flat} is calculated as:

$$\delta_{V_{\text{flat}}} = \left(\frac{1}{N} \sum_i^N \delta_{V_i}^2 + \delta_{V_a}^2 \right)^{\frac{1}{2}}, \quad (2)$$

where δ_{V_i} is the error associated with each of the N velocity points along the flat part of the rotation curve and δ_{V_a} is the dispersion around the average flat velocity value, which quantifies the degree of flatness of the curve. We note that the errors on inclination and position angles are already included in δ_{V_i} . Derived V_{flat} are listed in Table 1 (column 8). The grey lines and shaded regions in the rotation curves (rightmost panels) of Fig. 2 and of Fig. A1 denote the resulting V_{flat} and associated error, respectively.

The left-hand panel of Fig. 3 shows our determination of the stellar Tully–Fisher relation. Red squares at the high-mass end of the relation highlight our galaxy sample. As a comparison, we plot with cyan circles the relation from the Spitzer Photometry & Accurate Rotation Curves catalogue (SPARC, Lelli et al. 2016a), which is the largest collection of galaxies with high-quality photometry at $3.6\mu\text{m}$ and H I rotation curves currently available. We fit a linear function in the form $\log M_\star = \alpha \log V_{\text{flat}} + \beta$ to the entire sample and to galaxies with $\log(M_\star/M_\odot) > 10.5$ only, using an orthogonal distance regression technique to take into account errors in both variables. This allows us to quantify whether the relation changes slope at the high-mass end. The solid line in Fig. 3 denotes the best-fitting linear relation to the full sample, $\alpha = 4.25 \pm 0.19$ and $\beta = 0.80 \pm 0.23$, with an orthogonal scatter of $\sigma_\perp \simeq 0.15$ dex (grey band). We note that the grey band in Fig. 3 is wider than the errors of individual galaxies because the term δ_{V_a} (flatness deviation) dominates in equation (2). The dashed line is the best fit to the high-mass sample only, $\alpha = 3.51 \pm 0.22$ and $\beta = 2.61 \pm 0.21$. These best-fitting relations are consistent with each other and are in good agreement with recent determinations of the stellar Tully–Fisher relation (e.g. Reyes et al. 2011; Bekeraite et al. 2016; Bloom et al. 2017). From this we conclude that the relation seems to continue unbroken from dwarfs to super spirals. The two marginal outliers visible in the left-hand panel of Fig. 3 in our massive spiral sample are the two fastest rotators, 2MFGC08638 and 2MFGC12344.

The baryonic Tully–Fisher relation is considered the straightest and tightest of all Tully–Fisher relations (McGaugh et al. 2000; McGaugh 2005). Unfortunately, there are no measurements of the cold gas content of the galaxies in our sample, and the strong radio-frequency interference at the redshifted H I frequencies prevents such measurements. We can, however, have a rough estimate of gas masses in these galaxies via the Kennicutt–Schmidt relation (Schmidt 1959; Kennicutt 1998). In particular, we use the global star formation law recently calibrated by de los Reyes & Kennicutt (2019) in local spiral galaxies:

$$\log \left(\frac{\Sigma_{\text{SFR}}}{M_\odot \text{ yr}^{-1} \text{ kpc}^{-2}} \right) = 1.41 \log \left(\frac{\Sigma_{\text{gas}}}{M_\odot \text{ pc}^2} \right) - 3.84, \quad (3)$$

where Σ_{SFR} is the SFR density and Σ_{gas} is the cold gas surface density (atomic + molecular gas). We assumed a typical scatter of 0.3 dex for this relation. Σ_{SFR} was calculated by normalizing the SFR by the deprojected star-forming area πR_{25}^2 , where R_{25} is the radius at the 25 mag arcsec $^{-2}$ isophote in z -band. We then obtained the gas surface density Σ_{gas} via equation (3) and converted to gas mass M_{gas} by multiplying Σ_{gas} by the same area. This results in typical gas masses $\log(M_{\text{gas}}/M_\odot) \simeq 10.0$ – 10.9 and gas fractions $M_{\text{gas}}/M_\star \simeq 0.08$ – 0.20 . We note that these cold gas masses calculated via the Kennicutt–Schmidt law are also in good agreement with the expectations from




Figure 3. Tully–Fisher relations for massive spiral galaxies. The stellar Tully–Fisher relation (*left*) and the baryonic Tully–Fisher relation (*right*) are shown. We plot our sample of massive spirals (red squares) and galaxies from the SPARC catalogue (cyan circles) (Lelli, McGaugh & Schombert 2016a). Solid and dashed black lines are linear fits to all galaxies and to galaxies with stellar/baryonic masses larger than $3 \times 10^{10} M_{\odot}$, respectively. Best-fitting parameters for the relation in the form $\log M = \alpha \log V_{\text{flat}} + \beta$ are $(\alpha, \beta) = (4.25, 0.80)$ (full sample) and $(\alpha, \beta) = (3.51, 2.61)$ (high mass) for the stellar Tully–Fisher relation, and $(\alpha, \beta) = (3.60, 2.49)$ (full sample) and $(\alpha, \beta) = (3.26, 3.33)$ (high mass) for the baryonic Tully–Fisher relation. Grey bands denote the 1σ orthogonal scatter of data around the best-fitting relations for all galaxies, i.e. 0.15 dex for the stellar Tully–Fisher relation and 0.1 dex for the baryonic Tully–Fisher relation.

the empirical relation between stellar mass and HI mass in gas-rich spiral galaxies (e.g. Parkash et al. 2018; Naluminsa, Elson & Jarrett 2021). Finally, we calculate the total baryonic mass as $M_{\text{bar}} = M_{\star} + M_{\text{gas}}$.

The right-hand panel of Fig. 3 shows the baryonic Tully–Fisher relation for our galaxies (red squares) compared to the SPARC sample by Lelli et al. (2019). As before, the black solid line is a best fit to the entire mass range, $\alpha = 3.60 \pm 0.17$ and $\beta = 2.49 \pm 0.20$, which is virtually indistinguishable from previous determinations from SPARC (Lelli et al. 2016b, 2019). The black dashed lines are a best fit to galaxies with $\log(M_{\text{bar}}/M_{\odot}) \geq 10.5$ only, $\alpha = 3.26 \pm 0.14$ and $\beta = 3.33 \pm 0.19$. Our galaxies are perfectly compatible with the relation determined by lower mass galaxies, similar to what we find for the stellar Tully–Fisher relation (Fig. 3, left-hand panel). The only two galaxies that can be considered outliers of the baryonic Tully–Fisher relation are, again, the two fastest rotating galaxies in our sample. Although our gas mass estimates are rather crude, we are confident on our estimates of the total baryonic masses since stars always dominate the baryonic budget in giant spiral galaxies. Even if the gas masses were a factor of 2 larger/smaller than what we assume, our sample of galaxies would still be consistent with the scaling relations determined at lower masses.

6.2 Fall relation

For an axisymmetric disc rotating on cylinders about its symmetry axis, the specific stellar angular momentum within a radius R from the centre can be written as:

$$j_{\star}(< R) = \frac{J_{\star}(< R)}{M_{\star}(< R)} = \frac{\int_0^R \Sigma_{\star}(R') R'^2 V_{\star, \text{rot}}(R') dR'}{\int_0^R \Sigma_{\star}(R') R' dR'}, \quad (4)$$

where Σ_{\star} and $V_{\star, \text{rot}}$ are the mass surface density and the azimuthal velocity of the stellar component, respectively. Equation (4) is used to compute the specific angular momentum as a function of radius for our galaxies. We use the stellar surface density profiles derived in Section 3 and we assume $V_{\star, \text{rot}} = V_{\text{H}\alpha, \text{rot}}$. We expect any correction to the circular velocity for pressure support to be negligible for these highly flattened discs, i.e. $\sigma_{\star}/V_{\star, \text{rot}} \sim z_d/R_d \ll 1$, where z_d and R_d are the disc scale height and scale length, respectively (see also Mancera Piña et al. 2021a).

Fig. 4 shows cumulative j_{\star} profiles for our 43 galaxies calculated through equation (4). Black solid lines denote $j_{\star}(< R)$ within the last point of the measured rotation curve. Most profiles are converging within the optical disc but are not fully converged to the asymptotic value representing the total specific angular momentum of the galaxy. Therefore, we calculate j_{\star} at larger radii by extrapolating the rotation curves and surface brightness profiles beyond the last measured data point. Since all of our galaxies have reached the flat part of the rotation curve, as discussed in Section 5, we assume that the velocity remains constant at larger radii. For the surface brightness, we fit an exponential function to the outer regions of the optical disc (dashed lines in Fig. 1) and we assume that the exponential decline continues beyond the last measured point. Our extrapolation stops at the radius where the converging criterion used in Posti et al. (2018b) is met, i.e. $\Delta j_{\star}/j_{\star} < 0.1$ and $\Delta \log j_{\star}/\Delta \log R < 0.5$, where Δ is measured over the two outermost radii. Blue-dashed lines in Fig. 4 denote the extrapolated part of the converged j_{\star} profiles. Errors on the total specific angular momentum were calculated by bootstrapping 10k realization of each j_{\star} profile. To take into account a possible decline or increase of the rotation curve beyond the optical disc, we draw random extrapolations of the rotation curve by allowing a conservative maximum variation of 20 per cent from




Figure 4. Stellar specific angular momentum profiles for the 43 massive spiral galaxies in our sample. Black solid lines show the inner regions where both H α rotation velocity and stellar surface brightness are measured. These profiles are either fully converged or converging. Blue dashed lines show the extrapolation for a flat rotation curve and an exponentially declining surface density profile. The radius R is normalized to the disc scale length R_d in z -band and j_* to the specific angular momentum of a thin exponential disc with the same R_d and with a constant rotation curve (dotted line).

V_{flat} . Errors on the surface brightness were taken into account by drawing random extrapolations around the best-fitting exponential function. The 15.87th and the 84.13th percentiles of the resulting specific angular momentum distribution for each galaxy are taken as 1σ error. We find that typical uncertainties on j_* are of the order of 0.1 dex. The values of total j_* calculated for our sample are listed in Table 1 (column 10).

The Fall relation for our massive disc galaxies is presented in Fig. 5 (red squares). We compare it with the Fall relation at lower stellar masses determined by Posti et al. (2018b) using galaxies from the SPARC sample (cyan circles). The solid black line indicates the best-fitting relation $\log j_* = (0.64 \pm 0.11) \log M_* - (3.60 \pm 0.21)$ to the entire sample, with an orthogonal scatter of 0.19 dex (grey-shaded region), in good agreement with previous determinations (Fall 1983; Fall & Romanowsky 2013; Obreschkow & Glazebrook 2014; Cortese et al. 2016; Fall & Romanowsky 2018; Posti et al. 2018b; Murugesan et al. 2020; Mancera Piña et al. 2021a, b). A slope of 0.64 is very close to the value of $2/3$ predicted by tidal torque theories, indicating that f_j must be almost precisely a constant. The dashed line is the best fit to galaxies with $\log(M_*/M_\odot) \geq 10.5$, $\log j_* = (0.79 \pm 0.13) \log M_* - (5.23 \pm 0.23)$. As for the Tully–Fisher relations, our massive galaxies are broadly consistent with being on a single, unbroken relation, with the only significant outlier being, again, one of the two fastest rotators (2MFGC08638). We note that most of our galaxies lie slightly above the linear relation, which may suggest the beginning of a gradual curvature of the relation at the high-mass end. However, this slight bend towards larger j_* might be partially due to our secondary sample selection criterion (Section 2), which favoured galaxies with large disc sizes, hence with larger specific angular momenta ($j_* \propto R_d V_{\text{flat}}$).

7 DISCUSSION

Some previous studies have suggested that there might be a bend in the Tully–Fisher relation at the high-mass end (e.g. Peletier &




Figure 5. Fall relation ($j_* - M_*$) for massive spiral galaxies. Our sample is shown with red squares; cyan circles denote galaxies from the SPARC catalogue (Posti et al. 2018b). Solid and dashed black lines are linear fits in the form $\log j_* = \gamma \log M_* + \delta$ to all galaxies ($\gamma = 0.64$, $\delta = -3.60$) and to galaxies with $\log(M_*/M_\odot) \gtrsim 10.5$ ($\gamma = 0.79$, $\delta = -5.23$), respectively. The grey band marks the orthogonal scatter $\sigma_\perp = 0.19$ dex.

Willner 1993; Verheijen 2001; De Rijcke et al. 2007; Noordermeer & Verheijen 2007; Courtois et al. 2015; Ogle et al. 2019b). Most of these investigations focus on early-type or lenticular galaxies, which seem to lie on a relation that is offset from that of spiral galaxies (e.g. Williams, Bureau & Cappellari 2010; Cortesi et al. 2013) and/or appear to break at high-mass regime (e.g. Davis et al. 2016, however, see also den Heijer et al. 2015). The Tully–Fisher relation for massive discs is less certain, due to the scarcity of these systems. Noordermeer & Verheijen (2007) used a sample of 48 fairly massive galaxies ($M_* \lesssim 2 \times 10^{11} M_\odot$) with extended HI rotation curves and found that late-type discs (Spekkens & Giovanelli 2006) always lie on the stellar Tully–Fisher relation determined by less massive galaxies, while earlier-type discs (Noordermeer et al. 2007) seem to deviate from it. However, they also showed that this discrepancy is largely reduced when the asymptotic (flat) velocity is used instead of other velocity proxies, because earlier-type galaxies often have centrally peaked rotation curves that decline at large radii. The discrepancy disappears when using baryonic mass instead of stellar mass in the Tully–Fisher relation. Our galaxy sample, which extends to higher masses than that of Noordermeer & Verheijen (2007) and on average includes later-type discs with fairly flat rotation curves, confirms that the stellar and baryonic Tully–Fisher relations extend up to $M_* \simeq 6 \times 10^{11} M_\odot$ as unbroken power laws.

We use the analogous $M_h - V_h$ and $j_h - M_h$ relations for dark matter haloes as a reference for the observed Tully–Fisher and Fall relations, and we introduce the stellar mass fraction $f_M = M_*/M_h$, the angular momentum fraction $f_j = j_*/j_h$, and the ratio $f_V = V_{\text{flat}}/V_h$. With these three fractions, the Tully–Fisher and Fall relations become $M_* \propto f_M (V_{\text{flat}}/f_V)^3$ and $j_* \propto f_j (M_*/f_M)^{2/3}$. We use the statistical method of Posti et al. (2019b), on a galaxy sample consisting of their original sample plus our 43 massive spirals, to determine how the fractions f_M , f_j , and f_V vary with mass and to test for the presence

of bends in the observed scaling relations. The results of our new analysis are perfectly compatible with those of Posti et al. (2019b). In particular, we find that the data statistically prefer a model in which f_M is approximately proportional to $M_\star^{1/4}$, while both the angular momentum fraction f_j and the velocity fraction f_V are roughly constant with mass ($f_j \sim 0.7$, $f_V \sim 1$). Thus, on average, galactic discs have almost the same specific angular momentum as their dark matter haloes, independent of mass. This result is consistent with early models of disc formation (e.g. Fall & Efstathiou 1980; Dalcanton, Spergel & Summers 1997; Mo, Mao & White 1998), although the physical mechanisms responsible for it have not yet been fully elucidated (Romanowsky & Fall 2012; DeFelippis et al. 2017; Grand et al. 2019). On the other hand, a model with a prior imposed on f_M that follows the standard stellar-to-halo mass relation with turnover (e.g. Wechsler & Tinker 2018) is highly disfavoured by the observed scaling laws. Our statistical analysis confirms that f_M increases as a power law of M_\star for spiral galaxies up to the highest masses that we can measure, while, for lenticular and elliptical galaxies, f_M turns over at $M_\star \sim 5 \times 10^{10} M_\odot$ (Posti et al. 2019a; Posti & Fall 2021).

Ogle et al. (2019b), using a sub-sample of the galaxies analyzed in the present work (see Table 1), reported that the baryonic Tully–Fisher relation and the Fall relation break above a critical stellar mass $\log(M_\star/M_\odot) \simeq 11.5$, which seems to be in contrast with our findings. In this paper, we analyze a galaxy sample quite larger than Ogle et al.’s (43 versus 23), and we take advantage of a more sophisticated approach to derive galaxy kinematics. While we obtain rotation curves through a full modelling of the long-slit observations and we assume the V_{flat} as velocity estimator for the Tully–Fisher relations, Ogle et al. used the maximum rotation speed V_{max} measured directly from the flux-weighted centroid wavelength of the H α emission line. We stress that, although for some galaxies the V_{max} estimated by Ogle et al. (2019b) is slightly larger than our V_{flat} , for most galaxies their velocities are consistent with our within the errors. For the Fall relation, we calculated the specific angular momenta from the converged profiles $j_\star(R)$, while Ogle et al. used the estimator $j_\star = 2V_{\text{max}}R_d$ for pure exponential discs with scale radius R_d . Our $j_\star(R)$ profiles on average converge to the estimator for exponential discs quite well, as shown in Fig. 4. However, we found that the scale radii used in Ogle et al., which were taken from automated SDSS bulge-disc decomposition with fixed Sérsic index by Simard et al. (2011), are systematically larger than the values estimated here. This results in their estimates of j_\star being typically larger than ours. Ogle et al.’s (2019b) claim of a break in the scaling relations was mostly based on six outlier galaxies with very high-rotation velocities. We had to exclude two of these galaxies (OGC1304 and OGC0139) from our sample because a more attentive analysis showed that their current H α data are not good enough to determine an unambiguous rotation velocity, due to either asymmetric (OGC1304) or anomalous (OGC0139) H α emission (see Fig. A2). For two other galaxies, OGC0441 and 2MASXJ1123, we derived significantly smaller rotation speeds than Ogle et al.’s (384 versus 444 km s $^{-1}$ and 386 versus 436 km s $^{-1}$, respectively), and our measurements move these galaxies back on to the linear relations. In the case of OGC0441, the velocity discrepancy is mainly attributable to the different inclination angle, 44 $^\circ$ estimated in this work compared to 39 $^\circ$ assumed by Ogle et al. (2019b). The H α emission of 2MASXJ1123 is quite asymmetric, with the approaching side being less extended than the receding side and seemingly reaching higher velocities (see second row of Fig. 2); this might have affected the estimate of the systemic and maximum velocity in Ogle et al. (2019b). The last two galaxies, 2MFGC08638 and

2MFGC12344, are still outliers in the scaling relations determined in this paper, in agreement with Ogle et al.’s findings. However, we did not find evidence of any further extremely fast-rotating disc ($V_{\text{rot}} \gtrsim 400$ km s $^{-1}$) among the 20 massive spirals that we added to Ogle et al.’s (2019b) galaxy sample. Our analysis therefore leaves us with only two rapid rotators that appear to lie outside the Tully–Fisher and Fall relations, which is insufficient to test the claim that there is a break at $V_{\text{rot}} > 400$ km s $^{-1}$.

Massive discs are rare in the local Universe and fast rotators appear to be even rarer. Our current sample of 43 massive discs revealed only three galaxies with $V_{\text{flat}} > 400$ km s $^{-1}$, not enough to determine the shapes of the scaling relations in this range of extreme rotation velocities with any confidence. To detect or refute a bend at a statistically significant level, we would likely need ~ 10 – 20 galaxies with $V_{\text{flat}} > 400$ km s $^{-1}$. It might seem, at first glance, that this could be achieved with a relatively modest extension of the observing program described in this work. Unfortunately, however, it is difficult to know which galaxies are fast rotators until their rotation curves are measured. Because the space density of galaxies falls so rapidly with luminosity, size, and H I line width, it is necessary to observe many more galaxies that might have $V_{\text{flat}} > 400$ km s $^{-1}$ for every one that actually does exceed this limit. In practice, this means measuring the rotation curves of all the galaxies in a sample with selection criteria similar to those of the present sample, but ~ 5 – 10 times larger, amounting to ~ 200 – 400 galaxies. While not impossible, this would be a major undertaking.

The two galaxies in our sample with the highest rotational speeds, 2MFGC08638 and 2MFGC12344, are the only significant outliers in the scaling relations. The extreme rotation velocities (> 450 km s $^{-1}$) of these two galaxies place them on the high- V_{flat} side of the Tully–Fisher relations (Fig. 3) and the high- j_\star side of the Fall relation (Fig. 5). They have also exceptionally large stellar discs that extend up to 200 kpc in diameter in z band. These two galaxies are hard to reconcile with the relations determined at lower masses: either a significantly larger stellar mass or a significantly smaller asymptotic velocity would be required. Both these galaxies are highly inclined systems and we may expect some unaccounted dust absorption even at 3.4 μm , which may imply a M_\star/L_{W1} larger than the assumed value of 0.6. A mass-to-light ratio of $M_\star/L_{\text{W1}} \simeq 1.5$ would take these two galaxies back on the Tully–Fisher and Fall relations. Although such a high M_\star/L_{W1} cannot be completely ruled out on the basis of stellar population synthesis models (Fall & Romanowsky 2013; Into & Portinari 2013), it would imply stellar masses larger than $10^{12} M_\odot$, which would suspiciously make these two galaxies the most massive discs ever known. Another possibility is that these galaxies have rotation curves that sharply decline beyond the optical radius: a 20–30 per cent decrease in the asymptotic velocity would make them compatible with the Tully–Fisher relation. However, this kind of extreme decline has been rarely seen even in massive early-type discs (e.g. Noordermeer et al. 2007), and it seems unlikely to happen in late-type discs like 2MFGC08638 and 2MFGC12344. We also note that we ruled out the possibility of a scenario where the high-velocity H α emission is due to a peripheral ring of gas rather than to a filled galactic disc (see Section 5). H I data of the extended gas disc would be helpful to constrain better the asymptotic velocity of these two massive spiral galaxies.

Our findings suggest that most extremely massive spiral galaxies are following an evolutionary path similar to that of less massive spirals. These galaxies must have been continuing to assemble their stellar mass through secular evolution and minor mergers. Their current SFRs are of the order of 10 – $20 M_\odot \text{ yr}^{-1}$, which implies that these systems may have easily built stellar masses larger than

$10^{11} M_{\odot}$ throughout cosmic time, if we assume that, similarly to less massive spirals, they have been forming stars at a nearly constant or declining rate (Panter et al. 2007; Madau & Dickinson 2014). These galaxies must have experienced a relatively quiet evolution, without undergoing any major disruptive event, such as major mergers or strong AGN feedback, capable of transforming their disc-like morphology and of moving them off the scaling relations. Finally, our analysis indicates that star-forming galaxies have stellar/baryonic masses that scale with the dark matter haloes in a self-similar way up to the very high-mass end.

8 CONCLUSIONS

In this work, we used new $H\alpha$ -[N II] long-slit observations to study the kinematics of 43 extremely massive spiral galaxies with $\log(M_*/M_{\odot}) > 11$. We constructed 3D kinematic models of rotating discs; we extracted a long-slit mock observation from them and fitted it to our emission-line observations. This method allows us to deal with well-known biases due to instrumental response (PSF and spectrograph resolution) and to the high inclination angles of several galaxies in our sample, returning accurate rotation velocities and gas velocity dispersions. All galaxies analyzed show rotation curves that flatten within the optical radius, velocity dispersions of $25 - 30 \text{ km s}^{-1}$, typical of the ionized gas component, and $V_{\text{rot}}/\sigma_{\text{gas}} \gg 10$. Two galaxies in our sample (2MFGC08638 and 2MFG12344) are fast rotators ($V_{\text{flat}} > 450 \text{ km s}^{-1}$), with the most extreme galaxy, 2MFG12344, reaching an impressive rotation velocity of $\simeq 530 \text{ km s}^{-1}$ in the outer regions of the optical disc.

From the derived kinematics, we built the stellar and baryonic Tully–Fisher relations and the Fall relation. To be consistent with the latest determinations of these scaling laws from HI observations, we used the flat part of the rotation curve as velocity indicator for the Tully–Fisher relation and we estimated the total stellar specific angular momentum from the converged $j_*(R)$ profiles. In contrast with some previous results, we found that massive discs lie on both the Tully–Fisher and the Fall relations determined for less massive spiral galaxies, without any strong evidence for a bend or break at the high-mass end. The only marginal outliers are the two fastest rotators, which might be difficult to reconcile with the known scaling relations. While a larger population of such outliers could exist at $\log(M_*/M_{\odot}) > 11.5$, with the data currently available we ruled out the presence of a break at the high-mass end of the Tully–Fisher and Fall relations. The kinematics and scaling relations derived in this paper indicate that most of these giant discs are scaled up versions of regular spiral galaxies.

ACKNOWLEDGEMENTS

EMDT was supported by the US National Science Foundation under grant 1616177. LP acknowledges support from the Centre National d’Études Spatiales (CNES) and from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement no. 834148). This paper utilizes data from the Southern African Large Telescope (SALT), the Hale Telescope at Palomar Observatory, the Wide-field Infrared Survey Explorer (WISE), the Sloan Digital Sky Survey (SDSS), and the Legacy Surveys. WISE is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by NASA. Funding for the Sloan Digital Sky Survey IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the participating institutions. The Legacy Surveys consist of three

individual and complementary projects: the Dark Energy Camera Legacy Survey (DECaLS), the Beijing–Arizona Sky Survey (BASS), and the Mayall z-band Legacy Survey (MzLS), conducted using facilities supported by NSF’s OIR Lab, which is operated by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation. This research made use of Photutils, an Astropy package for detection and photometry of astronomical sources. This work relied on the NASA/IPAC Extragalactic Database, operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA.

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

- Andersen D. R., Bershadsky M. A., Sparke L. S., Gallagher J. S., II, Wilcots E. M., van Driel W., Monnier-Ragaigne D., 2006, *ApJS*, 166, 505
- Astropy Collaboration, 2013, *A&A*, 558, A33
- Astropy Collaboration, 2018, *AJ*, 156, 123
- Begeman K. G., 1987, PhD thesis, Kapteyn Institute, Groningen (The Netherlands)
- Bekeraite S. et al., 2016, *ApJ*, 827, L36
- Bloom J. V. et al., 2017, *MNRAS*, 472, 1809
- Bradley L., Sipocz B., Robitaille T., Erik T. et al., 2020, *astropy/photutils*: 1.0.0, <https://doi.org/10.5281/zenodo.4044744>
- Buckley D. A. H., Swart G. P., Meiring J. G., 2006, *Proc. SPIE*, 6267, 62670Z
- Bullock J. S., Dekel A., Kolatt T. S., Kravtsov A. V., Klypin A. A., Porciani C., Primack J. R., 2001, *ApJ*, 555, 240
- Burgh E. B., Nordsieck K. H., Kobulnicky H. A., Williams T. B., O’Donoghue D., Smith M. P., Percival J. W., 2003, *Proc. SPIE*, 4841, 1463
- Cattaneo A., Salucci P., Papastergis E., 2014, *ApJ*, 783, 66
- Cluver M. E. et al., 2014, *ApJ*, 782, 90
- Cluver M. E., Jarrett T. H., Dale D. A., Smith J. D. T., August T., Brown M. J. I., 2017, *ApJ*, 850, 68
- Cortese L. et al., 2016, *MNRAS*, 463, 170
- Cortesi A. et al., 2013, *MNRAS*, 432, 1010
- Courtois H. M., Zaritsky D., Sorce J. G., Pomarède D., 2015, *MNRAS*, 448, 1767
- Dalcanton J. J., Spergel D. N., Summers F. J., 1997, *ApJ*, 482, 659
- Davis T. A., Greene J., Ma C.-P., Pandya V., Blakeslee J. P., McConnell N., Thomas J., 2016, *MNRAS*, 455, 214
- de los Reyes M. A. C., Kennicutt R. C. Jr, 2019, *ApJ*, 872, 16
- De Rijcke S., Zeilinger W. W., Hau G. K. T., Prugniel P., Dejonghe H., 2007, *ApJ*, 659, 1172
- DeFelippis D., Genel S., Bryan G. L., Fall S. M., 2017, *ApJ*, 841, 16
- Dekel A., Silk J., 1986, *ApJ*, 303, 39
- den Heijer M. et al., 2015, *A&A*, 581, A98
- Dey A. et al., 2019, *AJ*, 157, 168
- Di Teodoro E. M., Fraternali F., 2015, *MNRAS*, 451, 3021
- Di Teodoro E. M. et al., 2018, *MNRAS*, 476, 804
- Dutton A. A., van den Bosch F. C., 2012, *MNRAS*, 421, 608
- Eisenstein D. J. et al., 2011, *AJ*, 142, 72
- Fall S. M., 1983, in Athanassoula E., ed., *Internal Kinematics and Dynamics of Galaxies*, Vol. 100. Reidel Publishing Co., Dordrecht (The Netherlands), p. 391
- Fall S. M., Efstathiou G., 1980, *MNRAS*, 193, 189
- Fall S. M., Romanowsky A. J., 2013, *ApJ*, 769, L26
- Fall S. M., Romanowsky A. J., 2018, *ApJ*, 868, 133
- Ferrero I. et al., 2017, *MNRAS*, 464, 4736
- Fouque P., Bottinelli L., Gouguenheim L., Paturel G., 1990, *ApJ*, 349, 1
- Fraternali F., Sancisi R., Kamphuis P., 2011, *A&A*, 531, A64
- Giovanelli R., Haynes M. P., Rubin V. C., Ford W. K. J., 1986, *ApJ*, 301, L7

- Grand R. J. J. et al., 2019, *MNRAS*, 490, 4786
- Huyer W., Neumaier A., 1998, *J. Glob. Optim.*, 14, 331
- Into T., Portinari L., 2013, *MNRAS*, 430, 2715
- Jarrett T. H., Chester T., Cutri R., Schneider S., Skrutskie M., Huchra J. P., 2000, *AJ*, 119, 2498
- Jarrett T. H., Cluver M. E., Brown M. J. I., Dale D. A., Tsai C. W., Masci F., 2019, *ApJS*, 245, 25
- Jedrzejewski R. I., 1987, *MNRAS*, 226, 747
- Karachentsev I. D., Mitronova S. N., Karachentseva V. E., Kudrya Y. N., Jarrett T. H., 2002, *A&A*, 396, 431
- Kennicutt R. C. Jr, 1998, *ApJ*, 498, 541
- Kent S. M., 1987, *AJ*, 93, 816
- Kent S. M., 1988, *AJ*, 96, 514
- Lapi A., Salucci P., Danese L., 2018, *ApJ*, 859, 2
- Lelli F., McGaugh S. S., Schombert J. M., 2016a, *AJ*, 152, 157
- Lelli F., McGaugh S. S., Schombert J. M., 2016b, *ApJ*, 816, L14
- Lelli F., McGaugh S. S., Schombert J. M., Desmond H., Katz H., 2019, *MNRAS*, 484, 3267
- Madau P., Dickinson M., 2014, *ARA&A*, 52, 415
- Mancera Piña P. E., Posti L., Fraternali F., Adams E. A. K., Oosterloo T., 2021a, *A&A*, 647, A76
- Mancera Piña P. E., Pezzulli G., Fraternali F., Fall S. M., Oosterloo T., Adams E. A. K., 2021b, *A&A*, 651, L15
- Martinsson T. P. K., Verheijen M. A. W., Westfall K. B., Bershadsky M. A., Schechtman-Rook A., Andersen D. R., Swaters R. A., 2013, *A&A*, 557, A130
- McGaugh S. S., 2005, *ApJ*, 632, 859
- McGaugh S. S., 2012, *AJ*, 143, 40
- McGaugh S. S., Schombert J. M., 2015, *ApJ*, 802, 18
- McGaugh S. S., Schombert J. M., Bothun G. D., de Blok W. J. G., 2000, *ApJ*, 533, L99
- Meidt S. E. et al., 2014, *ApJ*, 788, 144
- Mo H. J., Mao S., White S. D. M., 1998, *MNRAS*, 295, 319
- Mo H., van den Bosch F. C., White S., 2010, *Galaxy Formation and Evolution*. Cambridge University Press, Cambridge, UK
- Murugesan C., Kilborn V., Jarrett T., Wong O. I., Obreschkow D., Glazebrook K., Cluver M. E., Fluke C. J., 2020, *MNRAS*, 496, 2516
- Naluminsa E., Elson E. C., Jarrett T. H., 2021, *MNRAS*, 502, 5711
- Navarro J. F., Steinmetz M., 2000, *ApJ*, 538, 477
- Noordermeer E., Verheijen M. A. W., 2007, *MNRAS*, 381, 1463
- Noordermeer E., van der Hulst J. M., Sancisi R., Swaters R. S., van Albada T. S., 2007, *MNRAS*, 376, 1513
- Norris M. A., Meidt S., Van de Ven G., Schinnerer E., Groves B., Querejeta M., 2014, *ApJ*, 797, 55
- Obreschkow D., Glazebrook K., 2014, *ApJ*, 784, 26
- Ogle P. M., Lanz L., Nader C., Helou G., 2016, *ApJ*, 817, 109
- Ogle P. M., Lanz L., Appleton P. N., Helou G., Mazzarella J., 2019a, *ApJS*, 243, 14
- Ogle P. M., Jarrett T., Lanz L., Cluver M., Alatalo K., Appleton P. N., Mazzarella J. M., 2019b, *ApJ*, 884, L11
- Panther B., Jimenez R., Heavens A. F., Charlot S., 2007, *MNRAS*, 378, 1550
- Parkash V., Brown M. J. I., Jarrett T. H., Bonne N. J., 2018, *ApJ*, 864, 40
- Peebles P. J. E., 1969, *ApJ*, 155, 393
- Peletier R. F., Willner S. P., 1993, *ApJ*, 418, 626
- Persic M., Salucci P., Stel F., 1996, *MNRAS*, 281, 27
- Planck Collaboration XIII, 2016, *A&A*, 594, A13
- Ponomareva A. A., Verheijen M. A. W., Papastergis E., Bosma A., Peletier R. F., 2018, *MNRAS*, 474, 4366
- Posti L., Fall S. M., 2021, *A&A*, 649, A119
- Posti L., Pezzulli G., Fraternali F., Di Teodoro E. M., 2018a, *MNRAS*, 475, 232
- Posti L., Fraternali F., Di Teodoro E. M., Pezzulli G., 2018b, *A&A*, 612, L6
- Posti L., Fraternali F., Marasco A., 2019a, *A&A*, 626, A56
- Posti L., Marasco A., Fraternali F., Famaey B., 2019b, *A&A*, 629, A59
- Reyes R., Mandelbaum R., Gunn J. E., Pizagno J., Lackner C. N., 2011, *MNRAS*, 417, 2347
- Röck B., Vazdekis A., Peletier R. F., Knapen J. H., Falcón-Barroso J., 2015, *MNRAS*, 449, 2853
- Romanowsky A. J., Fall S. M., 2012, *ApJS*, 203, 17
- Schmidt M., 1959, *ApJ*, 129, 243
- Simard L., Mendel J. T., Patton D. R., Ellison S. L., McConnachie A. W., 2011, *ApJS*, 196, 11
- Skrutskie M. F. et al., 2006, *AJ*, 131, 1163
- Spekkens K., Giovanelli R., 2006, *AJ*, 132, 1426
- Torres-Flores S., Epinat B., Amram P., Plana H., Mendes de Oliveira C., 2011, *MNRAS*, 416, 1936
- Tully R. B., Fisher J. R., 1977, *A&A*, 500, 105
- van Albada T. S., Bahcall J. N., Begeman K., Sancisi R., 1985, *ApJ*, 295, 305
- Verheijen M. A. W., 2001, *ApJ*, 563, 694
- Wechsler R. H., Tinker J. L., 2018, *ARA&A*, 56, 435
- Williams M. J., Bureau M., Cappellari M., 2010, *MNRAS*, 409, 1330
- Wright E. L. et al., 2010, *AJ*, 140, 1868

SUPPORTING INFORMATION

Supplementary data are available at *MNRAS* online.

Figure A1. Surface photometry and kinematic analysis of 43 massive spiral galaxies.

Figure A2. The eight galaxies excluded from our kinematic analysis. [paper_ss_appendix.pdf](#)

Appendix A. Atlas of extremely massive spiral galaxies.

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.