

The histone demethylase KdmB is part of a trimeric protein complex and mediates virulence and patulin production in *Penicillium expansum*

Dianiris Luciano-Rosario¹, Omer Barda², Joanna Tannous³, Dean Frawley⁴, Özgür Bayram⁴, Edward Sionov², Dov Prusky², and Nancy P. Keller^{1,5}

¹Department of Plant Pathology, University of Wisconsin, Madison, Wisconsin, USA

²Institute of Postharvest and Food Sciences, The Volcani Center, Agricultural Research Organization, Rishon LeZion, Israel

³Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

⁴Faculty of Science and Engineering, National University of Ireland Maynooth, Kildare, Ireland

⁵Department of Medical Microbiology and Immunology, University of Wisconsin, Madison, Wisconsin, USA

*corresponding author

Abstract

Epigenetic modification of chromosome structure has increasingly been associated with alterations in secondary metabolism and sporulation defects in filamentous fungal pathogens. Recently, the epigenetic reader protein SntB was shown to govern virulence, spore production and mycotoxin synthesis in the fruit pathogen *Penicillium expansum*. Through immunoprecipitation-coupled mass spectrometry, we find that SntB is a member of a protein complex with KdmB, a histone demethylase and the essential protein RpdA, a histone deacetylase. Deletion of *kdmB* phenocopied some but not all characteristics of the Δ *sntB* mutant. KdmB deletion strains exhibited reduced lesion development on Golden Delicious apples and this was accompanied by decreased production of patulin in host tissue. In contrast to SntB loss, KdmB loss did not impact expression of three regulators (*laeA*, *pacC* and *creA*) of *P. expansum* virulence. However, Δ *kdmB* mutants were sensitive to several cell wall stressors which possibly contributed to the decreased virulence observed on apples. Slight differences in spore production and germination rates of Δ *kdmB* mutants *in vitro* did not impact overall diameter growth in culture.

Keywords: epigenetics, virulence, secondary metabolism, fungal biology, post-harvest disease

Introduction

Plant pathogenic filamentous fungi mediate interactions with their environments utilizing different strategies. Necrotrophic pathogens strategies include tissue acidification of the host tissues, cell wall degrading enzyme secretion, effector proteins delivery and secondary metabolite production and secretion (van Kan, 2006). Secondary metabolites are small bioactive molecules that are produced by diverse organisms including pathogenic filamentous fungi (Keller, 2019). These molecules are diverse in structure and function and mediate diverse ecological interactions. Many secondary metabolites have beneficial properties and have been utilized by humans as drugs or agricultural treatments, but others can have detrimental effects and are classified as mycotoxins.

Penicillium expansum is a post-harvest pathogen of pome fruits and the causative agent of blue mold disease. This necrotrophic pathogen produces the mycotoxins patulin and citrinin which pose a food safety threat to consumers of apples and their processed food products (Luciano-Rosario *et al.*, 2020). Patulin is difficult to remove from contaminated food products due to its heat and acidity resistance and thus the threshold levels imposed by international agencies like the World Health Organization WHO, FDA, and EU often lead to destruction of contaminated products (Ioï *et al.*, 2017). Other secondary metabolites produced by *P. expansum* include andrastins, roquefortine, chaetoglobosins, and communesins (Tannous *et al.*, 2018), although these compounds are only occasionally reported to contaminate apple products and their potential health impacts to consumers are not characterized (Olsen *et al.*, 2019; Andersen *et al.*, 2004; Larsen *et al.*, 1998; Tannous *et al.*, 2018). The production of secondary metabolites is regulated by diverse mechanisms at a molecular level. Endogenous regulatory mechanisms include regulation of secondary metabolite gene clusters – also known as biosynthetic gene clusters (BGCs) - by cluster-

specific transcription factors or by proteins that mediate chromatin accessibility of the transcriptional machinery.

In filamentous fungi, chromatin regulation couples BGC gene expression and thus secondary metabolite production with developmental processes such as virulence, sporulation, and resistance to stressors (Choi *et al.*, 2022; Strauss and Cánovas, 2021; Bachleitner *et al.*, 2021; Wu *et al.*, 2021). Typically, chromatin regulation occurs through histone post-translational modifications (PTMs) to histone tails (Pfannenstiel and Keller 2019). There are many possible PTMs such as acetylation, methylation, and phosphorylation among others and these modifications are created by protein complexes that physically interact with and modify histone tails. The different classes of histone tail modification proteins include “readers” which are proteins that identify a histone tail modification and mediate the response to the modification, “writers” or proteins that add modifications to histone tails and “erasers” or proteins that remove histone tail modifications (Pfannenstiel and Keller 2019).

The first reader protein to be identified in filamentous fungi was SntB, a homolog of Snt2 in *Saccharomyces cerevisiae* which was found to interact with an eraser enzyme, the histone deacetylase (HDAC) Rpd3 (Baker *et al.*, 2013). Through a random mutagenesis approach, SntB was identified in *Aspergillus nidulans* as a regulator of the mycotoxin sterigmatocystin (ST) (Pfannenstiel *et al.*, 2017). Homologs of SntB in filamentous pathogens have been shown to mediate virulence in *Fusarium oxysporum* (Denisov *et al.*, 2011) and *A. flavus* (Pfannenstiel *et al.*, 2019) where, in the latter species, loss of SntB also resulted in decreased aflatoxin synthesis but increased production of normally unexpressed metabolites. Recently, we described the role of

SntB in virulence, secondary metabolism and development of *P. expansum* where the $\Delta sntB$ mutant was altered in conidiation, germination rate, virulence and patulin production in apples (Tannous *et al.*, 2020).

Considering reader proteins partner with ‘erasers’ or ‘writers’ in yeast, efforts next focused on determining the SntB complex in filamentous fungi. Using a TAP-purification and mass-spec analysis, SntB was found to interact with the HDAC RpdA (homolog of Rpd3) and the histone demethylase KdmB in *A. nidulans* (Karahoda *et al.*, 2022). Further TAP-purification using KdmB showed it complexed with SntB and the “writer” enzyme EcoA thus establishing a multimeric KERS complex (Karahoda *et al.*, 2022). Deletion of KdmB presented considerable similarity to the SntB deletion phenotype in *A. nidulans* whereas RpdA deletion yielded a more severe phenotype and EcoA is an apparent essential protein and could not be deleted.

Here we sought to determine if a similar SntB complex could operate in *P. expansum* and, if so, if loss of any complex members impact fungal development and/or virulence of *P. expansum*. Through immunoprecipitation and mass spec analysis we found SntB interacts with KdmB and RpdA, similarly as in *A. nidulans*. Although unable to delete *rpda*, presumably due to lethality, we generated and assessed three $\Delta kdmB$ strains for fungal physiology, stress resistance, secondary metabolite production and virulence on apple. The mutant strains presented a similar phenotype as $\Delta sntB$ most notably with a decrease in virulence and mycotoxin synthesis on apple.

Materials and Methods

Background Strain and Culture Conditions

The experiments were conducted using the *P. expansum* Pe21 as the background strain (Hadas *et al.*, 2007). For all experiments, conidia were obtained from a fresh culture on Glucose Minimal Media (GMM) incubated for 5 days at 25°C. The conidia were harvested using a solution of sterile 0.01% Tween 80 and quantified using a hemacytometer.

$\Delta kdmB$ and 3XFLAG-tagged SntB knock-in Strains Construct Generation

To design and generate the transformation cassettes for the $\Delta kdmB$ strains, we used double joint PCR as described by Lim *et al.*, 2012. Briefly, 1.5kb upstream and downstream of the *kdmB* open reading frame (ORF) (PEXP_096530) were amplified using Pfu DNA polymerase (Agilent, Santa Clara, CA, United States). In addition, *A. fumigatus pyrG* was amplified in the same manner. Later, the obtained PCR products were gel extracted and combined in a (1:3:1) ratio for double joint PCR. The resulting PCR product was used as template DNA to amplify the construct which was purified using a G-50 column. The primer sequences are listed on Table S1A. A 3XFLAG-tagged SntB (PEXP_027800) knock-in strain was also generated in order to determine the subcellular localization of SntB and its interacting proteins. An epitope tagging of the protein was done with a FLAG tag at its C-terminus in the *P. expansum* Pe-21 WT strain. The 3X-Flag tag sequence is inserted upstream of the *sntB* stop codon through homologous recombination. A PCR fragment carrying the 3x-Flag sequence (GACTACAAAGACCATGACGGTGATTATAAAGATCATGATATCGATTACAAGGATGACGATGACAAG) and the TrpC terminator was amplified from plasmid psk529 (Hartmann *et al.*, 2010; Jimenez-Ortigosa *et al.*, 2012) using the primer pair FLAG:trpC(t)_F and FLAG:trpC(t)_R. This PCR product was fused to the hygromycin resistance gene *hph* under control of the *gpdA* promoter amplified from plasmid psk529 (Hartmann *et al.* 2010; Jimenez-Ortigosa *et al.*,

2012) using the primer pair Hyg_Flag F and Hyg_Flag R. The fragment containing the hygromycin resistance gene as a selectable marker and the 3X- flag sequence was used as the middle fragment in the double joint fusion PCR. The 5' fragment corresponds to the last 1kb of the *sntB* transcript missing the stop codon. This fragment was amplified using the primer pair PesntB_flagtag_5'F/R. The last fragment corresponds to the 3' flank of the *sntB* gene and was amplified using the primer pair PesntB_flagtag_3'F/R. The diagram of this construct is shown in Fig. S1C. The three fragments were fused together using a double joint PCR protocol described by Lim *et al.* (2012). All primers used to make the construct can be found in Table S1. In addition, we attempted to generate $\Delta rpdA$ and *rpdA*^{KD} (PEXP_031050) strains but were unsuccessful leading to infer that this locus is essential for *P. expansum* as it is for *Aspergillus nidulans* (Karahoda *et al.*, 2022)

Protoplast Isolation and PEG-mediated Transformation

To obtain transformants, we used gene replacement by homologous recombination as previously described by Greco *et al.*, 2019 with modifications described in Luciano-Rosario *et al.*, 2022. We used TDL 9.1, a $\Delta pyrG$ mutant, as a background strain for generating the $\Delta kdmB$ strains. The obtained transformants were screened and confirmed by PCR and Southern Blot. All the strains used in this study are listed in Table S1B.

For the 3XFLAG-tagged SntB knock-in strains, the protoplast transformation of *P. expansum* WT strain was conducted as described in Tannous *et al.* 2018. Transformants were screened on SMM medium supplemented with hygromycin at a concentration of 100 µg/mL. Ten transformants were randomly picked and subjected to southern blotting to confirm the correct tagging of *sntB* and the

single insertion of the construct. Only one transformant (labeled TJT23.1) was chosen for further study (Fig. S1D).

Physiology: Radial Growth, Conidiation, and Germination

For radial growth measurements, point inoculations were performed using 10^6 spores in GMM and the colony diameter was measured for 7 days. For conidiation assay, 10^6 spores were overlaid in GMM plates. Then, agar plugs were collected, homogenized and the spores quantified using the hemocytometer. For evaluating germination, 1mL of 1×10^5 spores/mL in liquid GMM were added to wells on a 24 well-plate. A total of 50 spores were tracked for 23 hours after inoculation and the percent germination was recorded every hour. Strains were assessed in triplicate.

Cell Wall Stressor Assay

GMM plates were prepared containing the assessed cell wall stressors: Calcofluor White (CFW), Congo Red (CR) and Sodium Dodecyl Sulfate SDS at the described final concentrations. Point inoculations at different spore concentrations (10^5 , 10^4 , and 10^3) were photographed 3 days post inoculation and compared to cultures in non-amended GMM plates.

Mycotoxin extraction from cultured media

A 10^6 fungal conidia/ml solution (100 μ l) was inoculated onto 55 mm petri dishes containing 10 ml of solid YES media (20 g bacto yeast extract, 150 g sucrose and 15 g bacto agar per liter). The plates were incubated at 28°C in the dark for 3-7 days as needed for sample collection. pH was measured directly in the agar cultures with a double pore slim electrode connected to a Sartorius PB-11 Basic Meter (Sartorius, Göttingen, Germany).

Patulin was extracted from 1 g of the homogenized medium containing fungal mycelium, which was mixed with 2 mL of HPLC grade ethyl acetate (BioLab, Jerusalem, Israel) and placed in an orbital shaker at 150 rpm for 30 min at room temperature. The sample was centrifuged for 10 min at 6000 g; the supernatant was transferred to a clean glass tube and was evaporated to dryness under a stream of gaseous nitrogen at 50 °C. The residue was redissolved in 1 mL of the mobile phase (0.02 M ammonium acetate:acetonitrile 9:1 v/v) and filtered through a 0.22 µM PTFE syringe filter prior to HPLC analysis.

Citrinin was extracted from 1 g of the homogenized YES medium with fungal mycelium using 1 mL of methanol (HPLC grade; BioLab, Jerusalem, Israel). The sample was shaken in an orbital shaker at 150 rpm for 30 min at room temperature and then centrifuged for 10 min at 6000 g. The supernatant was filtered through a 0.22 µM PTFE syringe filter and kept at –20 °C prior to HPLC analysis.

RNA isolation and qRT-PCR analyses

Mycelia grown on agar plates were harvested on day 3 post-inoculation, frozen in liquid nitrogen, lyophilized for 24 hours and kept at –80 °C until use. Total RNA was extracted from 100 mg of lyophilized tissue of the selected samples using the Hybrid-R RNA isolation kit (GeneAll, Seoul, South Korea) according to the manufacturer's protocol. The DNase and reverse-transcription reactions were performed on 1 µg of total RNA with the Maxima First-Strand cDNA Synthesis Kit (Thermo Scientific, Waltham, MA, United States) according to the manufacturer's instructions. The cDNA samples were diluted 1:10 (v/v) with ultrapure water.

The quantitative real-time PCR was performed using Fast SYBR green Master Mix (Applied Biosystems, Waltham, MA, USA) in a StepOnePlus Real-Time PCR System (Applied Biosystems, Waltham, MA, USA). The PCR conditions were as follows: 95 °C for 20 sec, followed by 40 cycles of 95 °C for 3 sec and 60 °C for 20 sec. The samples were normalized using *β-tubulin* as endogenous control and the relative expression levels were measured using the $2^{(-\Delta\Delta C_t)}$ analysis method. Results were analyzed with StepOne software v2.3. Primer sequences used for qRT-PCR analysis are listed in Table S1C.

Virulence and mycotoxin assessment in apples

Golden Delicious apples were obtained from a local supermarket. Fruits were subjected to surface sterilization using 1% sodium hypochlorite solution for 1 min and immediately rinsed in sterile distilled water. A 10 µL conidial suspension containing 10^7 conidia/mL was injected directly into the sterilized fruits at 2 mm depth. Following inoculation, the fruits were incubated in covered plastic containers at 28 °C for 16 days for symptom monitoring and sample collection. The diameter of the rotten spots was recorded at selected time points.

Patulin was extracted from 1 g of homogenized apple tissue samples using 2 mL ethyl acetate in a 15 mL glass tube. Each sample was vortexed for at least 5 min; then, 2 mL of 1.5% sodium bicarbonate solution was added, and the sample was vortexed vigorously again for 1 min. Next, 20 µL of acetic acid was added, and after shaking in an orbital shaker at 250 rpm for 5 min, the sample was left to stand for a few minutes at room temperature for phase separation. The upper phase was transferred to a new glass tube and evaporated to dryness under a stream of gaseous nitrogen at 50 °C. The samples were reconstituted in the mobile phase (0.02 M ammonium

acetate:acetonitrile 9:1 v/v) and filtered through a 0.22 µm PTFE syringe filter into a glass injection vial to prepare for the HPLC analysis.

Citrinin was extracted from 1 g of homogenized apple tissue using 2 mL of methanol. Each sample was then vortexed for 5 min and shaken using an orbital shaker at 250 rpm for 30 min before centrifuging at 6000 g for 10 min. The aliquots were filtered through a 0.22 µm PTFE membrane syringe filter into a glass injection vial and stored at -20 °C prior to HPLC analysis.

HPLC analysis

Both patulin and citrinin were quantitatively analyzed by injection of 20 µL filtered samples into a reverse-phase UHPLC system (Waters ACQUITY Arc, FTN-R, Milford, MA, USA) using a Kinetex 3.5 µm XB-C18 (150 × 4.6 mm) column (Phenomenex, Torrance, CA, USA). The column temperature was maintained at 30 °C, and the flow rate was 0.8 and 1 ml/min, respectively. For patulin, the separation was achieved with a mobile phase composed of 0.02 M ammonium acetate:acetonitrile (9:1 v/v); citrinin mobile phase consisted of 1% acetic acid in water and acetonitrile (50:50 v/v). The patulin peak was detected with a photodiode array (PDA) detector at 276 nm; citrinin was detected with a fluorescence detector (331 nm excitation, 500 nm emission). Both mycotoxins were quantified by comparing peak areas with calibration curves of the standard mycotoxins (Fermentek, Jerusalem, Israel).

Isolation of crude protein extracts for pull-down assay

The mycelia from vegetative cultures were frozen using liquid nitrogen and pulverised with a mortar and pestle. Protein crude extracts were prepared by re-suspending the pulverised mycelia

in 1ml protein extraction buffer (300mM NaCl, 50mM Tris-HCl pH 7.5, 10% glycerol, 1mM EDTA, 0.1% NP-40) that had been supplemented with 1mM DTT, 1X complete EDTA-free protease inhibitors (Roche), 1mM benzamidine, 0.5mM PMSF and 1X phosphatase inhibitors (1mM NaF, 0.5mM sodium orthovanadate, 8mM β -glycerol phosphate) immediately prior to use. Samples were mixed vigorously by vortexing and centrifuged at 13,000 RPM for 10-15 minutes at 4°C. 1ml of the crude protein supernatant was transferred to a new 1.5ml microcentrifuge tube.

Immunoprecipitation of FLAG-tagged proteins

For the immunoprecipitation of FLAG-tagged fusion proteins, 20 μ l anti-FLAG M2 magnetic beads (Sigma-M8823) were washed twice with 180 μ l protein extraction buffer, containing supplements. After each wash, the bead solution was placed on a magnetic rack and the protein extraction buffer was removed. The anti-FLAG M2 magnetic beads were then resuspended in 50 μ l protein extraction buffer and added to 1ml crude protein extract. This mixture was left to incubate on a rotator for 3 hours at 4°C. Samples were placed in a magnetic rack and the supernatant was discarded. Beads were washed twice with 1ml protein extraction buffer (without supplements) and were then washed for a third time with the same buffer containing 1mM DTT. All liquid was removed and the beads were stored at -80°C until further use.

Sample preparation for LC-MS/MS peptide identification

Isolated FLAG-tagged proteins were resuspended in 50mM ammonium bicarbonate. 1 μ l of 0.5M DTT was added and samples were incubated at 56°C for 20 minutes. 2.7 μ l of iodoacetamide (0.55M) was added and samples were incubated in the dark for 15 minutes. 1 μ l of 1% (w/v) ProteaseMAX (Promega) was added, followed by addition of 1 μ l trypsin (1 μ g/ μ l) (Promega).

Samples were left to incubate overnight at 37°C. The next day, 1µl of Trifluoroacetic acid (TFA) was added to each and samples were vortexed briefly and left to incubate for 5 minutes at room temperature. Beads were collected on a magnetic rack and the supernatant was transferred to new tubes. The supernatants were centrifuged at 13,000 RCF for 10 minutes and dried in a SpeedVac for 3 hours. Samples were stored at -20°C until further use.

Peptide samples were resuspended in 20 µl resuspension buffer (0.5% TFA) and sonicated for 3 minutes, followed by a brief centrifugation. ZipTip C₁₈ pipette tips (Millipore) were used to purify peptide samples prior to mass spectrometric analysis. To equilibrate the ZipTips, a wetting solution (0.1%, 80% acetonitrile) was aspirated 5 times, followed by aspiration of an equilibration buffer (0.1% TFA) 5 times. ZipTips were then used to pipette the peptide samples up and down 15-20 times. Then, the equilibration buffer was aspirated again 5 times, followed by elution of the peptides via aspiration of an elution buffer (0.1% TFA, 60% acetonitrile) 5 times into a new microcentrifuge tube. This solution was dried in a SpeedVac for 2 hours and peptide samples were stored at -20°C.

LC-MS/MS analysis and peptide identification

Immediately prior to loading, peptide samples were resuspended in 10µl Q-Exactive loading buffer (2% acetonitrile, 0.5% TFA) and 8µl were added to mass spectrometry vials (VWR). Digested peptides were separated using reverse phase liquid chromatography with an Ultimate 3000 NanoLC system (Dionex Corporation, Sunnyvale, CA, USA) followed by mass identification using a Q-Exactive mass spectrometer (Thermo Fisher Scientific). Samples were loaded by an autosampler onto a C18 trap column, which was switched on-line with an analytical Biobasic C18

Picofrit column (C18 PepMap, 75 μm id $\times$ 500 mm, 2 μm particle size, 100 Å pore size; Dionex). The peptides generated were eluted over 65 minutes. Full MS scans in the 300-1700 m/z range were recorded with a resolution of 140,000 (m/z 200) with a blocking mass set to 445.12003.

LC-MS identification of peptides were performed using the Proteome Discoverer Daemon 1.4 software (Thermo Fisher) and organism-specific taxon-defined protein databases. As controls, anti-FLAG magnetic beads were added to crude protein extracts from wild type strains. These samples were prepared for mass spectrometry analysis as described for the immunoprecipitated FLAG-tagged fusion proteins. Confirmation of protein interactions and unique peptides were determined by isolating only those that appear in the FLAG-tagged fusion protein purifications but do not appear in any of the wild type controls.

Results

KdmB interacts with SntB and RpdA in a trimeric complex

Fungal secondary metabolite BGCs, including the *pat* (patulin) and *cit* (citrinin) BGCs, are regulated by histone PTMs in response to various external stimuli. Following previous evidence that SntB loss in *P. expansum* resulted in decreased patulin and citrinin synthesis and decreased virulence in apples (Tannous *et al.*, 2020), and that SntB was a member of the KERS complex in *A. nidulans* (Karahoda *et al.*, 2022), we asked if we could identify any SntB interacting proteins and whether these proteins could also impact virulence and mycotoxin synthesis in *P. expansum*. In order to test this, we created a flag-tagged version of *sntB* to replace the native locus. One correct transformant, TJT23.1, was used for the protein immunoprecipitation experiments which were coupled with mass spectrometry analysis (Fig. S1D).

Four separate SntB-FLAG immunoprecipitations were analyzed by LC-MS/MS analysis. In all 4 replicates, SntB recruited the same top two proteins (Fig. 1A): a homolog of *A. nidulans* histone demethylase KdmB (PEXP_096530) and a homolog of *A. nidulans* histone deacetylase RpdA (PEXP_03150). The percentage of coverage and numbers of unique peptides for these two proteins were significantly high and were consistent across all replicates (Fig. 1A, Table S2). This provides strong evidence that SntB exhibits direct interactions with both KdmB and RpdA in *P. expansum*, forming a trimeric complex. The *P. expansum* KdmB and RpdA proteins exhibit 70.91% and 70.01% sequence similarity to the *A. nidulans* homologs respectively, implying a high level of conservation between the two species.

The *P. expansum* KdmB protein is a large multi-domain protein of 1,737 amino acids (aa), highly similar to *A. nidulans* KdmB, which is 1,717 aa (Fig. 1B). According to Uniprot (<https://www.uniprot.org>), *P. expansum* KdmB possesses the following domains: JmjN at aa position 71-112, ARID (A/B) at aa position 164-257, 2 PHD-type domains, one at aa position 476-526 and the other at aa position 1344-1393 and lastly a JmjC domain at aa position 618-784. The *P. expansum* RpdA protein is 620 aa, slightly smaller than *A. nidulans* which is 687 aa. Both proteins contain a single histone deacetylase domain. This domain is present at aa position 35-283 in *P. expansum*, while it exists at aa position 26-424 in *A. nidulans*. The SntB protein in *P. expansum* is a large, multidomain protein of 1,637 aa, similar to the *A. nidulans* homolog which is 1,596 aa (Karahoda *et al.*, 2022). *P. expansum* SntB contains the following domains: BAH at aa position 177-295, ELM2 at aa position 480-654, SANT at aa position 667-710 and two PHD-type domains at aa positions 942-992 and 1050-1182. The levels of sequence identity and domain

structure for these proteins between *P. expansum* and *A. nidulans* suggest that they perform similar roles and function as a complex in both species.

Interestingly, unlike what is reported in *A. nidulans* (Karahoda *et al.*, 2022) and our companion piece in *A. flavus* (Karahoda *et al.* in submission), the *P. expansum* EcoA homolog (PEXP_003320) was not detected in any of our pulldown data. The *P. expansum* EcoA protein exhibits 71.91% sequence similarity to the respective *A. nidulans* homolog. *P. expansum* EcoA is 376 aa, while the EcoA protein in *A. nidulans* is 401 aa (Karahoda *et al.*, 2022). *P. expansum* EcoA contains a ZF domain at aa position 132-168 and an acetyltransferase domain at aa position 305-373. These two domains are present in *A. nidulans* EcoA at similar positions, signifying a high degree of conservation between the two species. This may imply that this protein could be performing similar roles and could be associated with the KERS complex in both species. However, lack of EcoA in any pulldown data suggests that SntB does not interact with EcoA directly in *P. expansum*, which differs from what was observed for the KERS complex in *A. nidulans* (Karahoda *et al.*, 2022). However, it is still possible that EcoA exists as a member of this complex as it could interact with KdmB and/or RpdA and thus could be associated with SntB indirectly. The interaction of EcoA with the complex could also be weak or transient, existing only during certain stages of fungal development or under specific growth conditions or stresses.

ΔkdmB strains show enhanced germination initiation rate and decreased conidia production

To investigate the role of SntB interacting members in *P. expansum*, we aimed to generate gene deletion mutants for KdmB and RpdA. We successfully obtained *ΔkdmB* strains but could not obtain deletion nor knock-down strains for *rpda* suggesting that this is an essential gene in *P.*

expansum (Fig. S1A). As we were unable to generate a *kdmB* complementation strain, we analyzed three independent deletion strains for the following experiments (Fig. 1C).

Colony diameter, germination initiation rate and conidia production for the WT and three $\Delta kdmB$ sibling strains is shown in Fig. 2. We observed no difference in colony diameter when the strains were grown in GMM media and no obvious phenotypical difference between the $\Delta kdmB$ and WT strains in this media (Fig. 1D). This contrasted with $\Delta sntB$ that showed a decrease in diameter growth compared to WT (Tannous *et al.*, 2020). The $\Delta kdmB$ strains showed a one hour increase in germination initiation when compared to the WT strain although all the strains achieved 96% germination by the final recorded timepoint (Fig. 1E), a similar observation as reported for $\Delta sntB$ (Tannous *et al.*, 2020). In addition, we observed a decreased conidia production per area at days 4 and 5 for two out of the three $\Delta kdmB$ strains when compared to the WT strain (Fig. 1F).

$\Delta kdmB$ strains show susceptibility to cell wall stressors

Considering that Choi *et al.*, 2022 showed that the *A. fumigatus* $\Delta kdmB$ strain is more sensitive to H₂O₂ than the WT strain, we were curious to evaluate if *P. expansum* $\Delta kdmB$ strains were more sensitive to chemical stressors. We assessed the response of the WT and $\Delta kdmB$ strain to Calcofluor White (CFW), Congo Red (CR), and Sodium Dodecyl Sulfate (SDS). All the $\Delta kdmB$ strains showed increased susceptibility to CFW at 75 µg/mL and 100 µg/mL (Fig. 2A) but minimal sensitivity to CR and SDS (Fig. 2BC). The enhanced sensitivity to CFW suggests a possible alteration in chitin synthesis in the mutants.

Impaired virulence of $\Delta kdmB$ strains in Golden Delicious apples

To assess if *kdmB* is involved in virulence and blue mold disease, we performed an apple inoculation experiment and measured the lesion diameter for 7 days (Fig. 3AB). The $\Delta kdmB$ strains showed reduced lesion diameter over time when compared to the WT strain suggesting that *kdmB* plays a role in virulence. These results are consistent with those described for the SntB deleted strain, where a marked reduction in decay development in infected apples was also observed (Tannous *et al.*, 2020). Since acidification of host tissues is an essential virulence strategy that necrotrophic pathogens employ, and the $\Delta kdmB$ strains showed reduced lesion diameters on apples, we measured the resulting pH for these strains in the rotten apple tissue 16 days post inoculation. We found that the $\Delta kdmB$ strains are impaired in tissue acidification when compared to the WT strain (Fig. S2A). Interestingly, when measuring the pH in YES media, the same trends hold but to a decreased extent (Fig. S2B). In addition, as some secondary metabolites including patulin act as virulence factors for some pathogens (Snini *et al.*, 2016), we quantified patulin and citrinin in infected apples. The $\Delta kdmB$ strains showed decreased amounts of patulin and citrinin in apples (Fig. 3CD). This result was also shown in the mycotoxin producing media YES for patulin but not for citrinin (Fig. S2CD).

Epigenetic mutants often result in transcriptional changes of BGC genes. We evaluated the expression of select patulin and citrinin BGC genes to see if there was any correlation with expression and mycotoxin production. Expression of the backbone polyketide synthase gene *patK* required for patulin synthesis was decreased in the $\Delta kdmB$ compared to the WT strain when grown in YES media (Fig. 4A), possibly contributing to the decreased patulin amounts quantified in apple (Fig. 3CD). In contrast the citrinin polyketide synthase gene *citS* was increased in relative gene expression in the $\Delta kdmB$ strains when compared to the WT despite no increase in this metabolite

in apple tissue. Furthermore, we assessed transcriptional activity of additional regulators of secondary metabolism and virulence determinants. In general, we observed a trend towards a slight increase in relative gene expression in the $\Delta kdmB$ strains for the virulence factors *laeA*, *creA*, *pacC*, and *sntB* and for the *gox1* and *gox2* genes which mediate gluconic acid production (Kumar *et al.*, 2017; Tannous *et al.*, 2018; Chen *et al.*, 2018; Tannous *et al.*, 2020; Hadas *et al.*, 2007). Although these trends were noticed, the reported differences are not statistically significant. *xanC*, which encodes a transcription factor that regulates citrinin production (Wang *et al.*, 2021), showed a 5-fold increase in relative expression for the $\Delta kdmB$ strains when compared to WT, although as noted above, citrinin was not upregulated in the mutant strains.

Discussion

Fungal responses to the environment are governed by vast changes in transcriptionally active regions of the genome. Other than *S. cerevisiae* and *Schizosaccharomyces pombe*, the identity and function of chromatin modifying complexes is still in its infancy in fungi. However, recent advances such as the Hst1-Rfm1-Sum1 complex controlling virulence related genes in *C. glabrata* (Vázquez-Franco *et al.*, 2022) and the discovery of the KERS complex in *A. nidulans* governing fungal development and mycotoxin synthesis are providing new frameworks for continuous progress in this field. KERS is a heteromeric complex containing a ‘reader’ (SntB), two ‘eraser’ (RpdA and KdmB) and one ‘writer’ (EcoA) protein. A previous study showed SntB to be a key regulator of virulence, mycotoxin production and fungal development in the blue mold pathogen *P. expansum* (Tannous *et al.*, 2020). Here, as found in *A. nidulans*, we report that SntB recruits the Jarid1-type Jumonji domain demethylase KdmB and the class I HDAC RpdA (Fig. 1). In *A. nidulans*, KdmB and SntB form a bridging complex with KdmB interacting with EcoA and SntB

interacting with RpdA. Similarly, in an associated paper (Karahoda *et al. in submission*), tagging of KdmB with HA and GFP epitope tag in the plant pathogenic fungus *Aspergillus flavus* led to identification of EcoA, RpdA and SntB. However, in our four purifications we could not identify EcoA as an interaction partner of KdmB-RpdA-SntB complex. We speculate that if EcoA exists as a member of the KERS complex in *P. expansum*, it could be interacting with proteins other than SntB. Thus, further pulldowns of KdmB, RpdA and EcoA could help clarify the direct and indirect interactions of EcoA within this complex. These pulldown data could also provide information regarding the interaction partners of the complex, as well as the independent interactions of each individual tagged protein. This could provide information regarding possible roles of each protein that are independent of the KERS complex. Efforts to delete *kdmB* were successful but deletions were not possible for *rpda* and *ecoA* following that reported for *A. nidulans* and *A. flavus*. Our efforts therefore were to characterize *kdmB* deletion on the pathogenicity and mycotoxin synthesis of *P. expansum*.

KdmB has been studied in diverse filamentous fungi (Gacek-Matthews *et al.*, 2016; Hou *et al.*, 2020; Janevska *et al.*, 2018; Lukito *et al.*, 2019; Bachleitner *et al.*, 2019; Choi *et al.*, 2022). In these studies, in addition to establishing that KdmB demethylates H3K4me3, the data show that *kdmB* also plays a role in virulence, secondary metabolism, stress resistance and conidiation in a variety of fungi. For example, in *A. fumigatus*, an increased percent survival and decreased fungal burden in mice was reported when inoculated with a $\Delta kdmB$ strain as compared to WT (Choi *et al.*, 2022). In addition, the secondary metabolites gliotoxin and sterigmatocystin were reduced in $\Delta kdmB$ strains of *A. fumigatus* and *A. nidulans* respectively (Choi *et al.*, 2022; Gacek-Matthews *et al.*, 2016). Similarly, in the associated manuscript, *kdmB* and *rpda* mutants cannot produce the

mycotoxin aflatoxin in *A. flavus* (Karahoda *et al. in submission*). *Botrytis cinerea* $\Delta Bcjar1$ mutants showed reduced conidiation and tolerance to SDS and H₂O₂ (Hou *et al.*, 2020). While reduced conidiation was observed in *B. cinerea*, increased conidiation was reported for *F. fujikuroi* and *A. fumigatus* (Janevska *et al.*, 2018; Choi *et al.*, 2022).

We also found some alterations in physiological traits of $\Delta kdmB$ mutants. These mutants showed many similarities to *P. expansum* $\Delta sntB$ strains such as an increased germination initiation rate and decreased conidiation (Fig. 2 and Tannous *et al.*, 2020). These findings contrasted with $\Delta kdmB$ strains in *Fusarium fujikuroi* and *Aspergillus fumigatus* which presented increased conidia production (Janevska *et al.*, 2018; Choi *et al.*, 2022). While in the latter species *kdmB* may act as a negative regulator of conidiation, data suggest that in *P. expansum* it has an opposite role although we acknowledge that the cultural conditions may have a confounding impact on any conclusions on gene loss and physiology.

Since PTMs have been described to impact secondary metabolism in filamentous fungi, we hypothesized that *kdmB* would have a role in mediating regulation of secondary metabolism. There are many instances in which $\Delta kdmB$ strains have been shown to impact secondary metabolite production. For example, in *F. fujikuroi*, the $\Delta kdmB$ strain showed reduced bikaverin, fusarubins, fusarins and GA₃ production when compared to the WT strain (Janevska *et al.*, 2018). Similarly, paxilline and lolitrem B were reduced in *Epichloë festucae* $\Delta kdmB$ strains (Lukito *et al.*, 2019). A similar trend was described for *Fusarium graminearum*, *A. nidulans* and *A. fumigatus* in which $\Delta kdmB$ strains produced less deoxynivalenol (DON), fusarin C, zearalenone, fusarielin (Bachleitner *et al.*, 2019); sterigmatocystin (ST), emericellamide C, emericellamide D, orsellinic

acid (Gacek-Matthews et al., 2016); and gliotoxin respectively (Choi *et al.*, 2022). In concordance with these studies, we observed that *P. expansum* mutants showed reduced patulin and citrinin production in $\Delta kdmB$ strains in apples although gene expression largely did not follow suit. These results suggest that the decreased patulin and citrinin production observed in $\Delta kdmB$ infections of apples are independent from patulin and citrinin gene expression.

In addition, the $\Delta kdmB$ strains showed a reduced lesion diameter in apple inoculation assays when compared to the WT strain suggesting that *kdmB* plays a role in virulence as well. This result is very similar to the one reported for the $\Delta sntB$ strain. Along with the reduced production of patulin *in vivo*, which acts as a virulence factor for some apple cultivars, an additive effect may be possible due to the observed increased susceptibility to cell wall stressors for the $\Delta kdmB$ strains as the apple environment has a plethora of stressors from the apple immune response. Also, degradative enzyme production has been linked to deletion of chromatin modifying proteins (Wu *et al.*, 2021). We speculate that possibly there could be some changes in enzyme activity in these mutants as *P. expansum* mutants decreased secretion of plant cell degradative enzymes are less virulent on apple (Jurick *et al.*, 2020). Acidification of apple tissue by *P. expansum* is also an important virulence trait (Prusky *et al.*, 2004) and possibly the decreased acidification ability of the $\Delta kdmB$ strains could also be contributing to decreased lesion size.

The present data support an important role for KdmB in virulence of *P. expansum* and, via coupling with previous data of a *sntB* mutant (Tannous *et al.*, 2020), we hypothesize that much of the defects in virulence observed in both mutants are mediated by a KERS-like complex in the blue mold

fungus. Taken together, this work suggests this complex could be a potential target for disease control, possibly broadly applicable across filamentous ascomycetes.

Acknowledgements. This work was supported by Research Grant Award No. IS-5323-20C from BARD, the United States – Israel Binational Agricultural Research and Development Fund, to NPK and ES, by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under Award Number T32AI055397 and the University of Wisconsin-Madison Advanced Opportunity Fellowship, Science, and Medicine Graduate Research Scholars program to DL. ÖB was funded by Science Foundation Ireland, grant 21 FFP-P 10146. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or BARD. DF was funded by an IRC postgraduate scholarship (GOIPG/2018/35).

References

- Andersen, B., Smedsgaard, J., Frisvad, J.C., 2004. *Penicillium expansum*: consistent production of patulin, chaetoglobosins, and other secondary metabolites in culture and their natural occurrence in fruit products. *J Agric Food Chem* 52, 2421–2428. <https://doi.org/10.1021/jf035406k>
- Bachleitner, S., Sørensen, J.L., Gacek-Matthews, A., Sulyok, M., Studt, L., Strauss, J., 2019. Evidence of a Demethylase-Independent Role for the H3K4-Specific Histone Demethylases in *Aspergillus nidulans* and *Fusarium graminearum* Secondary Metabolism. *Front Microbiol* 10, 1759. <https://doi.org/10.3389/fmicb.2019.01759>
- Bachleitner, S., Sulyok, M., Sørensen, J.L., Strauss, J., Studt, L., 2021. The H4K20 methyltransferase Kmt5 is involved in secondary metabolism and stress response in phytopathogenic *Fusarium* species. *Fungal Genet Biol* 155, 103602. <https://doi.org/10.1016/j.fgb.2021.103602>
- Baker, L.A., Ueberheide, B.M., Dewell, S., Chait, B.T., Zheng, D., Allis, C.D., 2013. The yeast Snt2 protein coordinates the transcriptional response to hydrogen peroxide-mediated oxidative stress. *Mol Cell Biol* 33, 3735–3748. <https://doi.org/10.1128/MCB.00025-13>
- Chen, Y., Li, B., Xu, X., Zhang, Z., Tian, S., 2018. The pH-responsive PacC transcription factor plays pivotal roles in virulence and patulin biosynthesis in *Penicillium expansum*. *Environ Microbiol* 20, 4063–4078. <https://doi.org/10.1111/1462-2920.14453>
- Choi, Y.-H., Lee, M.-W., Shin, K.-S., 2022. The Lysine Demethylases KdmA and KdmB Differently Regulate Asexual Development, Stress Response, and Virulence in *Aspergillus fumigatus*. *J Fungi (Basel)* 8, 590. <https://doi.org/10.3390/jof8060590>
- Denisov, Y., Freeman, S., Yarden, O., 2011. Inactivation of Snt2, a BAH/PHD-containing transcription factor, impairs pathogenicity and increases autophagosome abundance in *Fusarium oxysporum*. *Mol Plant Pathol* 12, 449–461. <https://doi.org/10.1111/j.1364-3703.2010.00683.x>
- Gacek-Matthews, A., Berger, H., Sasaki, T., Wittstein, K., Gruber, C., Lewis, Z.A., Strauss, J., 2016. KdmB, a Jumonji Histone H3 Demethylase, Regulates Genome-Wide H3K4 Trimethylation and Is Required for Normal Induction of Secondary Metabolism in *Aspergillus nidulans*. *PLoS Genet* 12, e1006222. <https://doi.org/10.1371/journal.pgen.1006222>
- Greco, C., Pfannenstiel, B.T., Liu, J.C., Keller, N.P., 2019. Depsipeptide Aspergillicins Revealed by Chromatin Reader Protein Deletion. *ACS Chem. Biol.* 14, 1121–1128. <https://doi.org/10.1021/acscchembio.9b00161>
- Hadas, Y., Goldberg, I., Pines, O., Prusky, D., 2007. Involvement of Gluconic Acid and Glucose Oxidase in the Pathogenicity of *Penicillium expansum* in Apples. *Phytopathology* 97, 384–390. <https://doi.org/10.1094/PHYTO-97-3-0384>
- Hartmann, T., Dümig, M., Jaber, B.M., Szewczyk, E., Olbermann, P., Morschhäuser, J., Krappmann, S., 2010. Validation of a Self-Excising Marker in the Human Pathogen *Aspergillus fumigatus* by

- Employing the β -Rec/six Site-Specific Recombination System. *Appl Environ Microbiol* 76, 6313–6317. <https://doi.org/10.1128/AEM.00882-10>
- Hou, J., Feng, H.-Q., Chang, H.-W., Liu, Y., Li, G.-H., Yang, S., Sun, C.-H., Zhang, M.-Z., Yuan, Y., Sun, J., Zhu-Salzman, K., Zhang, H., Qin, Q.-M., 2020. The H3K4 demethylase Jar1 orchestrates ROS production and expression of pathogenesis-related genes to facilitate *Botrytis cinerea* virulence. *New Phytol* 225, 930–947. <https://doi.org/10.1111/nph.16200>
- Ioi, J.D., Zhou, T., Tsao, R., F Marcone, M., 2017. Mitigation of Patulin in Fresh and Processed Foods and Beverages. *Toxins (Basel)* 9, E157. <https://doi.org/10.3390/toxins9050157>
- Janevska, S., Güldener, U., Sulyok, M., Tudzynski, B., Studt, L., 2018. Set1 and Kdm5 are antagonists for H3K4 methylation and regulators of the major conidiation-specific transcription factor gene ABA1 in *Fusarium fujikuroi*. *Environ Microbiol* 20, 3343–3362. <https://doi.org/10.1111/1462-2920.14339>
- Jiménez-Ortigosa, C., Aïmanianda, V., Muszkieta, L., Mouyna, I., Alsteens, D., Pire, S., Beau, R., Krappmann, S., Beauvais, A., Dufrêne, Y.F., Roncero, C., Latgé, J.-P., 2012. Chitin Synthases with a Myosin Motor-Like Domain Control the Resistance of *Aspergillus fumigatus* to Echinocandins. *Antimicrob Agents Chemother* 56, 6121–6131. <https://doi.org/10.1128/AAC.00752-12>
- Karahoda, B., Pardeshi, L., Ulas, M., Dong, Z., Shirgaonkar, N., Guo, S., Wang, F., Tan, K., Sarikaya-Bayram, Ö., Bauer, I., Dowling, P., Fleming, A.B., Pfannenstiel, B.T., Luciano-Rosario, D., Berger, H., Graessle, S., Alhussain, M.M., Strauss, J., Keller, N.P., Wong, K.H., Bayram, Ö., 2022. The KdmB-EcoA-RpdA-SntB chromatin complex binds regulatory genes and coordinates fungal development with mycotoxin synthesis. *Nucleic Acids Research* 50, 9797–9813. <https://doi.org/10.1093/nar/gkac744>
- Keller, N.P., 2019. Fungal secondary metabolism: regulation, function and drug discovery. *Nat Rev Microbiol* 17, 167–180. <https://doi.org/10.1038/s41579-018-0121-1>
- Kumar, D., Barad, S., Chen, Y., Luo, X., Tannous, J., Dubey, A., Glam Matana, N., Tian, S., Li, B., Keller, N., Prusky, D., 2017. LaeA regulation of secondary metabolism modulates virulence in *Penicillium expansum* and is mediated by sucrose. *Mol Plant Pathol* 18, 1150–1163. <https://doi.org/10.1111/mpp.12469>
- Larsen, T.O., Frisvad, J.C., Ravn, G., Skaaning, T., 1998. Mycotoxin production by *Penicillium expansum* on blackcurrant and cherry juice. *Food Addit Contam* 15, 671–675. <https://doi.org/10.1080/02652039809374696>
- Lim, F.Y., Sanchez, J.F., Wang, C.C.C., Keller, N.P., 2012. Toward awakening cryptic secondary metabolite gene clusters in filamentous fungi. *Methods Enzymol* 517, 303–324. <https://doi.org/10.1016/B978-0-12-404634-4.00015-2>

- Luciano-Rosario, D., Keller, N.P., Jurick, W.M., 2020. *Penicillium expansum*: biology, omics, and management tools for a global postharvest pathogen causing blue mould of pome fruit. *Mol Plant Pathol* 21, 1391–1404. <https://doi.org/10.1111/mpp.12990>
- Lukito, Y., Chujo, T., Hale, T.K., Mace, W., Johnson, L.J., Scott, B., 2019. Regulation of subtelomeric fungal secondary metabolite genes by H3K4me3 regulators CclA and KdmB. *Mol Microbiol* 112, 837–853. <https://doi.org/10.1111/mmi.14320>
- Olsen, M., Lindqvist, R., Bakeeva, A., Leong, S.-L.L., Sulyok, M., 2019. Distribution of mycotoxins produced by *Penicillium* spp. inoculated in apple jam and crème fraîche during chilled storage. *Int J Food Microbiol* 292, 13–20. <https://doi.org/10.1016/j.ijfoodmicro.2018.12.003>
- Pfannenstiel, B.T., Greco, C., Sukowaty, A.T., Keller, N.P., 2018. The epigenetic reader SntB regulates secondary metabolism, development and global histone modifications in *Aspergillus flavus*. *Fungal Genet Biol* 120, 9–18. <https://doi.org/10.1016/j.fgb.2018.08.004>
- Pfannenstiel, B.T., Keller, N.P., 2019. On top of biosynthetic gene clusters: How epigenetic machinery influences secondary metabolism in fungi. *Biotechnol Adv* 37, 107345. <https://doi.org/10.1016/j.biotechadv.2019.02.001>
- Pfannenstiel, B.T., Zhao, X., Wortman, J., Wiemann, P., Throckmorton, K., Spraker, J.E., Soukup, A.A., Luo, X., Lindner, D.L., Lim, F.Y., Knox, B.P., Haas, B., Fischer, G.J., Choera, T., Butchko, R.A.E., Bok, J.-W., Affeldt, K.J., Keller, N.P., Palmer, J.M., 2017. Revitalization of a Forward Genetic Screen Identifies Three New Regulators of Fungal Secondary Metabolism in the Genus *Aspergillus*. *mBio* 8, e01246-17. <https://doi.org/10.1128/mBio.01246-17>
- Prusky, D., McEvoy, J.L., Saftner, R., Conway, W.S., Jones, R., 2004. Relationship Between Host Acidification and Virulence of *Penicillium* spp. on Apple and Citrus Fruit. *Phytopathology* 94, 44–51. <https://doi.org/10.1094/PHYTO.2004.94.1.44>
- Snini, S.P., Tannous, J., Heuillard, P., Bailly, S., Lippi, Y., Zehraoui, E., Barreau, C., Oswald, I.P., Puel, O., 2016. Patulin is a cultivar-dependent aggressiveness factor favouring the colonization of apples by *Penicillium expansum*. *Mol Plant Pathol* 17, 920–930. <https://doi.org/10.1111/mpp.12338>
- Strauss, J., Cánovas, D., 2021. Editorial SI FGB “Chromatin regulation and epigenetics.” *Fungal Genet Biol* 153, 103569. <https://doi.org/10.1016/j.fgb.2021.103569>
- Tannous, J., Barda, O., Luciano-Rosario, D., Prusky, D.B., Sionov, E., Keller, N.P., 2020. New Insight Into Pathogenicity and Secondary Metabolism of the Plant Pathogen *Penicillium expansum* Through Deletion of the Epigenetic Reader SntB. *Front Microbiol* 11, 610. <https://doi.org/10.3389/fmicb.2020.00610>
- Tannous, J., Keller, N.P., Atoui, A., El Khoury, A., Lteif, R., Oswald, I.P., Puel, O., 2018a. Secondary metabolism in *Penicillium expansum*: Emphasis on recent advances in patulin research. *Crit Rev Food Sci Nutr* 58, 2082–2098. <https://doi.org/10.1080/10408398.2017.1305945>

- Tannous, J., Kumar, D., Sela, N., Sionov, E., Prusky, D., Keller, N.P., 2018b. Fungal attack and host defence pathways unveiled in near-avirulent interactions of *Penicillium expansum* creA mutants on apples. *Mol Plant Pathol* 19, 2635–2650. <https://doi.org/10.1111/mpp.12734>
- van Kan, J.A.L., 2006. Licensed to kill: the lifestyle of a necrotrophic plant pathogen. *Trends in Plant Science* 11, 247–253. <https://doi.org/10.1016/j.tplants.2006.03.005>
- Vázquez-Franco, N., Gutiérrez-Escobedo, G., Juárez-Reyes, A., Orta-Zavalza, E., Castaño, I., De Las Peñas, A., 2022. *Candida glabrata* Hst1-Rfm1-Sum1 complex evolved to control virulence-related genes. *Fungal Genet Biol* 159, 103656. <https://doi.org/10.1016/j.fgb.2021.103656>
- Wang, W., Drott, M., Greco, C., Luciano-Rosario, D., Wang, P., Keller, N.P., 2021. Transcription Factor Repurposing Offers Insights into Evolution of Biosynthetic Gene Cluster Regulation. *mBio* 12, e0139921. <https://doi.org/10.1128/mBio.01399-21>
- Wu, H., Nakazawa, T., Morimoto, R., Shivani, null, Sakamoto, M., Honda, Y., 2021. Targeted disruption of *hir1* alters the transcriptional expression pattern of putative lignocellulolytic genes in the white-rot fungus *Pleurotus ostreatus*. *Fungal Genet Biol* 147, 103507. <https://doi.org/10.1016/j.fgb.2020.103507>