

**LA-UR-24-21031**

Accepted Manuscript

# Measuring neutron star radius with second and third generation gravitational wave detector networks

Bandopadhyay, Ananya

Kacanja, Keisi

Nitz, Alexander H.

Brown, Duncan A.

Somasundaram, Rahul

Provided by the author(s) and the Los Alamos National Laboratory (2024-11-01).

**To be published in:** Classical and Quantum Gravity

**DOI to publisher's version:** 10.1088/1361-6382/ad828a

**Permalink to record:**

<https://permalink.lanl.gov/object/view?what=info:lanl-repo/lareport/LA-UR-24-21031>



Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

PAPER • OPEN ACCESS

# Measuring neutron star radius with second and third generation gravitational wave detector networks






To cite this article: Ananya Bandopadhyay *et al* 2024 *Class. Quantum Grav.* **41** 225003

View the [article online](#) for updates and enhancements.

You may also like

- [Towards quantum gravity with neural networks: solving the quantum Hamilton constraint of  \$U\(1\)\$  BF theory](#)  
Hanno Sahlmann and Waleed Sherif
- [Boosted Kerr–Newman black holes](#)  
Rafael F Aranha and Rodrigo Maier
- [Hopfion-like solutions in de Sitter spacetime](#)  
Adam Grzela, Jacek Jezierski and Tomasz Smoka

# Measuring neutron star radius with second and third generation gravitational wave detector networks

Ananya Bandopadhyay<sup>1,\*</sup> , Keisi Kacanja<sup>1</sup> ,  
Rahul Somasundaram<sup>1,2</sup> , Alexander H Nitz<sup>1</sup>   
and Duncan A Brown<sup>1</sup> 

<sup>1</sup> Department of Physics, Syracuse University, Syracuse, NY 13244, United States of America

<sup>2</sup> Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, United States of America

E-mail: [abandopa@syr.edu](mailto:abandopa@syr.edu), [kkacanja@syr.edu](mailto:kkacanja@syr.edu), [rsomasun@syr.edu](mailto:rsomasun@syr.edu), [ahnitz@syr.edu](mailto:ahnitz@syr.edu)  
and [dabrown@syr.edu](mailto:dabrown@syr.edu)

Received 15 February 2024; revised 17 September 2024

Accepted for publication 1 October 2024

Published 15 October 2024



CrossMark

## Abstract

The next generation of ground-based interferometric gravitational wave detectors will observe mergers of black holes and neutron stars throughout cosmic time. A large number of the binary neutron star merger events will be observed with extreme high fidelity, and will provide stringent constraints on the equation of state of nuclear matter. In this paper, we investigate the systematic improvement in the measurability of the equation of state with increase in detector sensitivity by combining constraints obtained on the radius of a  $1.4M_{\odot}$  neutron star from a simulated source population. Since the measurability of the equation of state depends on its stiffness, we consider a range of realistic equations of state that span the current observational constraints. We show that a single 40 km Cosmic Explorer detector can pin down the neutron star radius for a soft, medium and stiff equation of state with a precision of 10 m within a decade, whereas the current generation of ground-based detectors like the

\* Author to whom any correspondence should be addressed.



Original Content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Advanced LIGO-Virgo network would take  $\mathcal{O}(10^5)$  years to do so for a soft equation of state.

Keywords: compact binaries, neutron stars, gravitational waves

## 1. Introduction

Cosmic Explorer is a groundbreaking next-generation gravitational wave detector, representing a significant leap in our capacity to explore a plethora of compact binary merger events. The development of Cosmic Explorer and other advanced gravitational wave detectors, such as Einstein Telescope [1, 2], will allow us to understand neutron stars and their internal structure. Cosmic Explorer will detect hundreds of signals with high signal-to-noise ratios (SNR), greater than 100 [3, 4]. With the improvement in sensitivity of the detectors, it will be possible to measure the radius of a neutron star with unprecedented accuracy. Inference from simulated populations performed by combining information from a large number of observed events, allows us to obtain improved constraints on the equation of state [5–10]. Previous works have demonstrated the feasibility of combining constraints from multiple observations using statistical tools, to obtain a tighter constraint on the equation of state.

In particular, Finstad *et al* [10] combined  $\mathcal{O}(300)$  signals in an Advanced LIGO-Virgo network and compared it to the constraints obtained by stacking observations for one year of operation of a single Cosmic Explorer detector, showing the order of magnitude improvement in measurement accuracy achieved by a singular Cosmic Explorer. However, Cosmic Explorer will operate as a network which calls for the need to systematically investigate different detector configurations in order to fully gauge the improvement in sensitivity that Cosmic Explorer would accomplish. Therefore, in this paper, we extend the work of Finstad *et al* [10] to systematically study the improvement in the measurability of the equation of state with the increase in network sensitivity from the current generation of Advanced LIGO-Virgo detectors at design sensitivity to the planned Cosmic Explorer detectors, which are expected to begin operations in 2030 s. We simulate an astrophysical population of neutron stars consistent with current observational constraints on the rate of binary neutron star mergers in the Universe [11]. We perform a full Bayesian parameter estimation to obtain a forecast on the expected number of years of observation at different detector sensitivities required to measure the radius of a neutron star accurate to 10 m, by simulating different realizations of the Universe from the above population. We present our results for an Advanced LIGO-Virgo network, a post O5 network comprised of three LIGO detectors at A<sup>#</sup> <sup>3</sup> sensitivity, a 20 km and a 40 km Cosmic Explorer detector <sup>4</sup>, and two other next generation detector networks. The first of these is comprised of a 40 km Cosmic Explorer along with two detectors at A<sup>#</sup> sensitivity, and the second consists of a 40 km Cosmic Explorer, a 20 km Cosmic Explorer and one detector at A<sup>#</sup> sensitivity. For each of the networks, we combine equation of state constraints (in terms of measurement of radius of a  $1.4M_{\odot}$  neutron star) from multiple events, observed across a given observing period for each detector (network). We study the measurability of the neutron star radius for three different equation of state models drawn from [12] that are calibrated with nuclear theory at lower densities and span the current observational constraints at densities relevant for neutron stars. We find that, while the current generation of ground-based gravitational wave detector network would require thousands of years to accurately measure the

<sup>3</sup> <https://dcc.ligo.org/LIGO-T2300041/public>.

<sup>4</sup> <https://dcc.cosmicexplorer.org/CE-T2000017>.


neutron star radius even for a stiff equation of state (i.e. relatively more measurable), a single 40 km Cosmic Explorer can do so within a year. Multi-detector networks including at least one Cosmic Explorer detector could constrain the radius to 10 m accuracy within a few years for even the softest equation of state considered here. Our result clearly demonstrates the potential of next generation detector networks to constrain the equation of state of dense nuclear matter to a higher accuracy than is possible for any existing experimental facility.

The rest of this article is organized as follows. In section 2, we describe the properties of the population of neutron stars that we are examining for our study. In section 3, we describe the Bayesian inference framework that is used to obtain combined constraints on the equation of state from the measurement of neutron star radius across an observed population. In section 4, we present the results of our analysis, i.e. the projected constraints on the radius of a  $1.4M_{\odot}$  neutron star for different detector networks, and conclude the study in section 5.

## 2. Neutron star population

To study the measurability of tidal deformability from the gravitational wave signals of inspiralling binary neutron star systems, we generate a population of binary neutron star systems with component masses drawn from a Gaussian centered at  $1.4M_{\odot}$  and having a standard deviation  $\sigma_m = 0.05M_{\odot}$ . The spins are chosen to be aligned with the orbital angular momentum, and are drawn from a Gaussian distribution with zero mean and  $\sigma_{\chi} = 0.02$ . The events are chosen to be isotropically distributed across sky locations, and the inclination and orientation of the binary systems are distributed uniformly on the sphere. The luminosity distances of the systems are drawn from a uniform distribution in the range  $d_L \in [20Mpc, 20Gpc]$ . Events at larger distances, i.e. having low values of signal-to-noise ratios, contain roughly the same amount of information about the measured parameters, thus drawing events uniformly in distance reduces the amount of computational resources spent on sampling over the parameters of a large number of ‘uninformative’ events occurring at higher distances in the simulated astrophysical population. It also gives us a greater selection of ‘informative’ events (occurring at nearby distances) to choose from, in constructing different realizations of the Universe, as compared to what we would have had if our simulated population of neutron stars was the true astrophysical population. We use the median merger rate for binary neutron stars  $\mathcal{R} = 320^{+490}_{-240} \text{Gpc}^{-3}\text{yr}^{-1}$  from [11] to normalize the astrophysical population, and reweight our neutron star population to a uniform in volume distribution. Our final distribution is uniform in volume. The uniform in volume assumption is reasonable for sources which contribute substantially to the equation of state measurement, since only the close by sources would be detected with signal-to-noise ratios high enough to extract equation of state information from them.

Since the measurability of the tidal deformability depends strongly on the equation of state, we choose a set of three equations of state representing a soft, a medium and a stiff one, for this study. These are then used to assign tidal deformabilities to the individual neutron stars in each simulated signal. They are chosen from a set of 2000 equations of state also used in [10], calibrated with chiral effective field theory calculations up to nuclear saturation density,  $n_{\text{sat}}$ . Beyond this point, their construction is based on general considerations of thermodynamic stability and causality, and employing a general extension scheme in the speed of sound  $c_s^2$  for densities between  $[n_{\text{sat}}, 12n_{\text{sat}}]$  [12]. Figure 1 shows the full set of 2000 equations of state, which we use as the prior for our parameter estimation runs, with the soft, medium and stiff equations of state labelled in blue, green and orange respectively. We note here that many of the equations of state that we use as our prior are already in conflict with current astrophysical



**Figure 1.** Neutron star mass-radius curves for the soft (blue), medium (orange) and stiff (green) equations of state used in our analysis. The curves shown in grey depict the full set of 2000 equations of state used as the prior for the Bayesian analysis. The equation of state set is constructed in such a way that the distribution of radii for a  $1.4M_{\odot}$  neutron star is roughly uniform.

constraints. However, as we go on to show, the posteriors obtained on the neutron star radius from the gravitational wave analysis, particularly for the next generation detectors, are strongly likelihood-informed, and the choice of a broad range of prior does not affect the results of our analysis.

Parameter estimation runs are performed for the simulated population separately for each equation of state. The waveform model used for generating the simulated signals is the IMRPhenomD\_NRTidal approximant. This is a frequency domain approximant that is based on a phenomenological inspiral-merger-ringdown model IMRPhenomD, originally constructed to model binary black hole systems with spins aligned with their angular momentum axis [13, 14]. Tidal effects are superposed on the binary black hole waveforms and calibrated with Numerical Relativity simulations of binary neutron star systems to obtain the binary neutron star waveform approximant [15, 16]. For high SNR signals, which will be detected abundantly by the next generation gravitational wave detectors, waveform systematics can become a dominant source of error in the extraction of parameters [9]. Earlier works have shown that tidal parameters can be recovered reliably using the IMRPhenomD\_NRTidal waveform for systems with small mass ratios and spins [17]. Stiffer equations of state have been shown to introduce a bias in the recovery of non-tidal parameters, which in turn would affect the measurement of the equation of state [17, 18]. Systems with unequal mass ratios and highly spinning systems also show an increased bias in the estimation of tidal parameters [19], thus indicating the need for improved waveform modelling for binary neutron star systems for the next generation of ground based detectors.

### 3. Injection study

To quantify the improvement in measurability of the equation of state through neutron star radius and tidal deformability measurements with increase in detector sensitivity, we perform a Bayesian Inference based parameter estimation for the injected signals, which are projected

onto simulated detector noise. We consider six different detector networks for the purpose of this study. The LIGO-Virgo network consists of the LIGO Handford, LIGO Livingston [20, 21] and Virgo detectors [22] simulated at design sensitivity [23]. The post-O5 network is also comprised of three detectors, LIGO Handford, LIGO Livingston and LIGO India, at  $A^\sharp$  sensitivity [24]. The two Cosmic Explorer detectors are considered as single detector networks, a 40 km Cosmic Explorer detector off the coast of Washington, and a 20 km Cosmic Explorer detector off the coast of Texas. Additionally, we consider two more next generation detector networks. The first of these is comprised of a 40 km Cosmic Explorer detector off the coast of Washington, and two detectors at Livingston and India, operating at  $A^\sharp$  sensitivity (henceforth referred to as  $CE40+2A^\sharp$ ). The final network consists of a 40 km Cosmic Explorer, a 20 km Cosmic Explorer and LIGO India operating at  $A^\sharp$  sensitivity (henceforth referred to as  $CE40+CE20+A^\sharp$ ). Since the site locations for the Cosmic Explorer detectors are yet to be determined, the aforementioned fiducial locations are chosen as being close enough to a wide range of potential sites for the detectors to be representative from the point of view of the science goals that we address here.

To assess the performance of each detector network, we simulate a population of binary neutron star sources and apply a lower bound of 30 on the signal-to-noise ratio (SNR) to determine the events for which parameter estimation runs are performed to infer source parameters. As noted in [10], the measurement of the neutron star radius  $R_{1,4}$  is dominated by the loudest signals, which come from the local Universe. At present, the only gravitational-wave event that has been used to successfully constrain the neutron star radius is GW170817, which had a signal-to-noise ratio of  $\sim 33$ . Considering the SNR threshold of 30 chosen for this work, any single event is either comparable to, or more informative than GW170817, as regards constraining the equation of state. At a population level, the contribution of the distant low-SNR sources to the measurement of equation of state parameters becomes negligible. For signals observed with lower signal-to-noise ratio than the threshold value, the error bars associated with radius measurement are large, implying that these signals contribute negligibly to the combined equation of state inference.

Parameter estimation for the injected signals is based on the Bayes theorem. Here we briefly review the theorem in the context of its application to gravitational wave astronomy. It states that for a set of parameters  $\vec{\theta}$  that can be used to construct a model  $M$  for the gravitational waveform, and given  $\vec{d}(t)$ , the gravitational wave strain data from the detectors,

$$p(\vec{\theta}|\vec{d}(t), M) = \frac{\pi(\vec{\theta}(t)|M) \mathcal{L}(\vec{d}(t)|\vec{\theta}, M)}{\mathcal{Z}}, \quad (1)$$

where  $p(\vec{\theta}|\vec{d}(t), M)$  is the data-informed posterior distribution of the parameter  $\vec{\theta}$ ,  $\pi(\vec{\theta}(t)|M)$  is our prior knowledge of the parameter vector  $\vec{\theta}$ , conditional on the model, and  $\mathcal{L}(\vec{d}(t)|\vec{\theta}, M)$  is the likelihood of obtaining data  $\vec{d}(t)$  given the waveform model  $M$  and the parameters  $\vec{\theta}$ . Assuming that each detector produces stationary, Gaussian noise that is uncorrelated between different detectors in the network, the likelihood function is given by [25]

$$\begin{aligned} \mathcal{L}(\vec{d}|\vec{\theta}, M) &= \exp\left(-\frac{1}{2} \sum_{i=1}^N \langle n_i | n_i \rangle\right) \\ &= \exp\left(-\frac{1}{2} \sum_{i=1}^N \langle d_i - h_i(\vec{\theta}) | d_i - h_i(\vec{\theta}) \rangle\right) \end{aligned} \quad (2)$$

where

$$\langle a|b \rangle = 4\mathbb{R} \int_{f_{\min}}^{f_{\max}} \frac{\tilde{a}^*(f) \tilde{b}(f)}{S_n(f)} \quad (3)$$

is the noise-weighted inner product of two functions.

$\mathcal{Z} = \int \pi(\vec{\theta}(t)|M) \mathcal{L}(\vec{d}(t)|\vec{\theta}, M) d\vec{\theta}$  is the Bayesian evidence, which acts as a normalization constant. We use ensemble samplers, like Markov Chain Monte Carlo samplers, that sample the parameter space of interest, and trace out the posterior probability distribution. For this work, we use the parallel-tempered emcee sampler [26], invoking it through the PyCBC Inference library [27], to estimate posterior distributions of all of the parameters used to model the binary neutron star signals, as described in section 2. We use the heterodyne likelihood model [28–30] in PyCBC for likelihood estimation, which evaluates the likelihood in discrete bins close to the peak value, to speed-up sampler convergence.

In the process of recovering the signal parameters through Markov-Chain Monte Carlo sampling, the sampling parameters are chosen to be the source frame chirp mass and mass ratio, component spins along the direction of the angular momenta, sky location, distance, geocentric time of coalescence, inclination, polarization angle, and the equation of state. We sample over the equations of state in the same procedure as [10]. The equations of state are indexed in increasing order of radii for a  $1.4M_{\odot}$  neutron star, and each index corresponds to a unique mass-radius (or equivalently, mass-tidal deformability) curve in the equation of state parameter space. The lower frequency cutoff for the signals analyzed are chosen to be higher than the seismic frequency limit for the detectors. For the LIGO-Virgo network at design sensitivity, we use a lower frequency cutoff of 20 Hz, for the detectors at A<sup>‡</sup> sensitivity, we use 10 Hz, and for the Cosmic Explorer detectors, the lower frequency cutoff is chosen to be 7 Hz. The high frequency cutoff for all signals is 2048 Hz.


Once the sampler converges, we obtain the full set of posterior samples for the parameters of each of the analyzed events. This represents our knowledge of the source parameters for the gravitational wave event, conditioned on the data. We then combine multiple events hierarchically, to obtain a hyperposterior for the neutron star radius, that represents our integrated knowledge of the equation of state from the observed population of neutron stars. To do this, we first convert the equation of state samples to radius samples for a  $1.4M_{\odot}$  neutron star (denoted as  $R_{1.4}$ ). This is done by using the mass-radius curve corresponding to each equation of state sample to determine the radius value for that sample. We then use a Gaussian Kernel Density Estimation [31] to obtain the posterior distribution on the radius for each event. The combined posterior for  $N$  events  $\{s_1, s_2, \dots, s_N\}$  is then obtained as

$$p(R_{1.4}|s_1, \dots, s_N) = p(R_{1.4})^{(1-N)} \prod_{i=1}^N p(R_{1.4}|s_i) \quad (4)$$

where  $p(R_{1.4})$  is the radius prior, uniform across all events.

## 4. Results

We simulate a realistic observing scenario for binary neutron star mergers by creating a population of neutron stars distributed uniformly in distance (details in section 2) and re-weighting our events to a population distributed uniformly in volume. We then make random draws from this population to construct different realizations of the Universe. We also randomize



**Figure 2.** Combined constraints on the measurement of  $R_{1.4}$  for the six different detector networks considered in this study. The blue, orange and green colors represent the constraints for the soft, medium and stiff equations of state respectively. The shaded regions in the figures represent the 90% credible intervals for the combined radius posteriors obtained using equation (4). The dotted lines depict the fractional error bars on radius measurement after a given number of years of observation. The dashed lines are the injected radius values.

the sequence of detection of the events and combine the posteriors for  $R_{1.4}$  using equation (4) to obtain the projected accuracy in the measurement of neutron star radius for a given observation period for each detector network. In figure 2, we show the projected constraints on the radius of a  $1.4M_{\odot}$  neutron star for a given random drawn population for the six detector networks described in section 3. For all of the detector networks, we find that the combined radius posterior converges towards the injected value. For the soft equation of state (depicted in blue


in the figure), the shifting of the confidence intervals above the horizontal line representing the injected value suggests a possible systematic bias that causes the radius to be over-estimated for all of the detector networks. This can be attributed, at least partially, to the discrete sampling in the equation of state index, which causes a mismatch between the equation of state prior and the population distribution. We return to a discussion of this bias in section 5. We find that a next generation detector configuration with one 40 km Cosmic Explorer, one 20 km Cosmic Explorer and one A<sup>#</sup> detector converges to the injected radius the fastest. Even one Cosmic Explorer detector converges faster to the radius than a three-detector Advanced LIGO-Virgo configuration.

As pointed out in [10], the loudest events have the most significant contribution towards lowering the fractional error bars on radius measurement. Thus, the exact time required to measure the radius of a neutron star to a certain accuracy will be affected by the order of occurrence of the events. To mitigate the effect of the ordering of events used in obtaining the projected constraints in figure 2, we draw different populations of the Universe for each detector network 500 times and plot the distribution of the number of years of operation it would take to constrain the radius of a  $1.4M_{\odot}$  neutron star with a precision of 10 m. These results are shown in figure 3 and a summary is also shown in table 1. From these, we can see that even for the stiffest equation of state, the current generation of ground based detectors like the Advanced LIGO-Virgo network would take several thousand years to measure the radius of a neutron star with a precision of 10 m, and this improves only slightly with the A<sup>#</sup> upgrade. For a very stiff equation of state which gives a radius of  $\sim 13$  km for a  $1.4M_{\odot}$  neutron star, the A<sup>#</sup> network may be able to constrain the radius with a precision of 10 m within 20 years, but this worsens significantly if the true equation of state is moderately soft. On the other hand, a single 40 km Cosmic Explorer detector will be able to constrain radius with a precision of 10 m within 12 years for the softest equation of state considered here, and within a year for the stiffest. In order to measure the equation of state for a neutron star in a viable time-frame, building Cosmic Explorer is absolutely essential.

## 5. Conclusion

In this paper, we have performed a comparative study of the potential of different second and third generation gravitational wave detector networks to precisely constrain the nuclear equation of state. We perform a full Bayesian inference on a simulated population of neutron stars to be observed at different network sensitivities. The equation of state models used in the analysis span the plausible range of stiffness allowed by current observational constraints for densities relevant for neutron stars, and are calibrated with chiral effective field theory at the low density regime. We combine constraints from individual events, and show that this procedure can be used to obtain a stringent constraint on the radius of a  $1.4M_{\odot}$  neutron star.

Our results demonstrate the systematic improvement in the measurability of the nuclear equation of state with improvement in sensitivity of gravitational detectors, from the current generation of Advanced LIGO-Virgo detectors, to the A<sup>#</sup> upgrade, and subsequently the next generation of detectors like Cosmic Explorer. The constraints obtained on the equation of state follow the expected hierarchy of measurability, with the stiffest equations of state giving the most stringent constraints, and the softer models leading to weaker constraints. Based on the current estimate for the merger rate from [11], we find that the Advanced LIGO-Virgo detectors operating at design sensitivity would take  $2 \times 10^{5+4 \times 10^4}_{-4 \times 10^4}$ ,  $5 \times 10^{4+1 \times 10^4}_{-1 \times 10^4}$  and  $7000^{+900}_{-900}$  years respectively to constrain the radius of a  $1.4M_{\odot}$  with a precision of 10 m, for the soft, medium and stiff equations of state respectively. With the detectors operating at A<sup>#</sup> sensitivity, the same



**Figure 3.** Distribution of the number of years of observation at different network sensitivities required to constrain the radius of a  $1.4M_{\odot}$  neutron star down with a precision of 10 m. The blue, orange and green histograms depict the expected distributions for the soft, medium and stiff equations of state respectively. For each equation of state (labeled by a different color) the leftmost and rightmost vertical dashed lines encompass the 90% confidence interval, and the central dashed line represents the median value of the distribution.

accuracy can be reached in  $300^{+50}_{-50}$ ,  $100^{+40}_{-40}$  and  $20^{+4}_{-4}$  years. However, with the next generation of ground based detectors like the Cosmic Explorer, the radius of a  $1.4M_{\odot}$  neutron star can be measured with a precision of 10 m well within a human lifetime, with a single 20 km detector

**Table 1.** A summary of the result of drawing 500 random realizations of the Universe from our simulated population for each detector network and each equation of state. For each realization of the Universe, the events are shuffled to randomize order of occurrence, and then used to determine the average number of years required to measure the radius of a  $1.4M_{\odot}$  neutron star with a precision of 10 m.

| Detector network         | Soft EoS   | Medium EoS                                       | Stiff EoS             |
|--------------------------|--|--|-----------------------|
| LIGO-Virgo               | $2 \times 10^{5+4 \times 10^4}_{-4 \times 10^4}$ | $5 \times 10^{4+1 \times 10^4}_{-1 \times 10^4}$ | $7000^{+900}_{-900}$  |
| 3 A <sup>#</sup>         | $300^{+50}_{-50}$                                | $100^{+40}_{-40}$                                | $20^{+4}_{-4}$        |
| CE20                     | $21^{+10}_{-10}$                                 | $15^{+7}_{-7}$                                   | $3^{+1}_{-1}$         |
| CE40                     | $12^{+3}_{-3}$                                   | $8^{+1}_{-1}$                                    | $1^{+0.6}_{-0.6}$     |
| CE40+2 A <sup>#</sup>    | $9^{+4}_{-4}$                                    | $6^{+2}_{-2}$                                    | $0.4^{+0.2}_{-0.2}$   |
| CE40+CE20+A <sup>#</sup> | $5^{+2}_{-2}$                                    | $3^{+1}_{-1}$                                    | $0.2^{+0.07}_{-0.07}$ |

doing it in  $21^{+10}_{-10}$ ,  $15^{+7}_{-7}$  and  $3^{+1}_{-1}$  years, and a 40 km detector requiring only  $12^{+3}_{-3}$ ,  $8^{+1}_{-1}$  and  $1^{+0.6}_{-0.6}$  years respectively for the soft, medium and stiff equations of state. The multi-detector network comprised of a 40 km Cosmic Explorer and 2 detectors at A<sup>#</sup> does it in  $9^{+4}_{-4}$ ,  $6^{+2}_{-2}$  and  $0.4^{+0.2}_{-0.2}$  years, whereas the network comprised of a 40 km Cosmic Explorer, a 20 km Cosmic Explorer and an A<sup>#</sup> takes  $5^{+2}_{-2}$ ,  $3^{+1}_{-1}$  and  $0.2^{+0.07}_{-0.07}$  years respectively to measure the radius with a precision of 10 m for the soft, medium and stiff equations of state. To assess the impact that a measurement accuracy of 10 m on  $R_{1.4}$  has on constraining the nuclear equation of state, one can overlay the confidence interval on the mass-radius curves constituting the prior (figure 1). This was done, for e.g. in [12], where using the same set of 2000 equations of state as the prior in their analysis for GW170817, the authors ruled out a significant fraction of the mass-radius curves at 90% confidence with a  $\sim 2$  km accuracy on the measurement of neutron-star radius for a single mass (as shown in figure 2 of [12]). With a 10 m precision, a much larger fraction of the mass-radius curves can be ruled out, thus yielding substantially stronger constraints on the equation of state than obtainable from other astrophysical observations, such as NICER or electromagnetic counterparts of binary neutron star mergers. Also, while the 10 m precision used here to constrain  $R_{1.4}$  is an arbitrary choice, and the exact limits on the required precision for neutron star radius will be set by nuclear theorists as their models continue to be upgraded with time, 10 m is a sufficiently high precision, that will allow us to measure subtle features of the equation of state, for e.g. a weak phase transition that is not manifested as a sharp discontinuity in the mass-radius curve. Thus, the next generation of ground based detectors will be instrumental in constraining the nuclear equation of state to a level of accuracy that can not be reached with any existing facilities.

In this paper, we have focussed our results on the measurement of the radius of a  $1.4M_{\odot}$  neutron star. In order to truly constrain the nuclear equation of state, one has to constrain the mass-radius curve (or equivalently, the mass-tidal-deformability curve), thus necessitating the measurement of radii of neutron stars along the full mass spectrum. However, with our approach of sampling directly over the equation of state, rather than in the tidal parameters such as  $\Lambda_{1,2}$ , we are not restricted to  $R_{1.4}$ , but can instead map the equation of state posteriors to the neutron star radius for any other mass. As the equation of state priors continue to be updated with constraints from nuclear theory and other astrophysical and terrestrial experiments, this approach will also allow us to place constraints on the equation of state at higher and lower densities, especially as we observe events involving neutron stars along the full mass spectrum. The posteriors obtained from the analysis of gravitational-wave signals detected from binary

neutron stars can also be combined with other astrophysical observations, such as pulsar timing and pulse profile modeling [32–34], to improve the combined constraints on the equation of state, as well as mitigate some of the systematic uncertainties involved in the models used to constrain the equation of state [35]. Finally, the advantage of representing the equation of state constraints as single parameter such as  $R_{1.4}$  is that it allows us to directly compare the posteriors on  $R_{1.4}$  to those on nuclear parameters obtained from terrestrial experiments, such as the neutron skin thickness of  $^{208}\text{Pb}$  [36, 37], to set tighter constraints on the equation of state.

Some of the source bias in the process of obtaining combined constraints by this method are pointed out in [10]. These biases include the imperfect knowledge of the mass distribution of neutron stars and the potential absence of an electromagnetic counterpart, posing challenges in distinguishing a binary neutron star system from a neutron star black hole system. In this study, we circumvent the bias caused by the imperfect knowledge of mass distribution by choosing the mass prior used in our analysis to be the same as the one used to simulate the neutron star population. However, in practical applications, as more and more signals from binary neutron star mergers are detected, the mass distribution used in the inference should also be simultaneously updated to mitigate this bias. Further, as seen in figure 2, for some of the equations of state, there is a systematic bias that causes the combined posteriors to deviate from the injected value. This can be interpreted as a finite-sampling effect, which we anticipate would be alleviated by sampling the equation of state in a continuous manner, instead of using a finite set of discrete equation of state samples as the prior. Solving the Tolman–Oppenheimer–Volkoff (TOV) equations that determine the neutron-star structure is a computational bottleneck that limits us from implementing a continuous sampling of the equation of state. Progress has been made in this direction through the use of machine learning algorithms to emulate solutions to the TOV equations, yielding results that are in good agreement with those obtained using the full TOV solver [38]. We leave it to a future work to investigate the improvement brought about through a continuous sampling of the equation of state. Another source of bias in obtaining constraints on the equation of state, particularly in terms of the neutron star radius, is that we do not measure radius directly from gravitational wave signals, but use the measurement of tidal deformability to obtain an estimate of the radius. Since the love number  $k_2$  (as defined in e.g. [39–41]), relating tidal deformability to radius, is also a function of radius, even an extremely precise measurement of the tidal deformability leads to a multimodal distribution for measured radius, which can ultimately tend to bias the median value of the measured radius away from its true value. Future works can extend this work to constrain the radius not only for a  $1.4M_{\odot}$  neutron star but for the whole mass range allowed by plausible equations of state. In conclusion, our work highlights the imperative need to build Cosmic Explorer in order to be able to accurately constrain the radius of a neutron star within a human lifetime.

### Data availability statement

The data that support the findings of this study are openly available at the following URL/DOI: <https://github.com/sugwg/bns-eos-nggw>.

### Acknowledgment

The authors acknowledge support from National Science Foundation Awards PHY-2011655 (AB and DAB), PHY-2207264 (DAB), and PHY-2309240 (AHN, KK and DAB). RS acknowledges support from the Nuclear Physics from Multi-Messenger Mergers (NP3M) Focused

Research Hub which is funded by the National Science Foundation under Grant Number 21-16686, and by the Laboratory Directed Research and Development program of Los Alamos National Laboratory under Project number 20220541ECR. This research was supported through computational resources provided by Syracuse University.

## ORCID iDs

Ananya Bandopadhyay  <https://orcid.org/0000-0002-5116-844X>

Keisi Kacanja  <https://orcid.org/0009-0004-9167-7769>

Rahul Somasundaram  <https://orcid.org/0000-0003-0427-3893>

Alexander H Nitz  <https://orcid.org/0000-0002-1850-4587>

Duncan A Brown  <https://orcid.org/0000-0002-9180-5765>

## References

- [1] Hild S, Chelkowski S, Freise A, Franc J, Morgado N, Flaminio R and DeSalvo R 2010 A Xylophone configuration for a third generation gravitational wave detector *Class. Quantum Grav.* **27** 015003
- [2] Sathyaprakash B *et al* 2012 Scientific objectives of einstein telescope *Class. Quantum Grav.* **29** 124013 Sathyaprakash B *et al* 2013 *Class. Quantum Grav.* **30** 079501
- [3] Abbott B P *et al* 2017 Exploring the sensitivity of next generation gravitational wave detectors *Class. Quantum Grav.* **34** 044001
- [4] Evans M *et al* 2021A Horizon study for cosmic explorer: science, observatories, and community (arXiv:2109.09882)
- [5] Lackey B D and Wade L 2015 Reconstructing the neutron-star equation of state with gravitational-wave detectors from a realistic population of inspiralling binary neutron stars *Phys. Rev. D* **91** 043002
- [6] Agathos M, Meidam J, Del Pozzo W, Li T G F, Tompitak M, Veitch J, Vitale S and Van Den Broeck C 2015 Constraining the neutron star equation of state with gravitational wave signals from coalescing binary neutron stars *Phys. Rev. D* **92** 023012
- [7] Hernandez Vivanco F, Smith R, Thrane E, Lasky P D, Talbot C and Raymond V 2019 Measuring the neutron star equation of state with gravitational waves: the first forty binary neutron star merger observations *Phys. Rev. D* **100** 103009
- [8] Pacilio C, Maselli A, Fasano M and Pani P 2022 Ranking love numbers for the neutron star equation of state: the need for third-generation detectors *Phys. Rev. Lett.* **128** 101101
- [9] Chatziioannou K 2022 Uncertainty limits on neutron star radius measurements with gravitational waves *Phys. Rev. D* **105** 084021
- [10] Finstad D, White L V and Brown D A 2023 Prospects for a precise equation of state measurement from advanced LIGO and cosmic explorer *Astrophys. J.* **955** 45
- [11] Abbott R *et al* 2021 Population properties of compact objects from the second ligo-virgo gravitational-wave transient catalog *Astrophys. J. Lett.* **913** L7
- [12] Capano C D, Tews I, Brown S M, Margalit B, De S, Kumar S, Brown D A, Krishnan B and Reddy S 2020 Stringent constraints on neutron-star radii from multimessenger observations and nuclear theory *Nat. Astron.* **4** 625–32
- [13] Husa S, Khan S, Hannam M, Pürrer M, Ohme F, Jiménez Forteza X and Bohé A 2016 Frequency-domain gravitational waves from nonprecessing black-hole binaries. I. New numerical waveforms and anatomy of the signal *Phys. Rev. D* **93** 044006
- [14] Khan S, Husa S, Hannam M, Ohme F, Pürrer M, Jiménez Forteza X and Bohé A 2016 Frequency-domain gravitational waves from nonprecessing black-hole binaries. II. A phenomenological model for the advanced detector era *Phys. Rev. D* **93** 044007
- [15] Barkett K *et al* 2016 Gravitational waveforms for neutron star binaries from binary black hole simulations *Phys. Rev. D* **93** 044064
- [16] Dietrich T, Bernuzzi S and Tichy W 2017 Closed-form tidal approximants for binary neutron star gravitational waveforms constructed from high-resolution numerical relativity simulations *Phys. Rev. D* **96** 121501

- [17] Dietrich T, Hinderer T and Samajdar A 2021 Interpreting binary neutron star mergers: describing the binary neutron star dynamics, modelling gravitational waveforms and analyzing detections *Gen. Relativ. Grav.* **53** 27
- [18] Dudi R, Pannarale F, Dietrich T, Hannam M, Bernuzzi S, Ohme F and Brüggmann B 2018 Relevance of tidal effects and post-merger dynamics for binary neutron star parameter estimation *Phys. Rev. D* **98** 084061
- [19] Samajdar A and Dietrich T 2018 Waveform systematics for binary neutron star gravitational wave signals: effects of the point-particle baseline and tidal descriptions *Phys. Rev. D* **98** 124030
- [20] Abbott B P *et al* 2016 GW150914: The Advanced LIGO Detectors in the Era of First Discoveries *Phys. Rev. Lett.* **116** 131103
- [21] Buikema A *et al* 2020 Sensitivity and performance of the Advanced LIGO detectors in the third observing run *Phys. Rev. D* **102** 062003
- [22] Acernese F *et al* 2015 Advanced Virgo: a second-generation interferometric gravitational wave detector *Class. Quantum Grav.* **32** 024001
- [23] Abbott B P *et al* 2018 Prospects for observing and localizing gravitational-wave transients with Advanced LIGO, advanced virgo and KAGRA *Living Rev. Rel.* **21** 3
- [24] Fritschel P *et al* 2022 Report from the Lsc post-o5 study group, LIGO, *Technical Report* T2200287 (<https://dcc.ligo.org/LIGO-T2200287/public>)
- [25] Wainstein L A and Zubakov V D 1970 Extraction of signals from noise
- [26] Foreman-Mackey D, Hogg D W, Lang D and Goodman J 2013 emcee: the MCMC Hammer *Publ. Astron. Soc. Pac.* **125** 306–12
- [27] Biwer C M, Capano C D, De S, Cabero M, Brown D A, Nitz A H and Raymond V 2019 PyCBC inference: a python-based parameter estimation toolkit for compact binary coalescence signals *Publ. Astron. Soc. Pac.* **131** 024503
- [28] Cornish N J 2021 Heterodyned likelihood for rapid gravitational wave parameter inference *Phys. Rev. D* **104** 104054
- [29] Zackay B, Dai L and Venumadhav T 2018 Relative binning and fast likelihood evaluation for gravitational wave parameter estimation vol 6 (arXiv:1806.08792 [astro-ph.IM])
- [30] Finstad D and Brown D A 2020 Fast parameter estimation of binary mergers for multimessenger follow-up *Astrophys. J. Lett.* **905** L9
- [31] Rosenblatt M 1956 Remarks on some nonparametric estimates of a density function *Ann. Math. Stat.* **27** 832–7
- [32] Reardon D J *et al* 2016 Timing analysis for 20 millisecond pulsars in the Parkes Pulsar Timing Array *Mon. Not. R. Astron. Soc.* **455** 1751–69
- [33] Riley T E *et al* 2019 A NICER view of PSR J0030+0451: millisecond pulsar parameter estimation *Astrophys. J. Lett.* **887** L21
- [34] Cromartie H T *et al* 2020 Relativistic Shapiro delay measurements of an extremely massive millisecond pulsar *Nat. Astron.* **4** 72–76
- [35] Landry P, Essick R and Chatziioannou K 2020 Nonparametric constraints on neutron star matter with existing and upcoming gravitational wave and pulsar observations *Phys. Rev. D* **101** 123007
- [36] Reed B T, Fattoyev F J, Horowitz C J and Piekarewicz J 2021 Implications of PREX-2 on the equation of state of neutron-rich matter *Phys. Rev. Lett.* **126** 172503
- [37] Essick R, Landry P, Schwenk A and Tews I 2021 Detailed examination of astrophysical constraints on the symmetry energy and the neutron skin of  $^{208}\text{Pb}$  with minimal modeling assumptions *Phys. Rev. C* **104** 065804
- [38] Reed B T, Somasundaram R, De S, Armstrong C L, Giuliani P, Capano C, Brown D A, and Tews I 2024 Towards accelerated nuclear-physics parameter estimation from binary neutron star mergers: emulators for the Tolman-Oppenheimer-Volkoff equations (arXiv:2405.20558)
- [39] Damour T and Nagar A 2009 Relativistic tidal properties of neutron stars *Phys. Rev. D* **80** 084035
- [40] Flanagan E E and Hinderer T 2008 Constraining neutron star tidal love numbers with gravitational wave detectors *Phys. Rev. D* **77** 021502
- [41] Hinderer T, Lackey B D, Lang R N and Read J S 2010 Tidal deformability of neutron stars with realistic equations of state and their gravitational wave signatures in binary inspiral *Phys. Rev. D* **81** 123016