

MASTER

NYO-3399-193

SU-1206-193

May 1969

APPLICATION OF A RELATIVISTIC RESONANCE FORMULA TO THE

$$e^+e^- \rightarrow \pi^+\pi^- \text{ EXPERIMENT*}$$

Michael Parkinson

Physics Department
Syracuse University
Syracuse, New York

ABSTRACT

The present data for the reaction $e^+e^- \rightarrow \pi^+\pi^-$ is analyzed with the aid of a general effective range resonance formula for the ρ meson. It is concluded that: (a) $m_\rho = 769 \pm 5$ MeV and $\Gamma_\rho = 109 \pm 10$ MeV provide a reasonable explanation of the present experimental points: (b) The full width at half maximum regardless of the peak height should give an excellent estimate of the ρ width. (c) It is hard to escape $|F_\pi(m_\rho^2)|^2 = \frac{m_\rho^2}{\Gamma_\rho^2}$. (d) The effects of final state interactions on resonance shape ought to be investigated.

* Work supported by the United States Atomic Energy Commission



Department of Physics
SYRACUSE UNIVERSITY
Syracuse, New York 13210

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

NYO-3399-193
SU-1206-193
May 1969

APPLICATION OF A RELATIVISTIC RESONANCE FORMULA TO THE

$$e^+e^- \rightarrow \pi^+\pi^- \text{ EXPERIMENT*}$$

Michael Parkinson
Physics Department
Syracuse University
Syracuse, New York

ABSTRACT

The present data for the reaction $e^+e^- \rightarrow \pi^+\pi^-$ is analyzed with the aid of a general effective range resonance formula for the ρ meson. It is concluded that: (a) $m_\rho = 769 \pm 5$ MeV and $\Gamma_\rho = 109 \pm 10$ MeV provide a reasonable explanation of the present experimental points: (b) The full width at half maximum regardless of the peak height should give an excellent estimate of the ρ width. (c) It is hard to escape $|\Gamma_\pi(m_\rho^2)|^2 = \frac{m_\rho^2}{\Gamma_\rho^2}$. (d) The effects of final state interactions on resonance shape ought to be investigated.

* Work supported by the United States Atomic Energy Commission

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

14

We present here an analysis of the pion form factor as measured in the $e^+e^- \rightarrow \pi^+\pi^-$ experiments of Orsay and Novosibirsk,⁽¹⁾ using a formula derived from a natural relativistic generalization of the Breit-Wigner formula for the ρ meson.⁽²⁾ This relativistic resonance formula, when tested against the $\pi N \rightarrow \pi\pi N$ data, appears to be quite satisfactory⁽³⁾ although we will have more to say on this subject later. For our purposes here, we need only the following expressions:

$$F_\pi(s) = D(0)/D(s) \quad , \quad s = (p_+ + p_-)^2 \quad (1)$$

$$D(s) = 1 - \frac{s-s_0}{\pi} \oint \frac{\mu(s') ds'}{(s'-s_0)^2 (s'-s)} - i \frac{\mu(s)}{s-s_0} \quad (2)$$

$$\mu(s') = \sum_{i=1}^N c_i^2 \rho_i(s') \theta(s'-t_i) \quad , \quad \theta(s) = \begin{cases} 1 & s > 0 \\ 0 & s < 0 \end{cases} \quad (3)$$

c_i , ρ_i , and t_i are given in Table I; i is the channel index.

A short discussion of the derivation of (1) - (3) will be given here; for a more complete discussion, see ref. 2.

Briefly then, for the l^- partial wave, we write the T matrix, related to S matrix by $S = 1 - 2ipT$, as $T^{-1} = K'^{-1} - i\rho$ where we have introduced the K' matrix; ρ is the matrix of relativistic phase space factors:

$$\rho_{ij} = \rho_i(s) \delta_{ij} \theta(s-t_i)$$

As defined, K' is not analytic, because ρ contains θ functions. We assume T is analytic because all currently acceptable theories have this property. Suppose we now define $R_{ij}(s) = r_i(s) \delta_{ij}$ where $r_i(s)$

$$r_i(s) = \frac{1}{\pi} \int_{t_i}^{\infty} \frac{\rho_i(s') ds'}{s' - s - i\epsilon} + \text{possible subtractions.}$$

Then $T^{-1} = K^{-1} - R$ defines an analytic K matrix which should be easier to approximate. At a resonance T becomes pure imaginary, which means $K^{-1} \rightarrow (K_0^{-1})(s_R - s)$, with $K_0 > 0$ so that we have a pole on the second sheet. K_0 is the matrix of coupling constants to the resonance: $(K_0)_{ij} = k_i k_j$. Thus $T = K_0(s_R - s - RK_0)^{-1} = K_0 / (s_R - s - \text{Tr}(RK_0))$ or $T = K_0 / \Delta(s)$ with $\Delta(s) = s_R - s - \text{Tr}(RK_0)$. In order to make $\text{Re} [\text{Tr}(RK_0)]$ vanish as $(s - s_R)^2$ when $s \rightarrow s_R$, we subtract it twice at the mass of the resonance. This formula, like all effective range formulas, usually has a pole on the real axis for $s < 0$; i.e., $\Delta(s_0) = 0$ for $s_0 < 0$. This pole must be removed to get the form factor. It can be shown that

$$\Delta(s) = (s - s_0) \left(1 - \frac{s - s_0}{\pi} \int_{-\infty}^{+\infty} \frac{\sum_{i=1}^N C_i \rho_i(s') \theta(s' - t_i)}{(s' - s_0)^2 (s' - s - i\epsilon)} ds' \right)$$

where C_i are proportional to the k_i and we display the zero at $s = s_0$ explicitly rather than that at $s = s_R$. Thus, $D(s) = \Delta(s) / (s - s_0)$ which leads to equations (1) - (3) if we elect to normalize $F(0) = 1$. Observe that D corresponds to the D function in the ND^{-1} method with the left hand cut replaced by a pole at $s = s_0$. Like the usual simple pole approximation at $s = s_R$, $D(s) \approx s_R - s - i\text{Im} \Gamma_R$ for $s \approx s_R$ but its behavior away from that point is not the same. It seems plausible that Eqn. (2) ought to be better than the usual simple pole approximation because it has more of the properties of the correct D function than the latter. For example, one significant property Eqn. (2) possesses is that the integral in D

gives the analytic continuation of the decay widths of the resonance into the various channels. That such an analytic continuation is needed in general was observed sometime ago,⁽⁴⁾ in the calculation of the mass shift to a nuclear state due to the opening and closing of a channel as the position of the channel threshold moves. For these reasons, we will use Eqn. (1) with the three channels that seem most significant: $\pi\pi$, $\pi\omega$, $\bar{N}N(+)$ when we attempt to fit the data. These three channels were quite adequate in accounting for the ρ meson in $\pi N \rightarrow \pi\pi N$ where it was found that: $\Gamma_\rho = 152 \pm 18$ MeV and $m_\rho = 768 \pm 2$ MeV.⁽³⁾ (4)

Let us now consider the $\pi\pi$ problem in the l^- partial wave. In Table I, we see estimated values for the C_i . With values such as these, $\text{Re } D(s)$ turns out to be approximately linear. We define m_ρ^2 and Γ_ρ as follows:

$$\text{Re } D(m_\rho^2) = 0 \quad (5)$$

$$m_\rho \Gamma_\rho = \frac{C_1^2 \rho_1(m^2)}{m^2 + |s_0|} \frac{1}{\text{Re } D'(m_\rho^2)}, \quad D'(s) = \frac{\partial D}{\partial s} \quad (6)$$

Plotted against $E = \sqrt{s}$, $D(0)/D(s)$ peaks for $E \approx m_\rho$ and has a full width at half maximum approximately equal to Γ_ρ . These facts lead us to a quantitative test that can be applied to the $e^+e^- \rightarrow \pi^+\pi^-$ data in order to see if Eqns. (1) - (3) can reasonably explain them. For we see that

$$|F_\pi(m_\rho^2)|^2 = \left(\frac{D(0)}{m_\rho^2 \text{Re } D'(m_\rho^2)} \right)^2 \frac{m_\rho^2}{\Gamma_\rho^2} \quad (7)$$

When $\text{Re } D(s)$ is approximately linear, because $D(s_0) = 1$ and $\text{Re } D(m_\rho^2) = 0$, we find that

$$D(0)/D(s_0) \approx \frac{m_\rho^2}{m_\rho^2 + |s_0|}$$

and

$$\operatorname{Re} D'(m_\rho^2) = \frac{1}{m_\rho^2 + |s_0|}$$

We do expect $D(s)$ to be roughly linear, and thus we expect the following relation to be true:

$$|F_\pi(m_\rho^2)|^2 \approx \frac{m_\rho^2}{\Gamma_\rho^2} \quad (8)$$

Comparing Eqn. (8) against experiment, we see that it is consistent at this point, but only because of the large error bars. This is particularly true of the Russian points. It will be interesting to see how Eqn. (8) fares in the future as the experimental results are sharpened.

For example, it may turn out the D is sufficiently non-linear so that Eqn. (8) will not hold. For example, if we set $g_{\rho NN} = g_{\rho \pi \omega} = 0$ or if we take $g_{\rho \pi \omega} = 1.5$ we get sufficient non-linearity to affect Eqn. (7), quite substantially in the latter case. Neither of these examples are particularly realistic in the present scheme of things, but nevertheless even in these cases, we find that a very good estimate of Γ_ρ , the ρ meson decay width or inverse lifetime, is the full width at half maximum in the form factor, regardless of the height of the peak. This suggests the desirability of obtaining very accurate data points around the half maximum on each side of the peak as a way to measure the ρ width precisely and rather independently of the exact dynamics reflected, for example, in the thresholds and coupling constants which enter into Eqn. (2).

Application of Eqns. (1) - (3) to Experimental Data

In Figure (1), one sees the results of fitting Eqn. (1) to the present experimental data (see App. 1 for details). It is quite reassuring to see that the $\pi \rho \omega$ and ρNN coupling constants are consistent with the values we expect from other sources of information.⁽⁵⁾ Although the χ^2 obtained is not so

good, (5% confidence level) observe that the Russian point at 855 MeV and the Orsay point at 886 MeV are in disagreement, and alone contribute a χ^2 of 10. Furthermore, Orsay Pt. at 940 MeV contributes another 6. These large contributions indicate either an experimental or theoretical problem to be resolved in the future. From Eqn. (4) and Eqn. (5), we obtain:

$$\begin{aligned} m_{\rho} &= 5.49 \pm 0.02 m_{\pi} = 769 \pm 3 \text{ MeV} \\ \Gamma_{\rho} &= 0.778 \pm 0.04 m_{\pi} = 109 \pm 6 \text{ MeV} \end{aligned} \quad (9)$$

Note that the curve in Fig. 1 satisfies Eqn. (7) very closely, despite the fact that $D(s)$ deviates from linearity, as reflected in the values $D(0) = 0.136$ and $\text{Re}D'(m_{\rho}^2) = 4.47 \times 10^{-3} m_{\pi}^{-2}$. Nevertheless, $D(0)/m_{\rho}^2 \text{Re}D'(m_{\rho}^2) = 1.01$ which shows the insensitivity of Eqn. (7) to possible nonlinearity. (The linear predictions for $D(0)$ and $D'(m_{\rho}^2)$ are $D(0) = 0.0732$ and $\text{Re}D'(m_{\rho}^2) = 2.44 \times 10^{-3} m_{\pi}^{-2}$).

The above values indicate, of course, a problem, when compared with those of Eqn. (4). Inasmuch as the sample of ρ 's in the $e^+e^- \rightarrow \pi^+\pi^-$ experiment are "cleaner" than those in $\pi N \rightarrow \pi\pi N$, one is inclined to take Eqn. (9) more seriously.

This kind of difficulty is not new. Although the success of the hypothesis that the ρ meson is universally coupled to the isospin current has been great, we have, in fact, a list of annoying discrepancies between theory and experiment, some of which are as follows:

- (1) The ρ meson width never seems to be quite the same from experiment to experiment. This effect became so pronounced that at one point the Rosenfeld tables⁽⁶⁾ ceased giving a width for the ρ meson and even now merely gives 125 ± 20 MeV, which would appear to be fairly safe. In the author's opinion, the probable cause for these differences is to be found in an accurate many body calculation for those states ($\pi\pi N$, $\pi\pi\pi$, $N3\pi$, $N4\pi$, etc.) where ρ mesons

are seen, but rescattering effects, three body effects, or final state interactions are significant. One would guess that these effects broaden the peak.

(2) The nucleon isovector form factor appears to be given by something like

$$\frac{1}{(q^2 + m_\rho^2)^2}$$

instead of

$$\frac{1}{q^2 + m_\rho^2}$$

as ρ meson dominance most naturally suggests. This is even more puzzling when one discovers in Eqn. (2) that the $N\bar{N}$ contribution is the most important single factor in producing the zero at $s = m_\rho^2$, which is to say that the $N\bar{N}$ state is the most important factor in producing the ρ meson as a resonance. As already pointed out in ref. (2), this seems to indicate that the ρ meson is best considered an $N\bar{N}$ bound state rather than a $\pi\pi$ resonance. This is the same as saying that in a world made of nucleons, which do not bind to make pions, the ρ meson would exist as a stable particle, but that since we do have pions, it decays into them.

(3) There seems to be no ρ meson production from $^1S_0 N\bar{N}$ ⁽⁷⁾. This is very mysterious and is the cause, in a recent paper applying the Veneziano model to $\bar{p}n \rightarrow \pi^+\pi^-\pi^-$,⁽⁸⁾ for adding an ad hoc factor killing the ρ meson. If indeed the thesis of ref. (8) is correct, and the $\bar{p}n \rightarrow \pi^+\pi^-\pi^-$ amplitude is essentially just the $\pi^- \rightarrow \pi^+\pi^-\pi^-$ amplitude with the initial π^- having a mass of $2m_N$, then we cannot avoid seeing ρ mesons. We do not see them; therefore, the annihilation amplitude cannot be the $\pi\pi$ scattering amplitude with one heavy π meson. Furthermore, things seem to be even much more complicated than this, because any model the author knows of says that there ought to be ρ mesons produced. There are resonance bands at an effective mass of m_ρ in the $\bar{p}n \rightarrow \pi^+\pi^-\pi^-$ Dalitz plot, but they are uniformly populated which is not what one naively

expects of a ρ meson. Perhaps again an adequate understanding of three body effects would show that we have ρ mesons, but that our naive expectations are significantly altered.

These difficulties indicate at least three significant points that ought to be investigated in the world of the ρ mesons and there are others more fundamental: e.g., the original question raised in 1960⁽⁹⁾ as to how the ρ meson acquires a mass if it is a Yang-Mills field. So, the ρ meson, historically the first of a new short-lived breed of meson (so short-lived in fact that serious questions were raised as to whether it deserved to be called a particle at all) still remains both an interesting and not very well understood member of our zoo.

Comparison of This Analysis With Other Work

There have been other analyses similar in nature to that undertaken here. Those of Ref. (10) and (11) are contained in Eqn. (1) in the following way: in Ref. (10), our Eqn. (2) was used with only the $\pi\pi$ channel and in Ref. (11), our Eqn. (2) was used with only the $\pi\pi$ channel on which a cutoff was placed in addition. In Ref. (12), a great variety of formulae are presented. Their Eqns. (12b) and (13) with $R = 1/m_\rho$ ought to agree very closely with the results here. And in fact, the curve presented here is right in the middle of the various curves they calculated. They did not, however, use $R = 1/m_\rho$ favoring $R = 0$ instead. Two arguments can be presented for $R \approx 1/m_\rho$:

- (a) Nonrelativistic theory predicts $R \approx$ interaction radius $\approx 1/m_\rho$.
- (b) The relativistic theory of Ref. (2) and here gives a phase shift close to that of the nonrelativistic formula where $R \approx 1/m_\rho$.

CONCLUSIONS

The analysis presented here suggests that:

$$(a) \quad \Gamma_\rho = 109 \text{ MeV} \pm 10 \text{ MeV}$$

$$m_\rho = 769 \text{ MeV} \pm 5 \text{ MeV}$$

The error bars are conservative estimates.

(b) A measurement of the full width at half maximum should provide a good estimate of the rho meson width, regardless of the height of the peak.

(c) It is very unlikely to find anything much different from

$$\left| F_\pi(m_\rho^2) \right|^2 = \frac{m_\rho^2}{\Gamma_\rho^2}$$

(d) Studies of final state interactions in $\pi N \rightarrow \pi\pi N$ and $N\bar{N} \rightarrow 3\pi$ should be undertaken in order to understand their effect on the observed resonance peaks in these interactions.

The author would like to thank Prof. J. J. Sakurai for a discussion of the $e^+e^- \rightarrow \pi^+\pi^-$ situation, and Prof. J. E. Augustin for communicating his latest results prior to publication.

APPENDIX

Here we present in some detail the method and functions for implementing in a numerical calculation equations (1) - (3) in the text.

In Table I we see the necessary ρ functions (phase space functions), thresholds, masses and approximate values of the coupling constants.

As shown in Ref. 2, we need the following functions (we take this opportunity to correct a remaining error in Ref. 2, which is the absence of an exponent of 2 on the $s - m_{\rho}^2$ in the fourth term of the equation for H_3 and on the $s - m_{\rho}^2$ under the integral sign):

$$f(s, t, u, \Lambda) = P \int_t^{\Lambda} \frac{ds'}{[(s'-u)(s'-t)]^{1/2}} \frac{1}{s'-s} \quad (0 < u < t) \quad (A.1)$$

$$f'(s, t, u, \Lambda) = \frac{1}{(s-t)(s-u)} \left[\frac{[(\Lambda-t)(\Lambda-u)]^{1/2}}{s-\Lambda} + \left(\frac{t+u}{2} - s \right) f(s, t, u, \Lambda) \right] \quad (A.2)$$

$$\begin{aligned} H_1(s, t) &= P \int_t^{\infty} \frac{(s'-t)^2}{(s'(s'-t))^{1/2}} \frac{ds'}{(s'-s_0)^2 (s'-s)} = \frac{(s-t)^2}{(s-s_0)^2} f(s, t, 0, \infty) \\ &+ \frac{\partial}{\partial s_0} \left[\frac{(s_0-t)^2}{s_0-s} \right] f(s_0, t, 0, \infty) + \frac{(s_0-t)^2}{s_0-s} f'(s_0, t, 0, \infty) \end{aligned} \quad (A.3)$$

$$\begin{aligned} H_2(s, t) &= P \int_t^{\infty} \frac{(s'-t)(s')}{[s'(s'-t)]^{1/2}} \frac{ds'}{(s'-s_0)^2 (s'-s)} = \frac{(s-t)s}{(s-s_0)^2} f(s, t, 0, \infty) \\ &+ \frac{\partial}{\partial s_0} \left[\frac{(s_0-t)(s_0)}{s_0-s} \right] f(s_0, t, 0, \infty) + \frac{(s_0-t)(s_0)}{s_0-s} f'(s_0, t, 0, \infty) \end{aligned} \quad (A.4)$$

$$\begin{aligned} H_3(s, t, u, \Lambda) &= P \int_t^{\Lambda} \frac{(s'-t)^2 (s'-u)^2}{[(s'-t)(s'-u)]^{1/2}} \frac{ds'}{s'(s'-s_0)^2 (s'-s)} = \\ &= \cosh^{-1} \frac{\Lambda - \frac{1}{2}(t+u)}{\frac{1}{2}(t-u)} - \frac{t^2 u^2}{s s_0^2} f(0, t, u, \Lambda) \end{aligned}$$

$$\begin{aligned}
 & + \frac{\partial}{\partial s_0} \left[\frac{(s_0 - t)^2 (s_0 - u)^2}{s_0 (s_0 - s)} \right] f(s_0, t, u, \Lambda) \\
 & + \frac{(s_0 - t)^2 (s_0 - u)^2}{(s_0 - s) (s_0)} f'(s_0, t, u, \Lambda) + \frac{(s - t)^2 (s - u)^2}{s (s - s_0)^2} f(s, t, u, \Lambda) \quad (A.5)
 \end{aligned}$$

(Note change in H_3 as compared to Ref. 2).

We further need

$$\begin{aligned}
 f''(s, t, u, \Lambda) = & \frac{1}{(s-t)(s-u)} \left\{ \left(\frac{t+u}{2} - s \right) f'(s, t, u, \Lambda) - \right. \\
 & - f(s, t, u, \Lambda) \left[1 + \left(\frac{t+u}{2} - s \right) \left(\frac{1}{s-t} + \frac{1}{s-u} \right) \right] \\
 & \left. - \frac{\sqrt{(\Lambda-t)(\Lambda-u)}}{s-\Lambda} \left[\frac{1}{s-t} + \frac{1}{s-u} + \frac{1}{s-\Lambda} \right] \right\}
 \end{aligned}$$

where the primes on f indicate partial differentiation with respect to s .

The reason we need f'' is to be found when we consider the singular points $s = 0$ and $s = s_0$. For these points, we do the following:

$$(1) \text{ Define } h_1(s) = (s-t)^2 f(s, t, 0, \infty).$$

Then

$$\begin{aligned}
 H_1(s_0, t) &= \frac{1}{2} \frac{\partial^2}{\partial s_0^2} h_1(s_0) \\
 &= f(s_0, t, 0, \infty) + (s_0 - t) f'(s_0, t, 0, \infty) + \frac{1}{2} (s_0 - t)^2 f''(s_0, t, 0, \infty) \quad (A.7)
 \end{aligned}$$

$$(2) \text{ Define } h_2(s) = s(s-t) f(s, t, 0, \infty).$$

Then

$$\begin{aligned}
 H_2(s_0, t) &= \frac{1}{2} \frac{\partial^2}{\partial s_0^2} h_2(s_0) \\
 &= f(s_0, t, 0, \infty) + (2s_0 - t) f'(s_0, t, 0, \infty) + \frac{1}{2} s_0 (s_0 - t) f''(s_0, t, 0, \infty) \quad (A.8)
 \end{aligned}$$

(3) Define $h_3(s) = (s-t)^2 (s-u)^2 f(s, t, u, \Lambda)$.

$$\begin{aligned} H_3(0, t, u, \Lambda) &= \frac{\partial}{\partial x} \left[\frac{h_3(x)}{(x-s_0)^2} \right]_{x=0} + \frac{\partial}{\partial s_0} \left(\frac{h_3(s_0)}{s_0^2} \right) \\ &= p(0, s_0) f'(0, t, u, \Lambda) + p'(0, s_0) f(0, t, u, \Lambda) \\ &\quad + p(s_0, 0) f'(s_0, t, u, \Lambda) + p'(s_0, 0) f(s_0, t, u, \Lambda) \end{aligned} \quad (A.9)$$

where

$$p(b, a) = \frac{(b-t)^2 (b-u)^2}{b-a} \quad (A.10)$$

and

$$p'(b, a) = p(b, a) \left[\frac{2}{b-t} + \frac{2}{b-u} - \frac{1}{b-a} \right] \quad (A.11)$$

Also,

$$\begin{aligned} H_3(s_0, t, u, \Lambda) &= - \frac{h_3(0, t, u, \Lambda)}{s_0^3} + \frac{1}{2} \frac{\partial^2}{\partial s_0^2} \left[\frac{h_3(s_0, t, u, \Lambda)}{s_0} \right] \\ &= - \frac{t^2 u^2 f(0, t, u, \Lambda)}{s_0^3} + \frac{1}{2} \left[p(s_0, 0) f''(s_0, t, u, \Lambda) \right. \\ &\quad \left. + 2p'(s_0, 0) f'(s_0, t, u, \Lambda) + p''(s_0, 0) f(s_0, t, u, \Lambda) \right] \end{aligned} \quad (A.12)$$

where

$$p''(b, a) = p(b, a) \left\{ \left(\frac{2}{b-t} + \frac{2}{b-u} - \frac{1}{b-a} \right)^2 + \frac{1}{(b-u)^2} - \frac{2}{(b-t)^2} - \frac{2}{(b-u)^2} \right\}$$

($H_1(s, t)$ and $H_2(s, t)$ are well behaved at $s = 0$)

The need for $s = 0$ is clear because $F(s) = D(0)/D(s)$. The need for $s = s_0$ is less obvious, but occurs because it is good to supply a minimizing routine with derivatives. One parameter being s_0 , we need $\frac{\partial D}{\partial s_0}$ and find that

$$\frac{\partial}{\partial s_0} \left[(s-s_0) H_1(s, t) \right] = H_1(s, t) - 2H_1(s_0, t) \text{ so that } H_1(s_0, t) \text{ is required.}$$

The formulas (A.1) to (A.13) were used to define $\psi(s)$ function and its derivatives to the author's version of Davidon's variable metric ⁽¹³⁾ which provided the results shown in Fig. 1 and discussed in the text. In Eqns. (A.5), (A.9) and (A.12), Λ was taken as $200 m_{\pi}^2$.

REFERENCES

1. V. L. Auslander, G. I. Budker, Iu. N. Pestov, V. A. Sidorov, A. N. Skrinsky, and A. G. Khabakhapashev: Phys. Letters 25B, 433(67) and Novosibirsk preprint 243 (68);
J. E. Augustin, J. C. Bizot, J. Buon, J. Haissihski, D. Lalanne, P C. Marin, H. Nguyen Ngoc, J. Perez-y- Jorba, F. Rumpf, E. Silva and S. Tarernier: Phys. Rev. Letters 20, 126 (68); Phys. Letters 28B, 508 (69); Private communication.
2. J. S. Ball and M. Parkinson, Phys. Rev. 162, 1509 (67).
3. Tomas Brunilla, Matts Roos and Jan Pisut: Cern preprint Th. 972 (Dec. 2 (68)).
4. J. B. Ehrman, Phys. Rev. 81, 412 (51);
R. G. Thomas, Phys. Rev. 88, 1109 (52);
For a modern discussion, see S. C. Frautschi, Phys. Letters 8, 141 (64).
5. The residues at the ρ pole are the crucial thing. We find these to be:
 $g_{\rho NN} = 2.06 \pm 0.03$; $g_{\rho \pi \omega} = 0.31 \pm 0.15$ compared to the expected 2.11 and 0.45. Both are in agreement with the expected values. The value 2.11 for $g_{\rho NN}$ is based on ρ dominance of the nucleon isovector form factor and ρ universality; so, the agreement here is better than we have a right to expect.
6. Naomi Barash-Schmidt, Angela Barbaro-Galteri, Leroy R. Price, Arthur H. Rosenfeld, Paul Sading, Charles G. Wohl: Lawrence Radiation Laboratory preprint UCRL 8030 Revised, *Jan. 69*
7. C. Baltay, P. Franzini, N. Gelfand G. Lutjens, J. Severiens, J. Steinberger, D. Tycko, D. Zanello: Phys. Rev. 140, B1039 (65);
N. Gelfand, Phys. Rev. 169, 1077 (68);
P. Anninos, L. Gray, P. Hagerty, T. Kalogeropoulos, S. Zenone, R. Bizzari, G. Ciapetti, M. Gaspero, I. Laasko, S. Lichtman, G. C. Monetti: Phys. Rev. Letters 20, 402 (68).

8. C. Lovelace, Phys. Letters 28B, 264 (68).
9. J. J. Sakurai, Annals of Physics 11, 1 (60).
10. G. J. Gounaris and J. J. Sakurai, Phys. Rev. Letters 21, 244 (68);
G. J. Gounaris, Fermi Institute preprint 68-86 (Dec. 1968).
11. M. T. Vaughn and K. C. Wali, Phys. Rev. Letters 21, 938 (68).
12. Matts Roos and Ján Pišút, Cern preprint TH. 1000.
13. W. C. Davidon, Argonne National Laboratory preprint ANL 5990 (1959).

TABLE I

Channel	Particles	Masses	Thresholds		Coupling (a) Constants	Functions ρ_i
			t	u		
1	$\pi\pi$	$m_\pi=1$	4.0	0	$C_1 \approx 0.90$	$\left[\frac{(s-t_1)^3}{16s} \right]^{1/2}$
2	$NN(+ -)$	$m_N=6.72$	180.6	0	$C_2 \approx 3.0$	$\left[\frac{(s-t_2)(s)}{4} \right]^{1/2}$
3	$\pi\omega$	$m_\omega=5.61$	43.7	21.2	$C_3 \approx 0.60$	$\left[\frac{(s-t_3)^3(s-u_3)^3}{16s} \right]^{1/2}$

(a) These are varied in order to fit the data. The approximate values shown are those expected from other considerations: $C_1 \approx 0.90$ corresponds to a 100-125 MeV width, depending on s_p and $D'(m^2)$, C_2 is inferred from universal ρ coupling and vector meson dominance of the electromagnetic form factor, and C_3 comes from the $\omega \rightarrow 3\pi$ decay width. See ref. 2 for more details.

TABLE CAPTION

The relevant data for insertion into Eqns. (1) - (3). We have used the values and conventions of Ref. 2 except for a permutation of channels. The (+ -) refers to the helicities of the nucleons. The integral over channel 3 has been cutoff sharply at $200 m_\pi^2$.

FIGURE CAPTION

The curve was fitted using eqns. (1) - (3) (see appendix for details) with four parameters, obtaining $s_0 = 380 \pm 8 \text{ m}_\pi^2$, $c_1 = 1.15 \pm 0.01$, $c_2 = 3.0 \pm 0.05$, $c_3 = 0.45 \pm 0.20 \text{ m}_\pi^{-1}$. The χ^2 was 25; see text for discussion.

