BNL - 49518 Informal Report UC-721

# EFFECTS OF MICROBIAL PROCESSES ON GAS GENERATION UNDER EXPECTED WIPP REPOSITORY CONDITIONS:

**Annual Report Through 1992** 

A. J. Francis and J. B. Gillow

Prepared for:
Sandia National Laboratories
Waste Isolation Pilot Plant Gas Generation Program
Albuquerque, New Mexico

under

U.S. DOE Contract No. 67-8602

by

Brookhaven National Laboratory Upton, NY 11973

Under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy.



# EFFECTS OF MICROBIAL PROCESSES ON GAS GENERATION UNDER EXPECTED WIPP REPOSITORY CONDITIONS:

**Annual Report Through 1992** 

A. J. Francis and J. B. Gillow

Prepared for:
Sandia National Laboratories
Waste Isolation Pilot Plant Gas Generation Program
Albuquerque, New Mexico
under
U.S. DOE Contract No. 67-8602
by
Brookhaven National Laboratory
Upton, NY 11973

#### **ABSTRACT**

Microbial processes involved in gas generation from degradation of the organic constituents of transuranic waste under conditions expected at the Waste Isolation Pilot Plant (WIPP) repository are being investigated at Brookhaven National Laboratory. These laboratory studies are part of the Sandia National Laboratories - WIPP Gas Generation Program. Gas generation due to microbial degradation of representative cellulosic waste was investigated in short-term (< 6 months) and long-term (> 6 months) experiments by incubating representative paper (filter paper, paper towels, and tissue) in WIPP brine under initially aerobic (air) and anaerobic (nitrogen) conditions. Samples from the WIPP surficial environment and underground workings harbor gas-producing halophilic microorganisms, the activities of which were studied in short-term experiments. The microorganisms metabolized a variety of organic compounds including cellulose under aerobic, anaerobic, and denitrifying conditions. In long-term experiments, the effects of added nutrients (trace

amounts of ammonium nitrate, phosphate, and yeast extract), no nutrients, and nutrients plus excess nitrate on gas production from cellulose degradation. Results to date (up to 200 days of incubation) show that: (i) gas production was not detected in abiotic control samples; (ii) cellulose incubated without nutrients produced limited but sustained gas production; (iii) the addition of a nutrient solution enhanced the biodegradation of cellulose as evidenced by an increase in the production of total gas, carbon dioxide, and nitrous oxide; (iv) in the presence of excess nitrate, gas production was the highest and nitrous oxide accumulated to varying amounts; (v) the addition of bentonite increased the background carbon dioxide concentration and appeared to stimulate microbial activity; and (vi) in addition to total gas and carbon dioxide production, cellulose degradation in nutrient-amended samples was evidenced by the gradual bleaching of brown paper towel, the formation of gas bubbles, the formation of paper pulp, and the appearance of a red color at the bottom of the sample bottles, indicating the growth of halophilic microorganisms. Estimates of the total gas production on the basis of initial results ranged from 0.001 mL g<sup>-1</sup> cellulose day<sup>-1</sup> to 0.039 mL g<sup>-1</sup> cellulose day<sup>-1</sup>.

# **CONTENTS**

1.0	INTE	RODUCTION	1
	1.1 E	Background	2
	1.2 E	Biochemistry of Cellulose Degradation	3
	1	1.2.1 Biodegradation of Cellulose	3
	1	1.2.2 Metabolic Diversity of Halophilic Microorganisms	8
2.0	EXP	PERIMENTAL APPROACH	10
3.0	SHO	PRT-TERM EXPERIMENTS	13
	3.1 F	Rationale	13
	3.2 N	Methodology	13
	3	3.2.1 Sample Collection	13
	3	3.2.2 Direct Counts of Bacteria	14
		3.2.3 Activity Measurements	14
	3	3.2.4 Denitrification Studies	15
	3.3 I	Results and Discussion	18
		3.3.1 Direct Counts of Bacteria	18
		3.3.2 Activity Measurements	18
		3.3.3 Denitrification Studies	28
	3.4 5	Summary, Short-Term Experiments	37
4.0	LON	NG-TERM EXPERIMENTS	41
		Objective	41
		Rationale	42
		4.2.1 Aerobic (Sealed) Treatments	43
		4.2.2 Anaerobic Treatments	44
		Materials and Methods	45
		4.3.1 Cellulose	45
		4.3.2 WIPP Brine	50
		4.3.3 Bentonite	50
		4.3.4 Inoculum	50
		Sample Treatments	53
		4.4.1 Anaerobic Sample Preparation	53
		4.4.2 Aerobic Sample Preparation	53
		1.1.2 Gas Analyses	54

# **CONTENTS** (Continued)

5.0 Results and Discussion (Long-Term Inundated Experiment)	54
5.1 Aerobic Treatments	55
5.1.1 Total Gas Production	55
5.1.2 Carbon Dioxide Production	58
5.1.3 Nitrous Oxide Production	61
5.2 Anaerobic Treatments	63
5.2.1 Total Gas	63
5.2.2 Carbon Dioxide Production	66
5.2.3 Nitrous Oxide Production	70
6.0 SUMMARY	70
7.0 REFERENCES	78
APPENDIX A Details of the Short-Term Activity Measurements	A-1
APPENDIX B Gas Analyses	<b>B-</b> 1
APPENDIX C Details of Long-Term Experiment	C-1
APPENDIX D Gas Production Data (Gross and Net) for the Long-Term Inundated Experiment	D-1
APPENDIX E Mixed Inoculum Activity Measurements	E-1

# 1.0 INTRODUCTION

Transuranic (TRU) wastes contain alpha-emitting transuranium nuclides with half-lives greater than twenty years and concentrations greater than 100 nCi per gram. TRU wastes are generated from nuclear weapons production and other related nuclear processing procedures. The wastes include adsorbed liquids, sludges, organics, and cemented materials containing the following radionuclides: <sup>232</sup>Th, <sup>233</sup>U, <sup>235</sup>U, <sup>236</sup>U, <sup>237</sup>Np, <sup>238</sup>Pu, <sup>239</sup>Pu, <sup>240</sup>Pu, <sup>241</sup>Pu, <sup>242</sup>Pu, <sup>241</sup>Am, <sup>244</sup>Cm, <sup>252</sup>Cf, and a variety of metals. Typically, TRU waste is classified as either contact-handled (CH), which does not require shielding, or remote-handled (RH), which requires shielding because of the hazard of gamma radiation exposure. The Waste Isolation Pilot Plant (WIPP) is a mined, geologic repository developed to demonstrate that radioactive transuranic wastes generated in defense-related activities can be safely and permanently disposed of underground. The WIPP is a U.S. Department of Energy facility located in southeastern New Mexico, about 2150 ft (656 m) below the surface, in a bedded salt, evaporite Permian formation. A major long-term concern is the potential for gas generation from the corrosion of Fe and Fe-based alloys, microbial degradation of cellulosic, plastic, and rubber materials, and radiolysis of brine and waste material by alpha-emitting radionuclides (Molecke, 1979 and Brush, 1990). Gas generation can cause pressurization and the formation of fractures which could allow the radionuclides to migrate away from the disposal site.

Anoxic corrosion and microbiological activity are the two most important processes that may generate appreciable amounts of gas. The current estimate of gas production due to anoxic corrosion is 900 moles per drum of waste (Brush, 1991). Caldwell et al. (1979) reported that microbial gas production due to biodegradation of TRU waste could be significant. Recently, Lappin et al. (1989) estimated such production rates at 1 mole of gas per drum of waste per year for 600 years. Brush 1991; and Brush et al., (1990) proposed a minimum and maximum range at 0 to 5 moles per drum per year, as did the earlier estimates by Molecke (1979).

Laboratory studies are under way to determine the rate and extent of gas production due to radiolysis, corrosion, and microbial activity, to support the Sandia National Laboratories-WIPP Gas Generation Program efforts to assess the long-term performance of the WIPP repository. Studies of the effects of microbial processes have been underway at Brookhaven National Laboratory since 1991. These laboratory studies are funded under Sandia National Laboratories contract no. 67-8602. \*This report summarizes the progress and status of the WIPP-funded work at BNL. The microbiological data obtained from the laboratory studies from initiation through 1992 are presented.

# 1.1 Background

The WIPP waste repository is located 2150 ft (656 m) below ground surface, with 56 rooms planned or under construction in a bedded salt formation. About 6,800 drums of waste in 55-gallon (208-L) containers will be placed in each room of 3,640-m³ capacity. Each drum will contain, on average, about 10 kg of cellulosic waste (approx. 70,000 kg of cellulosic per room), 70% of which is paper (Brush, 1990). The rest of the potentially biodegradable portion of the waste consists of plastic and rubber, and other organic compounds. Wastes consisting of inorganic process sludges from secondary waste treatment, containing a total of ~3 million moles of nitrate and a much smaller amount of phosphate, will also be emplaced in the WIPP (Brush, 1990; Brush et al., 1991).

Microorganisms can enter the WIPP from several sources, including: (i) association with the TRU waste; (ii) the surface environment via the mine ventilation systems and human intrusion; and (iii) resident populations in the salt crystals and brine formations. Alpha radiation from TRU waste is not expected to have significant effects on microbial activity (Barnhart et al., 1980; Francis, 1990). Previous studies of low-level radioactive wastes and waste leachates have shown that microbes in the wastes can metabolize organic carbon compounds (Francis et al., 1980a,b; Francis, 1985). Halotolerant and halophilic microorganisms (10<sup>1</sup> to 10<sup>5</sup> colony forming units/mL) including aerobic, nitrate-reducing, and anaerobic bacteria were detected in the WIPP surficial environment and underground

workings (work of R. Vreeland, West Chester University, to be published). Cellulose-degrading extreme halophiles from the underground workings also have been isolated (work of R. Vreeland, West Chester University, to be published). Introduced microorganisms, as well as resident or indigenous halotolerant and halophilic bacteria, can metabolize organic compounds and nitrate in the waste, and may generate metabolic byproducts, such as organic acids, alcohols, carbon dioxide, nitrous oxide, nitrogen, hydrogen sulfide, and methane.

# 1.2 Biochemistry of Cellulose Degradation

# 1.2.1 Biodegradation of Cellulose

The cellulosic portion of the TRU waste will approximately be comprised of the following (Brush, 1990): paper (70%), cloth (4%), plywood (10%), and lumber (16%) (untreated: 10% and, treated: 6%). In addition, the waste contains plastic materials (primarily polyethylene and polyvinyl chlorides) and rubber materials (primarily neoprene and hypalon), their characteristics may be altered by alpha-irradiation, which may enhance their biodegradability and potential for gas generation.

Cellulose, hemicellulose, and lignin make up the three major components of plant vascular material, lignocellulose. Lignin is a highly branched, constitutionally undefined aromatic polymer that makes up 15 to 38% of hardwood and softwood trees. It is considered highly resistant to biodegradation, although thermochemically modified lignin has been shown to biodegrade (Colberg and Young, 1982).

Cellulose is an unbranched polymer of several thousand D-glucose units linked together by  $\beta$ -1,4 glucosicic bonds. The strength of the polymer is derived from the multitude of hydrogen bonds, with concentrations of hydrogen bonds in microcrystalline regions and fewer bonds in amorphous regions. Cellulose is insoluble; therefore, hydrolysis is a

prerequisite to microbial degradation. Hydrolysis of cellulose results in the formation of cellobiose, which is then hydrolyzed to glucose (see Figure 1).

Biodegradation of cellulose by white-rot fungus Trichoderma reesei and the bacteria Cellulomonas has provided insights into the enzymology, the mechanisms of action, and the pathways of cellulose degradation. Several extracellular enzymes are involved in the breakdown of cellulose. The cellulase enzymes, consisting of exoglucanase (exoenzyme) and endoglucanase (endoenzyme), break the cellulose chains into various smaller fragments, starting with: (i) different 1,4- $\beta$ -endoglucanases that attack the 1,4- $\beta$ -linkages, randomly depolymerizing internal units; (ii) 1,4- $\beta$ -exoglucanases that remove cellobiose from the nonreducing chain end of the molecule, and (iii) 1,4-\beta-glucosidases (cellobiase), that hydrolyze cellobiose to glucose (Priest, 1984). Amorphous regions of cellulose are degraded by both the endo- and exo-glucanases separately. Synergistic action of the two enzymes is necessary for degrading crystalline cellulose (Poulsen and Peterson, 1992). These enzymes (produced by a variety of aerobic and anaerobic bacteria, fungi and protozoa) coordinate to hydrolyze cellulose into soluble components, which then are converted into a variety of end products. Bacteria, including aerobes such as Cellulomonas sp. and Cellvibrio gilvus (Bott and Kaplan, 1991), and anaerobes, such as Clostridium sp. (Benoit et al., 1992), Clostridium thermocellum (Lynd et al., 1989), Acetovibrio celluloyticus (Laube and Martin, 1981), and Ruminococcus albus (Pavlostathis et al., 1988) produce extracellular cellulase enzymes in the presence of cellulosics. These enzymes are induced by the presence of substrate (Hrmova et al., 1991) and are attenuated by soil and other absorptive materials (Hope and Burns, 1985). Close proximity of the cell to the substrate is necessary for degradation. A purified enzyme extract of T. reesei was shown to effectively degrade cellulose (Priest, 1984), and non-oxygen labile endoglucanase from Clostridium thermocellum was shown to strongly absorb to native cellulose (Ng et al., 1977). The rate of enzyme induction also depends upon the presence of the necessary nutrients, nitrogen and phosphorus (Skujins, 1976). Upon induction, hydrolysis by the cellulases is the rate-limiting step which, once achieved, follows first-order kinetics (Pavlostathis, 1991). The available surface area is an important determinant of the

Figure 1. Cellulose Degradation Pathway.

rate of digestion of cellulose. Studies of cellulytic bacteria in the rumen demonstrated the importance for degradation of adherence of microbial cells to the cellulose surface, with fermentation rates correlated to surface area (Weimer et al., 1990). The primary hydrolysis products of cellulose are cellobiose and glucose, which then are converted to organic acids, carbon dioxide, hydrogen, and methane by various microbial processes.

Glucose, generated from cellulose, is readily used by a variety of microorganisms. The specific process depends on the availability of electron acceptors such as oxygen, nitrate, sulfate, and CO<sub>2</sub>. In the presence of oxygen, carbon dioxide and water are formed during the oxidation of glucose.

$$C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O + energy$$

As oxygen is consumed, the alternate electron acceptors are used: nitrate, iron (III) oxides and hydroxides, manganese (IV) oxides and hydroxides, sulfate, and carbon dioxide. In the case of nitrate, dissimilatory reduction transforms nitrate to ammonium (dissimilatory nitrate reduction to ammonium (DNRA)) or to nitrous oxide, and then nitrogen (denitrification):

$$NO_3 \rightarrow NO_2 \rightarrow NH_4^+$$
 (DNRA)  
 $NO_3 \rightarrow NO_2 \rightarrow N_2O \rightarrow N_2$  (Denitrification)

Both processes are affected by the concentration of oxygen; the organisms catalyzing these transformations are microaerophiles or facultative anaerobes, capable of metabolism under low oxygen conditions or in its absence. Denitrification will slow down or cease with higher oxygen concentrations which inhibit the production of specific enzymes (Tiedje, 1988). Denitrifiers use several substrates, such as glucose and low molecular weight organic acids and alcohols. The use of nitrate as an alternate electron acceptor may be significant in the WIPP because of the presence of nitrate in the waste, predominantly from process sludges.

In the absence of oxygen and nitrate, anaerobic microorganisms will dissipate electrons via fermentation of the carbohydrate:

$$C_6H_{12}O_6 \rightarrow 2C_3H_4O_3$$
 (pyruvic acid) + 4H<sup>+</sup>  
4H<sup>+</sup> + 2C<sub>3</sub>H<sub>4</sub>O<sub>3</sub>  $\rightarrow$  2C<sub>3</sub>H<sub>6</sub>O<sub>3</sub> (lactic acid)

Fermentation products, such as low molecular weight organic acids and alcohols are available for preferential use by denitrifiers, sulfate reducers, and methanogens. Brines from the WIPP contain 160 to 300 mM sulfate (Brush, 1990); sulfate-reduction could be significant in the presence of metabolizable carbon. It occurs under reducing conditions (Eh -150 to -200mV, pH = 7.00), resulting in a change in Eh (-250mV) with growth and sulfide formation (Postgate, 1984):

2 lactate + 
$$SO_4$$
  $\rightarrow$  2 acetate +  $2CO_2$  +  $2H_2O$  +  $S$ 

Sulfate reduction results in the formation of H<sub>2</sub>S and insoluble metal sulfides. Sulfate-reducing bacteria (SRB) convert lactate, pyruvate, alcohols, amino acids, and acids of the tricarboxylic acid cycle to acetate and CO<sub>2</sub>. Glucose and other sugars seldom seem to be used directly by SRB.

The presence of CO<sub>2</sub>, H<sub>2</sub>, organic acids, and a low Eh generated by these anaerobic microbial processes provide a conducive environment for the growth of methanogenic bacteria. Methanogens can use (i) acetate; (ii) methanol; or (iii) carbon dioxide and hydrogen and produce methane:

(i) 
$$CH_3COO^- + H^+ \rightarrow CH_4 + CO_2$$
  
(ii)  $4CH_3OH \rightarrow 3CH_4 + CO_2 + 2H_2O$   
(iii)  $4H_2 + CO_2 \rightarrow CH_4 + 2H_2O$ 

Approximately 70% of the methane formed in sludge and freshwater sediments is due to reaction (i), whereas halophiles predominantly carry out reaction (ii).

Additionally, iron(III) reduction may be a significant process in the WIPP because of the presence of metallic and oxidized forms of iron (5 x 10<sup>6</sup> moles). Iron reduction involves the oxidation of organic carbon concomitant with the reduction of iron, whereby iron is used as the electron acceptor in the absence of oxygen, resulting in the dissolution of Fe(III) to Fe(II). Manganese reduction results in the formation of soluble Mn(II) from Mn(IV). Soluble uranyl ions can be reduced to insoluble U(IV) by anaerobic bacteria (Francis et al., 1991; Lovely, 1992). Corrosion caused by microbes could transform metal ions at a passive surface, resulting in metal sulfide precipitates (Kearns et al., 1992). The use of hydrogen solely from passivation can result in the formation of methane by methanogens, accelerating cathodic depolarization and increasing corrosion (Lorowitz et al., 1992). Figure 2 shows microbiologically mediated redox processes.

# 1.2.2 Metabolic Diversity of Halophilic Microorganisms

Halobacteria isolated from hypersaline environments can metabole a wide variety of organic compounds under aerobic and anaerobic conditions. Most of the extreme halophiles are archaebacteria; that is, they are a distinct group of microorganisms, apart from the eubacteria that make up the majority of prokaryotes, with an ancient lineage composed of other types of organisms adapted to extreme environments, such as alkaliphiles, thermophiles, and methanogens. Moderate halophiles grow best in an environment containing 0.5 to 2.5 M NaCl, while extreme halophiles grow best in 2.5 to 5.2 M NaCl (Kusher and Kamekur, 1988; Ventosa, 1988). Brines in the WIPP repository consist of 5.1-5.3 M chloride and 1.83-4 M sodium, 0.63-1.44 M magnesium, and 0.04-0.30 M potassium (Brush, 1990; Molecke, 1983).

Halophiles grow anaerobically by: (i) fermenting glucose, fructose, glycerol, citrate, and lactate (Javor, 1984); (ii) reducing nitrate to nitrogen gas using a variety of carbohydrates (Tomlinson et al., 1986); (iii) degrading chitin; and (iv) producing methane from methylamines, CO<sub>2</sub> and H<sub>2</sub> (Zhilina and Zavarzin, 1990). In hypersaline environments acetate-utilizing methanogens have been difficult to isolate (Zhilina and Zavarzin, 1990).




Figure 2. Electron-free energy for biologically mediated redox sequence with organic carbon (CH<sub>2</sub>O) acting as the electron donor. Calculated for standard conditions at pH 7. (CH<sub>2</sub>O) represents one-sixth of glucose (i.e. -153 kJ mol<sup>-1</sup>), from Zehnder, A.J., and W. Stumm. 1988. "Geochemistry and Biogeochemistry of Anaerobic Habitats," Biology of Anaerobic Microorganisms. Ed. A. J. Zehnder. New York, NY: John Wiley and Sons. 19.

Sulfate reducer also have been difficult to isolate, although saltmarsh sediments harbor abundant SRB populations (Dicker and Smith, 1985). The WIPP site contains a variety of halotolerant and halophilic bacterial populations. Isolates from underground brine seepages and salt crystals, and from brine and sediment from surficial lakes near the WIPP site, revealed a great diversity of colony characteristics when grown on solid media (work of R. Vreeland, West Chester University, to be published). These isolates were shown to use amino acids, glucose, and cellulose.

# 2.0 EXPERIMENTAL APPROACH

Laboratory experiments were designed to determine the potential gas generation due to biodegradation of cellulose under conditions expected in the WIPP repository (Figure 3). The experiments were divided into short-term (<6 months) and long-term (>2 years). In the short-term experiments, we examined the influence of electron donors and acceptors on the activities of specific microbial processes relevant to the WIPP disposal environment. In the long-term experiments, we measured gas generation due to biodegradation of cellulose under realistic conditions expected in the WIPP repository after the waste was in place. The conditions include humid and inundated, and initially aerobic and anaerobic environments. The effects of addition of nutrients and bentonite also was investigated.




Figure 3. Experimental Approach

#### 3.0 SHORT-TERM EXPERIMENTS

# 3.1 Rationale

The two main objectives of the short-term experiments were: (i) to determine gas production due to the activity of aerobes and anaerobes in the presence of specific electron donors (cellulose, glucose, succinate) and acceptors (oxygen, nitrate, sulfate) under hypersaline conditions; and (ii) to evaluate the suitability of inocula for use in the long-term experiments.

Anaerobic microbial processes were emphasized in the short-term experiments because they are expected to have the greatest impact on the long-term performance of the WIPP repository. These experiments provided an opportunity to specify aspects of the long-term investigations by manipulating the experimental variables. The microbes tested include those present in the brine collected from WIPP underground workings, surficial sediment slurries from the surrounding lakes at the WIPP site, and in axenic (pure) or mixed cultures isolated from these sources. In addition, analytical methods were tested and standardized during this phase.

# 3.2 Methodology

# 3.2.1 Sample Collection

Sediment and water samples from Nash Draw (near the WIPP site) and muck pile salt, rib salt, and brine (from the WIPP underground workings) were collected from August 21 to 25, 1991. Core samples of mud from Laguna Cinco, Quatro, Tres, and Surprise Springs, all in Nash Draw, were obtained using sterile iron pipes. Air was excluded by driving the core deep into the mud and capping it while submersed. Corrosion of iron end-caps also prevented contamination by oxygen. Lake brine samples were collected in sterile polyethylene containers from the four lakes as well as from Lindsay Lake, also in Nash

Draw. Mud and brine from all the lakes were also collected with sterile glass serum bottles, which were then stoppered and crimped to exclude air. Salt from the WIPP underground was collected using sterile spatulas and sterile containers. G-Seep brine from the WIPP underground was collected in sterile polyethylene bottles by Glen Barker, SNL.

The samples were shipped to BNL within two days after collection. The mud samples were extruded in a nitrogen-filled glove box and transferred to sterile serum bottles, fitted with butyl rubber stoppers, and stored at 4°C. Brine from Lindsay Lake and the WIPP site was also stored at 4°C. Viable bacteria in these samples were counted by Russell Vreeland, West Chester University, and the total number of bacteria and microbial activity were determined at BNL.

#### 3.2.2 Direct Counts of Bacteria

Samples were shaken on a wrist-action shaker for 45 minutes to disperse the contents. One mL was removed with a sterile needle and syringe, dispensed into a snap-cap vial and preserved with 5% (v/v) glutaraldehyde. Double-stranded DNA specific stain 4'6-diamidino-2 phenylindole dihydrochloride (DAPI, Polysciences, Inc.) was added to the sample and it was incubated for seven minutes in the dark. The sample was filtered through a 0.2  $\mu$ m black membrane filter (Poretics Corp.), and then placed on a slide and examined under oil-immersion at 1875 x magnification. Slides were prepared in triplicate for each sample, the blue-fluorescing cells were counted directly using a calibrated grid eyepiece under epifluorescent light. The DAPI stain differentiated the cells from salt grains: DNA-containing material was blue and the salt yellow (Coleman, 1980).

# 3.2.3. Activity Measurements

Production of gas by aerobes, anaerobes, and denitrifiers in a mixed inoculum of WIPP salt and brine and Nash Draw brine and sediment was determined. The carbon sources tested included glucose  $(C_6H_{12}O_6)$ , cellulose  $([C_6H_{12}O_6]_n)$ , and succinate  $(C_4H_6O_4)$ .

Two hundred g of S2180, W30 WIPP muck-pile salt were dissolved in sterile deionized water and diluted to 1 L. To a 450 mL aliquot of the solution, 10 mL of Laguna Cinco mud slurry and 40 mL of Laguna Cinco lake water were added. The mixture was kept in an anaerobic glove box. Nine and a half mL of the mixture were added with a sterile syringe to 20 mL sterile serum bottles containing 0.5 mL of a sterile concentrated nutrient stock solution containing 5 mg yeast extract, 10 mg potassium phosphate, 125 µmol ammonium nitrate, and 275 \(mu\)mol glucose (See Appendix A). Aerobic samples were incubated with an initial headspace of air in sealed containers, whereas anaerobic samples were incubated in a nitrogen atmosphere after purging the samples several times with nitrogen. The serum bottles were fitted and sealed with butyl rubber stoppers. Denitrification activity was determined by adding a nutrient solution containing 5 mg yeast extract, 10 mg potassium phosphate, 125  $\mu$ mol nitrate added as ammonium nitrate, 99  $\mu$ mol nitrate added as potassium nitrate, and 185  $\mu$ mol of succinate. These samples contained a total of 224  $\mu$ mol of nitrate and were prepared anaerobically. A set of aerobe, anaerobe, and denitrifier treatments were also prepared by including approximately 0.5 g filter paper instead of the respective carbon sources, glucose and succinate.

Six samples were prepared for each treatment, two of which were treated with 1 mL of 10% formaldehyde to serve as abiotic controls. Brine samples without nutrients were also prepared, and two were treated with formalin to determine endogenous activity. The headspace gas of all the samples was analyzed after 48, 83, and 147 days of incubation at 30°C, and the total volume of gas, carbon dioxide, and nitrous oxide in the headspace determined (see Appendix B).

# 3.2.4 Denitrification Studies

In the absence of oxygen and in the presence of metabolizable organic carbon, some aerobic bacteria can use nitrate as an alternate electron acceptor. This process, called

denitrification, or the reduction of nitrate, converts 80% or more of the available nitrate-N to nitrogen gas (Tiedje, 1988). Nitrate is reduced to nitrogen by denitrifying bacteria via the following steps:

$$NO_3^- \rightarrow NO_2^- \rightarrow N_2O \rightarrow N_2$$

Nitrate is also reduced by dissimilatory nitrate reduction to ammonium, but with either incomplete conversion to nitrogen or a far lesser yield of nitrogen (Tiedje, 1988). Nitrous oxide is an intermediate product and does not generally accumulate, although some halophiles accumulate nitrous oxide under certain conditions (Tomlinson et al., 1986).

Denitrification was observed in Nash Draw sediment slurry, G-seep, and a pure culture by the acetylene blockage technique (Yoshinari and Knowles, 1976). Acetylene inhibits the conversion (reduction) of  $N_2O$  to  $N_2$ , resulting in the stoichiometric accumulation of  $N_2O$  in the headspace (Balderston et al., 1976):

$$NO_3^- \rightarrow NO_2^- \rightarrow N_2O - \int \int \rightarrow N_2$$
  
 $C_2H_2$ 

Acetylene was injected into the duplicate samples of each treatment to give a final concentration of 10%. Nitrous oxide in the headspace of each sample was measured with a gas chromatograph equipped with a <sup>63</sup>Ni electron capture detector (see Appendix B). Samples were analyzed at 0, 21, and 43 hours, and at appropriate intervals thereafter.

#### 3.2.4.1 DENITRIFICATION IN SEDIMENT SLURRY SAMPLE

Laguna Cinco sediment (collected on August 22, 1991 and stored anoxically at 4°C for two months), and a fresh sample from the same site (collected on December 10, 1991) were examined for denitrifier activity. The fresh sample was examined within 48 hours of collection. The sample was mixed well, and a 1 mL slurry (August:  $0.56 \pm 0.01$  g dry sediment; December:  $0.59 \pm 0.01$  g dry sediment) was pipetted into sterile 20 mL serum

bottles in a glove box filled with nitrogen. One mL of filter-sterilized (0.22  $\mu$ m) brine (20% w/v WIPP halite) were added to the slurries resulting in a final volume of 2 mL. The slurry contained the following nutrients: (i) no additions (unamended); (ii) 3  $\mu$ mol succinate (carbon-amended); (iii) 300 nmol nitrate (nitrate-amended); and (iv) 3  $\mu$ mol succinate and 300 nmol nitrate (carbon-and nitrate-amended at a C:N ratio of 10:1). Succinate was used as the carbon source to discourage the growth of fermentative organisms. Triplicate samples of each treatment were prepared. The pH of each treatment was measured at the beginning and end of the experiment. Formaldehyde-treated samples served as controls. The serum bottles containing samples were sealed with rubber stoppers in the glove box in a nitrogen atmosphere and incubated at 30°C. Denitrification was determined by the acetylene blockage technique.

#### 3.2.4.2 DENITRIFICATION IN G-SEEP

G-Seep brine collected on December 10, 1991 from the WIPP underground workings was prepared within 48 hours of collection to assay for denitrification activity. Five mL aliquots of G-Seep brine were transferred into sterile serum bottles in an anaerobic glove box. One mL of filter-sterilized (0.22  $\mu$ m) brine (20% w/v WIPP halite) with nutrients was added. The G-Seep samples contained the following nutrients: (i) no additions (unamended); (ii) 1.5  $\mu$ mol succinate (carbon-amended); (iii) 150 nmol nitrate (nitrate-amended); and (iv) 1.5  $\mu$ mol succinate and 150 nmol nitrate (carbon- and nitrate-amended). Formaldehyde-treated samples served as controls. Triplicate samples of each treatment were prepared, sealed with butyl rubber stoppers in the glove box in a nitrogen atmosphere, and incubated at 30°C. Denitrification was determined by the acetylene blockage technique. Samples were analyzed after 0, 3, 30, and 60 days.

# 3.3 RESULTS AND DISCUSSION

# 3.3.1 Bacterial Population

Table 1 shows the numbers of bacteria, both viable and non-viable in brine samples from Nash Draw and in G-Seep from the WIPP underground. The direct counts of bacteria in Nash Draw samples range from  $5.5 \times 10^6$  to  $1.0 \times 10^7$  cells/mL. The G-Seep brine contained  $7.2 \times 10^4$  to  $3.2 \times 10^6$  cells/mL.

# 3.3.2 Activity Measurements

#### 3.3.2.1 **AEROBIC**

Table 2 shows the total gas, carbon dioxide, and nitrous-oxide production in the mixed inoculum slurry samples incubated initially under aerobic conditions with glucose, ammonium nitrate, yeast extract, and potassium phosphate. In the control samples, there was a slight increase in carbon dioxide due to abiotic reactions, but no nitrous oxide was detected. In amended samples, the concentration of both carbon dioxide and nitrous oxide increased.

# 3.3.2.2 ANAEROBIC GLUCOSE METABOLISM

Mixed inoculum incubated anaerobically with glucose, ammonium nitrate, yeast extract, and potassium phosphate showed little activity (Table 3). There was no significant production of carbon dioxide in amended samples. The reason for the lack of anaerobic metabolism of glucose in these samples is not known.

Table 1. Bacterial Populations in Nash Draw and G-Seep Samples

Brine Source	Number of Cells/mL
Nash Draw	
Surprise Springs	5.5 x 10 <sup>6</sup>
Laguna Cinco	6.8 x 10 <sup>6</sup>
Lindsay Lake	7.0 x 10 <sup>6</sup>
Laguna Tres South	9.0 x 10 <sup>6</sup>
Laguna Quatro	$1.0 \times 10^7$
WIPP Underground Workings	
G-Seep #9	7.2 x 10 <sup>4</sup>
G-Seep #23	$3.4 \times 10^6$

Table 2. Aerobic Glucose Metabolism by Mixed Inoculum.

	Incubation Time (Days)									
Treatment*	48 Gas Produced**			83 Gas Produced**			Gas Produced**			
	Total (ml)	CO <sub>2</sub> (µmoles)	N <sub>2</sub> Ο (μmoles)	Total (ml)	CO <sub>2</sub> (µmoles)	N <sub>2</sub> O · (μmoles)	Total (ml)	CO <sub>2</sub> (µmoles)	N <sub>2</sub> O (μmoles)	
Control (n=2) [Formalin treated]	0.80 ± 0.10	14.0 ± 0.6	nd	1.31 ± 0.57	15.8 ± 0.2	nd	0.79 ± 0.60	20.1 ± 0.1	nd	
Amended (n=4)	$0.20 \pm 0.20$	76.9 ± 6.2	7.20 ± 4.58	$0.83 \pm 0.48$	83.0 ± 15.3	8.75 ± 6.14	$0.55 \pm 0.43$	80.4 ± 14.8	8.27 ± 5.71	

<sup>\*</sup> Each sample contained 125 µmoles of ammonium nitrate, 278 µmoles of glucose, 5 mg yeast extract, and 10 mg potassium phosphate

20

<sup>\*\*</sup> Dissolved gas concentration not included nd = not detected

Table 3. Anaerobic Glucose Metabolism by Mixed Inoculum.

Treatment*	48 Gas Produced**			Incubation Time (Days)  83  Gas Produced**			147 Gas Produced**			
	Total (ml)	CO2 (µmoles)	N <sub>2</sub> Ο (μmoles)	Total (ml)	CO <sub>2</sub> (µmoles)	N2O (μmoles)	Total (ml)	CO 2 (µmoles)	<u>N₂O</u> (μmoles)	
Control (n=2) [Formalin treated]	0.60 ± 0.50	12.6 ± 0.3	nd	1.03 ± 0.23	13.5 ± 0.2	nd	0.13 ± 0.14	16.2 ± 0.1	nd	
Amended (n=4)	nd	$5.6 \pm 0.15$	nd	$0.34 \pm 0.20$	7.72 ± 0.18	$0.478 \pm 0.030$	-0.12 ± 0.12	8.00 ± 0.20	2.32 ± 1.59	

<sup>\*</sup> Each sample contained 125 μmoles of ammonium nitrate, 278 μmoles of glucose, 5 mg yeast extract, and 10 mg potassium phosphate \*\* Dissolved gas concentration not included

#### 3.3.2.3 CELLULOSE DEGRADATION

## 3.3.3.3.1 Aerobic

Table 4 shows cellulose degradation in samples incubated initially under aerobic conditions with filter paper, ammonium nitrate, yeast extract, and potassium phosphate. Control samples showed no activity. A significant increase in carbon dioxide was observed after 83 days in amended samples. One sample produced 136  $\mu$ mol of carbon dioxide, and the pressure increased to 4.47 psi. The paper in the bottle showed disintegration, and dissolution. Blackening of specific samples was noted after 147 days, indicative of sulfate reduction.

## 3.3.2.3.2 Anaerobic

Samples incubated anaerobically with filter paper, ammonium nitrate, yeast extract, and potassium phosphate showed a significant increase in activity after 48 days (Table 5). Nitrous oxide was detected in amended samples, but not in the controls. An increase in carbon dioxide was observed after 83 days. There was a large variation in gas production of anaerobic samples, one sample produced  $103 \mu mol$  of carbon dioxide and the pressure reached 5.12 psi. The filter paper in this sample also disintegrated.

# 3.3.2.3.3 Cellulose Degradation in the Presence of Excess Nitrate

Mixed inoculum incubated anaerobically with filter paper, ammonium nitrate, yeast extract, potassium phosphate, and excess nitrate produced less nitrous oxide than the succinate-amended samples (Table 6). Only 20% of the nitrate was converted to nitrous oxide in the samples with acetylene. Carbon dioxide production was not significant until 147 days.

Table 4. Cellulose Degradation by Mixed Inoculum Under Aerobic Conditions.

		48		Incub	oation Time (Da 83	ys)		147	
Treatment*	Gas Produced**			Gas Produced**			Gas Produced**		
	Total (ml)	CO <sub>2</sub> (µmoles)	<u>N2O</u> (μmoles)	Total (ml)	<u>CO</u> 2 (μmoles)	N2O (μmoles)	Total (ml)	<u>CO</u> 2 (μmoles)	N <sub>2</sub> O (μmoles)
Control (n=2) [Formalin treated]	0.90 ± 0.20	10.9 ± 0.5	nd	1.65 ± 0.49	12.7 ± 0.1	nd	1.00 ± 0.43	14.7 ± 0.0	nd
Amended (n=4)***	nd	32.0 ± 1.9	7.29 ± 1.84	0.95 ± 0.61	57.1 ± 22.8	4.12 ± 1.38	nd	$63.9 \pm 20.7$	5.97 ± 2.12

<sup>\*</sup> Each sample contained 125 μmoles of ammonium nitrate, 0.5 grams of filter paper, 5 mg yeast extract, and 10 mg potassium phosphate

7

<sup>\*\*</sup> Dissolved gas concentration not included

<sup>\*\*\*</sup> Samples exhibited disintegration of the filter paper at 83 days

Table 5. Cellulose Degradation by Mixed Inoculum Under Anaerobic Conditions.

		48		Incubation Time (Days) 83				147	
Treatment*		Gas Produced	**		Gas Produced	**	***************************************	Gas Produced*	*
	Total (ml)	CO <sub>2</sub> (µmoles)	N <sub>2</sub> O (µmoles)	Total (ml)	<u>CO 2</u> (μmoles)	<u>N2O</u> (μmoles)	Total (ml)	CO <sub>2</sub> (µmoles)	<u>N<sub>2</sub>O</u> (μmoles)
Control (n=2) [Formalin treated]	$0.60 \pm 0.00$	11.3 ± 0.1	nd	1.02 ± 0.22	12.5 ± 0	nd	0.17 ± 0.17	14.3 ± 0.0	nd
Amended (n=4)									
1	nd	8.85	9.29	0.12	10.7	5.79	0.05	13.1	nd
2***	nd	6.83	10.7	3.48	65.0	nd	1.90	95.6	nd
3***	nd	9.12	nd	3.76	103	nd	0.77	90.8	nd
<b>4***</b>	nd	9.73	9.43	2.18	53.0	nd	3.81	105	nd
Mean	nd	8.63 ± 0.54	$7.36 \pm 2.14$	$2.39 \pm 0.72$	57.9 ± 16.5	1.45 ± 1.25	1.63 ± 0.71	76.1 ± 18.4	nd

Each sample contained 125 μmoles of ammonium nitrate, 0.5 grams of filter paper, 5 mg yeast extract, and 10 mg potassium phosphate
 Dissolved gas concentration not included
 At 83 days, samples exhibited disintegration of filter paper

Table 6. Cellulose Degradation by Mixed Inoculum in the Presence of Excess Nitrate.

				Incul	oation Time (Da	ıys)			
		48			83			147	
Treatment*		Gas Produced	**		Gas Produced	**		Gas Produced	**
	Total (ml)	CO <sub>2</sub> (µmoles)	N <sub>2</sub> O (μmoles)	Total (ml)	<u>CO</u> <sub>2</sub> (μmoles)	N2O (µmoles)	Total (ml)	<u>CO 2</u> (μmoles)	<u>N2O</u> (μmoles)
Control(n=2) [Formal	in treated]								
(w/acetyl.)	nd	8.93	nd	1.03	11.9	nđ	0.73	12.7	nd
(w/o acetyl.)	nd	9.90	nd	0.24	10.9	nd	-0.44	12.0	nd
Amended (n=4)									
(w/acetyl., n=2)	nd	$5.24 \pm 0.40$	23.6 ± 1.2	$0.38 \pm 0.27$	20.3 ± 11.4	$19.5 \pm 2.1$	1.39 ± 1.97	56.7 ± 36.2	14.4 ± 1.7
(w/o acetyl., n=2)	nd	$8.00 \pm 0.95$	5.59 ± 4.00	$0.06 \pm 0.02$	8.13 ± 0.41	$4.94 \pm 0.78$	$3.62 \pm 0.88$	78.0 ± 5.2	nd

Each sample contained 224 μmoles of nitrate, 0.5 grams of filter paper, 5 mg yeast extract, and 10 mg potassium phosphate
 Dissolved gas concentration not included

Cellulose degradation by mixed inoculum was observed in samples incubated under aerobic and anaerobic conditions and in the presence of excess nitrate. Disintegration of the filter paper in aerobic and anaerobic samples was noted after 83 days, indicating cellulose degradation. Analysis of these samples showed an increase in total gas, carbon dioxide, and nitrous-oxide production (Tables 4 and 5). The filter paper exhibited areas of thinning and clearing at 83 days in samples containing excess nitrate. At 147 days, the filter paper had fully disintegrated and carbon dioxide content had increased (Table 6). Aerobic samples containing cellulose showed little increase in carbon dioxide at 147 days; one sample turned black, possibly indicating the onset of sulfate reduction. This sample was checked after 220 days and the presence of hydrogen sulfide was confirmed by gas chromatography. Anaerobic samples also produced gas at 147 days, and one sample showed blackening (Sample 2, Table 5).

#### 3.3.2.4 DENITRIFICATION

Mixed inoculum samples incubated anaerobically with succinate, ammonium nitrate, yeast extract, potassium phosphate, and excess nitrate exhibited denitrification activity. Complete conversion of nitrate (224  $\mu$ mol) to nitrous oxide (129  $\mu$ mol) was noted in the presence of acetylene after 48 days of incubation (Table 7). The accumulation of nitrous oxide in amended samples without acetylene was much less than in the samples containing acetylene, indicating that nitrous oxide was converted to nitrogen. Control samples showed no activity. Additional studies on denitrification by WIPP sediment slurry brine and an axenic culture isolated from the brine are described next.

#### 3.3.3 Denitrification Studies

## 3.3.3.1 DENITRIFICATION IN SEDIMENT

Microbial denitrification was analyzed in freshly collected sediment and stored sediment. Nitrous oxide was both being produced and converted to  $N_2$  at the same time (acetylene was

Table 7. Denitrification Activity by Mixed Inoculum.

				Incub	ation Time (Da	ys)			
		48			83			147	
Treatment*		Gas Produced	**		Gas Produced	**		Gas Produced*	*
	Total (ml)	CO <sub>2</sub> (µmoles)	N <sub>2</sub> Ο (μmoles)	Total (ml)	CO <sub>2</sub> (µmoles)	N <sub>2</sub> O (μmoles)	Total (ml)	CO 2 (µmoles)	<u>N<sub>2</sub>O</u> (μmoles)
Control(n=2] [Formal	in treated]								
w/acetyl.)	nd	8.93	nd	1.37	10.2	nd	1.08	11.8	nd
w/o acetyl.)	nd	8.26	nd	nd	11.5	nd	nd	12.9	nd
Amended (n=4)									
w/acetyl., n=2)	nđ	$3.43 \pm 0.24$	129 ± 2	$0.50 \pm 0.20$	$6.41 \pm 0.05$	114 ± 5	-0.12 ± 0.24	na	na
(w/o acetyl., n=2)	nd	51.0 ± 36.1	$6.97 \pm 2.12$	1.07 ± 0.20	30.4 ± 21.5	12.1 ± 2.0	$1.01 \pm 0.39$	na	nd

27

nd = not detected

na = not analyzed

Each sample contained 224 μmoles of nitrate, 185 μmoles of succinate, 5 mg yeast extract, and 10 mg potassium phosphate Dissolved gas concentration not included

not added to the samples) as shown in Figures 4A and 4B. The nitrous oxide concentrations reported in these studies do not include N<sub>2</sub>O dissolved in the brine solution. Nitrous oxide was not detected in the control sample, which had been treated with formaldehyde. In unamended samples and samples amended with succinate, ~5 nmol of N<sub>2</sub>O was detected at about 45 hours of incubation, and N<sub>2</sub>O was not detected in the headspace thereafter. In samples amended with nitrate, the N<sub>2</sub>O concentration reached its maximum (30 nmol gdw<sup>-1</sup>) at about 200 hours and then disappeared rapidly by conversion to N<sub>2</sub> (Figure 4A). In the succinate and nitrate amended samples, much less N<sub>2</sub>O was detected than in the nitrate-amended samples. Freshly collected samples showed little accumulation of N<sub>2</sub>O in the headspace (Figure 4B), most probably due to rapid and complete denitrification.

The rates of denitrification in the unamended and amended (succinate and nitrate) sediment samples were determined by the acetylene blockage technique. Addition of acetylene inhibits the reduction of N<sub>2</sub>O to N<sub>2</sub> and allows N<sub>2</sub>O to accumulate in the headspace which is analyzed by gas chromatography. In Figures 5A and 5B and Table 8a and 8b, denitrification in stored (2 months) and freshly collected (assayed within 48 hours) samples are compared. In the stored samples, Figures 5A and 5B, denitrifying activity initially in the unamended sample was minimal. After lag phase, denitrification proceeded at about 0.14 nmol N<sub>2</sub>O gdw<sup>-1</sup>. The rate is presented in Table 9a. The sediment contained sufficient indigenous carbon and nitrate to produce 23 nmoles of N<sub>2</sub>O gdw<sup>-1</sup>. Denitrification in the stored, carbon- amended samples was similar to that in the unamended samples (0.22 nmol N<sub>2</sub>O gdw<sup>-1</sup> h<sup>-1</sup>). Denitrification activity in the stored sample was stimulated by the addition of nitrate. After an initial lag, the rate of denitrification was 1.36 nmol nitrous oxide gdw<sup>-1</sup> h<sup>-1</sup>.

A stimulatory effect on the rate and extent of denitrification was seen in both stored and fresh samples when carbon and nitrate were added. Denitrification proceeded at 1.78 nmol N<sub>2</sub>O gdw<sup>-1</sup> h<sup>-1</sup>, finally producing 209 nmol nitrous oxide gdw<sup>-1</sup> in the stored sample, translating to denitrification of 80% of the added nitrate. In the freshly collected sediment, without added carbon or nitrate, 205 nmoles of nitrous oxide gdw<sup>-1</sup> were produced at a rate




Figure 4. Denitrification by Laguna Cinco sediment; (A) stored sample, (B) fresh sample




Figure 5. Denitrification determined by the acetylene blockage technique in (A) stored sample, (B) fresh sample

Table 8a. Denitrification in Laguna Cinco Sediment (Stored 2 Months)

Time	Control*	None	Addition Carbon	Nitrate	Carbon & Nitrate
(Hours)			nmol nitrous oxide	**/sample	
0	nd	nd	nd	nd	nd
21		$1.96 \pm 0.58$	2.16	$1.50 \pm 0.11$	0.514
42.5	nd	$3.44 \pm 0.64$	na	3.21	$3.86 \pm 0.22$
67		8.16	$0.00 \pm 0.00$	$9.64 \pm 0.74$	9.64
155	nd	$20.7 \pm 1.4$	$19.6 \pm 0.1$	$129 \pm 5$	$166 \pm 6$
183		$22.3 \pm 3.0$	$19.4 \pm 0.9$	$183 \pm 0$	$194 \pm 2$
207		na	na	$192 \pm 0$	$209 \pm 2$
231		na	na	$168 \pm 4$	$191 \pm 9$
303	nd	$23.3 \pm 1.5$	$18.3 \pm 0.2$	$171 \pm 8$	$191 \pm 5$

Table 8b. Denitrification in Laguna Cinco Sediment (Assayed Within 48 Hours)

Time	Control*	None	Addition Carbon	Nitrate	Carbon & Nitrate
(Hours)			nmol nitrous oxide	**/sample	***********************
0	nd	nd	nd	nd	nd
45		na	na	$13.6 \pm 0.1$	$5.58 \pm 3.94$
66	nd	$26.5 \pm 1.1$	$29.0 \pm 1.0$	30.5	30.5
91.5		$43.0 \pm 0.3$	$44.2 \pm 4.6$	$39.4 \pm 1.2$	$45.2 \pm 0.5$
114	nd	$68.6 \pm 3.1$	$61.6 \pm 1.8$	$65.6 \pm 2.1$	$79.6 \pm 1.2$
138		$86.0 \pm 3.7$	$78.1 \pm 1.5$	$110 \pm 3$	$141 \pm 5$
165		$142 \pm 5$	$115 \pm 3$	$164 \pm 12$	$233 \pm 12$
309		$188 \pm 11$	$194 \pm 2$	$357 \pm 21$	$436 \pm 18$
405	nd	$205 \pm 1$	$202 \pm 8$	$372 \pm 3$	$458 \pm 15$
-	nd	• • • • • • • • • • • • • • • • • • • •		$372 \pm 3$	458

<sup>\*</sup> Formalin treated samples

<sup>\*\*</sup> Dissolved gas concentration not included

nd = none detected

na = not analyzed

Table 9a. Denitrification Rate (Stored Sample)\*

Addition	Rate			
(nmol nitrou	itrous oxide/g dry weight/hour)			
None (Unamended)	0.14			
Carbon	0.22			
Nitrate	1.36			
Carbon and Nitrate	1.78			

<sup>\*</sup>Rate after lag phase, from 67 to 155 hours (see Table 8a).

Table 9b. Denitrification Rate (Fresh Sample)\*

Addition	Rate				
(nmol nitrou	(nmol nitrous oxide/g dry weight/hour)				
None (Unamended)	1.17				
Carbon	0.87				
Nitrate	1.93				
Carbon and Nitrate	3.00				

<sup>\*</sup>Rate after lag phase, from 66 to 165 hours (see Table 8b).

of 1.44 nmol nitrous oxide gdw<sup>-1</sup> h<sup>-1</sup>, as shown in Table 9b. These samples are not carbon-limited. Addition of nitrate, however, increased N<sub>2</sub>O production, suggesting that these samples are nitrate-limited. In the absence of carbon or nitrate limitations in the freshly collected sediment, denitrification proceeded at 3.00 nmol N<sub>2</sub>O gdw<sup>-1</sup> h<sup>-1</sup>, and resulted in the production of 458 nmol nitrous oxide gdw<sup>-1</sup> (Table 8b); all of the total nitrate present was reduced.

These experiments show that microorganisms present in mud from Laguna Cinco denitrified 70-80% of the added nitrate in nitrate-amended samples and 80%-100% of the nitrate in carbon and nitrate samples. We added 540 nmol nitrate gdw<sup>-1</sup> to each sample; therefore, complete conversion of the nitrate to nitrous oxide should yield 270 nmol N<sub>2</sub>O gdw<sup>-1</sup>. About 70% of the added nitrate was reduced to nitrous oxide in the stored sample, or 192 nmoles N<sub>2</sub>O gdw<sup>-1</sup> was produced. Freshly collected sediment with added nitrate produced 372 nmoles N<sub>2</sub>O gdw<sup>-1</sup>; 81% of the total nitrate present was reduced. This sediment contained a large quantity of indigenous nitrate.

Denitrification in unamended samples provides information about the nutrient conditions in the sediment. If all the available indigenous nitrate was converted to nitrous oxide in the carbon-amended sample, then the stored sediment contained about 40 nmoles of nitrate gdw<sup>-1</sup>, and the fresh sediment contained 410 nmol gdw<sup>-1</sup>. Similar conversions of nitrate to nitrous oxide using a pure culture were used to determine sub-ppb concentrations of nitrate in lake waters (Christensen and Tiedje, 1988). Thus, the assay can be used not only to detect the presence and activities of denitrifying organisms, but also to detect easily metabolizable low-molecular-weight organic carbon compounds in the environment (Francis et al., 1989).

The presence of metabolizable carbon and nitrogen compounds in the WIPP surficial environments has important implications. For example, a steady mix of microbial populations can be actively maintained when presented with an adequate supply of a

limiting nutrient. From the standpoint of the long-term experiment, we have shown in part that the organisms are active in the proposed inoculum.

The results also show that active denitrification under anaerobic conditions occurred in Laguna Cinco mud collected in August 1991 and stored for two months, and in mud collected in December 1991, and assayed within 48 hours. Therefore, the stored and fresh samples contain viable organisms that can be used as an inoculum for the long-term experiment.

#### 3.3.3.2 DENITRIFICATION IN G-SEEP

Figure 6 and Table 10 show denitrification in G-Seep brine. In the carbon and nitrogen amended samples  $\sim 84\%$  of the added nitrate (150 nmoles) was converted to  $N_2O$  (62.7 nmoles) after two months of incubation. The dissolved  $N_2O$  in the brine was not determined; hence the  $N_2O$  values reported are not corrected for  $N_2$  solubility. Lack of  $N_2O$  production in the unamended and carbon amended samples suggests that G-Seep brine is nitrate-limited. However, in the nitrate amended samples, 45% of the added nitrate was converted to  $N_2O$ , indicating metabolizable carbon is present which eventually became limiting. These results also suggest that the microbes in the brine were able to readily metabolize the added carbon and nitrate via denitrification.

### 3.3.3.3 DENITRIFICATION BY AN AXENIC PURE CULTURE

A pure culture of a denitrifying bacterium was isolated from the sediment slurry sample during the initial denitrification experiments. The isolate was grown in the following medium: sodium succinate, 5 g; potassium nitrate, 1 g; yeast extract, 0.5 g; WIPP salt (20% w/v), 1000 mL; pH 6.85. The isolate, designated as BWFG-1, was a gram-negative rod, facultative anaerobe, and grew rapidly within 48 hours in the liquid medium. On solid medium, the culture produced circular, convex, light-orange colonies with entire margins.




Figure 6. Denitrification in G-Seep Brine. [In C&N amended samples, 84% of the added nitrate was converted to nitrous oxide]

Table 10. Denitrification in G-Seep Brine

			Amendment*		
Time	Control**	Unamended	Carbon	Nitrate	Carbon & Nitrate
(Days)		nn	noles nitrous oxide**	*/sample	
0	nd	nd	nd	nd	nd
3.25	nd	nd	nd	nd	nd
30	nd	$3.30 \pm 0.01$	$2.53 \pm 0.43$	$24.2 \pm 0.8$	3 55.9 ± 4.4
60	nd	$5.72 \pm 0.72$	$7.58 \pm 2.98$	$34.0 \pm 0.9$	$62.7 \pm 4.0$

Samples injected with acetylene to accumulate nitrous oxide

nd = none detected

36

<sup>\*</sup> Amendment: Carbon = 1.5 umol succinate, Nitrate = 150 nmol

<sup>\*\*</sup> Formalin treated samples

<sup>\*\*\*</sup> Dissolved gas concentration not included

BWFG-1 is an archaebacterium. It was confirmed by hybridization with fluorescently labeled oligodeoxynucleotide probes complementary to the 16S ribosomal RNA segment specific for archaebacteria (DeLong et al., 1989). Extreme halophiles were previously described in the literature as being archaebacteria, placing them in a distinct phylogenetic group of organisms that contains other genera from extreme environments, such as methanogens, thermoacidophiles, and alkalophiles (Ross et al., 1981).

The rate of denitrification by BWFG-1 was determined by adding 2.5 mL of a 24 h culture to 40 mL of medium in the presence of acetylene. Control samples included uninoculated medium, inoculated samples treated with 0.5% v/v formaldehyde, and inoculated samples without acetylene. Triplicate samples were incubated anaerobically at 30°C. The number of bacterial cells were counted using the DAPI method.

Production of nitrous oxide by the pure culture of (25  $\mu$ mol h<sup>-1</sup>) is shown in Figure 7 and Table 11. The bacterium denitrified nitrate at a rate 2.5  $\mu$ mol h<sup>-1</sup>. About 72% of the added nitrate (392  $\mu$ mol) was converted to N<sub>2</sub>O (142  $\mu$ mol) in about three days; these N<sub>2</sub> values do not include the amount of N<sub>2</sub>O dissolved in the growth medium. There was no accumulation of nitrous oxide in samples incubated without acetylene, indicating complete reduction of nitrate to N<sub>2</sub>. In Figure 8, the direct counts of the bacteria during the course of denitrification are presented. A marked increase in the number of cells corresponding to nitrous oxide production was observed. The pH of the growth medium increased from 6.85 to 8.00 after three days, and turbidity measured spectrophotometrically at 600 nm was 0.08 at 0h, 0.58 at 29h, and 0.80 at 50h.

# 3.4 Summary, Short-Term Experiments

1. Direct microscopic examination of brine from Nash Draw lakes and from G-Seep showed that between 10<sup>4</sup> to 10<sup>7</sup> cells/mL bacteria are present.




Figure 7. Denitrification by a pure culture of bacteria isolated from the WIPP site. [72% of added nitrate was converted to nitrous oxide at a rate of  $2.5 \,\mu$ moles/h]

Table 11. Denitrification by a Pure Culture Isolated from the WIPP Environment: Nitrous Oxide Production Over Time

Time (Hours)	Inoculated (w/o acetyle		Inocula (w/ acet		Inoc.* (w/o acet.)	Inoc.* (w/acet.)	Uninoc. (w/o acet.)	Uninoc. (w/acet.)
				µmoles nitro	us oxide**/sam	ple		
0	nd		nd		nd	nd	nd	nd
3	$0.001 \pm$	0.001	0.011	$\pm 0.034$	nd	nd	nd	nd
4.5	$0.007 \pm$	0.002	0.245	$\pm 0.081$	nd	nd	nd	nd
6	$0.009 \pm$	0.002	0.434	± 0.090	nd	nd	nd	nd
7	$0.010 \pm$	0.003	0.563	± 0.016	nd	nd	nd	nd
9.25	$0.009 \pm$	0.003	0.990	± 0.027	nd	nd	nd	nd
13	$0.014 \pm$	0.003	2.33	± 0.03	nd	nd	nd	nd
23	$0.019 \pm$	0.003	15.2	± 0.9	nd	nd	nd	nd
29	$0.187 \pm$	0.084	23.0	± 0.5	nd	nd	nd	nd
36	1.50 ±	0.38	63.8	± 2.6	nd	nd	nd	nd
50	$0.086 \pm$	0.059	130	± 2	nd	nd	nd	nd
78	$0.140 \pm$	0.030	142***	± 1	nd	nd	nd	nd

Defined culture medium (per sample): 740 umoles succinate, 392 umoles nitrate, 20% w/v WIPP salt.

ND = none detected

<sup>\* =</sup> Sample treated with formalin (0.5%)

<sup>\*\* =</sup> Dissolved gas concentration not included

<sup>\*\*\* = 72%</sup> of nitrate added converted to nitrous oxide




Figure 8. Growth of denitrifying bacteria.

- 2. Cellulose was degraded by a mixed culture derived from samples consisting of Nash Draw sediment slurry, salt crystals, and G-Seep brine in the presence of added nutrients (nitrate, phosphate, and yeast extract). Cellulose degradation was confirmed by an increase in carbon dioxide production and disintegration of filter paper.
- 3. Storage of sediment and lake water at 4°C for about two months did not significantly affect the activity of microbes in the samples.
- 4. The denitrification assay is a useful method to rapidly determine the activity of dentrifying microbes in WIPP samples. The assay also confirmed the presence of metabolizable carbon in the sediment and in WIPP brine.
- 5. Denitrifiers were detected in G-Seep, although their source was not identified.
- 6. An axenic culture of archaebacteria was isolated from the WIPP site denitrified nitrate at a rate of 2.5  $\mu$ mol h<sup>-1</sup>. The characteristics and growth rate of this culture have been elucidated for future studies to examine the influence of environmental variables on specific microbial processes in the WIPP repository.
- 7. Short-term experiments have provided useful information on microbial activity under accelerated test conditions that are relevant to the WIPP repository; further work will include examination of the other anaerobic processes such as fermentation, sulfate reduction, and methanogenesis.

## **4.0 LONG-TERM EXPERIMENTS**

## 4.1 Objective

The objective of the long-term study is to determine the rate and extent of gas generation over the long term (>2 years) from cellulose biodegradation under humid and inundated conditions, in the presence and absence of added nutrient.

### 4.2 Rationale

The TRU waste that will be placed in the WIPP repository contains an average of about 10 kg of cellulosic material per drum, approximately 70% of which is paper (Brush 1990a). Initially, the repository will be ventilated, but the addition of backfill (salt, or bentonite/salt mixture to fill void spaces around waste containers) will seal the drums inside the disposal rooms. Initially, it will also be dry, but after sealing the repository, humid conditions will develop. The ambient humidity is expected to be 18 to 27 g/m<sup>3</sup> (about 74% relative humidity, RH), and the temperature about 30°C (Brush, 1990). Microenvironments of condensed liquid brine may exist under humid conditions. Diffusion of water vapor through high-efficiency particulate, HEPA, air filters on waste containers will result in humid conditions inside the containers. Eventually, corrosion or rupturing of the containers due to salt creep-room closure will expose the waste to salt and backfill. Process sludges from other breached waste containers are expected to be leached by the brine. This is presumed to be the major source of nitrate and phosphate (nutrients) in the repository. The accumulation of potentially intruding brines from the surrounding Salado Formation will most likely begin after sealing the rooms, which will inundate them. In the event of potential, inadvertent human intrusion, fluids may also seep in from the Castille Formation into the Salado Formation. The atmosphere inside the disposal environment will become anaerobic in the short-term (months to years) due to consumption of oxygen by corrosion, radiolysis, and microbial processes acting on the waste materials. Microenvironments of trapped air that contain oxygen will continue to exist after sealing. Radiolysis of organic wastes will deplete oxygen, whereas radiolysis of nitrate-bearing sludges will release oxygen. Radiolysis of brines may also produce some oxygen.

A succession of microbial processes will occur under the changing environmental conditions inside the repository. Environment changes from aerobic to anaerobic, humid to inundated (and possibly back to humid), and asaline to saline will stimulate the activities of (i) microbes initially present in the waste, and (ii) resident and indigenous halotolerant

or halophilic bacteria in the brine and salt. To examine the influences of various microbial processes on gas generation, sample prepared were treated to simulate the following scenarios.

#### 4.2.1 Aerobic

During the early stages of waste emplacement, the environment will be aerobic but will become anaerobic over a period of time after closure because of corrosion, aerobic microbial activity, and radiolytic processes. In these long-term experiments, the cellulose samples will be placed in serum bottles, sealed with air, and incubated. The conditions will be initially aerobic and become anaerobic with time due to consumption of oxygen by aerobes, thus paving the way for anaerobic microbial activity.

### 4.2.1.1 SCENARIO 1

After emplacement and sealing of waste containers in WIPP disposal rooms, the intact and nearly intact containers will be isolated from backfill and brine. The humidity inside the disposal rooms is expected to be 18 to 27 g/m³ (about 74% RH), and humidity inside the containers is expected to reach equilibrium with the room environment. The cellulose will be in an asaline, humid, aerobic environment for possibly months, several years, or up to a few decades, as water vapor diffuses through the waste drum particulate filters. Microorganisms capable of cellulose degradation and gas production under these conditions will be active probably under a sub-optimal moisture content.

#### 4.2.1.2 SCENARIO 2

Room closure and corrosion will breach many of the containers and expose the waste material to backfill, salt, and brine. The cellulose is expected to contact the salt and backfill material, and microbial degradation of the cellulose is expected to occur under saline, humid conditions.

#### 4.2.1.3 SCENARIO 3

Influx of intruding brines from the Salado Formation, brine capillary rise through the backfill, and brine dissolution will all tend to inundate some portion or all of the disposal rooms with brine. Brine inundation will accelerate the onset of anoxic conditions as any residual air pockets are flooded. Process sludge TRU wastes contain significant quantities of nitrate and lesser quantities of phosphate. The breach of these sludge containers and inundation by brine will then transport the nitrate and will bring nonhalophilic, halotolerant and halophilic microbes into contact with cellulose.

Inundation of the WIPP waste by brine by the above mentioned or other processes will accelerate the activities of halophilic and halotelerant microbes. In particular, dentrification activity under microaerophilic and anaerobic conditions could be significant and may contribute to the total quantity and to the proportion of gases produced (N<sub>2</sub>, N<sub>2</sub>O, and CO<sub>2</sub>).

#### 4.2.2 Anaerobic

At least a portion of the WIPP repository wastes will be anaerobic at the start (within their containers possibly due to radiolysis and microbial action at the initial stages) and remain anaerobic thereafter. Under these conditions, short-term (i.e., operational phase) and in the long-term, degradation of cellulosic waste by anaerobic microorganisms could be significant.

### 4.2.2.1 SCENARIO 4

Some of the cellulose in the disposal environment may become anaerobic before any significant aerobic microbial activity. Cessation of air flow from closure of the disposal rooms, and oxic corrosion plus radiolysis, may bring about anoxic conditions in a humid environment. If the cellulose is exposed to salt under humid conditions, halotolerant or halophilic microbes that can grow in humid and anoxic environments may be involved in

degrading of cellulose. With the onset of anoxic conditions, alternate electron acceptors such as nitrate and sulfate will be used by microbes in the degradation of cellulose and its product intermediates.

### 4.2.2.2 SCENARIO 5

With the onset of brine intrusion in the disposal rooms, inundation will be more likely to cause anaerobiosis by forcing out any residual trapped air. Cellulose in contact with brine may undergo degradation by halophilic and halotolerant microbes present in the brine and waste. Because of the breaching of the waste containers, it is likely that nitrate originating in the sludges will be transported by the brine. It may come in contact with cellulosic wastes and enhance the degradation of cellulose.

Scenarios 3 and 5 will be examined in the long-term inundated experiment. Scenarios 1, 2 and 4 will be examined in a long-term humid experiment in CY1993. Figures 9 through 12 give the complete treatment matrix for the long-term inundated experiments.

### 4.3 Materials and Methods

#### 4.3.1 Cellulosics

Simulated TRU cellulosic waste material was composed of four types of paper: (i) filter paper, (ii) white paper towel, (iii) brown paper towel, and (iv) Kimwipes<sup>•</sup> (lintless tissue wipes). These papers are typical cellulosic wastes from laboratory and process activities. The four paper types were cut into strips in a large paper shredder, and then cut into 1 cm x 1 cm squares.

Samples of shredded paper types, each weighing 1.25 g, were thoroughly mixed and added to 160 mL acid washed (10% HCl), sterile (autoclaved at 120°C, 20 psi for 20 min) serum bottles.




Figure 9. Long-term inundated experiment treatment matrix (aerobic samples).




Figure 10. Long-term inundated experiment treatment matrix (aerobic samples containing bentonite).




Figure 11. Long-term inundated experiment treatment matrix (anaerobic samples).




Figure 12. Long-term inundated experiment treatment matrix (anaerobic samples containing bentonite).

#### 4.3.2 WIPP Brine

Fifteen liters of G-Seep #9 brine were provided by SNL and stored at 4°C until use.

### 4.3.3 Bentonite

Bentonite clay in two one-L containers was provided by SNL. The bentonite was a granular MX-80 Volclay bentonite available from the American Colloid Company of Belle Fourche, SD (Table 12).

#### 4.3.4 inoculum

The microbial inoculum used in these studies were obtained from the following three sources (Table 13): (i) mud and brine from Nash Draw: collected on December 12, 1991 and stored at 4°C, the mud was stored anoxically in serum bottles, (mud samples were filtered through sterile cotton in a nitrogen-filled glove box to remove large particulates); (ii) brine from the WIPP underground workings: 200 mL of G-Seep were collected on December 12, 1991; and (iii) asaline inoculum from laboratory contamination: 2.5 g of dust containing asaline microorganisms was gathered from labs in Bldg. 318 at BNL.

The mud, brine, and dust samples were then mixed together in a sterile beaker in a nitrogen-filled glove box. The total volume of the mixed inoculum was 583 mL. The activity of the mixed inoculum was examined by incubation under aerobic and anaerobic conditions in the presence of metabolizabled substrate. The results are presented in Appendix E.

Table 12. Composition of Bentonite\*.

Chemical Composition	(NaCa) <sub>.35</sub> (Al <sub>1.60</sub> Fe <sub>.15</sub> (Si <sub>3.90</sub> Al <sub>.10</sub> ) 0 <sub>10</sub> (OH)		
Montmorillonite Content	90 Percent		
Typical Chemical Analysis	Silica Alumina Iron (Ferric) Iron (Ferrous) Magnesium Sodium and Potassium Calcium Crystal Water Trace Elements		63.02% SiO <sub>2</sub> 21.08% Al <sub>2</sub> O <sub>3</sub> 3.25% Fe <sub>2</sub> O <sub>3</sub> 0.35% FeO 2.67% MgO 2.57% Na <sub>2</sub> O  0.67% CaO 5.64% H <sub>2</sub> O 0.72%
Exchangeable Ions (Milli-equivalents/100 g)	Sodium Calcium Magnesium	55-65 15-25 10-15	
Moisture Content	10% Maximum as S	Shipped	
рН	8.5 - 10.5		

<sup>\*</sup> Data provided by the American Colloid Company, Skokie, IL.

Table 13. Composition of Mixed Inoculum.

Source .	Mud Slurry (mL)	Brine (mL)	
Laguna Quatro Mud and Brine	60	40	
Laguna Cinco Mud and Brine	35	40	
Laguna Tres South Mud and Brine	13	40	
Lindsey Lake Mud and Brine	50	40	
Surprise Springs Mud and Brine	25	40	
G-Seep Brine		200	
Total	183	400	

## 4.4 Sample Treatments

The treatments consisted of (a) 100 mL of brine, and (b) 100 mL of brine and 5 g mixed cellulosic papers. The samples were incubated with and without nutrients, which consisted of yeast extract (0.05%), potassium phosphate dibasic (0.1%), and ammonium nitrate (0.1%). Some samples also received excess nitrate in the form of potassium nitrate (0.5%).

## 4.4.1 Anaerobic Sample Preparation

The serum bottles containing the mixed cellulosic papers were flushed with nitrogen and placed inside an anaerobic, nitrogen-containing glove box for 24 hours before inoculation to remove any trapped air. G-Seep brine (10 L) was removed from storage at 4°C and equilibrated overnight at room temperature. One hundred mL of the brine solutions (with and without nutrients or excess nitrate) were added to sample bottles with and without bentonite containing either no cellulose, cellulose, or glucose. Bentonite (6 g) was added to separate sample bottles inside the glove box to determine its influence on gas production. The samples were gently mixed to distribute the bentonite.

The microbial inoculum prepared from various sources was continually mixed and 4 mL was added to specific samples (3.8% V/V inoculum). The samples were gently mixed (to blend the inoculum) and then capped with butyl rubber stoppers. Control samples received 3 mL of 37% formaldehyde to give a final concentration of 1% formaldehyde.

## 4.4.2 Aerobic Sample Preparation

Aerobic (sealed) samples were prepared as described above with the following exceptions: 1) brine solutions were not purged with ultra high-purity (UHP)  $N_2$ , 2) the mixed inoculum was removed from the glove box; 3) brine was added to the bottles, inoculated, and capped with butyl rubber stoppers outside of glove box, thereby sealing air

in the headspace. Appendix C has a detailed description of all the treatments (aerobic and anaerobic) and the number of replicate samples. All samples were placed in a  $30 \pm 2^{\circ}$ C incubator.

# 4.4.3 Gas Analyses

The headspace gas of select samples was analyzed for total gas production, carbon dioxide, and nitrous oxide at time 0 (January 29, 1992) and thereafter at monthly intervals. Control samples were analyzed less frequently. The methods used for the headspace gas analyses are presented in Appendix B.

## 5.0 RESULTS AND DISCUSSION

The treatments consist of cellulose samples which were (i) uninoculated, (ii) inoculated with a mixed inoculum, (iii) inoculated and amended with nutrients, (yeast extract (0.05%), potassium phosphate (0.1%), and ammonium nitrate (0.1%)), and (iv) inoculated with nutrients plus excess nitrate (0.5%) potassium nitrate).

The results presented for aerobic and anaerobic samples represent the amount of gas produced per gram of cellulose (Figures 13 through 24). A detailed description of the procedure used to calculate the results are given in Appendix D. Tables 1 through 12 in Appendix D present data on a per sample basis, and Tables 13 through 24 in Appendix D present data on a per gram cellulose basis. Gas production rates on a per gram cellulose per day basis are presented in Table 14, and on per drum of waste per year basis in Table 15.

### 5.1 Aerobic Treatments

#### 5.1.1 Total Gas Production

Figure 13 shows the total gas produced in samples incubated with an initial atmosphere of air (aerobic). The formalin-treated control samples showed no increase in total gas production, and in fact showed a slight decrease. Likewise, uninoculated and inoculated samples which received no nutrients showed a slight decrease in total gas (-0.18 mL g<sup>-1</sup> cellulose and -0.34 mL g<sup>-1</sup> cellulose respectively (Table 13, Appendix D)). The decrease in total gas may be due, in part, to sampling. A decrease in gas production was more evident in inoculated samples because of oxygen consumption, indicating the start of microbial activity, (oxygen was not analyzed in these samples but is planned for the future). In the nutrient-amended inoculated samples, an initial decrease in gas volume (-0.27 mL g<sup>-1</sup> cellulose at 45 days) was followed by an increase after 69 days to 0.86 mL g<sup>-1</sup> cellulose at 200 days at a rate of 0.008 g<sup>-1</sup> cellulose day<sup>-1</sup>. This rate was calculated from linear slope of gas production from 69 to 200 days. Excess nitrate stimulated the rate of gas production (0.023 mL g<sup>-1</sup> cellulose day<sup>-1</sup> after 69 days) resulting in a total of 4.42 mL g<sup>-1</sup> cellulose at 200 days. This stimulatory effect was also evidenced by the lack of a long lag-phase (see Figure 13) and was a result of the metabolism of dissolved carbon in the presence of nitrate.

Total gas production in aerobic samples containing bentonite is presented in Figure 14. Uninoculated and inoculated samples, with no added nutrients, did not produce gas. Inoculated samples containing nutrients produced 4.38 mL of gas g<sup>-1</sup> cellulose after 200 days (Table 14, Appendix D), at a rate of 0.028 mL g<sup>-1</sup> cellulose day<sup>-1</sup>. In the presence of excess nitrate, the total production increased to 6.07 g<sup>-1</sup> cellulose at 200 days at a rate of 0.034 g<sup>-1</sup> cellulose day<sup>-1</sup>. Enhanced total gas production was seen in samples containing bentonite, and was apparently due to a combination of abiotic and biotic factors, which are evident upon examining carbon dioxide evolution in the presence of bentonite.




Figure 13. Total gas produced in samples incubated with an initial atmosphere of air.




Figure 14. Total gas produced in samples containing bentonite incubated with an initial atmosphere of air.

Extrapolation of the gas production rates from mL per g cellulose per day to mol per drum of waste per year is accomplished with the following conversion factors: for an assumed average drum of transuranic waste, with about 10 Kg of cellulosic materials, a total gas generation rate of 0.01 mL g<sup>-1</sup> cellulose day<sup>-1</sup> corresponds to a gas generation rate of 1.6 mol gas per drum per year.

## 5.1.2 Carbon Dioxide Production

Uninoculated samples produced 4.00  $\mu$ mol of CO<sub>2</sub> g<sup>-1</sup> cellulose at 200 days, which was slightly less than the formalin treated controls (7.62  $\mu$ mol g cellulose<sup>-1</sup>), as shown in Figure 15 (and Table 15, Appendix D). However, inoculated samples produced 8.30  $\mu$ mol of CO<sub>2</sub> g<sup>-1</sup> cellulose, slightly higher than uninoculated and formalin-treated controls, due to the onset of microbial activity. Inoculated samples containing nutrients produced 40.8  $\mu$ mol carbon dioxide g<sup>-1</sup> cellulose at 200 days, at a rate of 0.283  $\mu$ mol g<sup>-1</sup> cellulose day<sup>-1</sup>. In the presence of excess nitrate, 95.6  $\mu$ mol carbon dioxide g<sup>-1</sup> cellulose were produced at a rate of 0.484  $\mu$ mol g<sup>-1</sup> cellulose day<sup>-1</sup>, more than twice that of samples without excess nitrate. Evidence of the growth of halophilic bacteria was noted in nutrient-amended samples by a red/pink color at the bottom of the bottles. This red coloration, characteristic of halophiles, is caused by the presence of bacteriorhodopsin, a 50-carbon carotenoid pigment. This coloration was not seen in formalin-treated controls or unamended samples.

The addition of bentonite resulted in the production of a significant amount of abiotically produced carbon dioxide in samples without cellulose. Table 5, Appendix D shows that carbon dioxide increased from 17.7  $\mu$ mol sample<sup>-1</sup> to 40.0  $\mu$ mol sample<sup>-1</sup> without cellulose, inoculum and nutrients (sample 4(NC)-a), compared to the same treatment without bentonite. The latter treatment showed a slight increase, from 1.38 to 2.18  $\mu$ mol sample<sup>-1</sup> (see Table 2, Appendix D). Formalin-treated control samples also showed the same trend. The net effect was an increase in abiotically produced carbon dioxide (approximately 40  $\mu$ mol sample<sup>-1</sup>, (Table 5) by the addition of bentonite. Carbon dioxide production was insignificant in uninoculated unamended samples, (Figure 16). The




Figure 15. Carbon dioxide produced in samples incubated with an initial atmosphere of air.




Figure 16. Carbon dioxide produced in samples containing bentonite incubated in an initial atmosphere of air.

inoculated unamended samples with bentonite produced  $21.5 \,\mu\text{mol g}^{-1}$  cellulose at 200 days (Figure 16), whereas the samples without bentonite produced  $8.30 \,\mu\text{mol g}^{-1}$  cellulose (Table 15, Appendix D). Nutrient-amended inoculated samples produced  $69.8 \,\mu\text{mol}$  carbon dioxide  $g^{-1}$  cellulose, whereas nutrient-amended inoculated samples with excess nitrate produced  $116 \,\mu\text{mol g}^{-1}$  cellulose at 200 days. After an initial lag of 69 days, carbon dioxide was produced at a rate of 0.533 and 0.869  $\,\mu\text{mol g}^{-1}$  cellulose day<sup>-1</sup> in nutrient amended and nutrient amended plus excess nitrate samples, respectively. These rates of carbon dioxide production are higher than the same treatments without bentonite (0.283 and 0.484  $\,\mu\text{mol}$  g cellulose<sup>-1</sup> day<sup>-1</sup>, respectively). The buffering effect of CaCO<sub>3</sub>, as well as minerals and trace elements including Fe, Al, Si and exchangeable cations (Na<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>) and anions (Cl<sup>-</sup>, SO<sub>4</sub><sup>2-</sup>) present in the bentonite (Wanner et. al., 1992), may enhance microbial activity. Bentonite also provides an attachment site for microorganisms that may favor growth.

Extrapolation of the carbon dioxide production rates from  $\mu$ mol per  $g^{-1}$  cellulose day<sup>-1</sup> to mol drums<sup>-1</sup> of waste year<sup>-1</sup> is accomplished with the following conversion factor: for an assumed average drum of transuranic waste, with about 10 Kg of cellulosic materials, a carbon dioxide gas generation rate of 1.0  $\mu$ mol CO<sub>2</sub>  $g^{-1}$  cellulose day<sup>-1</sup> corresponds to 3.7 mol of gas drum<sup>-1</sup> year<sup>-1</sup>.

## 5.1.3 Nitrous-Oxide Production

Nutrient-amended samples contained 250  $\mu$ mol nitrate g<sup>-1</sup> cellulose, while nutrient-amended samples plus excess nitrate contained 1240  $\mu$ mol nitrate g<sup>-1</sup> cellulose. Acetylene was not added to samples and therefore the nitrous oxide is both being produced and reduced to N<sub>2</sub> in these samples. Nitrous oxide was not detected in uninoculated or inoculated samples without amendments, indicative of the lack of microbial activity (see Figure 17). Nitrous oxide accumulated in the inoculated nutrient-amended samples, with production peaking at 24.4  $\mu$ mol g<sup>-1</sup> cellulose at 132 days and then declining to 1.76  $\mu$ mol g<sup>-1</sup> cellulose at 164 days (Table 17, Appendix D). Nitrous-oxide was produced at a rate of 0.674  $\mu$ mol g<sup>-1</sup> cellulose day<sup>-1</sup> from 69 to 104 days. In the presence of excess nitrate, nitrous




Figure 17. Nitrous oxide produced in samples incubated with an initial atmosphere of air.

oxide was produced at the rate of 0.835  $\mu$ mol g<sup>-1</sup> cellulose day<sup>-1</sup>, reaching 115  $\mu$ mol g<sup>-1</sup> cellulose at 200 days.

In the presence of bentonite, nitrous oxide was not detected in uninoculated and inoculated unamended samples (see Table 18, Appendix D and Figure 18). The addition of nutrients to inoculated stimulated nitrous oxide production from 69 to 104 days at 1.00  $\mu$ mol g<sup>-1</sup> cellulose day<sup>-1</sup>. Thereafter, N<sub>2</sub>O did not accumulate in the headspace, probably because of depletion of available nitrate, or rapid conversion of N<sub>2</sub>O to nitrogen gas. In the presence of excess nitrate, nitrous oxide was produced at a rate of 0.589  $\mu$ mol g<sup>-1</sup> cellulose day<sup>-1</sup>, reaching a maximum (82.7  $\mu$ mol g<sup>-1</sup> cellulose) at 200 days. The continued accumulation of nitrous oxide at 200 days was probably due to the abundance of available nitrate. Addition of bentonite did not result in a substantial accumulation of N<sub>2</sub>O in the headspace, suggesting that N<sub>2</sub>O was rapidly converted to N<sub>2</sub> as soon as it was formed. Samples with and without bentonite exhibited the same trend (Figures 17 and 18), although less N<sub>2</sub>O was detected in samples containing bentonite.

### 5.2 Anaerobic Treatments

### 5.2.1 Total Gas Production

Figure 19 shows the total gas produced in samples incubated under anaerobic conditions in the presence of nitrogen. Uninoculated samples showed a slight loss of gas of about 3.20 mL sample<sup>-1</sup>, presumably due to sampling. Inoculated samples without nutrients produced 0.59 mL g<sup>-1</sup> cellulose at 200 days (see Table 19, Appendix D). After a lag of about 45 days, total gas production increased in inoculated samples containing nutrients, which produced 2.27 mL g<sup>-1</sup> cellulose at a rate of 0.021 mL g<sup>-1</sup> cellulose day<sup>-1</sup>. With nutrients plus excess nitrate, 5.44 mL of gas were produced at a rate of 0.039 mL g<sup>-1</sup> cellulose day<sup>-1</sup>.




Figure 18. Nitrous oxide produced in samples containing bentonite incubated with an initial atmosphere of air.




Figure 19. Total gas produced in anaerobic samples.

Figure 20 shows total gas produced in anaerobic samples containing bentonite. Uninoculated unamended, samples showed a net loss at 200 days to -0.28 mL g<sup>-1</sup> cellulose (Table 20, Appendix D), due to a combination of sampling and data correction. inoculated unamended, samples produced 0.81 mL g<sup>-1</sup> cellulose after a lag of 69 days, at a rate of 0.007 mL g<sup>-1</sup> cellulose day. In nutrient-amended samples, the addition of bentonite had no significant effect and it was similar to samples without bentonite (Figure 19). In contrast the addition of bentonite increased gas production in aerobic-amended samples. Inoculated amended samples produced 1.92 mL of gas at a rate of 0.013 g<sup>-1</sup> cellulose day<sup>-1</sup>, and inoculated samples with excess nitrate produced 3.52 mL of gas at a rate of 0.025 g<sup>-1</sup> cellulose day<sup>-1</sup>.

## 5.2.2 Carbon Dioxide Production

Uninoculated samples produced only about 3.59  $\mu$ mol carbon dioxide g<sup>-1</sup> cellulose over 200 days (Figure 21), but in the presence of inoculum, carbon dioxide increased to 5.47  $\mu$ mol g<sup>-1</sup> cellulose (Table 21, Appendix D). After a lag of 69 days, inoculated samples amended with nutrients produced 26.0  $\mu$ mol carbon dioxide at a rate of 0.198  $\mu$ mol g<sup>-1</sup> cellulose day<sup>-1</sup> (Table 21, Appendix D). Addition of excess nitrate stimulated carbon dioxide production to 61.4  $\mu$ mol at the rate of 0.422  $\mu$ mol g<sup>-1</sup> cellulose day<sup>-1</sup>.

Figure 22 shows carbon dioxide production in anaerobic samples containing bentonite. Uninoculated unamended samples produced  $0.22~\mu\text{mol g}^{-1}$  cellulose at 200 days, while inoculated unamended samples produced  $8.28~\mu\text{mol g}^{-1}$  cellulose. The addition of bentonite enhanced the background (abiotic) carbon dioxide concentration by about  $40.0~\mu\text{mol}$  ( Table 22, Appendix D). Inoculated samples plus nutrients produced  $31.8~\mu\text{mol}$  carbon dioxide  $g^{-1}$  cellulose over 200 days at a rate of  $0.236~\mu\text{mol}$   $g^{-1}$  cellulose day<sup>-1</sup>. Inoculated samples with nutrients plus excess nitrate produced  $35.0~\mu\text{mol}$  carbon dioxide  $g^{-1}$  cellulose over 200 days at a rate of  $0.252~\mu\text{mol}$   $g^{-1}$  cellulose day<sup>-1</sup>.




Figure 20. Total gas produced in anaerobic samples containing bentonite.




Figure 21. Carbon dioxide produced in anaerobic samples.




Figure 22. Carbon dioxide produced in anaerobic samples containing bentonite.

## 5.2.3 Nitrous Oxide Production

The rate of nitrous oxide accumulation in anaerobic samples is shown in Figure 23. Nitrous oxide was not detected in either uninoculated or inoculated samples (Table 23, Appendix D). In the inoculated amended samples, nitrous oxide accumulated to 15.5  $\mu$ mol after 100 days and remained relatively unchanged up to 200 days. In the presence of excess nitrate, nitrous oxide was produced after 69 days at a rate of 0.602  $\mu$ mol g<sup>-1</sup> cellulose day<sup>-1</sup>, reaching a concentration of 79  $\mu$ mol at 200 days. In contrast, nitrous oxide accumulation was higher in aerobic samples (see Figure 15) than anaerobic samples.

Figure 24 shows nitrous oxide production in anaerobic samples containing bentonite. Nitrous oxide was not detected in uninoculated and inoculated samples (see Table 24, Appendix D). In inoculated samples with nutrients, only trace amounts of nitrous oxide were detected at 104 and 164 days. In the inoculated samples containing excess nitrate, nitrous oxide was produced at a rate of  $0.647 \mu \text{mol g}^{-1}$  cellulose day<sup>-1</sup> after 104 days and reached 62.1  $\mu \text{mol g}^{-1}$  cellulose at 200 days. In comparison to treatments without bentonite, less nitrous oxide accumulated when bentonite present. This may be caused by either an enhancement of denitrification activity (enabling the complete conversion of nitrous oxide to nitrogen without accumulation of nitrous oxide) or suppression of denitrification activity.

### 6.0 SUMMARY

Gas generation from microbial degradation of a mixture of cellulosic waste was investigated. Cellulosic waste consisting of a mixture of filter paper, paper towels, and Kimwipes were incubated in the presence of WIPP brine with and without a mixed inoculum, nutrients, or nutrients plus excess nitrate and bentonite. Abiotic (control) samples were treated with formalin and showed no microbial activity. Nitrogen (anaerobic) or aircontaining (aerobic) samples with cellulose showed an increase in total gas, CO<sub>2</sub> and N<sub>2</sub>O when inoculated with a mixed inoculum without any added nutrients, or nutrients plus excess nitrate, and bentonite. In particular, samples which received nutrients plus excess nitrate




Figure 23. Nitrous oxide produced in anaerobic samples.




Figure 24. Nitrous oxide produced in anaerobic samples containing bentonite.

produced much more gas, CO<sub>2</sub>, and N<sub>2</sub>O than samples which did not. Bentonite increased the background level of CO<sub>2</sub> concentration due to abiotic reactions and it also appears that bentonite has a stimulatory effect on aerobic microbial activity.

Table 14 summarizes the rate and extent of gas production due to the presence of cellulose for the period between 69 and 200 days (131 days). Before 69 days, gas production in most treatments was not directly attributable to the presence of cellulose because the samples without cellulose also produced gas due to carry over of nutrients in the mixed inoculum, and metabolism of added nutrients. After 69 days, gas production in samples with cellulose exceeded those without cellulose. Gas production rates were calculated from the linear slope between 69 and 200 days. Negative values in the table denote a loss in gas volume. The negative rate reported for carbon dioxide in uninoculated treatments is the result of inactivity in these samples, and should be interpreted as "zero".

The total volume of gas produced in air-containing (aerobic) samples was highest in the presence of bentonite. Gas was produced at a rate of 0.028 mL g<sup>-1</sup> cellulose day<sup>-1</sup> in inoculated, nutrient-amended samples, and at 0.034 mL g<sup>-1</sup> cellulose day<sup>-1</sup> in inoculated, nutrient-amended samples containing excess nitrate. The highest amount of gas was produced in the presence of excess nitrate (6.07 mL g<sup>-1</sup> cellulose).

In anaerobic samples, gas production was highest in the absence of bentonite and was the lowest when it was present. Gas was produced at a rate of 0.021 mL g<sup>-1</sup> cellulose day<sup>-1</sup> in inoculated nutrient-amended samples, and 0.039 ml g<sup>-1</sup> cellulose day<sup>-1</sup> in inoculated nutrient amended samples containing excess nitrate.

The concentration of carbon dioxide was highest in aerobic treatments in the presence of bentonite. In samples containing excess nitrate,  $116 \mu mol CO_2 g^{-1}$  cellulose was produced at a rate of 0.869  $\mu mol g^{-1}$  cellulose day<sup>-1</sup>. Inoculated unamended samples of this treatment also showed the highest amount of carbon dioxide, with 21.5  $\mu mol g^{-1}$  cellulose produced over 200 days. In the absence of bentonite, aerobic samples that were inoculated with

Table 14. Summary of Rate and Net Gas Production in Samples Containing Cellulose

	Total Volum	e of Gas	Carbon Dio	oxide	Nitrous Ox	ride
Sample	Rato* (mi/g cell./doy)	Total Produced at 200 Days (ml/g cell.)	Rate* (µmol/g cell./day)	Total Produced at 200 Days (µmol/g cell.)	Rate* (umol/g cell./doy)	Total Produced at 200 Days (µmol/g cell.)
Aerobe						
Uninoculated	0.001	-0.18	-0.001	4.00		ND
Inoculated	-0.001	-0.34	0.033	8.30	<b></b>	ND
Inoc. + Nutrients	0.008	0.86	0.283	40.8	0.674****	24.4
Inoc. + Nut. + Nitr.	0.023***	4.42	0.484****	95.6	0.835	115
Aerobe + Bentonite						
Uninoculated	0.003	0.00	-0.016	2.32		0.048
Inoculated	0.001	-0.08	0.134	21.5		ND
Inoc. + Nutrients	0.028	4.38	0.533	69.8	1.00****	31.8****
Inoc. + Nut. + Nitr.	0.034	6.07	0.869	116	0.589	82.7
Anaerobe						
Uninoculated	-0.004	-0.09	-0.003	3.59		ND
Inoculated	0.003	0.59	0.016	5.47	••	ND
Inoc. + Nutrients	0.021	2.27	0.198	26.0	0.564****	15.8****
Inoc. + Nut. + Nitr.	0.039	5.44	0.422	61.4	0.602	79.0
Anaerobe + Bentonite						
Uninoculated	-0.003	-0.28	-0.005	0.22		ND
Inoculated	0.007	0.81	0.057	8.28		ND
Inoc. + Nutrients	0.013	1.92	0.236	31.8		ND
Inoc. + Nut. + Nitr.	0.025	3.52	0.252	35.0	0.647*****	62.1****

<sup>\*</sup> Rate calculated from 69 days (end of lag phase) to 200 days (131 days) except where noted; rate assumed linear and averaged over available data.

Negative values denote gas volume loss or decrease in concentration over time.

ND - not detected

<sup>\*\*</sup>For an assumed average drum of transuranic waste, with 10 kg of cellulosic material, the following rate conversion factors are applicable:

 $<sup>0.01 \</sup>text{ ml/g/day} = 1.6 \text{ moles/drum/year}$ 

<sup>1.0</sup>  $\mu$ mole/g/day = 3.7 moles/drum/year

<sup>\*\*\*</sup> Lag phase not present, gas production started at T=0

<sup>\*\*\*\*</sup> Nitrous oxide reached maximum at 104 days, rate is over 35 days

<sup>\*\*\*\*\*</sup> Nitrous oxide reached maximum concentration at 132 days, rate is over 32 days.

<sup>\*\*\*\*\*\*</sup> Lag phase lasted 104 days

nutrients produced 40.8  $\mu$ mol CO<sub>2</sub> g<sup>-1</sup> cellulose, whereas the samples containing excess nitrate produced 95.6  $\mu$ mol CO<sub>2</sub> g<sup>-1</sup> cellulose. Aerobic samples produced more carbon dioxide than anaerobic samples. In the absence of bentonite, anaerobic samples produced 0.422  $\mu$ mol CO<sub>2</sub> g<sup>-1</sup> cellulose day<sup>-1</sup> with a total yield 61.4  $\mu$ mol CO<sub>2</sub> g<sup>-1</sup> cellulose in the presence of excess nitrate. However, in the presence of bentonite, 35.0  $\mu$ mol g<sup>-1</sup> cellulose was produced in anaerobic samples containing excess nitrate. This is about one third of that produced in aerobic samples. Therefore, initially aerobic processes were more efficient in producing CO<sub>2</sub>.

In the presence of cellulose and nutrients, there is significantly greater amount of gas production than in samples without nutrients. In the absence of nutrients, microbial activity was minimal. Aerobic, denitrifying, and anaerobic primarily the fermentative activities were the predominant microbial processes noted to date.

Production of nitrous-oxide correlated with the presence of excess nitrate (1240  $\mu$ mol g<sup>-1</sup> cellulose), and 115  $\mu$ mol g<sup>-1</sup> cellulose was produced in aerobic samples without bentonite. Nitrous oxide did not accumulate in nutrient-amended samples which contained 0.1% nitrate (250  $\mu$ mol g<sup>-1</sup> cellulose); it quickly disappeared after about 30 days. Bentonite did not stimulate the accumulation of nitrous-oxide, but instead, was correlated with a lower accumulation of N<sub>2</sub>O.

The long-term inundated experiments showed enhanced halophilic bacterial activity in the presence of cellulose under aerobic and anaerobic conditions. Up to 200 days, gas production was highest in nutrient-amended including excess nitrate treatments containing an initial concentration of oxygen; this was enhanced by the addition of bentonite.

Table 15 presents gas production data for total gas and carbon dioxide scaled up to drum<sup>-1</sup> of waste year<sup>-1</sup>. Nitrous oxide is not presented because the production of this gas, while significant as a marker of microbial activity, is subject to conversion to nitrogen which affects the production rate. The rates drum<sup>-1</sup> of waste year<sup>-1</sup> were calculated on the basis

Table 15. Summary of Gas Production Rates Expressed as per g Cellulose/Day and Scaled up to per Drum/Year.

	Total Volume	Total Volume of Gas Car		rbon Dioxide	
Sample	Rate*	Calculated Amount Produced per Waste Drum**	Rate*	Calculated Amount Produced per Waste Drum**	
	(mi/g cell./day)	(moles/drum/year)	(µmol/g cell./day)	(moles/drum/year)	
Aerobe					
Uninoculated	0.001	0.16	-0.001		
Inoculated	-0.001	-0.16	0.033	0.12	
Inoc. + Nutrients	0.008	1.28	0.283	1.05	
Inoc. + Nut. + Nitr.	0.023***	3.68	0.484***	1.79	
Aerobe + Bentonite					
Uninoculated	0.003	0.48	-0.016		
Inoculated	0.001	0.16	0.134	0.5	
Inoc. + Nutrients	0.028	4.48	0.533	1.97	
Inoc. + Nut. + Nitr.	0.034	5.44	0.869	3.21	
Anaerobe					
Uninoculated	-0.004	-0.64	-0.003		
Inoculated	0.003	0.48	0.016	0.06	
Inoc. + Nutrients	0.021	3.36	0.198	0.73	
Inoc. + Nut. + Nitr.	0.039	6.24	0.422	1.56	
Anaerobe + Bentonite					
Uninoculated	-0.003	-0.48	-0.005		
Inoculated	0.007	1.12	0.057	0.21	
Inoc. + Nutrients	0.013	2.08	0.236	0.87	
Inoc. + Nut. + Nitr.	0.025	4.00	0.252	0.93	

<sup>\*</sup> Rate calculated from 69 days (end of lag phase) to 200 days (131 days) except where noted; rate assumed linear and averaged over available data.

Negative values denote gas volume loss or decrease in concentration over time

<sup>\*\*</sup>For an assumed average drum of transuranic waste, with 10 kg of cellulosic material, the following rate conversion factors were used:

 $<sup>0.01 \</sup>text{ ml/g/day} = 1.6 \text{ moles/drum/year}$ 

<sup>1.0</sup> µmole/g/day = 3.7 moles/drum/year

<sup>\*\*\*</sup> Lag phase not present, gas production started at T=0

an assumed average drum of transuranic waste with about 10 Kg of cellulosic materials. A total gas generation rate of 0.01 mL of gas g<sup>-1</sup> cellulose day<sup>-1</sup> corresponds to a gas generation rate of 1.6 mol of gas drum<sup>-1</sup> year<sup>-1</sup>. A carbon dioxide gas generation rate of 1.0  $\mu$ mol CO<sub>2</sub> g<sup>-1</sup> cellulose day<sup>-1</sup> corresponds to a gas generation rate of 1.6 mol gas drum<sup>-1</sup> year<sup>-1</sup>.

The data contained in this report is a summarization of work in progress, (a status report) and should not be interpreted as final values. Most of the long-term studies are still in progress. Gas production rates will undoubtedly be modified after long-term data, up to about two years or longer, are obtained and analyzed. The preliminary data included herein, and resultant gas production rates, should only be used for preliminary interpretations and tentative conclusions. Further data and interpretation from this microbial degradation-gas generation study will be documented in the future.

#### 7.0 REFERENCES

- Balderston, W.L., B. Sherr, and W.J. Payne. 1976. "Blockage by Acetylene of Nitrous Oxide Reduction in *Pseudomonas perfectomarinus*," <u>Applied Environmental Microbiology</u>. Vol. 31, No.4, 504-508.
- Barnhart, B.J., E.W. Campbell, E. Martinez, D.E. Caldwell, and R. Hallett. 1980. Potential Microbial Impact on Transuranic Wastes Under Conditions Expected in the Waste Isolation Pilot Plant (WIPP), Annual Report, October 1, 1978 September 30, 1979. LA-8297-PR. Los Alamos National, NM: Los Alamos Scientific Laboratory.
- Benoit, L., C. Cailliez, E. Petitdemange, and J. Gitton. 1992. "Isolation of Cellulolytic Mesophilic Clostridia from a Municipal Solid Waste Digestor," *Microbial Ecology*. Vol. 23, No.2, 117-125.
- Black, J., X. Tong, and D. Grbic-Galic. 1991. Laboratory Studies of Microbial Degradation of Transuranic (TRU) Waste, Fine Gress Report January 1, 1989 April 30, 1991. Stanford, CA: Environmenta eering and Science, Department of Civil Engineering, Stanford University. Copy on file in the Sandia WIPP Central Files, Waste Management and Transportation Library, Sandia National Laboratories, Albuquerque, NM).
- Bott, T.L., and L.A. Kaplan. 1991. "Selection of Surrogates for a Genetically Engineered Microorganism with Cellulolytic Capability for Ecological Studies in Streams," Canadian Journal of Microbiology. Vol. 37, No. 11, 848-857.
- Brush, L.H. 1990. Test Plan for Laboratory and Modeling Studies of Repository and Radionuclide Chemistry for the Waste Isolation Pilot Plant. SAND90-0266. Albuquerque, NM: Sandia National Laboratories.

- Brush, L.H. 1991. "Appendix A: Current Estimates of Gas Production Rates, Gas Production Potentials, and Expected Chemical Conditions Relevant to Radionuclide Chemistry for the Long-Term WIPP Performance Assessment," Preliminary Comparison with 40 CFR Part 191, Subpart B for the Waste Isolation Pilot Plant, December 1991. Vol. 3: Reference Data. WIPP Performance Assessment Division. Eds. R.P. Rechard, A.C. Peterson, J.D. Schreiber, H.J. Iuzzolino, M.S. Tierny, and J.S. Sandha. SAND91-0893/3. Albuquerque, NM: Sandia National Laboratories. A-25 through A-36.
- Brush, L.H., D. Grbic-Galic, D.T. Reed, X. Tong, R.H. Vreeland, and R.E. Westerman. 1990. "Preliminary Results of Laboratory Studies of Repository Chemistry for the Waste Isolation Pilot Plant," Scientific Basic for Nuclear Waste Management XIV, Boston, MA, November 26-29, 1990. Eds. T.A. Abrajano, Jr. and L.H. Johnson. SAND90-1031C. Pittsburgh, PA: Materials Research Society. Vol. 212, 893-900.
- Caldwell, D.E., R.C. Hallet, M.A. Molecke, E. Martinez, and B.J. Barnhart. 1988. Rates of CO<sub>2</sub> Production from the Microbial Degradation of Transuranic Wastes Under Simulated Geologic Isolation Conditions. SAND87-7170. Albuquerque, NM: Sandia National Laboratories.
- Christensen, S., and J.M. Tiedje. 1988. "Sub-Parts-Per-Billion Nitrate Method: Use of an N<sub>2</sub>O-Producing Denitrifier to Convert NO<sub>3</sub>- or <sup>15</sup>NO<sub>3</sub>- to N<sub>2</sub>O," Applied Environmental Microbiology. Vol. 54, No. 6, 1409-1413.
- Colberg, P.J., and L.Y. Young. "Biodegradation of Lignin-Derived Molecules Under Anaerobic Conditions," Canadian Journal of Microbiology. Vol. 28, No.7, 886-889.
- Coleman, A.W. 1980. "Enhanced Detection of Bacteria in Natural Environments by Fluorochrome Staining of DNA," *Limnology and Oceanography*. Vol. 25, No. 5, 948-951.

- DeLong, E.F., G.S. Wickham, and N.R. Pace. 1989. "Phylogenetic Stains: Ribosomal RNA-Bbased Probes for the Identification of Single Cells," *Science*. Vol. 243, No.4896, 1360-1363.
- Dicker, H.J., and D.W. Smith. "Effects of Organic Amendments on Sulfate Reduction Activity, H<sub>2</sub> Consumption, and H<sub>2</sub> Production in Salt Marsh Sediments," *Microbial Ecology*. Vol. 11, No. 4, 299-315.
- Francis, A.J., S. Dobbs, and R.F. Doering. 1980a. "Biogenesis of Tritiated and Carbon-14 Methane from Low-Level Radioactive Waste," *Nuclear Chemical Waste Management*. Vol. 1, No. 1, 153-159.
- Francis, A.J., S. Dobbs, and B.J. Nine. 1980b. "Microbial Activity of Trench Leachates from Shallow-Lland, Low-Level Radioactive Waste Disposal Sites," *Applied and Environmental Microbiology*. Vol. 40, No. 1, 108-113.
- Francis, A.J. 1985. Low-level Radioactive Wastes in Subsurface Soils," Soil Reclamation Processes: Microbiological Analyses and Applications. Eds. R.L. Tate III and D.A. Klein, New York, NY: Marcel Decker, Inc., 279-331.
- Francis, A.J., J.M. Slater, and C.J. Dodge. 1989. "Denitrification in Deep Subsurface Sediments," *Geomicrobiology Journal*. Vol. 7, No. 1-2, 103-116.
- Francis, A.J. 1990. "Microbial Dissolution and Stabilization of Toxic Metals and Radionuclides in Mixed Wastes," *Experientia*. Vol. 46, No. 8, 840-851.
- Francis, A.J., C.J. Dodge, J.B. Gillow, and J.E. Cline. 1991. "Microbial Transformations of Uranium in Wastes," *Radiochimica Acta*. Vol. 52/53, pt. 2, 311-316.

- Hope, C.F.A., and R.G. Burns. 1985. "The Barrier-Ring Plate Technique for Studying Extracellular Enzyme Diffusion and Microbial Growth in Model Soil Environments," *Journal of General Microbiology*. Vol. 131, pt. 5, 1237-1243.
- Hrmová, M., E. Petráková and P.Belly. 1991. "Induction of Cellulose- and Xylan-Degrading Enzyme Systems in *Aspergillus terreus* by Homo- and Hetero-Disaccharides Composed of Glucose and Xylose," *Journal of General Microbiology*. Vol. 137, pt. 3, 541-547.
- Javor, B.J. 1984. "Growth Potential of Halophilic Bacteria Isolated from Solar Salt Environments: Carbon Sources and Salt Requirements," Applied and Environmental Microbiology. Vol. 48, No.2, 352-360.
- Kearns, J.T., C.R. Clayton, G.P. Halala, J.B. Gillow, and A.J. Francis. 1992. "Application of XPS to the Study of MIC," *Materials Performance*. Vol 31, No. 10, 48-51.
- Kushner, D.J., and M. Kamekura. 1988. "Chapter 5: Physiology of Halophilic Eubacteria," Halophilic Bacteria. Ed. F. Rodriquez-Valera. Boca Raton, FL: CRC Press, Inc. Vol I, 110.
- Lappin, A.R., R.L. Hunter, D.P. Garber, and P.B. Davies, eds. 1989. Systems Analysis,

  Long-Term Radionuclide Transport, and Dose Assessments, Waste Isolation Pilot Plant
  (WIPP), Southeastern New Mexico; March 1989. SAND89-0462, Albuquerque, NM:
  Sandia National Laboratories.
- Laube, V.M., and S.M. Martin. 1981. "Conversion of Cellulose to Methane and Carbon Dioxide by Triculture of Acetovibrio cellulyticus, Desulfovibrio sp., and Methanosarcina barkeri,' Applied and Environmental Microbiology, Vol. 42, No. 3, 413-420.

- Lovely, D.R., E.J. Phillips, Y.A. Gorby, and E.R. Landa. 1992. "Microbial Reduction of Uranium," *Nature*. Vol. 350, 413-416.
- Lorowitz, W.H., D.P. Nagle, Jr., and R.S. Tanner. 1992. "Anaerobic Oxidation of Elemental Metals Coupled to Methanogenesis by *Methanobacterium thermoautotrophicum*," Environmental Science and Technology. Vol. 26, No. 8, 1606-1610.
- Lynd, L.R., H.E. Grethlein, and R.H. Wolkin. 1989. "Fermentation of Cellulosic Substrates in Batch and Continuous Culture by *Clostridium thermocellum*," *Applied and Environmental Microbiology*. Vol. 55, No. 12, 3131-3139.
- Molecke, M.A. 1979. Gas Generation from Transuranic Waste Degradation: Data Summary and Interpretation. SAND79-1245. Albuquerque, NM: Sandia National Laboratories.
- Molecke, M.A. 1983. A Comparison of Brines Relevant to Nuclear Waste Experimentation. SAND83-0516. Albuquerque, NM: Sandia National Laboratories.
- Ng, T.K., P.J. Weimer, and J.G. Zeikus. 1977. "Cellulolytic and Physiological Properties of Clostridium thermocellum," Archives Microbiology. Vol. 114, No. 1, 1-7.
- Pavlostathis, S.G., and E. Giraldo-Gomez. 1991. "Kinetics of Anaerobic Treatment: A Critical Review," Critical Reviews in Environmental Control. Vol. 21, No. 5-6, 411-490.
- Pavlostathis, S.G., T.L. Miller, and M.J. Wolin. 1988. "Fermentation of Insoluble Cellulose by Continuous Cultures of Ruminococcus albus," Applied and Environmental Microbiology. Vol. 54, No. 11, 2655-2659.

- Postgate, J.R. 1984. The Sulphate-Reducing Bacteria. 2nd ed. New York, NY: Cambridge University Press. 56-100.
- Poulsen, O.M., and L.W. Petersen. 1992. "Degradation of Microcrystalline Cellulose: Synergism Between Different Endoglucanases of Cellulomonas sp. ATCC 21399," Biotechnology and Bioengineering. Vol. 39, No.1, 121-123.
- Priest, F.G. 1984. Extracellular Enzymes. Wokingham Berkshire, England: Van Nostrad Reinhold (UK) Co. Ltd. 42-43.
- Rodriquez-Valera, F., ed. 1988. *Halophilic Bacteria*. Boca Raton, FL: CRC Press, Inc. Vol I, 72.
- Ross, H.N.M., M.D. Collins, B.J. Tindall, and W.D. Grant. 1981. "A rapid procedure for the Detection of Archaebacterial Lipids in Halophilic Bacteria," *Journal of General Microbiology*. Vol. 123, pt. 1, 75-80.
- Skujins, J. 1976. "Extracellular Enzymes in Soil," CRC Critical Reviews in Microbiology. Vol. 4, No. 4, 383-421.
- Tiedje, J.M. 1988. "Ecology of Denitrification and Dissimilatory Nitrate Reduction to Ammonium," *Biology of Anaerobic Microorganisms*. Ed. A.J.B. Zehnder, New York, NY: John Wiley and Sons. 179-186.
- Tomlinson, G.A., L.L Jahnke, and L.I. Hochstein. 1986. "Halobacterium denitrificans sp. nov., an Extremely Halophilic Denitrifying Bacterium," International Journal of Syst. Bacteriology. Vol. 36, No. 1, 66-70.

- Ventosa, A. 1988. "Chapter 3: Taxonomy of Moderately Halophilic Heterotrophic Eubacteria," *Halophilic Bateria*. Ed. F. Rodriquez-Valera. Boca Raton, FL: CRC Press, Inc. Vol. I, 72.
- Vreeland, R. et al., 1991. West Chester University, to be published.
- Wanner H., P. Wersin, and N. Sierro. 1992. Thermodynamic modelling of beutonite-ground interaction and implications for near field chemistry in a repository for spent fuel. SKB Technical Report 92-37. Stockholm, Sweden: Swedish Nuclear Fuel and Waste Management, Co. 1-6.
- Weimer, P.J., J.M. Lopez-Guisa, and A.D. French. 1990. "Effect of Cellulose Fine Structure on Kinetics of its Digestion by Mixed Ruminal Microorganisms in Vitro," Applied and Environmental Microbiology. Vol. 56, No. 8, 2421-2419.
- Yoshinari, T., and R. Knowles. 1976. "Acetylene Inhibition of Nitrous Oxide Reduction by Denitrifying Bacteria," *Biochemical Biophysical Research Communications*. Vol. 69, No. 3, 705-710.
- Zehnder, A.J., and W. Stumm. 1988. "Geochemistry and Biogeochemistry of Anaerobic Habitats," *Biology of Anaerobic Microorganisms*. Ed. A. J. Zehnder. New York, NY: John Wiley and Sons. 19.
- Zhilina, T.N., and G.A. Zavarzin. 1990. "Extremely Halophilic Methylotrophic, Anaerobic Bacteria," FEMS Microbiology Reviews. Vol. 87, No. 3-4, 315-322.

## APPENDIX A: DETAILS OF THE SHORT-TERM ACTIVITY MEASUREMENTS

## APPENDIX A: DETAILS OF THE SHORT-TERM ACTIVITY MEASUREMENTS

The short-term experiments were developed to determine the activity of specific groups of organisms (aerobes, anaerobes, and denitrifiers) by using WIPP salt from the underground and Nash Draw brine as a basal medium and inoculum.

A concentrated stock solution of nutrients (20x) was prepared and 0.5mL dispensed into 20-mL serum bottles. Medium for anaerobes (glucose fermenters) and denitrifiers was dispensed and sealed inside a nitrogen-filled glove box. Medium for aerobes was prepared outside of the glove box in air. The samples were capped with butyl rubber stoppers and aluminum crimps and autoclaved (120°C, 20 psi, 15 minutes).

The nutrient solution was added prior to the addition of inoculum to achieve a final concentration of nutrients in the samples as follows:

### Aerobe and Anaerobe Series

glucose	5.0 g/L
yeast extract	0.5 g/L
potassium phosphate	1.0 g/L
ammonium nitrate	1.0 g/L
inoculum	9.5 mL/bottle
pH=6.8	

## Denitrifier Series

sodium succinate	5.0 g/L
yeast extract	0.5 g/L
potassium phosphate (dibasic)	1.0 g/L
ammonium nitrate	1.0 g/L
potassium nitrate	1.0 g/L
inoculum	9.5 mL/bottle
pH = 6.8	

To determine denitrification, 2 mL of acetylene was injected into the headspace and nitrous oxide production was determined by gas chromatography.

Cellulose degradation was investigated by replacing the carbon source with 0.5 g Whatman #1 filter paper.

A large volume of inoculum was prepared by dissolving 200 g of S2180, W30 muck pile salt from the WIPP underground workings into 1 L of sterile water. A 450 mL aliquot was poured into a sterile beaker and 10 mL of Laguna Cinco mud slurry and 50 mL of Laguna Cinco brine from Nash Draw were added. This was done inside the anaerobic glove box. A total of 510 mL of inoculum was prepared; 9.5 mL of this inoculum was added to each bottle of sterile nutrient medium through a sterile needle and syringe to attain a final volume of 10 mL in each bottle. The pressure was equalized after the addition of inoculum with a sterile syringe and 0.22  $\mu$ m filter.

A total of 6 samples were prepared per series. Two of the 6 were treated with 1 mL of 10 % formalin to serve as a control. Six samples were also prepared without nutrient additions, 2 treated with formalin.

Samples were incubated at 30°C and analyzed for gas production (total gas, carbon dioxide and nitrous oxide) at specific time periods.

## **APPENDIX B: GAS ANALYSIS**

## APPENDIX B: GAS ANALYSIS

## **Total Gas**

Headspace pressure was measured with a Wallace & Tiernan® digital pressure model 661-D/A035 gauge calibrated to National Institute of Standards and Technology (NIST) standards.

## Carbon Dioxide

Carbon dioxide was analyzed using a gas chromatograph equipped with a thermal conductivity detector. Instrument conditions are listed below:

Column	SS 12' x 1/8" Porapak QS
Column temp (°C)	100
Carrier gas	He
Carrier flow (mL/min)	35
Injector temp (°C)	150
Manifold temp (°C)	210
Detector temp (°C)	250
Detector current (mA)	225
Detection limit (nmol/mL)	1

Instrument calibrated with gas standards traceable to NIST.

## **Nitrous Oxide**

Nitrous oxide was analyzed using a gas chromatograph equipped with an electron capture detector (63Ni). Instrument conditions are listed below:

Column	SS 12' x 1/8" Porapak QS
Column temp (°C)	70
Carrier gas	$N_2$
Carrier flow (mL/min)	30
Injector temp (°C)	270
Detector temp (°C)	270
Detector current (nA)	2
Detection limit (pmol/mL)	100

Instrument calibrated with gas standards traceable to NIST.

## APPENDIX C: DETAILS OF THE LONG-TERM EXPERIMENT

### APPENDIX C: DETAILS OF THE LONG-TERM EXPERIMENT

The following is a detailed description of the samples prepared for the long-term inundated experiment, including the number and chemical composition of treatments.

## Sample Preparation

Detergent (Alconox®) and acid-washed (10% HCl) 160 mL serum bottles were rinsed with deionized water. They were then dried in a drying oven, covered with aluminum foil and autoclaved, thus completing preparation of the bottles.

Four paper types were used for the experiment:

Whatman #1 filter paper Brown paper towel White paper towel Kimwipes

The papers were reduced to approximately 1 cm by 1 cm squares.

The bottles were filled with the processed paper. The papers were mixed together prior to filling and 1.25 g of each paper type were added to each bottle for a total of 5 grams of paper per bottle.

## **Anaerobic Samples**

Ten liters of G-Seep #9 were removed from storage at 4°C and equilibrated overnight to room temperature. Storage at 4°C was necessary in order to prevent microbial activity in the storage containers; which could possibly pre-enrich the brine with specific microbes.

Sixty filled bottles (w/paper) were arranged and prepared for treatment as follows:

36 w/paper: unamended (G-Seep w/o additions)

18 w/o bentonite

18 w/ bentonite

12 w/paper: amended (G-Seep w/ nutrients)

6 w/o bentonite

6 w/ bentonite

12 w/paper: amended w/ excess nitrate (G-Seep w/ nutrients and excess nitrate)

6 w/o bentonite

6 w/ bentonite

Thirty-two bottles w/o paper were arranged and prepared for treatment as follows:

16 unamended (G-Seep w/o additions)

8 w/o bentonite

8 w/ bentonite

8 amended (G-Seep w/ nutrients)

4 w/o bentonite

4 w/ bentonite

8 amended w/ excess nitrate (G-Seep

w/ nutrients and excess nitrate)

4 w/o bentonite

4 w/ bentonite

A total of 92 bottles were prepared for anaerobic treatment.

## **Aerobic Samples**

Sixty filled bottles (w/ paper) were arranged and prepared for treatment as follows:

36 w/paper: unamended (G-Seep w/o additions)

18 w/o bentonite

18 w/ bentonite

12 w/paper: amended (G-Seep w/ nutrients)

6 w/o bentonite

6 w/ bentonite

12 w/paper: amended w/ excess nitrate (G-Seep w/ nutrients and

excess nitrate)

6 w/o bentonite

6 w/ bentonite

Thirty-two bottles w/o paper were arranged and prepared for treatment as follows:

16 unamended (G-Seep w/o additions)

8 w/o bentonite

8 w/ bentonite

8 amended (G-Seep w/ nutrients)

4 w/o bentonite

4 w/ bentonite

## 8 amended w/ excess nitrate (G-Seep w/ nutrients and excess nitrate) 4 w/o bentonite

4 w/ bentonite

A total of 92 bottles were prepared for aerobic treatment.

## **Nutrient Additions**

The following lists the quantities added and final concentrations of nutrients in the samples. This list applies to both the aerobic and anaerobic samples with and without paper.

(a) Amended: The following quantities of nutrients were used for the amended treatments:

	grams/L	$\mathbf{m}\mathbf{M}$
Ammonium nitrate	1	12.5
Potassium phosphate		
(dibasic)	1	7.35
Yeast extract (0.05%)	0.5	

Each 100 mL sample contained the following final concentration of nutrients:

	grams/100mL	μmoles
Ammonium nitrate	0.1	1250
Potassium phosphate (dibasic)	0.1	735
Yeast extract (0.05%)	0.05	

(b) Amended and excess nitrate added: Potassium nitrate was added in addition to the ammonium nitrate:

	grams/L	$\mathbf{m}\mathbf{M}$
Ammonium nitrate	1	12.5
Potassium phosphate		
(dibasic)	1	7.35
Potassium nitrate	5	49.5
Yeast extract (0.05%)	0.5	

Each 100 mL sample contained the following final concentration of nutrients:

	grams/100mL	μmoles
Ammonium nitrate Potassium phosphate	0.1	1250
(dibasic)	0.1	735
Potassium nitrate	0.5	4950
Yeast extract (0.05%)	0.05	

(c) <u>Glucose added</u> (instead of paper): Samples were prepared without paper with a glucose addition to determine the ability of the inoculum to grow in the amended samples. The treatment was composed of the following:

	grams/L	mM
Glucose	5	27.7
Ammonium nitrate	1	12.5
Potassium phospate		
(dibasic)	1	7.35
Yeast extract (0.05%)	0.5	

Each 100 mL sample contains the following quantity of nutrients:

	grams/100mL	μmoles
Glucose	0.5	2770
Ammonium nitrate Potassium phosphate	0.1	1250
(dibasic)	0.1	735
Yeast extract (0.05%)	0.05	

(d) Glucose added (instead of paper) and excess nitrate: Samples were prepared without paper with a glucose addition and additional nitrate. The treatment was composed of the following:

	grams/L	mM
Glucose	5	27.7
Ammonium nitrate	1	12.5
Potassium phospate (dibasic)	1	7.35
Potassium nitrate	5	49.5
Yeast extract (0.05%)	0.5	***

Each 100 mL sample contained the following final concentration of nutrients:

	grams/100mL	umoles
Glucose	0.5	2770
Ammonium nitrate	0.1	1250
Potassium phosphate		
(dibasic)	0.1	735
Potassium nitrate	0.5	4950
Yeast extract (0.05%)	0.05	

## Sample Volumes

The final sample volume (displacement of liquid plus cellulose or liquid w/o cellulose) and the headspace volume of each treatment is as follows:

Cellulose Treatments	sample vol. (mL	hdsp. vol (mL*
U (uninoculated) I (inoculated) UC (uninoculated	110 114	50 46
control)	113	47
IC (inoculated control)	117	43
"No paper" Treatments	sample vol. (mL	hdsp. vol (mL*
NU (uninoculated) NI (inoculated)	sample vol. (mL 100 104	hdsp. vol (mL* 60 56
NU (uninoculated)	100	60

<sup>\*</sup>Headspace vol. (hdsp. vol) calculated by subtracting sample volume from volume of bottle (160ml).

Acetylene was not injected into the headspace of any of the samples in order to prevent pertubation of the headspace gas.

# APPENDIX D: GAS PRODUCTION DATA (GROSS AND NET) FOR THE LONG-TERM INUNDATED EXPERIMENT

## APPENDIX D: GAS PRODUCTION DATA (GROSS AND NET) FOR THE LONG-TERM INUNDATED EXPERIMENT

#### i. Data Reduction

Total gas, carbon dioxide, and nitrous oxide production by aerobic and anaerobic samples up to 200 days incubation is presented in Appendix D, Tables 1-12, on a per sample basis. This data was used to prepare Tables 13-24 in which total gas, carbon dioxide, and nitrous oxide production in aerobic and anaerobic treatments is presented on a per gram cellulose basis. Figures 13-24 are based on the data from Tables 13-24. The data in tables 13-24 have been corrected for gas production in the absence of cellulose by subtracting measured gas values in respective treatments prepared without cellulose. As an example, total gas production in aerobic samples was corrected (Table 13) by subtracting total gas production in treatments without cellulose (designated "NC", Appendix D, Table 1) from samples with cellulose (designated "C") at each time period. The resultant number was divided by the total amount of cellulose in each sample bottle (5 grams) in order to arrive at a value that represents the total gas produced per gram of cellulose. Tables 13 to 24 represent the gas produced strictly due to the presence of cellulose in the samples, and are corrected for gas produced due to metabolism of dissolved organic carbon present in the brine or in the nutrient addition as measured in specific control treatments. Carbon dioxide was produced in certain samples in the absence of cellulose, specifically prior to 69 days. After this period, gas production reported is that which is above and beyond the control treatments. Significant quantities of N<sub>2</sub>O were not detected in the absence of cellulose. Gas produced at time 0 was not subtracted from later values, as may seem obvious in order to normalize the starting values at 0. This was not done because time 0 measurements were taken 3 days after sample preparation, therefore gas present at time 0 is due to evolution of dissolved gases and headspace equilibration. These processes contribute to the overall gas production.

Figures 13 to 24 present data from Table 13 to 24. All data in these figures are presented as gas produced per gram cellulose, and have been corrected for gas production in the absence of

cellulose. The data plotted are the mean values of three samples and error bars represent the standard error of the mean. In the legend: "uninoculated" refers to samples that have received no mixed inoculum or nutrients; "inoculated" refers to samples that have received mixed inoculum but no nutrients; "inoculated + nutrients" refers to samples that have received both mixed inoculum and nutrients (amended), and; "inoculated + nutrients + nitrate" refers to samples that have received (i) mixed inoculum, (ii) nutrients, and (iii) excess nitrate in the form of potassium nitrate (amended plus excess nitrate).

Data for treatments prepared with glucose to determine activity of the mixed inoculum are presented in Figures 25-30. Data for gas analysis of these samples, along with formalin treated controls for all cellulose and no-cellulose treatments, are presented in Appendix D, Tables 1(a) to 4(c).

Analysis of the samples will continue at select intervals (as determined by gas producing activity) past 200 days and up to 800 days. The analyses will then stop if activity ceases, as indicated by cessation of gas production, or be prolonged if gas producing activity continues in the samples. If this occurs in select samples only then these samples will be reserved for long term monitoring, and analyses of inactive samples will stop.

Table 1. Gross Data for the Long-Term Inundated Experiment: Total Volume of Gas Produced in Aerobic Samples.

Treatments*	Sample	Volume of Gas Produced (ml)														
[Brine]	Designation							ubation Time (	Days)							
			0		45		69	104		13	32		164	-	200	
Unamended/Uninoculated (1)																
Formalin treated, w/o cellulose	1(NC)-f	4.22	± 0.13	-0.35	± 0.13	-0.06	± 0.45	-0.26 ± 0.06	-0.13	±	0.13		NA	-1.44	± 0.0	
Formalin treated, with cellulose	1(C)-f	5.44	± 0.32	4.09	± 0.16	4.19	± 0.10	2.08 ± 1.06	1.37	±	0.93		NA	0.67	± 0.9	
As is, w/o cellulose	1(NC)-a	4.39	± 0.03	3.71	± 0.03	3.84	± 0.03	1.90 ± 0.61	1.33	±	0.85		NA	-0.61	± 0.7	
As is, with cellulose	1(C)-a	4.63	± 0.10	2.28	± 0.95	2.04	± 0.92	$0.34 \pm 0.68$	0.34	±	0.61		NA	-1.53	± 0.5	
Inamended/Inoculated (2)													***			
Formalin treated, w/o cellulose	2(NC)-f	4.96	± 0.02	2.88	± 0.04	3.50	± 0.00	2.49 ± 0.04	2.45	±	0.14		NA	0.07	± 0.2	
Formalin treated, with cellulose	2(C)-f	4.89	± 0.06	2.72	± 0.38	1.76	± 0.56	0.91 ± 0.41	0.41	±	0.53		NA		± 0.4	
As is, w/o cellulose	2(NC)-a	3.56	± 0.02	2.90	± 0.15	2.21	± 0.30	$0.53 \pm 0.11$	-0.91	±	0.04		NA	-2.86	± 0.0	
As is, with cellulose	2(C)-a	3.16	± 0.03	2.97	± 0.22	1.44	± 0.16	-0.44 ± 0.22	-1.28	±	0.25		NA	-4.57	± 0.3	
Amended/Inoculated (3)													, <u>, , , , , , , , , , , , , , , , , , </u>			
As is, w/o cellulose	3(NC)-a	3.20	± 0.04	1.37	± 0.04	0.30	± 0.00	0.11 ± 0.04	-1.49	±	0.15	-2.48	± 0.38	-4.53	± 0.4	
as is, with cellulose	3(C)-a		± 0.16		± 0.69		$\pm$ 0.31	1.91 ± 1.53	1.44	±	1.19	0.34	± 0.63	-0.22	± 0.4	
as is, with glucose	3(G)-a	3.73	± 0.08	2.29	± 0.15	1.83	± 0.38	$-1.79 \pm 0.34$	-1.22	±	1.37	1.87	± 2.63	-1.83	± 1.30	
excess nitrate, w/o cellulose	3(NC)-x		± 0.08		± 0.46		± 0.08	-0.78 ± 0.30	-1.14	±	0.04		± 0.18		± 0.2	
excess nitrate, with cellulose	3(C)-x		± 0.06		± 0.72		± 3.04	$11.9 \pm 4.4$	15.5	±	5.3	18.3	± 6.1		± 4.0	
Excess nitrate, with glucose	3(G)-x	3.52	$\pm 0.02$	2.74	$\pm 0.50$	0.88	± 0.76	$-1.49 \pm 0.27$	-1.30	±	0.04	-0.86	± 0.72	2.90	± 2.02	

Table 2. Gross Data for the Long-Term Inundated Experiment: Production of Carbon Dioxide\* in Aerobic Samples

Treatments	Sample			Ca	rbon Dioxide	(µmoles/sample)		
[Brine]	Designation				Incubation Ti	me (Days)		
		0	45	69	104	132	164	200
Unamended/Uninoculated (1)								
Formalin treated, w/o cellulose	1(NC)-f	2.59 ± 0.01	2.50 ± 0.01	$3.40 \pm 0.40$	$3.32 \pm 0.00$	3.61 ± 0.00	NA	3.68 ± 0.0
Formalin treated, with cellulose	1(C)-f	$30.5 \pm 2.7$	40.8 ± 0.1	$44.4 \pm 0.2$	$40.6 \pm 0.8$	$41.8 \pm 0.3$	NA	$41.8 \pm 0.5$
As is, w/o cellulose	1(NC)-a	1.38 ± 0.01	1.49 ± 0.01	$0.94 \pm 0.00$	1.66 ± 0.01	1.83 ± 0.06	NA	2.18 ± 0.0
As is, with cellulose	1(C)-a	$13.8 \pm 0.5$	$21.1 \pm 0.3$	22.0 ± 0.1	21.3 ± 0.1	23.1 ± 0.1	NA	$22.2 \pm 0.1$
Unamended/Inoculated (2)								
Formalin treated, w/o cellulose	2(NC)-f	4.97 ± 0.45	4.38 ± 0.29	4.70 ± 0.11	4.88 ± 0.3	5.02 ± 0.30	NA	4.94 ± 0.27
Formalin treated, with cellulose	2(C)-f	$34.5 \pm 0.8$	$35.6 \pm 0.0$	$38.7 \pm 0.7$	$34.3 \pm 0.6$	$35.2 \pm 0.1$	NA	33.2 ± 1.1
As is, w/o cellulose	2(NC)-a	$2.30 \pm 0.02$	2.97 ± 0.03	2.92 ± 0.09	5.88 ± 0.08	6.56 ± 0.11	NA	7.01 ± 0.11
As is, with cellulose	2(C)-a	12.1 ± 0.3	19.7 ± 0.8	22.6 ± 0.9	$30.8 \pm 0.9$	40.9 ± 1.0	NA	48.5 ± 1.4
Amended/Inoculated (3)								
As is, w/o cellulose	3(NC)-a	2.87 ± 0.03	27.8 ± 1.2	72.9 ± 2.6	113 ± 5	130 ± 2	150 ± 1	142 ± 1
As is, with cellulose	3(C)-a	$2.80 \pm 0.10$	$50.9 \pm 1.4$	$91.8 \pm 5.9$	$215 \pm 37$	$278 \pm 25$	$333 \pm 21$	$346 \pm 27$
As is, with glucose	3(G)-a	$2.41 \pm 0.03$	$4.03 \pm 0.13$	$30.8 \pm 2.1$	94.9 ± 7.5	167 ± 32	$325 \pm 108$	323 ± 95
Excess nitrate, w/o cellulose	3(NC)-x	2.79 ± 0.03	17.2 ± 1.9	48.9 ± 0.2	92.2 ± 0.59	115 ± 3	131 ± 2	126 ± 2
Excess nitrate, with cellulose	3(C)-x	$2.60 \pm 0.10$	$51.6 \pm 0.0$	$210 \pm 21$	$399 \pm 18$	$533 \pm 13$	$612 \pm 20$	$604 \pm 30$
Excess nitrate, with glucose	3(G)-x	$2.51 \pm 0.01$	$4.05 \pm 0.25$	$24.1 \pm 5.9$	$72.0 \pm 7.7$	$121 \pm 4$	$211 \pm 6$	296 ± 20

<sup>\*</sup>Dissolved carbon dioxide concentrations not included.

Table 3. Gross Data for the Long-Term Inundated Experiment: Production of Nitrous Oxide\* in Aerobic Samples

Treatments	Sample			Nitrous Oxide (µmoles/sample) Incubation Time (Days)														
[Brine]	Designation		_							Incuba		•	Days)					
			0	69		104		132			164			200				
Unamended/Uninoculated (1)																		
Formalin treated, w/o cellulose	1 (NC)-f	0.000	£ 0.000	0.003	±	0.002	0.000	±	0.000	0.000	±	0.000		NA	ı	0.001	±	0.001
Formalin treated, with cellulose	1(C)-f	0.000 :	£ 0.000	0.003	±	0.001	0.003	±	0.001	0.013	±	0.001		NA	i	0.005	±	0.002
As is, w/o cellulose	1(NC)-a	0.000	e 0.000	0.162	±	0.110	0.000	±	0.000	0.000	±	0.000		NA		0.051	±	0.035
As is, with cellulose	1(C)-a	0.000	₺ 0.000	0.003	±	0.001	0.006	±	0.001	0.015	±	0.001		NA		0.008	±	0.000
Unamended/Inoculated (2)																		
Formalin treated, w/o cellulose	2(NC)-f	0.000	0.000	0.025	±	0.018	0.000	±	0.000	0.000	±	0.000		NA		0.000	±	0.000
Formalin treated, with cellulose	2(C)-f	0.000	0.000	0.001	±	0.000	0.002	±	0.000	0.014	±	0.002		NA		0.004	±	0.000
As is, w/o cellulose	2(NC)-a	0.003	0.000	0.001	±	0.001	0.000	±	0.000	0.016	±	0.011		NA		0.062	±	0.007
As is, with cellulose	2(C)-a	0.014	0.000	0.002	±	0.001	0.000	±	0.000	0.000	±	0.000		NA		0.009	±	0.006
Amended/Inoculated (3)																		
As is, w/o cellulose	3(NC)-a	0.000 ±	0.000	0.061	±	0.043	0.087	±	0.061	0.024	±	0.003	0.000	±	0.000	0.050	±	0.033
As is, with cellulose	3(C)-a	0.004 ±	0.000	0.182	± 1	0.147	118	±	48	122	±	39	8.81	±	5.98	29.2	±	25.2
As is, with glucose	3(G)-a	0.000 ±	0.000	0.056	±	0.010	0.252	±	0.122	5.64	±	3.40	114	±	80	105	±	74
Excess nitrate, w/o cellulose	3(NC)-x	0.000 ±		0.001	<u>`</u>	0.000	0.004	±	0.002	0.006	±	0.004	0.006	±	0.001	0.023	±	0.006
excess nitrate, with cellulose	3(C)-x	$0.000 \pm$			±	8.8	245	±	11	326	±	18	484	_	67	577	±	129
Excess nitrate, with glucose	3(G)-x	$0.000 \pm$	0.000	0.000	± (	0.000	2.97	±	2.10	2.99	±	2.08	11.2	±	7.3	66.0	±	41.2

<sup>\*</sup>Dissolved nitrous oxide concentrations not included.

Table 4. Gross Data for the Long-Term Inundated Experiment: Total Volume of Gas Produced in Aerobic Samples

Treatments*	Sample		Volume of Gas Produced (ml)														
[Brine/Bentonite]	Designation						Inc	ubation	Time (Da	ys)							
	<del></del>		0		45	_	69		104		132			164		200	
Unamended/Uninoculated (4)																	
Formalin treated, w/o cellulose	4(NC)-f	5.93	± 0.04	1.24	± 2.02	1.43	± 1.74	1.09	± 1.05	-0.31	+	0.00		NA	-2.17	± 0.0	
Formalin treated, with cellulose	4(C)-f	6.59	± 0.16	4.38	± 0.13		± 0.13		± 0.13	0.10	±	0.03		NA		± 0.0	
As is, w/o cellulose	4(NC)-a	5.43	± 0.04	3.63	± 0.53	4.78	± 0.78	2.65	± 0.45	1.92	±	0.20		NA	-0.82	± 0.2	
As is, with cellulose	4(C)-a	5.17	± 0.10	3.74	± 0.10	2.99	± 0.20	1.67	± 0.24	1.26	±	0.20		NA		± 0.2	
Inamended/Inoculated (5)																	
Formalin treated, w/o cellulose	5(NC)-f	5.08	± 0.36	2.56	± 0.83	1.19	± 1.73	0.76	± 0.94	0.54	±	0.65		NA	-2.02	± 0.30	
Formalin treated, with cellulose	5(C)-f	5.65	± 0.12	3.63	± 0.03	2.72	± 0.00	2.22	± 0.00	0.88	±	0.15		NA		± 0.12	
As is, w/o cellulose .	5(NC)-a		± 0.30	3.62	± 0.04	2.78	± 0.08	1.64	± 0.15	1.14	±	0.11		NA	-2.06	± 0.34	
As is, with cellulose	5(C)-a	3.79	± 0.06	3.13	± 0.09	1.88	± 0.19	0.16	± 0.66	-1.94	±	0.47		NA	-2.47	± 0.78	
Amended/Inoculated (6)																	
As is, w/o cellulose	6(NC)-a		± 0.04	0.84	± 0.84	1.52	± 0.84	1.56	± 0.80	0.11	±	0.53	-1.49	± 0.34	-3.47	± 0.15	
s is, with cellulose	6(C)-a		± 0.16	3.00	± 0.06	5.07	± 0.75	10.7	± 1.8	14.9	±	1.5	18.9	± 1.1	18.4	± 1.0	
s is, with glucose	6(G)-a	1.60	± 0.04	1.10	± 0.69	0.69	± 0.50	-0.34	± 0.19	-1.07	±	0.27	-1.62	± 0.19	-1.52	± 0.61	
xcess nitrate, w/o cellulose	6(NC)-x		± 0.08		± 0.04		± 0.08		± 0.00		±	0.08	-1.30	± 0.04	-3.05	± 0.08	
xcess nitrate, with cellulose			± 0.06		± 0.09		± 0.09		± 1.2	18.7	±	1.5	24.4	± 0.9	27.3	± 0.2	
xcess nitrate, with glucose	6(G)-x	2.29	± 0.23	2.51	± 1.71	0.69	± 0.57	-0.38	± 0.5	-0.53	±	0.11	-1.49	± 0.15	-1.94	± 0.69	

Table 5. Gross Data for the Long-Term Inundated Experiment: Production of Carbon Dioxide\* in Aerobic Samples

Treatments	Sample			Ca	rbon Dioxide	(µmoles/sample)		
[Brine/Bentonite]	Designation				Incubation Ti	me (Days)		
		0	45	69	104	132	164	200
Unamended/Uninoculated (4)								
Formalin treated, w/o cellulose	4(NC)-f	50.3 ± 0.4	88.8 ± 1.0	98.8 ± 0.4	101 ± 1	115 ± 2	NA	121 ± 4
Formalin treated, with cellulose	4(C)-f	71.1 ± 1.2	$95.3 \pm 0.2$	$110 \pm 0$	$114 \pm 0$	127 ± 1	NA	137 ± 1
As is, w/o cellulose	4(NC)-a	17.7 ± 0.3	$38.8 \pm 0.4$	44.5 ± 0.4	40.2 ± 0.3	42.7 ± 0.2	NA	40.0 ± 0.1
As is, with cellulose	4(C)-a	25.3 ± 1.5	$47.6 \pm 0.3$	66.9 ± 11.8	49 ± 0.7	51.8 ± 0.5	NA	51.6 ± 0.1
Unamended/Inoculated (5)								
Formalin treated, w/o cellulose	5(NC)-f	37.3 ± 10.1	59.6 ± 15.7	66.1 ± 18.4	77.4 ± 18.1	76.0 ± 26.5	NA	79.0 ± 33.0
Formalin treated, with cellulose	5(C)-f	61.9 ± 0.9	$83.5 \pm 0.0$	$94.0 \pm 0.2$	$93.2 \pm 0.6$	$100 \pm 0$	NA	$102 \pm 0$
As is, w/o cellulose	5(NC)-a	13.5 ± 2.9	36.7 ± 0.7	41.3 ± 0.2	38.7 ± 0.7	41.4 ± 0.7	NA	37.6 ± 0.7
As is, with cellulose	5(C)-a	23.7 ± 0.2	43.6 ± 0.4	$61.3 \pm 4.0$	80.3 ± 2.1	101 ± 3	NA	145 ± 6
Amended/Inoculated (6)								
As is, w/o cellulose	6(NC)-a	11.3 ± 0.1	103 ± 3	169 ± 3	185 ± 2	201 ± 2	203 ± 0	181 ± 1
As is, with cellulose	6(C)-a	$8.60 \pm 0.00$	$72.4 \pm 1.0$	$156 \pm 5$	$247 \pm 11$	358 ± 19	$492 \pm 6$	$530 \pm 6$
As is, with glucose	6(G)-a	8.50 ± 2.80	67.1 ± 0.2	73.1 ± 0.4	72.9 ± 0.2	$75.8 \pm 0.4$	79.5 ± 1.0	75.7 ± 1.7
Excess nitrate, w/o cellulose	6(NC)-x	$10.5 \pm 0.4$	84.6 ± 0.0	153 ± 1	171 ± 3	$139 \pm 37$	184 ± 4	159 ± 4
Excess nitrate, with cellulose	6(C)-x	$8.9 \pm 0.1$	$68.0 \pm 1.2$	$164 \pm 3$	$307 \pm 23$	499 ± 75	711 ± 46	$741 \pm 30$
Excess nitrate, with glucose	6(G)-x	$9.40 \pm 0.90$	$65.0 \pm 2.0$	$71.4 \pm 1.2$	$71.2 \pm 0.6$	$76.5 \pm 0.2$	$80.1 \pm 1.5$	$81.1 \pm 5.0$

<sup>\*</sup>Dissolved carbon dioxide concentrations not included.

Table 6. Gross Data for the Long-Term Inundated Experiment: Production of Nitrous Oxide\* in Aerobic Samples

Treatments	Sample			<del> </del>	<del></del>			N	itrous Ox			es/samp	le)				
[Brine/Bentonite]	Designation	0		6	9		10	4	incuba	13	Time (D	ays) 	16	54		20	0
Unamended/Uninoculated (4)																	
Formalin treated, w/o cellulose	4(NC)-f	0.000 ±	0.000	0.002	0.002	0.213	±	0.146	0.010	±	0.007		NA	\	0.000	±	0.000
Formalin treated, with cellulose	4(C)-f	0.000 ±	0.000	0.008	0.002	0.001	±	0.001	0.019	±	0.000		NA	١	0.006	<b>±</b> .	0.001
As is, w/o cellulose	4(NC)-a	0.000 ±	0.000	0.006	0.001	0.003	±	0.002	0.017	±	0.000		NA	\	0.005	±	0.003
As is, with cellulose	4(C)-a	0.000 ±	0.000	0.731	0.489	0.188	±	0.054	0.222	±	0.070		NA	١	0.245	±	0.101
Inamended/Inoculated (5)																	
Formalin treated, w/o cellulose	5(NC)-f	0.000 ±	0.000	0.070	0.049	0.009	±	0.002	0.166	±	0.106		NA	١	0.076	±	0.049
Formalin treated, with cellulose	5(C)-f	0.000 ±	0.000	0.012 ±	0.001	0.011	±	0.002	0.025	±	0.001		NA	1	0.009	±	0.002
As is, w/o cellulose	5(NC)-a	0.000 ±	0.000	0.038	0.024	0.195	±	0.065	0.210	±	0.066		NA	\	0.161	±	0.050
As is, with cellulose	5(C)-a	0.000 ±	0.000	0.010	0.006	0.000	±	0.000	0.000	±	0.000		NA	1	0.064	±	0.027
Amended/Inoculated (6)																	
As is, w/o cellulose	6(NC)-a	0.000 ±	0.000	0.002 ±	0.001	1.21	±	0.36	6.06	±	4.21	4.96	±	3.51	4.27	±	2.55
As is, with cellulose	6(C)-a	0.000 ±	0.000	31.6 ±		160	±	53	132	±	8	0.528	±	0.459	0.000	±	0.000
as is, with glucose	6(G)-a	0.000 ±	0.000	0.003 ±	0.000	0.003	±	0.000	0.028	±	0.001	0.007	±	0.001	0.015	±	0.004
xcess nitrate, w/o cellulose	6(NC)-x	0.000 ±	0.000	0.000 ±	0.000	14.6	±	2.7	13.9	±	5.2	13.8	±	5.4	11.6	±	4.9
xcess nitrate, with cellulose	6(C)-x	$0.000 \pm$	0.000	27.1 ±		142		22	217	±	41	354	±	41	425	±	35
excess nitrate, with glucose	6(G)-x	$0.000 \pm$	0.000	0.005 ±	0.001	0.010	±	0.001	0.022	+	0.003	0.092	±	0.063	0.011	±	0.001

<sup>\*</sup>Dissolved nitrous oxide concentrations not included.

Table 7. Gross Data for the Long-Term Inundated Experiment: Total Volume of Gas Produced in Anaerobic Samples

Treatments*	Sample								as Produc							
[Brine]	Designation							ubation	Time (Da	ys)						
			0		45		69		104		13	32		164		200
Unamended/Uninoculated (7)																
Formalin treated, w/o cellulose	7(NC)-f	2.99	± 0.10	2.65	± 0.10	2.01	± 0.03	1.09	± 0.06	1.66	±	0.00		NA	0.06	± 0.0
Formalin treated, with cellulose	7(C)-f	4.28	± 0.06	3.49	± 0.06	3.10	± 0.03		± 0.03		±	0.03		NA		
As is, w/o cellulose	7(NC)-a	3.20	± 0.34	2.96	± 0.07	0.99	± 0.14	2.24	± 0.00	2.07	±	0.10		NA	0.65	± 0.3
As is, with cellulose	7(C)-a	3.37	± 0.27	3.47	± 0.10	2.99	± 0.07	1.33	± 0.34	2.28	±	1.17		NA		± 0.1
Unamended/Inoculated (8)																
Formalin treated, w/o cellulose	8(NC)-f	3.79	± 0.04	2.31	± 0.07	1.73	± 0.18	0.76	± 0.11	1.55	±	0.11		NA	-0.07	+ 01
Formalin treated, with cellulose	8(C)-f	3.89	± 0.03	3.39	± 0.09	2.54	± 0.15		± 0.00		±	0.06		NA		± 0.0
As is, w/o cellulose	8(NC)-a	3.66	± 0.08	-1.37	± 0.99	1.41	± 0.23	1.87	± 0.08	2.10	±	0.08		NA	-0.50	± 0.0
As is, with cellulose	8(C)-a	3.47	± 0.06	3.63	± 0.13	2.53	± 0.16	1.78	± 0.13	2.22	±	0.06		NA		± 0.6
Amended/Inoculated (9)																
As is, w/o cellulose	9(NC)-a	3.77	± 0.04	3.35	± 0.08	6.63	± 0.30	7.96	± 1.33	9.07	±	0.88	8.41	± 0.48	8.04	± 0.4
As is, with cellulose	9(C)-a		± 0.09	3.44	± 0.16	4.04	± 0.03	11.3	± 0.5	16.7	±	1.3	19.2	± 1.2	19.4	± 0.6
as is, with glucose	9(G)-a	2.97	± 0.08	2.59	± 0.15	2.36	± 0.15	1.64	± 0.15	1.52	±	0.27	0.18	± 0.16	-1.26	
excess nitrate, w/o cellulose	9(NC)-x		± 0.04	2.86	± 0.08	4.30	± 0.23	6.59	± 0.76	9.18	±	0.23	8.65	± 0.67	6.93	± 0.2
Excess nitrate, with cellulose	9(C)-x		± 0.13	2.28	± 0.38	5.76	± 1.47	16.6	± 2.9	14.2	±	3.9	28.9	± 6.2	34.1	± 7.1
excess nitrate, with glucose	9(G)-x	3.01	± 0.11	0.61	± 0.15	0.19	± 0.11	0.46	± 0.04	0.46	+	0.61		± 2.08	1.14	

Table 8. Gross Data for the Long-Term Inundated Experiment: Production of Carbon Dioxide\* in Anaerobic Samples

Treatments	Sample			Ca		(µmoles/sample)	)	
[Brine]	Designation		_		Incubation Ti	` • /		
	·	0	45	69	104	132	164	200
Unamended/Uninoculated (7)								
Formalin treated, w/o cellulose	7(NC)-f	2.26 ± 0.01	1.92 ± 0.03	2.20 ± 0.20	2.24 ± 0.04	2.33 ± 0.01	NA	2.39 ± 0.01
Formalin treated, with cellulose	7(C)-f	$31.7 \pm 0.5$	$38.0 \pm 0.1$	$38.0 \pm 0.i$	$35.0 \pm 0.2$	$36.4 \pm 0.1$	NA	$36.3 \pm 0.1$
As is, w/o cellulose	7(NC)-a	1.12 ± 0.01	$0.58 \pm 0.13$	$0.90 \pm 0.00$	1.37 ± 0.01	1.35 ± 0.01	NA	1.46 ± 0.03
As is, with cellulose	7(C)-a	$13.0 \pm 0.4$	$19.3 \pm 0.1$	$20.5 \pm 0.1$	19.5 ± 0.1	$20.5 \pm 0.2$	NA	19.4 ± 0.2
Inamended/Inoculated (8)								
Formalin treated, w/o cellulose	8(NC)-f	4.42 ± 0.01	$3.88 \pm 0.08$	$3.89 \pm 0.13$	$3.88 \pm 0.04$	$3.99 \pm 0.03$	NA	3.97 ± 0.03
Formalin treated, with cellulose	8(C)-f	$32.1 \pm 0.7$	$34.3 \pm 0.4$	$33.5 \pm 0.2$	$31.2 \pm 0.0$	$32.5 \pm 0.5$	NA	$32.8 \pm 0.4$
As is, w/o cellulose	8(NC)-a	1.96 ± 0.02	$1.37 \pm 0.01$	$2.29 \pm 0.02$	$2.75 \pm 0.01$	$2.86 \pm 0.01$	NA	2.74 ± 0.01
As is, with cellulose	8(C)-a	$12.5 \pm 0.2$	$18.4 \pm 0.2$	19.0 ± 0.1	$17.8 \pm 0.7$	$22.7 \pm 0.5$	NA	30.1 ± 1.7
Amended/Inoculated (9)								
As is, w/o cellulose	9(NC)-a	3.02 ± 0.00	6.86 ± 0.02	58.9 ± 3.1	95.9 ± 9.5	109 ± 4	119 ± 3	120 ± 1
As is, with cellulose	9(U)-a	$2.70 \pm 0.00$	$25.8 \pm 0.2$	$42.5 \pm 1.7$	$132 \pm 3$	$200 \pm 7$	$240 \pm 3$	$250 \pm 4$
As is, with glucose	9(G)-a	$2.25 \pm 0.01$	$3.36 \pm 0.04$	$2.96 \pm 0.04$	$3.50 \pm 0.05$	$3.88 \pm 0.06$	4.26 ± 0.09	4.31 ± 0.07
excess nitrate, w/o cellulose	9(NC)-x	$3.07 \pm 0.03$	5.17 ± 0.28	$35.3 \pm 5.3$	94.5 ± 4.4	120 ± 1	125 ± 1	123 ± 1
Excess nitrate, with cellulose	9(C)-x	$5.40 \pm 0.00$	$26.6 \pm 0.2$	65.8 ± 17.1	$193 \pm 33$	$249 \pm 32$	$352 \pm 40$	$430 \pm 41$
Excess nitrate, with glucose	9(G)-x	$1.81 \pm 0.27$	$2.88 \pm 0.19$	$2.65 \pm 0.25$	$8.64 \pm 3.62$	$23.6 \pm 13.3$	$53.6 \pm 31.4$	74.9 ± 40.9

<sup>\*</sup>Dissolved carbon dioxide concentrations not included.

Table 9. Gross Data for the Long-Term Inundated Experiment: Production of Nitrous Oxide\* in Anaerobic Samples

Treatments	Sample								N	itrous Ox	ride	(µmol	es/samp	le)				
[Brine]	Designation									Incuba	tior	ı Time (D	Days)					
			0		69	<del>}</del>		10	)4		13	32		16	54		20	00
Unamended/Uninoculated (7)																		
Formalin treated, w/o cellulose	7(NC)-f	0.000	± 0.00	0.000	±	0.000	0.000	±	0.000	0.000	±	0.000		NA	\	0.000	±	0.000
Formalin treated, with cellulose	7(C)-f	0.000	± 0.00	0.009	±	0.007	0.004	±	0.001	0.007	±	0.005		NA	١	0.002	±	0.002
As is, w/o cellulose	7(NC)-a	0.000	± 0.00	0.003	±	0.002	0.001	±	0.000	0.000	±	0.000		NA	\	0.000	±	0.000
As is, with cellulose	7(C)-a	0.000	± 0.00	0.002	±	0.002	0.003	±	0.002	0.000	±	0.000		NA	<b>\</b>	0.002	±	0.002
Unamended/Inoculated (8)																		
Formalin treated, w/o cellulose	8(NC)-f	0.000	± 0.00	0.000	±	0.000	0.000	±	0.000	0.000	±	0.000		NA	1	0.000	±	0.000
Formalin treated, with cellulose	8(C)-f	0.000	± 0.00	0.002	±	0.001	0.002	±	0.000	0.014	±	0.001		NA	1	0.008	±	0.000
As is, w/o cellulose	8(NC)-a	0.000	± 0.000	0.000	±	0.000	0.000	±	0.000	0.000	±	0.000		NA	1	0.000	±	0.000
As is, with cellulose .	8(C)-a	0.000	± 0.000	0.001	±	0.001	0.000	±	0.000	0.000	±	0.000		NA	1	0.000	±	0.000
Amended/Inoculated (9)																		
As is, w/o cellulose	9(NC)-a	0.000	± 0.000	0.376	±	0.262	35.5	±	25.1	7.55	±	5.34	0.000	±	0.000	0.359	±	0.235
As is, with cellulose	9(C)-a	0.000	± 0.000			0.145	13.2	_	10.1	86.5	±	41.3	89.3	±	43.0	77.8		42.2
As is, with glucose	9(G)-a	0.000	± 0.000	0.002	±	0.001	0.000	±	0.000	0.005	±	0.003	0.005	±	0.000	0.015	±	0.010
Excess nitrate, w/o cellulose	9(NC)-x	0.000				0.021	5.13	±	2.48	0.014	±	0.002	0.000	±	0.000	0.000		0.000
excess nitrate, with cellulose	9(C)-x	0.000	± 0.000			0.422	85.7	±	50.2	196	±	75	280	±	160	395	_	228
excess nitrate, with glucose	9(G)-x	0.000	± 0.000	0.028	±	0.020	3.84	±	2.72	0.523	±	0.006	0.539	±	0.381	13.8	±	0.5

<sup>\*</sup>Dissolved nitrous oxide concentrations not included.

Table 10. Gross Data for the Long-Term Inundated Experiment: Total Volume of Gas Produced in Anaerobic Samples.

Treatments*	Sample								as Produce			<del></del>				
[Brine/Bentonite]	Designation							ubation	Time (Day	ys)						
			0		45		69		104		13	32		164		200
Unamended/Uninoculated (10)																
Formalin treated, w/o cellulose	10(NC)-f	4.65	± 0.23	4.61	± 0.47	2.37	± 1.47	3.41	± 0.19	4.07	±	0.04		NA	2.33	± 0.00
Formalin treated, with cellulose	10(C)-f	4.96	± 0.06	4.64	± 0.03	4.22	± 0.00	2.94	± 0.16	3.39	±	0.16		NA	2.05	± 0.10
As is, w/o cellulose	10(NC)-a	4.24	± 0.33	4.04	± 0.04	3.51	± 0.69	2.86	± 0.08	3.27	±	0.20		NA	1.55	± 0.0
As is, with cellulose	10(C)-a	3.84	± 0.48	3.84	± 0.14	3.33	± 0.27	2.01	± 0.41	2.18	±	0.48		NA	0.14	± 0.4
Unamended/Inoculated (11)																
Formalin treated, w/o cellulose	11(NC)-f	5.16	± 0.07	5.01	± 0.14	3.89	± 0.22	2.85	± 0.25	3.57	±	0.25		NA	1.23	± 0.22
Formalin treated, with cellulose	11(C)-f	4.65	± 0.03	4.53	± 0.00	3.66	± 0.03	1.73	± 1.14	2.16	±	1.08		NA	0.94	± 1.20
As is, w/o cellulose	11(NC)-a	4.23	± 0.11	3.62	± 0.08	3.47	± 0.19	2.06	± 0.19	2.44	±	0.30		NA	0.34	± 0.34
As is, with cellulose	11(C)-a	4.38	± 0.03	4.19	± 0.06	3.19	± 0.16	2.88	± 0.22	3.88	±	0.19		NA	4.38	± 0.38
Amended/Inoculated (12)																
As is, w/o cellulose	12(NC)-a	3.85	± 0.08	4.53	± 0.15	6.97	± 0.15	9.90	± 0.08	10.9	±	0.1		± 0.3		± 0.80
As is, with cellulose	12(C)-a	3.32	± 0.25	4.29	$\pm 0.09$	7.92	± 0.44	16.8	± 0.4	19.8	±	0.4		± 0.3	19.5	± 0.4
As is, with glucose	12(G)-a	3.50	± 0.04	5.52	± 0.34	9.49	± 2.40	24.7	± 3.3	33.9	±	0.6	38.8	± 1.8	37.8	± 2.3
Excess nitrate, w/o cellulose	12(NC)-x		± 0.00		± 0.15		± 0.69		± 0.3	12.4		0.0		± 0.1		± 0.1
Excess nitrate, with cellulose	12(C)-x		± 0.09		± 0.16		± 0.31		± 0.4	20.8		0.5		± 0.7		± 1.4
Excess nitrate, with glucose	12(G)-x	3.09	± 0.08	2.29	± 0.19	2.21	± 0.46	23.4	± 0.6	52.3	±	6.7	77.4	± 20.2	80.7	± 21.1

Table 11. Gross Data for the Long-Term Inundated Experiment: Production of Carbon Dioxide\* in Anaerobic Samples

Treatments	Sample			Ca	rbon Dioxide	(µmoles/sample	)	
[Brine/Bentonite]	Designation				Incubation Ti	me (Days)		
		0	45	69	104	132	164	200
Unamended/Uninoculated (10)								
Formalin treated, w/o cellulose	10(NC)-f	55.8 ± 0.6	82.9 ± 1.1	85.5 ± 0.8	86.4 ± 1.7	92.8 ± 2.0	NA	94.9 ± 1.1
Formalin treated, with cellulose	10(C)-f	$65.5 \pm 1.0$	$85.5 \pm 0.2$	88.6 ± 0.4	$81.9 \pm 0.3$	$88.7 \pm 0.9$	NA	$90.2 \pm 0.3$
As is, w/o cellulose	10(NC)-a	16.4 ± 0.4	$37.7 \pm 0.0$	38.4 ± 0.1	38.1 ± 0.3	39.8 ± 0.3	NA	40.0 ± 0.4
As is, with cellulose	10(C)-a	26.6 ± 0.5	$42.6 \pm 0.2$	$43.0 \pm 0.4$	$41.3 \pm 0.2$	$43.1 \pm 0.5$	NA	41.1 ± 0.2
Inamended/Inoculated (11)								
Formalin treated, w/o cellulose	11(NC)-f	53.0 ± 0.2	76.9 ± 0.6	80.2 ± 0.4	76.5 ± 0.0	83.7 ± 0.6	NA	83.5 ± 0.2
Formalin treated, with cellulose	11(C)-f	64.4 ± 1.3	$76.3 \pm 0.5$	78.6 ± 0.2	75.1 ± 1.9	78.6 ± 1.6	NA	80.4 ± 0.8
As is, w/o cellulose	11(NC)-a	$20.8 \pm 0.0$	$35.0 \pm 0.1$	$36.3 \pm 0.1$	$35.2 \pm 0.2$	19.6 ± 11.9	NA	36.2 ± 0.2
As is, with cellulose	11(C)-a	$30.1 \pm 0.6$	$38.1 \pm 0.2$	$40.5 \pm 0.1$	48 ± 2.5	59.9 ± 0.7	NA	77.6 ± 1.0
Amended/Inoculated (12)								
As is, w/o cellulose	12(NC)-a	14.0 ± 0.0	60.2 ± 0.2	99.8 ± 4.7	150 ± 4	16.6 ± 0.4	177 ± 1	179 ± 0
As is, with cellulose	12(C)-a	$12.0 \pm 0.8$	$55.0 \pm 0.3$	$104 \pm 5$	$209 \pm 2$	$260 \pm 8$	$295 \pm 10$	$338 \pm 10$
As is, with glucose	12(G)-a	15.2 ± 0.6	76.0 ± 2.4	144 ± 28	$375 \pm 36$	590 ± 15	691 ± 20	786 ± 25
Excess nitrate, w/o cellulose	12(NC)-x	13.9 ± 0.3	69.2 ± 0.0	121 ± 3	166 ± 1	186 ± 2	196 ± 2	196 ± 2
excess nitrate, with cellulose	12(C)-x	$10.3 \pm 0.6$	$57.4 \pm 1.7$	$122 \pm 3$	195 ± 5	$264 \pm 6$	$309 \pm 7$	$371 \pm 14$
Excess nitrate, with glucose	12(G)-x	$15.1 \pm 0.0$	$69.4 \pm 1.2$	$111 \pm 4$	$422 \pm 12$	$916 \pm 35$	1370 ± 19	1610 ± 43

<sup>\*</sup>Dissolved carbon dioxide concentrations not included.

Table 12. Gross Data for the Long-Term Inundated Experiment: Production of Nitrous Oxide\* in Anaerobic Samples

Treatments	Sample								N	Incuba		umol) Time (D	es/sampl	le)				
[Brine/Bentonite]	Designation		0		69			10	4		13	•		16	4		20	00
Unamended/Uninoculated (10)																		
Formalin treated, w/o cellulose	10(NC)-f	0.000 :		0.000	_	0.000		±	0.000	0.000	_	0.000		NA		0.000	_	0.00
Formalin treated, with cellulose	10(C)-f	0.000	£ 0.000	0.000	±	0.000	0.001	±	0.000	0.000	±	0.000		NA		0.000	±	0.00
As is, w/o cellulose	10(NC)-a	0.000 :	£ 0.000	0.004	±	0.003	0.102	±	0.065	1.040	±	0.490		NA	ı	1.73	±	0.10
As is, with cellulose	10(C)-a	0.000	₺ 0.000	0.005	±	0.002	0.008	±	0.004	0.021	±	0.009		NA		0.031	±	0.011
Inamended/Inoculated (11)																		
Formalin treated, w/o cellulose	11(NC)-f	0.000 :	£ 0.000	0.000	±	0.000	0.000	±	0.000	0.000	±	0.000		NA	ı	0.002	±	0.00
Formalin treated, with cellulose	11(C)-f	0.000	£ 0.000	0.000	±	0.000	0.006	±	0.002	0.024	±	0.003		NA		0.000	±	0.00
As is, w/o cellulose	11(NC)-a	0.000 :	e 0.000	0.001	±	0.001	0.000	±	0.000	0.037	±	0.026		NA		0.007	±	0.00
As is, with cellulose	11(C)-a	0.000	0.000	0.001	±	0.001	0.000	±	0.000	0.000	±	0.000		NA		0.000	±	0.00
Amended/Inoculated (12)																		
As is, w/o cellulose	12(NC)-a	0.000	0.000	0.016	±	0.006	0.534	±	0.378	0.000	±	0.000	0.000	±	0.000	3.35		2.19
As is, with cellulose	12(C)-a	0.000	0.000	0.111	±	0.056	2.34	±	1.92	0.000		0.000	0.301	±	0.246	0.000		0.00
As is, with glucose	12(G)-a	0.000	0.000	1.56	±	1.10	40.1	±	28.3	0.782	±	0.557	0.016	±	0.011	0.000	±	0.00
excess nitrate, w/o cellulose	12(NC)-x	0.000	0.000	0.007	±	0.001	0.081	±	0.057	0.000		0.000	0.003	±	0.002	0.481		0.34
excess nitrate, with cellulose	12(C)-x	0.004	0.000			2.39		±	0.018	47.1	±	17.8	154	±	20	311		23
xcess nitrate, with glucose	12(G)-x	0.005	0.000	0.077	±	0.049	2.31	±	1.59	261	±	184	616	±	428	585	±	406

<sup>\*</sup>Dissolved nitrous oxide concentrations not included.

Table 13. Total Volume of Gas Produced in Aerobic Samples in the Presence of Cellulose\*

Treatments	Sample				Gas Produced/Gran	n Cellulose		
[Brine]	Desig.	0	45	Incubation 69	on Time (Days) 104	132	164	200
Unamended/U	Ininoculate							
Formalin	1(C)-f	0.24 ± 0.06	0.89 ± 0.03	0.85 ± 0.02	0.42 ± 0.21	0.27 ± 0.19	NA	0.42 ± 0.18
As is	1(C)-a	0.05 ± 0.02	-0.29 ± 0.19	-0.36 ± 0.18	-0.31 ± 0.14	-0.20 ± 0.12	NA	-0.18 ± 0.10
Unamended/I	noculated	(2)						
Formalin	2(C)-f	-0.01 ± 0.01	-0.03 ± 0.08	-0.35 ± 0.11	-0.32 ± 0.08	-0.41 ± 0.11	NA	-0.11 ± 0.09
As is	2(C)-a	-0.08 ± 0.01	0.01 ± 0.04	-0.15 ± 0.03	-0.11 ± 0.04	0.00 ± 0.05	NA	-0.34 ± 0.07
Amended/Ino	culated (3)							
As is	3(C)-a	-0.12 ± 0.03	-0.27 ± 0.14	-0.25 ± 0.06	$0.36 \pm 0.31$	$0.29 \pm 0.24$	$0.07 \pm 0.13$	0.86 ± 0.08
Exc. nitrate	3(C)-x	-0.02 ± 0.01	0.02 ± 0.14	1.39 ± 0.61	2.38 ± 0.87	3.10 ± 1.06	4.02 ± 1.21	4.42 ± 0.80

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose NA = not analyzed

0-1

Table 14. Total Volume of Gas Produced in Aerobic Samples in the Presence of Cellulose and Bentonite\*

Treatments	Sample				Gas Produced/Gran	n Cellulose		
[Brine/Bent.]	Designation	l		Incubation	on Time (Days)			
		0	45	69	104	132	164	200
Unamended/U	ninoculated	1 (4)						
Formalin	4(C)-f	$0.13 \pm 0.03$	$0.63 \pm 0.03$	$0.47 \pm 0.03$	$0.15 \pm 0.03$	$0.02 \pm 0.01$	NA	0.04 ± 0.01
As is	4(C)-a	-0.05 ± 0.02	$0.02 \pm 0.02$	-0.36 ± 0.04	-0.2J ± 0.05	-0.13 ± 0.04	NA	$0.00 \pm 0.05$
<i>Unamended/Ir</i> Formalin As is	5(C)-f 5(C)-a	0.11 ± 0.02 0.03 ± 0.01	0.21 ± 0.01 -0.10 ± 0.02	0.31 ± 0.00 -0.18 ± 0.04	0.29 ± 0.00 -0.30 ± 0.13	$0.07 \pm 0.03$ -0.23 \pm 0.09	NA NA	0.16 ± 0.02 -0.08 ± 0.16
Amended/Inoc	, ,	0.25 ± 0.02	0.42 ± 0.01	0.71 ± 0.15	192 ± 025	206 ± 0.20	407 ± 0.22	4.28 ± 0.20
43 13	6(C)-a	$-0.25 \pm 0.03$	$0.43 \pm 0.01$	$0.71 \pm 0.15$	$1.82 \pm 0.35$	$2.96 \pm 0.30$	$4.07 \pm 0.22$	4.38 ± 0.20
Exc. nitrate	6(C)-x	$0.30 \pm 0.01$	$0.85 \pm 0.02$	$1.56 \pm 0.02$	$2.23 \pm 0.24$	$3.74 \pm 0.29$	$5.15 \pm 0.18$	6.07 ± 0.04

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose NA = not analyzed

Table 15. Production of Carbon Dioxide in Aerobic Samples in the Presence of Cellulose\*

Sample					ellulose)		
Desig.	0	45	69	104	132	164	200
ninoculated	1(1)						
1(C)-f	5.58 ± 0.54	7.66 ± 0.02	8.20 ± 0.04	7.46 ± 0.16	7.64 ± 0.06	NA	7.62 ± 0.10
1(C)-a	2.48 ± 0.10	3.92 ± 0.06	4.21 ± 0.02	3.93 ± 0.02	4.25 ± 0.02	NA	4.00 ± 0.02
2(C)-f	5.91 ± 0.16	6.24 ± 0.00	6.80 ± 0.14	5.88 ± 0.12	6.04 ± 0.02	NA NA	5.65 ± 0.22 8.30 ± 0.23
	1.90 ± 0.00	3.33 1 0.10	3.54 1 0.16	4.96 1 0.16			
	ND	462 ± 0.29	279 ± 119	20.4 + 7.4	20.6 + 5	366 + 42	40.8 ± 5.4
.)(C)-a	NU	4.62 ± 0.28 6.88 ± 0.00	32.2 ± 4.2	61.4 ± 3.6	83.6 ± 2.6	96.2 ± 4.0	95.6 ± 6.0
	Desig.  Ininoculated  1(C)-f  1(C)-a	Desig.  0  Ininoculated (1)  1(C)-f	Desig.  0 45  Ininoculated (1)  1(C)-f 5.58 ± 0.54 7.66 ± 0.02  1(C)-a 2.48 ± 0.10 3.92 ± 0.06  Designoculated (2)  2(C)-f 5.91 ± 0.16 6.24 ± 0.00  2(C)-a 1.96 ± 0.06 3.35 ± 0.16  Evaluated (3)	Incubated (1)  1(C)-f	Design   Inculation Time (Days)	Inculation Time (Days) 69 104 132  Ininoculated (1)  1(C)-f 5.58 ± 0.54 7.66 ± 0.02 8.20 ± 0.04 7.46 ± 0.16 7.64 ± 0.06  1(C)-a 2.48 ± 0.10 3.92 ± 0.06 4.21 ± 0.02 3.93 ± 0.02 4.25 ± 0.02  Inoculated (2)  2(C)-f 5.91 ± 0.16 6.24 ± 0.00 6.80 ± 0.14 5.88 ± 0.12 6.04 ± 0.02  2(C)-a 1.96 ± 0.06 3.35 ± 0.16 3.94 ± 0.18 4.98 ± 0.18 6.87 ± 0.20  Inculated (3)  3(C)-a ND 4.62 ± 0.28 3.78 ± 1.18 20.4 ± 7.4 29.6 ± 5	Inculation Time (Days) 69 104 132 164  Ininoculated (1)  1(C)-f 5.58 ± 0.54 7.66 ± 0.02 8.20 ± 0.04 7.46 ± 0.16 7.64 ± 0.06 NA  1(C)-a 2.48 ± 0.10 3.92 ± 0.06 4.21 ± 0.02 3.93 ± 0.02 4.25 ± 0.02 NA  Inculated (2)  2(C)-f 5.91 ± 0.16 6.24 ± 0.00 6.80 ± 0.14 5.88 ± 0.12 6.04 ± 0.02 NA  2(C)-a 1.96 ± 0.06 3.35 ± 0.16 3.94 ± 0.18 4.98 ± 0.18 6.87 ± 0.20 NA  Inculated (3)  Inculated (3)  3(C)-a ND 4.62 ± 0.28 3.78 ± 1.18 20.4 ± 7.4 29.6 ± 5 36.6 ± 4.2

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose; dissolved gas concentration not included

ND - not detected

NA - not analyzed

Table 16. Production of Carbon Dioxide in Aerobic Samples in the Presence of Cellulose and Bentonite\*

Treatments	Sample				Dioxide (µmoles/ tion Time (Days)	gran cenalose)		
[Brine/Bent.]	Desig.	0	45	69	104	132	164	200_
Unamended/U	ninoculated	i (4)						
Formalin	4(C)-f	4.16 ± 0.24	1.30 ± 0.04	2.24 ± 0.00	2.60 ± 0.00	2.40 ± 0.20	NA	3.20 ± 0.20
As is	4(C)-a	1.52 ± 0.30	1.76 ± 0.06	4.48 ± 2.36	1.76 ± 0.14	1.82 ± 0.10	NA	2.32 ± 0.02
Unamended/Iv	roculated (5	5)						
Formalin	5(C)-f	4.92 ± 0.18	4.78 ± 0.00	5.58 ± 0.04	3.16 ± 0.12	4.80 ± 0.00	NA	4.60 ± 0.00
As is	5(C)-a	2.04 ± 0.04	1.38 ± 0.08	4.00 ± 0.80	8.32 ± 0.42	11.9 ± 0.6	NA	21.5 ± 1.2
Amended/Inoc	culated (6)							
As is	6(C)-a	ND	ND	ND	12.4 ± 2.2	31.4 ± 3.8	57.8 ± 1.2	69.8 ± 1.2
Exc. nitrate	6(C)-x	ND	ND	2.2 ± 0.6	27.2 ± 4.6	72.0 ± 15.0	105 ± 9	116 ± 6

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose; dissolved gas concentration not included

ND - not detected

NA - not analyzed

100

Table 17. Production of Nitrous Oxide in Aerobic Samples in the Presence of Cellulose\*

Treatments	Sample					N:	itrous Oxi			s/gram (	elk	dose)						
[Brine]	Desig.		0		69		incuba	<b>tion</b> 10	Time (D	ays)	1.3	32		16	4		20	0
				<del></del>				- 10										
Unamended/l	I ninoculat	ed (1)																
Formalin	1(C)-f		ND		ND		0.001	ŧ	0.000	0.003	±	0.000		NA		0.001	Ì	0.000
As is	1(C)-a		ND		ND		0.001	±	0.000	0.003	±	0.000		NA			ND	
Formalin As is	2(C)-f 2(C)-a		ND ± 0.000		ND ND			ND ND		0.003	±	0.000		NA NA		0.001	± ND	
Amended/Ino	culated (3)	1				<del>, , , , , , , , , , , , , , , , , , , </del>												
As is	3(C)-a	0.001	± 0.000	0.024	±	0.029	23.6	±	9.6	24.4	±	7.8	1.76	±	1.20	5.83	±	5.04
				5.64			49.0			65.2			96.8			115	±	26

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose; dissolved gas concentration not included

ND - not detected

NA - not analyzed

Table 18. Production of Nitrous Oxide in Aerobic Samples in the Presence of Cellulose and Bentonite\*

Treatments	Sample _			Ni	trous Oxi		s/gram a	cellulose)				
[Brine/Bent.]	Desig.	0		69	Incuba	tion Time (D	ays)	132	1	64	-	200
Unamended/U	ninoculated (	4)										
Formalin	4(C)-f	ND	0.001	± 0.000		ND	0.002	± 0.000	N	A	0.001	± 0.000
As is	4(C)-a	ND	0.145	± 0.098	0.037	± 0.011	0.041	± 0.014	N	A	0.048	± 0.020
Inamended/Ir	oculated (5)											
Formalin	5(C)-f	ND	ı	ND		ND		ND	N	A		ND
As is	5(C)-a	. ND	0.000 :	± 0.001		ND		ND	N	Α		ND
Amended/Inoc	ulated (6)											
As is	6(C)-a	ND	6.32	± 2.64	31.8	± 10.6	25.2	± 1.6	N	D		ND
Exc. nitrate	6(C)-x	ND	5.42	± 1.30	25.5	± 4.4	40.6	± 8.2	68.0 ±	8.2	82.7	± 7.0

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose; dissolved gas concentration not included

ND - not detected

NA - not analyzed

7.7

Table 19. Total Volume of Gas Produced in Anaerobic Samples in the Presence of Cellulose\*

Treatments	Sample				f Gas Produced/Gra tion Time (Days)	am Cellulose		
(Brine)	Designation	0	45	69	104	132	164	200
Unamended/U	Ininoculated	(7)						
Formalin	7(C)-f	0.26 ± 0.01	0.17 ± 0.01	0.22 ± 0.01	0.17 ± 0.01	$0.14 \pm 0.01$	NA	0.26 ± 0.00
As is	7(C)-a	0.03 ± 0.05	0.10 ± 0.02	0.40 ± 0.01	-0.18 ± 0.07	0.04 ± 0.23	NA	-0.09 ± 0.03
Unamended/1	noculated (8	")						
Formalin	8(C)-f	0.02 ± 0.01	0.22 ± 0.02	0.16 ± 0.03	$0.16 \pm 0.00$	$0.12 \pm 0.01$	NA	0.22 ± 0.00
As is	8(C)-a	-0.04 ± 0.01	1.00 ± 0.03	0.22 ± 0.03	-0.02 ± 0.03	0.02 ± 0.01	NA	0.59 ± 0.13
Amended/Ino	culated (9)							
As is	9(C)-a	-0.08 ± 0.02	0.02 ± 0.03	-0.52 ± 0.01	0.66 ± 0.09	$1.52 \pm 0.25$	2.15 ± 0.24	2.27 ± 0.12
Exc. nitrate	9(C)-x	0.01 ± 0.03	-0.12 ± 0.08	0.29 ± 0.29	2.00 ± 0.58	2.24 ± 0.78	4.04 ± 1.24	5.44 ± 1.42

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose NA = not analyzed

Table 20. Total Volume of Gas Produced in Anaerobic Samples in the Presence of Cellulose and Bentonite\*

Treatments	Sample				Bas Produced/Gran	n Cellulose		
[Brine/Bent.]	Designation	0	45	Incubation 69	on Time (Days) 104	132	164	200
Unamended/U	ninoculated	i (10)						
Formalin	10(C)-f	0.06 ± 0.01	0.01 ± 0.01	0.37 ± 0.00	-0.09 ± 0.03	-0.14 ± 0.03	NA	-0.06 ± 0.03
As is	10(C)-a	-0.08 ± 0.10	-0.04 ± 0.03	-0.04 ± 0.05	-0.17 ± 0.08	-0.22 ± 0.10	NA	-0.28 ± 0.09
Unamended/Ii	noculated (1	(1)						
Formalin	11(C)-f	-0.10 ± 0.01	-0.10 ± 0.00	-0.05 ± 0.01	-0.22 ± 0.23	-0.28 ± 0.22	NA	-0.06 ± 0.2
As is	11(C)-a	0.03 ± 0.01	0.11 ± 0.01	-0.06 ± 0.03	$0.16 \pm 0.04$	0.29 ± 0.04	NA	0.81 ± 0.0
Amended/Ino	culated (12)							
As is	12(C)-a	-0.11 ± 0.05	-0.05 ± 0.02	0.19 ± 0.09	1.39 ± 0.09	1.78 ± 0.08	1.44 ± 0.07	1.92 ± 0.0
Exc. nitrate	12(C)-x	-0.06 ± 0.02	-0.09 ± 0.03	0.23 ± 0.06	0.78 ± 0.09	1.68 ± 0.10	2.19 ± 0.14	3.52 ± 0.2

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose NA = not analyzed

Table 21. Production of Carbon Dioxide in Anaerobic Samples in the Presence of Cellulose\*

Treatments	Sample				pioxide (µmoles/g	gram cellulose)		
[Brine]	Desig.				ion Time (Days)	400	161	200
		0	45 .	69	104	132	164	200
Inamended/U	ninoculated	1(7)						
Formalin	7(C)-f	5.89 ± 0.10	7.22 ± 0.02	7.16 ± 0.02	6.55 ± 0.04	$6.81 \pm 0.02$	NA	6.78 ± 0.02
As is	7(C)-a	2.38 ± 0.08	3.74 ± 0.02	3.92 ± 0.02	3.63 ± 0.02	3.83 ± 0.04	NA	3.59 ± 0.04
Jnamended/1	noculated (8	3)						
Formalin	8(C)-f	5.54 ± 0.14	$6.08 \pm 0.08$	5.92 ± 0.04	5.46 ± 0.00	5.70 ± 0.10	NA	5.77 ± 0.00
As is	8(C)-a	2.11 ± 0.04	3.41 ± 0.04	3.34 ± 0.02	3.01 ± 0.14	3.97 ± 0.10	NA	5.47 ± 0.3
Amended/Ino	culated (9)							
As is	9(C)-a	ND	3.79 ± 0.04	ND	7.22 ± 0.60	18.2 ± 1.4	24.2 ± 0.6	26.0 ± 0.8
Exc. nitrate	9(C)-x	0.47 ± 0.00	4.29 ± 0.04	6.10 ± 3.42	19.7 ± 6.6	25.8 ± 6.4	45.4 ± 8.0	61.4 ± 8.2

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose; dissolved gas concentration not included

ND - not detected

NA - not analyzed

Table 22. Production of Carbon Dioxide\* in Anaerobic Samples in the Presence of Cellulose and Bentonite

ON		<b>ON</b>										
		UIT	020	09.0 ±	± 08.δ	00.1 ±	9.21	2.1 ±	22.6	<b>₽.I</b> .±	32.0	82 ±
QN		<b>GN</b>	18.0	00.1 ±	± 8.11	<b>₽</b> .0 ±	89.84	9°I ∓	23.6	<b>0.2</b> ±	31.8	₽. 2.0
21.0 ± 8	29.0	<b>40.0</b> ±	18.0	20.0 ±	₹ 95°7	0\$.0 ±	90.8	<b>≯1.0</b> ±		AN	82.8	02.0 ±
97.0 ± 8		QN		П	3N	QN		QΝ		ΑN		QN
1.0 ± p	86.0	<b>40.0</b> ±	26.0	80.0 ±	Ŧ <b>1</b> 9.0	<b>≯</b> 0.04	99.0	01.0 ±		AN	22.0	<b>≯0.0</b> ±
2.0 ± 4	22.0	<b>≯</b> 0.0 ∓	29.0	80.0 ±	)N	QN		<b>ON</b>		ΔN		ПN
						•						
0		SÞ		Incubatio		10¢ : (Ds\2)		132		191		200
2.0 ± p 1.0 ± p	86.0	T T	<b>+0.0 +0.0</b>	28.0 <b>+</b> 0.0 28.0 <b>+</b> 0.0	ND   ND   ND   ND   ND   ND   ND   ND	80.0 ± 26.0 +0.0 140.0 80.0 ± 26.0 +0.0	Incubation Time (Days)   104   104   104   104   104   104   104   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   1004   100	Incubation Time (Days)   104   269   104   104   104   104   104   104   104   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105   105	Incubation Time (Days)   132   69   104   132   134   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135	Incubation Time (Days)   132   69   104   132   132   134   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135   135	Name	Name

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose; dissolved gas concentration not included WA - not analyzed

Table 23. Production of Nitrous Oxide in Anaerobic Samples in the Presence of Cellulose\*

Treatments	Sample		<u> </u>		es/gram cellulose)		
[Brine]	Desig.	_		Incubation Time (I		464	200
		0	69	104	132	164	200
U <b>namended/U</b>	ninoculated (	7)					
Formalin	7(C)-f	ND	0.002 ± 0.001	0.001 ± 0.000	0.001 ± 0.001	NA	ND
As is	7(C)-a	ND	ND	ND	ND	NA 	ND
Unamended/I	noculated (8)						
Formalin	8(C)-f	ND	ND	ND	$0.003 \pm 0.000$	NA	0.002 ± 0.000
As is	8(C)-a	ND	ND	ND	ND	NA	ND
Amended/Ino	culated (9)						
As is	9(C)-a	ND	ND	ND	15.8 ± 8.3	17.9 ± 8.6	15.5 ± 8.4
Exc. nitrate	9(C)-x	ND	0.158 ± 0.084	16.1 ± 10.0	39.2 ± 15.0	56.0 ± 32.0	79.0 ± 45.6

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose; dissolved gas concentration not included ND - not detected

Table 24. Production of Nitrous Oxide\* in Anaerobic Samples in the Presence of Cellulose and Bentonite

Treatments	Sample		Ni		es/gram cellulose)		
Brine/Bent.]	Desig.			Incubation Time (D			000
			69	104	132	164	200
Inamended/U	ninoculated (	10)			•		
Formalin	10(C)-f	ND	ND	ND	ND	NA	ND
As is	10(C)-a	ND	ND	ND	ND	NA	ND
Inamended/Ii	noculated (11)	)					
Formalin	11(C)-f	ND	ND	$0.001 \pm 0.000$	$0.005 \pm 0.001$	NA	ND
As is	11(C)-a	ND	ND	ND	ND	NA 	ND
Amended/Inoc	ulated (12)						
As is	12(C)-a	ND	0.019 ± 0.011	0.361 ± 0.384	ND	$0.060 \pm 0.049$	ND
Exc. nitrate	12(C)-x 0	.001 ± 0.000	0.759 ± 0.478	0.000 ± 0.000	9.42 ± 3.56	30.8 ± 4.0	62.1 ± 4.6

<sup>\*</sup> All values have been corrected with specific controls for gas production in the absence of cellulose; dissolved gas concentration not included ND - not detected NA - not analyzed

. .

# APPENDIX E: MEASUREMENTS OF MIXED INOCULUM ACTIVITY

To determine the activity of the mixed inoculum, brine samples were incubated with glucose with an initial air and  $N_2$  atmosphere. Total gas,  $CO_2$  and  $N_2O$  were periodically monitored.

# **Aerobic Samples**

### **Total Gas Production**

Samples containing glucose, nutrients, excess nitrates with and without bentonite did not show an increase in total gas production (Figure 25, Tables 1(a)-2(a) Appendix D).

## **Carbon Dioxide Production**

Production of carbon dioxide was evident after 132 days in amended and in amended plus excess nitrate samples without bentonite. After 200 days, amended samples produced 323  $\mu$ mol of CO<sub>2</sub>, whereas amended plus excess nitrate samples produced 296  $\mu$ mol CO<sub>2</sub>. Samples with bentonite did not produce significant amounts of CO<sub>2</sub> beyond the initial background level (~70-80  $\mu$ mol) (Figure 26, Tables 1(b)-2(b) Appendix D).

### **Nitrous Oxide Production**

Nitrous oxide was detected at 132 days in aerobic samples without bentonite (Figure 27, Tables 1(c)-2(c) Appendix D). Less  $N_2O$  was detected in amended samples with excess nitrate, and in samples containing bentonite,  $N_2O$  was not detected.

# **Anaerobic Samples**

# **Total Gas Production**

Amended and amended samples with excess nitrate showed no increase in total gas production. However, in the presence of bentonite, production in amended samples increased after 45 day of incubation and reached to 37.8 mL at 200 days (Figure 28, Tables 3(a)-4(a) Appendix D). Amended samples plus excess nitrate produced more gas (80.65 mL of gas at 200 days) than the other samples.

## Carbon Dioxide Production

Production of carbon dioxide was not detected in the amended samples, whereas the amended samples with excess nitrate produced a small amount of  $CO_2$ . Carbon dioxide production in samples containing bentonite was significant (Figure 29, Tables 3(b)-4(b) Appendix D). Amended samples containing excess nitrate produced 1610  $\mu$ mol cí  $CO_2$  at 200 days while the basic amended samples produced 786 moles. Carbon dioxide production was much higher in the glucose/bentonite samples than in the cellulose/bentonite samples, indicating the potential of the microorganisms to produce significant amounts of  $CO_2$  under hypersaline conditions when a simple sugar is present. In addition, bentonite seems to enhance the overall gas production.

### **Nitrous Oxide Production**

Accumulation of nitrous oxide was detected only in samples containing bentonite (Figure 30, Tables 3(c)-4(c) Appendix D). With excess nitrate, the  $N_2O$  concentration reached 585  $\mu$ mol and then started to decline.

# Summary

We examined glucose metabolism by the mixed inoculum used in the long-term experiments. Total gas production in samples incubated under aerobic conditions was not evident, but anaerobic samples produced significant amounts of gas, especially the samples with bentonite plus excess nitrate. Production of carbon dioxide in the aerobic samples was observed only in the amended and excess-nitrate samples without bentonite. Bentonite enhanced the activity of anaerobes. Carbon-dioxide production in anaerobic samples with excess nitrate reached a higher amount than any of the totals reached thus far in the long-term inundated experiment. Substantial amounts of N<sub>2</sub>O also accumulated in the headspace of these samples.




Figure 25. Total gas produced in samples containing glucose incubated with an initial atmosphere of air.




Figure 26. Carbon dioxide produced in samples containing glucose incubated with an initial atmosphere of air.




Figure 27. Nitrous oxide produced in samples containing glucose incubated with an initial atmosphere of air.




Figure 28. Total gas produced in anaerobic samples containing glucose.




Figure 29. Carbon dioxide produced in anaerobic samples containing glucose.




Figure 30. Nitrous oxide produced in anaerobic samples containing glucose.

# DATE FILMED 5/5/94