






**AIM**

**Association for Information and Image Management**




1100 Wayne Avenue, Suite 1100  
Silver Spring, Maryland 20910  
301/587-8202



Centimeter



Inches



MANUFACTURED TO AIM STANDARDS  
BY APPLIED IMAGE, INC.

**1 of 1**

RECEIVED  
JUN 22 1993  
OSTI

## Measuring The Sea Quark Polarization

Yousef Makdisi

(for the RHIC Spin Collaboration \*)

RHIC Project, Brookhaven National Laboratory, Upton NY 11973

### ABSTRACT

Spin is a fundamental degree of freedom and measuring the spin structure functions of the nucleon should be a basic endeavor for hadron physics. Polarization experiments have been the domain of fixed target experiments. Over the years large transverse asymmetries have been observed where the prevailing QCD theories predicted little or no asymmetries, and conversely the latest deep inelastic scattering experiments of polarized leptons from polarized targets point to the possibility that little of the nucleon spin is carried by the valence quarks.

The possibility of colliding high luminosity polarized proton beams in the Brookhaven Relativistic Heavy Ion Collider (RHIC) provides a great opportunity to extend these studies and systematically probe the spin dependent parton distributions specially to those reactions that are inaccessible to current experiments. This presentation focuses on the measurement of sea quark and possibly the strange quark polarization utilizing the approved RHIC detectors.

Submitted to the Proceedings of the Workshop on Future Directions in Particle  
and Nuclear Physics at Multi-GeV Hadron Facilities.

March 4-6, 1993 Brookhaven National Laboratory, Upton, NY, 11973. USA.

### DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**MASTER**

---

*Work done under contract no. DE-AC02-76CHO-0016 with the U.S. Department of Energy. Accordingly, the U.S. Government retains a non-exclusive, royalty-free licence to publish or reproduce the published form of this contribution, or allow others to do so, for U.S. Government purposes.*

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED *ok*

# Measuring The Sea Quark Polarization

Yusef Makdisi

(for the RHIC Spin Collaboration \*)

*RHIC Project, Brookhaven National Laboratory, Upton NY 11973*

## ABSTRACT

Spin is a fundamental degree of freedom and measuring the spin structure functions of the nucleon should be a basic endeavor for hadron physics. Polarization experiments have been the domain of fixed target experiments. Over the years large transverse asymmetries have been observed where the prevailing QCD theories predicted little or no asymmetries, and conversely the latest deep inelastic scattering experiments of polarized leptons from polarized targets point to the possibility that little of the nucleon spin is carried by the valence quarks.

The possibility of colliding high luminosity polarized proton beams in the Brookhaven Relativistic Heavy Ion Collider (RHIC) provides a great opportunity to extend these studies and systematically probe the spin dependent parton distributions specially to those reactions that are inaccessible to current experiments. This presentation focuses on the measurement of sea quark and possibly the strange quark polarization utilizing the approved RHIC detectors.

## 1. Introduction

Deep inelastic scattering experiments of longitudinally polarized leptons from longitudinally polarized targets have been the main vehicle, in the past decade, for measuring the nucleon spin structure functions. The Yale/SLAC [1] experiment and later the NMC [2] experiment at CERN provided the first detailed measurement of the first moment of the proton spin structure function

$$\Gamma_1^P = \int g_1^P(x) dx = 0.126 \pm 0.010 \text{ (stat)} \pm 0.015 \text{ (syst)}$$

First results from the SMC [3] experiment on a similar measurement of the deuteron spin structure function give

$$\Gamma_1^d = 0.023 \pm 0.020 \text{ (stat)} \pm 0.015 \text{ (syst)}$$

and determine the value for the neutron to be

$$\Gamma_1^n = -0.08 \pm 0.04 \text{ (stat)} \pm 0.04 \text{ (syst)}$$

These results suggest that, contrary to intuition, the contribution of the valence quarks to the overall spin of the nucleon is exceedingly small. What follows then is either the gluons or the sea quarks are highly polarized or both. Note that if the gluons are not polarized then the strange quarks will carry significant polarization, thus the need to measure the contributions of those partons inside the nucleon.

Deep inelastic experiments are not adept at differentiating between quark flavors and provide little sensitivity to the gluon contribution due to the lack of coupling between leptons and gluons. A proposal from

---

\* Work done under contract no. DE-AC02-76CHO-0016 with the U.S. Department of Energy.

the RHIC Spin Collaboration [4] details the physics potential and goals of a polarized RHIC collider utilizing two RHIC detectors STAR [5] and PHENIX [6]. Polarized gluon distribution functions will be studied through direct photon production as well as inclusive single and dijet production (see H. Spinka these proceedings). The Drell Yan process is our best candidate to probe the sea quark distributions and a recent work by Bourrely and Soffer [7] suggest  $W^{+/-}$  production as another potential mechanism.

## 2. Physics and Rates

The RSC proposal will use two longitudinally polarized colliding proton beams to measure the double spin asymmetry  $A_{LL}$  defined as the difference between the measured cross sections with parallel and anti parallel beam helicities divided by the sum. This is related to the sum of the respective parton asymmetries of all the sub processes that contribute to the cross section.

$$A_{LL} d\sigma = \sum_{ij} 1/(1+\delta_{ij}) \int dx_a dx_b [ \Delta f_i^a(x) \cdot \Delta f_j^b(x) \cdot \hat{a}_{LL}^{ij} d\sigma_{ij} + (i \leftrightarrow j) ]$$


These measurements result in sums of products of two partonic distributions. For example, in Drell Yan the product is that of  $[\Delta q(x)/q(x)] \cdot [\Delta \bar{q}(x)/\bar{q}(x)]$ . The sea quark distribution is then extracted if the associated valence quark distribution is known.

The proposal assumes 100 days of running at an overall efficiency of 50%, a luminosity of  $2 \times 10^{32} \text{ cm}^{-2} \text{ sec}^{-1}$  at center of mass energy of 500 GeV, and beam polarization of 70%.

### 1. The Drell Yan Process

Lepton pairs from Drell Yan production are detected in both STAR and PHENIX through the process  $pp \Rightarrow \gamma^* + X \Rightarrow l + l^- + X$ . Monte Carlo simulations [8] of the production rates were carried out for both detectors using PYTHIA with the EHLQ1 set of parton structure functions. STAR will detect electron positron pairs using the proposed electromagnetic barrel and endcap calorimeters with acceptance in  $|\eta| \leq 2.0$  and  $\phi = 2\pi$ ; PHENIX will use its barrel calorimeter with coverage of  $|\eta| \leq 0.35$  and  $\phi = \pi$  along with the dimuon endcap that covers  $1.1 \leq \eta \leq 2.5$  and  $\phi = 2\pi$ . Lepton pair production can proceed through several partonic processes, qq annihilation, qq Bremsstrahlung, and qg production, figure 1.

Figure 1. Contributions to the dilepton rates from various partonic processes PHENIX ( $\mu\mu$ ),  $\sqrt{S}=200 \text{ GeV}$ , and  $M_{\mu\mu}=2-3 \text{ GeV}/c^2$ .



A cut on the sum of the  $q$  and  $\bar{q}$  momenta to equal that of the virtual photon eliminates the Bremsstrahlung events and a cut on transverse momentum  $p_T < 1 \text{ GeV}/c$  of the produced pairs assures a sample where the  $q\bar{q}$  annihilation process dominates. Table I shows the event rates available from each detector for different center of mass collision energies.

$M_{ee}, \text{ GeV}/c^2$ ( $\sqrt{s}=200$ )	5-9	9-12	12-15	15-20	20-25
Without Endcaps	33,000	21,000	8,500	5,500	1,900
With Endcaps	71,000	50,000	21,000	13,500	4,300

$M_{\mu\mu}, \text{ GeV}/c^2$	2-5	5-9	9-12
Events ( $\sqrt{s}=200$ )	96,000	23,000	4,000
Events ( $\sqrt{s}=500$ )	240,000	79,000	19,000

Table I. Dilepton rates for both STAR and PHENIX

Good  $e/\pi$  rejection of the order of 100:1 for each lepton is required. In addition, STAR will run with a threshold of  $E_T > 4 \text{ GeV}/c^2$  in order to achieve manageable trigger rates. These simulations do not include contributions from open charm production that has a substantial cross section and the combinatorics from which present a potential background to the Drell Yan measurement.

A rate of 10,000 events per bin allows a 2% statistical error in the asymmetry measurement  $A_{LL}$  when the beam polarization is taken into account. The above table shows that both STAR and PHENIX can achieve such sensitivities for dilepton masses up to 25 and 15  $\text{GeV}/c^2$  respectively. Larger values of  $A_{LL}$  are expected if the kinematic variables are chosen to ensure that the interacting quark from one proton carries an appreciable polarization ( $x_1 > 0.1$ ). Similarly, the momentum fraction carried by the struck antiquark should be away from zero where one expects no polarization. Representative kinematic coverages for both detectors are shown in figure 2. The STAR detector has the better coverage due to its large geometrical acceptance. The kinematic acceptance in PHENIX can be improved by running at lower center of mass energies. However, this is likely to result lower sensitivity.




Figure 2. Diplots of the kinematic regions covered by the PHENIX and STAR detectors respectively. The axes refer to the momentum fractions carried by the interacting partons.

## 2. Measuring the sea polarization via $W^+ / W^-$ production

Bourrely and Soffer [7] suggest using the parity violating asymmetries  $A_L^{PV}$  and  $A_{LL}^{PV}$  in  $W^+$  and  $W^-$  production to get another handle on the sea quark polarization. They predict a large sensitivity to sea quark polarization in  $W^-$  production as well as a strong dependence on rapidity. The effect is most pronounced at  $y = -1$ . Their same calculation for  $A_{LL}$  in  $W^-$  production also reveals a strong dependence on sea quark polarization with a sensitivity that is relatively flat between  $-1 < y < 1$  which is quite desirable for the STAR and PHENIX geometries. The  $W^+$  measurement is used for normalization purposes, Figure 3.




Figure 3. Predictions for Parity violating asymmetries in  $W$  production. The solid lines with sea quark polarization, and the dashed lines with zero polarization.

There is appreciable acceptance for  $W$  production in STAR and PHENIX. The rates were calculated using the same PHYTHIA simulation package described earlier. Table II shows the expected number of events for running at center of mass energy of 500 GEV. These rates provide for sensitivities at the level of a few percent.

Boson	STAR barrel & Endcaps	PHENIX Barrel
$W^+$	80,000	16,000
$W^-$	30,000	6,000

Table II.  $W$  production for STAR and PHENIX

### 3. Transverse quark polarization

The proposed polarized RHIC facility allows for the beam polarization to be rotated so that transverse double spin asymmetries could be measured. These measurements are sensitive to the transverse spin structure functions of the partons within the nucleon that have never been measured. Theoretical calculations [9] are underway to assess the differences between longitudinal and transverse spins. Only the quarks are expected to participate as the gluons cannot carry transverse spin.

Large transverse single spin asymmetries have been observed in inclusive pion production experiments at transverse momenta up to 4 GeV/c, Figure 4. Perturbative QCD calculations expected these effects to be quite small and to vanish at high momentum transfer. However, perturbative calculations to orders next to leading twist do show some promise in explaining these single spin asymmetries.




Figure 4. The single spin asymmetry in inclusive pion production versus transverse momentum.

It would be interesting to extend these measurements to higher transverse momenta. STAR and PHENIX will measure neutral pion production up to  $p_t$  of 20 GeV/c but lack any particle identification for high momentum charged hadrons. A third RHIC detector, The Forward Angle and Mid rapidity Spectrometer [10] can achieve pion, kaon, and proton separation for momenta up to 25 GeV/c albeit with small acceptance. The case for kaons is particularly interesting as these hadrons carry strange valence quarks. These could provide an interesting indication of the polarization of the strange quarks from the parent protons.

## References

\* The RHIC Spin Collaboration: M. Biddo, D. Grosnick, D. Hill, D. Underwood, A. Yokosawa-ANL; G. Bunce, A. Carroll, E. Courant, R. Fernow, Y.Y. Lee, D. Lowenstein, Y. Makdisi, L. Ratner, T. Roser, M. Tannenbaum - BNL; J. B. Carroll, T. Hallman, G. J. Igo- UCLA; N. Tornqvist- Helsinki; N. Akchurin, Y. Onel- Iowa; S.Y. Lee- Indiana; S. Hiramatsu, Y. Mori, H Sato- KEK; L. Madansky, R. Welsh- Johns Hopkins; H. Enyo, K. Imai, A. Masaïke- Kyoto; J. Soffer- Marseille; R. Jaffe- MIT; A Bravar, M. Conte, A. Penzo, M Pusterla, P. Schiavon- Genova, Padova, Trieste; J. Collins, S. Heppelmann, G. Ladinsky, E. Minor, R. Robinett- Penn. State; Y. Arestov, B. Chuiko, A. Davidenko, A. Derevschikov, O. Grachov, A. Likhoded, A. Messhanin, S. Nurushev, V. Rykov, A. Ufimtsev, A. Vasiliev- IHEP Protvino; A. Efremov- Dubna; M. Werlen- Lausanne.

- [1] SLAC E-80, M.J. Alguard et al., Phys. Rev. Lett. 37, (1976) 1261; *ibid.* 41 (1978) 70; SLAC E-130 G.Baum et al., Phys Rev. Lett. 45 (1980) 2000; *ibid.* 51 (1983) 1135.
- [2] EMC, J. Ashman et al., Phys. Lett. B206 (1988) 364; J. Ashman et al., Nucl. Phys. B328 (1989) 1.
- [3] SMC, B. Adeva et al, Measurement of the Spin Dependent Structure Function  $g_1(x)$  of The Deuteron. CERN-PPE/93-47, 8-Mar-1993. Submitted to Phys. Lett. B.
- [4] M. Biddo et al., BNL Proposal on Spin Physics Using the RHIC Polarized Collider, 14-Aug-1993.
- [5] M. E. Biddo et al., The STAR conceptual Design Report , 15-June-1992.
- [6] J. C. Gregory et al. PHENIX Conceptual Design Report , 29-Jan-1993
- [7] C. Bourrely and J. Soffer, Parton Distributions and Parity Violating Asymmetries in  $W^{+/-}$  and Z Production at RHIC. Preprint CPT-93/ P.2865, Jan 1993.
- [8] A. Derevschikov and V. Rykov, Notes on the Drell Yan Pairs, Z and W production in STAR and PHENIX at RHIC, RSC-BNL/IHEP -4 report. Aug 1992, unpublished.
- [9] R. L. Jaffe and X. Ji, Phys. Rev. Lett. 67 (1991) 552.
- [10] D. Beavis et al., Forward Angle and Mid rapidity Hadron Spectrometer at RHIC. Letter of Intent # 8.

**DATE  
FILMED**

8 / 31 / 93

**END**

