



Fermi National Accelerator Laboratory

FNAL/C--92/174

DE92 018636

FERMILAB-Conf-92/174

RECEIVED BY OSTI  
AUG 3 1992

# Update on Hadroproduced Charm at TPL

Keith Thorne

*Fermi National Accelerator Laboratory  
P.O. Box 500, Batavia, Illinois 60510*

June 1992

Presented at the *XXVIIth Recontres de Moriond, QCD and High Energy interactions*, Les Arcs, France, March 22-28, 1992

MASTER

## **Disclaimer**

*This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.*

# Update on Hadroproduced Charm at TPL

Keith Thorne

*Fermi National Accelerator Laboratory*

*Batavia, IL USA*

Representing the E-769\* and E-791† Collaborations

## Abstract

Two experiments have now been run at Fermilab using the Tagged Photon Laboratory (TPL) spectrometer with an incident hadron beam to study heavy quark physics. Results (preliminary) from the first experiment, E-769, on charm hadroproduction dependence on the target atomic number,  $x_F$  and  $p_t$  are presented. The next experiment, E-791, just completed data-taking with an upgraded spectrometer and data-acquisition system to collect a high-statistics sample of charm decays. Preliminary plots and estimates of final sample size are presented

---

Centro Brasileiro de Pesquisas Físicas\*†, University of California at Santa Cruz†, University of Cincinnati†, Fermilab\*†, Illinois Institute of Technology†, University of Mississippi\*†, Ohio State University†, Princeton University†, Universidade Federal do Rio de Janeiro†, Tel Aviv University†, University of Toronto\*, Tufts University\*†, University of Wisconsin\*†, Yale University\*†

## 1 Introduction

The Tagged Photon Laboratory spectrometer is of a now conventional open geometry design for fixed-target heavy quark physics. Following a small nuclear target are up to 17 silicon microstrip detectors for finding decay vertices from heavy quark states, two magnets and 37 drift and proportional wire chambers for momentum determination, two gas threshold Cerenkov detectors for particle identification, electromagnetic and hadronic calorimeters, and muon counters behind a shielding wall. The previous experiment, E-691, used a tagged photon beam to produce heavy quark particles. The beamline has since been converted to a high-energy hadron beam. This allowed for both the study of charm hadroproduction, and an expected increase in the total number of charm decays reconstructed.

The common philosophy in these experiments is to use an open, efficient trigger for charm, delaying the application of more restrictive criteria until the off-line analysis. Such a technique both saves such event selection until after all the detector calibrations are made, and also uses cheaper, general-purpose computers, rather than expensive, single-purpose trigger processors. The trigger used requires a certain minimum amount of transverse energy be deposited in the calorimeters. To use this method, a high-rate data acquisition system is required to log the data onto tape.

## 2 E-769 Results

The initial experiment, E-769, concentrated on the study of charm hadroproduction. They used 250 GeV/c positive and negative hadron beams, with the beam particle type tagged by a differential Cerenkov counter and a transition radiation detector. This tagging was used to enhance the number of triggers from the kaon component of the beam. A total of 370 million triggers were recorded, with about 4,000 charm events being fully reconstructed.

The dependence of the charm cross-section with the atomic weight ( $A$ ) of the target material provides information on the distance scale of the production. E-769 used a target made up of foils of beryllium, aluminum, copper and tungsten to study this scaling, which is fit to  $A^\alpha$ . Their preliminary result <sup>1)</sup> for the  $D^+$  is  $\alpha = 1.04 \pm 0.08$  and for the  $D^0$  it is  $\alpha = 0.99 \pm 0.10$ , both from negative  $\pi^-$  data (Figure 1). These are consistent with a hard-scattering process, which requires  $\alpha = 1$ . They are now studying the  $x_F$  dependence of this A-dependence.

The experiment has also measured the basic  $x_F$  and  $p_t$  dependence of the D meson production. They fit the differential cross-section  $d\sigma/dx_F$  as  $(1 - x_F)^n$ . For the  $D^+$  they find  $n = 3.8 \pm 0.4$ , and that  $n = 4.13 \pm 0.32$  for the  $D^0$ . As seen in Figure 2, this dependence agrees with the Nason, Dawson, Ellis<sup>2)</sup> model for charm quark production, but disagrees when fragmentation models based on  $e^+e^-$  data are convoluted<sup>3)</sup> with the quark production. This implies that the D mesons carry away nearly all the momentum of the charm quark. The differential cross-section as a function of  $p_t$  was first fit to  $e^{-bp_t^2}$ . The results were  $b = 0.98 \pm 0.07$  for  $D^+$  and  $b = 0.95 \pm 0.09$  for  $D^0$ . As is shown in Figure 3, at high  $p_t$  the cross-section is fit better as  $e^{-b'p_t}$ .

E-769 has seen good signals for all the beam particle types ( $\pi$ ,  $K$  and  $p$ ). They




Figure 1: A-dependence of  $D^+$  and  $D^0$  production from 250 GeV/c  $\pi^-$  1)




Figure 2:  $x_F$  dependence of  $D^+$  production. The solid line is a fit to  $(1 - x_F)^n$ . The dashed line is the Nason, Dawson, Ellis prediction <sup>2)</sup> for charm quark production, which is in good agreement. The dotted line is with fragmentation functions from  $e^+e^-$  data convoluted with it <sup>3)</sup>.




Figure 3:  $p_t$  dependence of  $D^+$  production from 250 GeV/c  $\pi^-$ . The dashed line is a fit to  $e^{-bp_t^2}$ . The solid line, a fit to  $e^{-b'p_t}$ , is in better agreement with the data at large  $p_t$ .




Figure 4: Distribution of reconstructed primary vertices from E-791 data, showing cleanly-separated target profiles

are now studying the relative D meson cross-sections, and their  $x_F$  and  $p_t$  dependence.

### 3 E-791 Results

E-791 also used the high-energy hadron beam, but this time with an emphasis on charm decay physics. E-791 used the experience of E-769 to make upgrades which would enhance the precision and charm yield of the apparatus. The beam energy was raised to 500 GeV, and more silicon microstrip planes were added to improve vertexing. The biggest change was the large increase in the data-taking rate. The readout deadtime was reduced to  $50\mu s$  using fast digitizing electronics, and a highly-parallel DA system was built to record data at 9 Megabytes/s. This enabled us to record 20 billion events during the 1991 fixed-target run at Fermilab.

Preliminary analyses of some E-791 data have been made. A plot of reconstructed primary vertices (See Figure 4) reveals that the 5 target discs (1 platinum, 4 diamond) are cleanly separated. This indicates how well we can separate true secondary decay vertices between the targets from secondary interactions within the targets. We have run 100 tapes (0.5 % of our sample) through a preliminary reconstruction and have observed  $D^+ \rightarrow K\pi\pi$ ,  $D^0 \rightarrow K\pi$ , and  $D^* \rightarrow D^0\pi$  signals without using particle identification from the Cerenkov detectors. From these signals, we now estimate that we will have at least 100,000 fully reconstructed charm decays from our total sample.

### References

1. Gay, C. Ph.D. Dissertation, University of Toronto, Toronto, Ont., CANADA (unpublished)
2. Nason, P., Dawson, S., Ellis, R., *Nucl. Phys.* **263B**(1986) 36
3. Wu, Z. Ph.D. Dissertation, Yale University, New Haven, Ct., USA (unpublished)

**END**

**DATE  
FILMED**

9/04/92

