

10
11/28/94 48 S(1)

Conf-4409102-

SLAC-PUB-6699
OREXP-94-6
September 1994
(E/T)

Electroweak Coupling Measurements from Polarized Bhabha Scattering at SLD

The SLD Collaboration*

*Stanford Linear Accelerator Center
Stanford University, Stanford, California 94309*

Represented by

Kevin T. Pitts†

*Department of Physics, University of Oregon
Eugene, OR 97403*

Abstract

The cross section for Bhabha scattering ($e^+e^- \rightarrow e^+e^-$) with polarized electrons at the center of mass energy of the Z^0 resonance has been measured with the SLD experiment at the SLAC Linear Collider (SLC) during the 1992 and 1993 runs. The first measurement of the left-right asymmetry in Bhabha scattering ($A_{LR}^{e^+e^-}(\theta)$) is presented. From $A_{LR}^{e^+e^-}(\theta)$ the effective weak mixing angle is measured to be $\sin^2\theta_W^{\text{eff}} = 0.2245 \pm 0.0049 \pm 0.0010$. When combined with the measurement of A_{LR} , the effective electron couplings are measured to be $v_e = -0.0414 \pm 0.0020$ and $a_e = -0.4977 \pm 0.0045$.

*Presented at the 11th International Symposium on High Energy Spin Physics (SPIN 94),
Bloomington, Indiana, September 15-22, 1994.*

This work was supported by Department of Energy contracts: DE-FG02-91ER40676 (BU), DE-FG03-92ER40701 (CIT), DE-FG03-91ER40818 (UCSB), DE-FG02-91ER40672 (Colorado), DE-FG02-91ER40877 (Illinois), DE-FG03-91ER40861 (Indiana), DE-AC03-76SF00098 (LBL), DE-FG02-92ER40718 (Massachusetts), DE-AC02-74ER03069 (MIT), DE-FG02-92ER40226 (Oregon), DE-AC03-76SF00818 (SLAC), DE-FG08-91ER40627 (Texas), DE-AC02-74ER03081 (Wisconsin), DE-FG02-92ER40704 (Yale); National Science Foundation grants: PHY-91-13426 (UCSC), PHY-88-21230 (Columbia), PHY-92-05239 (Cincinnati), PHY-88-17930 (Rutgers), PHY-88-19316 (Vanderbilt), PHY-92-03212 (Washington); the UK Science and Engineering Research Council (Daresbury and RAL); the Istituto Nazionale di Fisica Nucleare of Italy (Bologna, Ferrara, Frascati, Pisa, Padova, Perugia); the Natural Sciences and Engineering Research Council of Canada (British Columbia, Victoria, TRIUMF); and the Japan-UE Cooperative Research Project in High Energy Physics (KEK, Nagoya, Tohoku).

*current address: Fermilab, P.O. Box 500, Batavia, IL 60510

-5

The SLD Collaboration has recently performed the most precise single measurement of the effective electroweak mixing angle, $\sin^2\theta_W^{\text{eff}}$, by measuring the left-right cross section asymmetry (A_{LR}) in Z boson production at the Z^0 resonance [1]. The left-right cross section asymmetry is a measure of the initial state electron coupling to the Z^0 , which allows all visible fermion final states to be included in the measurement. For simplicity, the e^+e^- final state (Bhabha scattering) is omitted in the A_{LR} measurement due to the dilution of the asymmetry from the large QED contribution of the t-channel photon exchange. Here, two new results are presented: the first measurement of the left-right cross section asymmetry in polarized Bhabha scattering ($A_{LR}^{e^+e^-}(|\cos\theta|)$), and measurements of the effective electron coupling parameters based on a combined analysis of the A_{LR} measurement [1] and the Bhabha cross section and angular distributions. The vector coupling measurement is the most precise yet presented [2].

In the Standard Model, measuring the left-right asymmetry yields a value for the quantity A_e , a measure of the degree of parity violation in the neutral current, since:

$$A_{LR} = A_e = \frac{2v_e a_e}{v_e^2 + a_e^2} = \frac{2[1 - 4\sin^2\theta_W^{\text{eff}}]}{1 + [1 - 4\sin^2\theta_W^{\text{eff}}]^2}, \quad (1)$$

where the effective electroweak mixing parameter is defined as $\sin^2\theta_W^{\text{eff}} = \frac{1}{4}(1 - v_e/a_e)$, and v_e and a_e are the effective vector and axial vector electroweak coupling parameters of the electron. The partial width for Z^0 decaying in e^+e^- is dependent on the coupling parameters:

$$\Gamma_{ee} = \frac{G_F M_Z^3}{6\sqrt{2}\pi} (v_e^2 + a_e^2)(1 + \delta_e), \quad (2)$$

where $\delta_e = \frac{3\alpha}{4\pi}$ is the correction for final state radiation. G_F is the Fermi coupling constant and M_Z is the Z^0 boson mass. By measuring A_e and Γ_{ee} , the above equations can be utilized to extract v_e and a_e .

Event selection is calorimetry-based and makes use of the distinct topology of the e^+e^- final state. The efficiency and contamination for the wide angle events are calculated from Monte Carlo simulations. Corrections are applied as a function of scattering angle to account for angle-dependent changes in response.

MASTER

2

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Table 1: Number of accepted events for the 1992 run. ($\langle \mathcal{P}_e \rangle = 22.4\%$)

region	left-handed	right-handed	$A_{LR}^{e^+e^-}(\text{raw})$
$0.0 < \cos\theta_{CM} < 0.70$	157	137	0.068 ± 0.058
$0.70 < \cos\theta_{CM} < 0.94$	208	205	0.0073 ± 0.049
$0.94 < \cos\theta_{CM} < 0.98$	305	318	-0.021 ± 0.040
$0.998 < \cos\theta_{CM} < 0.9994$	12,395	12,353	0.0017 ± 0.0064

 Table 2: Number of accepted events for the 1993 run. ($\langle \mathcal{P}_e \rangle = 63.0\%$)

region	left-handed	right-handed	$A_{LR}^{e^+e^-}(\text{raw})$
$0.0 < \cos\theta_{CM} < 0.70$	864	702	0.103 ± 0.0253
$0.70 < \cos\theta_{CM} < 0.94$	1,039	946	0.047 ± 0.022
$0.94 < \cos\theta_{CM} < 0.98$	1,566	1,479	0.029 ± 0.018
$0.998 < \cos\theta_{CM} < 0.9996$	93,727	94,319	-0.0032 ± 0.0023

Tables I and II show the number of events accepted, by beam helicity, for the 1992 and 1993 SLC runs. The raw asymmetry is defined as:

$$\bar{A}_{LR}^{e^+e^-}(\theta) = \langle \mathcal{P}_e \rangle A_{LR}^{e^+e^-}(\theta) = (N_L - N_R)/(N_L + N_R),$$

where $N_L(N_R)$ is the number of events tagged with a left-(right-) handed electron beam as a function of the $|\cos\theta|$, where θ is the center-of-mass scattering angle for the e^+e^- system after initial state radiation. Aside from the charge ambiguity which is unresolved by the calorimeter measurement, the center-of-mass scattering angle is derived trivially from the measured electron and positron laboratory scattering angles. The angular regions in the table are chosen to emphasize the different regimes of the $e^+e^- \rightarrow e^+e^-$ distribution: for $|\cos\theta| < 0.7$ the s-channel Z^0 decay dominates; from 0.7 to 0.94 the s-channel Z^0 decay, the t-channel photon exchange and the interference between those two interactions all contribute; for $|\cos\theta| > 0.94$, the t-channel photon exchange dominates. The region of $0.998 < |\cos\theta| < 0.9996$ is that which is covered by the small angle silicon/tungsten luminosity mon-

itor (LUM). The expected asymmetry ($A_{LR}^{e^+e^-}(\theta)$) is largest at $\cos\theta = 0$, and may be approximately written as $A_{LR}^{e^+e^-}(\theta) = A_e(1 - f_t(|\cos\theta|))$, where $f_t(|\cos\theta|)$ represents the t-channel contribution. For the region $|\cos\theta| < 0.7$, $\langle f_t \rangle \simeq 0.12$. The expected asymmetry falls to very small values ($\sim 10^{-4}$) in the small angle region where the t-channel photon exchange dominates.




Figure 1: Differential angular distribution for $e^+e^- \rightarrow e^+e^-$. The points are the corrected data, the dashed line is the fit.

To extract Γ_{ee} and A_e , the data are fit to the differential e^+e^- cross section using the maximum likelihood method. Two programs are used to calculate the differential e^+e^- cross section: EXPOSTAR [4] and, as a cross check, DMIBA [5]. The EXPOSTAR program calculates the differential cross sections within the framework of the Standard Model. The DMIBA program calculates the differential e^+e^- cross section in a model independent manner. To extract the maximal amount of information from the differential polarized Bhabha scattering distribution, the fit is performed over the entire angular region accepted by the liquid argon calorimeter (LAC), where $|\cos\theta| < 0.98$. No t-channel subtraction is performed. All ten lowest order terms in the

cross section are included in the fit: the four pure s-channel and t-channel terms for photon and Z^0 exchange, and the six interference terms [6]. The fit also includes initial state radiation. Since the measurement is calorimetric it is insensitive to final state radiation.

The partial width Γ_{ee} is extracted from the data in two ways: (1) using the full fit to the differential cross section for $|\cos\theta| \leq 0.98$, and (2) measuring the cross section in the central region ($|\cos\theta| < 0.6$) where the systematic errors are smaller, yielding a more precise measurement. For the fits we use $M_Z = 91.187$ GeV/c² and $\Gamma_Z = 2.489$ GeV/c² [7]. Figure 1 shows the fit to the full $e^+e^- \rightarrow e^+e^-$ distribution, which yields $\Gamma_{ee} = 83.14 \pm 1.03$ (stat) ± 1.95 (sys) MeV. The 2.4% systematic error is dominated (2.1%) by the uncertainty in the efficiency correction factors in the angular region $0.6 < |\cos\theta| < 0.98$, where the LAC response is difficult to model due to materials from interior detector elements [3].




Figure 2: Left-right asymmetry, $\hat{A}_{LR}^{e^+e^-}(|\cos\theta|)$ for polarized $e^+e^- \rightarrow e^+e^-$. The points are the corrected data, the solid curve is the fit.

A more precise determination of Γ_{ee} was performed using only the central

region of the LAC ($|\cos\theta| < 0.6$) and the small angle region in the LUM [8]. The program MIBA [9] is then used to calculate Γ_{ee} based on the total measured cross section within the defined fiducial region. From this method, we find:

$$\Gamma_{ee} = 82.89 \pm 1.20 \text{ (stat)} \pm 0.89 \text{ (sys)} \text{ MeV.}$$

The loss in statistical precision of the limited fiducial region is more than compensated by the improvement in the systematic error. The 1.1% systematic error is dominated by the accuracy of the detector simulation (0.74%) and the uncertainty in the absolute luminosity (0.52%).

To extract A_e from the Bhabha events, the right- and left-handed differential $e^+e^- \rightarrow e^+e^-$ cross sections are fit directly for v_e and a_e using EXPOSTAR. This yields

$$A_e = 0.202 \pm 0.038 \text{ (stat)} \pm 0.008 \text{ (sys).}$$

Figure 2 shows the measured left-right cross section asymmetry for $e^+e^- \rightarrow e^+e^-$ ($A_{LR}^{e^+e^-}(|\cos\theta|)$) compared to the fit. The measurement of A_e is limited by the statistical uncertainty. The 3.8% systematic is dominated by a 3.2% uncertainty in the angle-dependent response correction factors. The polarization uncertainty contributes 1.7% and asymmetry factors from the SLC contribute 0.06% as discussed in Refs. [1] and [3].

The results for Γ_{ee} and A_e from above may now be used in equations 1 and 2 to extract the effective vector and axial vector couplings to the Z^0 : $v_e = -0.0507 \pm 0.0096 \text{ (stat)} \pm 0.0020 \text{ (sys)}$, $a_e = -0.4968 \pm 0.0039 \text{ (stat)} \pm 0.0027 \text{ (sys)}$, where lower energy e^+e^- annihilation data have been utilized to assign $|v_e| < |a_e|$, and $v_e e$ scattering data have been utilized to establish $v_e < 0$ and $a_e < 0$ [10]. Figure 3 shows the one standard deviation (68%) contour for these electron vector and axial vector coupling measurements. Most of the sensitivity to the electron vector coupling and, hence, $\sin^2\theta_W^{\text{eff}}$ arises from the measurement of A_e , while the sensitivity to the axial vector coupling arises from Γ_{ee} . Also shown are standard model calculations using the program ZFITTER [11].

The effective electroweak mixing angle represented by these vector and axial vector couplings is:

$$\sin^2\theta_W^{\text{eff}} = 0.2245 \pm 0.0049 \text{ (stat)} \pm 0.0010 \text{ (sys).}$$

We reiterate that this measurement derives strictly from the Bhabha events.




Figure 3: One standard deviation (68%) contour in the a_e , v_e plane. The large ellipse is for $e^+e^- \rightarrow e^+e^-$, the smaller ellipse includes the measurement of A_{LR} . The hatched region shows the Standard Model calculation as a function of the mass of the top quark. The width of the hatched region is the variation due to the uncertainty in the Higgs mass.

The SLD Collaboration has published a more precise measurement of A_e from the left-right cross section asymmetry (A_{LR}) measurement [1]. Combining the Bhabha results with the SLD measurement of A_{LR} gives:

$$v_e = -0.0414 \pm 0.0020 \quad a_e = -0.4977 \pm 0.0045,$$

the most precise measurement of the electron vector coupling to the Z^0 published to date. The v_e , a_e contour including the A_{LR} measurement is also shown in Figure 3, demonstrating the increased sensitivity in v_e from A_{LR} .

We thank the personnel of the SLAC accelerator department and the technical staffs of our collaborating institutions for their outstanding efforts on our behalf.

References

- [1] SLD Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **73**, 25 (1994).
- [2] SLD Collaboration, K. Abe *et al.*, SLAC-PUB-6605, August 1994.
- [3] K.T. Pitts, Ph.D. Thesis, University of Oregon, SLAC Report 446 (1994).
- [4] D. Levinthal, F. Bird, R.G. Stuart and B.W. Lynn, Z. Phys. C **53**, 617 (1992).
- [5] P. Comas and M. Martinez, Z. Phys. C **58**, 15 (1993).
- [6] M. Greco, Nucl. Phys. **B177**, 97 (1986).
- [7] The LEP Collaborations, Report No. CERN-PPE-93-157, August, 1993.
- [8] J.M. Yamartino, Ph.D. Thesis, MIT, SLAC Report 426, February 1994.
- [9] M. Martinez and R. Miquel, Z. Phys. C **53**, 115 (1992).
- [10] S.L. Wu, Phys. Rep. **107**, 59 (1984).
- [11] D. Bardin *et al.*, Report No. CERN-TH-6443-92, May 1992.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

*The SLD Collaboration

K. Abe,⁽²⁸⁾ I. Abt,⁽¹⁴⁾ T. Akagi,⁽²⁶⁾ W.W. Ash,⁽²⁶⁾ D. Aston,⁽²⁶⁾
N. Bacchetta,⁽²¹⁾ K.G. Baird,⁽²⁴⁾ C. Baltay,⁽³²⁾ H.R. Band,⁽³¹⁾
M.B. Barakat,⁽³²⁾ G. Baranko,⁽¹⁰⁾ O. Bardon,⁽¹⁶⁾ T. Barklow,⁽²⁶⁾
A.O. Bazarko,⁽¹¹⁾ R. Ben-David,⁽³²⁾ A.C. Benvenuti,⁽²⁾ T. Bierenz,⁽²⁶⁾
G.M. Bilei,⁽²²⁾ D. Bisello,⁽²¹⁾ G. Blaylock,⁽⁷⁾ J.R. Bogart,⁽²⁶⁾ T. Bolton,⁽¹¹⁾
G.R. Bower,⁽²⁶⁾ J.E. Brau,⁽²⁰⁾ M. Breidenbach,⁽²⁶⁾ W.M. Bugg,⁽²⁷⁾
D. Burke,⁽²⁶⁾ T.H. Burnett,⁽³⁰⁾ P.N. Burrows,⁽¹⁶⁾ W. Busza,⁽¹⁵⁾
A. Calcaterra,⁽¹³⁾ D.O. Caldwell,⁽⁶⁾ D. Calloway,⁽²⁶⁾ B. Camanzi,⁽¹²⁾
M. Carpinelli,⁽²³⁾ R. Cassell,⁽²⁶⁾ R. Castaldi,⁽²³⁾ A. Castro,⁽²¹⁾
M. Cavalli-Sforza,⁽⁷⁾ E. Church,⁽³⁰⁾ H.O. Cohn,⁽²⁷⁾ J.A. Coller,⁽³⁾
V. Cook,⁽³⁰⁾ R. Cotton,⁽⁴⁾ R.F. Cowan,⁽¹⁶⁾ D.G. Coyne,⁽⁷⁾ A. D'Oliveira,⁽⁸⁾
C.J.S. Damerell,⁽²⁵⁾ S. Dasu,⁽²⁶⁾ R. De Sangro,⁽¹³⁾ P. De Simone,⁽¹³⁾
R. Dell'Orso,⁽²³⁾ M. Dima,⁽⁹⁾ P.Y.C. Du,⁽²⁷⁾ R. Dubois,⁽²⁶⁾ B.I. Eisenstein,⁽¹⁴⁾
R. Elia,⁽²⁶⁾ D. Falciai,⁽²²⁾ C. Fan,⁽²⁾ M.J. Fero,⁽¹⁶⁾ R. Frey,⁽²⁰⁾ K. Furuno,⁽²⁰⁾
T. Gillman,⁽²⁵⁾ G. Gladding,⁽¹⁴⁾ S. Gonzalez,⁽¹⁶⁾ G.D. Hallewell,⁽²⁶⁾
E.L. Hart,⁽²⁷⁾ Y. Hasegawa,⁽²⁸⁾ S. Hedges,⁽⁴⁾ S.S. Hertzbach,⁽¹⁷⁾
M.D. Hildreth,⁽²⁶⁾ J. Huber,⁽²⁰⁾ M.E. Huffer,⁽²⁶⁾ E.W. Hughes,⁽²⁶⁾
H. Hwang,⁽²⁰⁾ Y. Iwasaki,⁽²⁸⁾ P. Jacques,⁽²⁴⁾ J. Jaros,⁽²⁶⁾ A.S. Johnson,⁽³⁾
J.R. Johnson,⁽³¹⁾ R.A. Johnson,⁽⁸⁾ T. Junk,⁽²⁶⁾ R. Kajikawa,⁽¹⁹⁾
M. Kalelkar,⁽²⁴⁾ I. Karliner,⁽¹⁴⁾ H. Kawahara,⁽²⁶⁾ H.W. Kendall,⁽¹⁶⁾
M.E. King,⁽²⁶⁾ R. King,⁽²⁶⁾ R.R. Kofler,⁽¹⁷⁾ N.M. Krishna,⁽¹⁰⁾ R.S. Kroeger,⁽¹⁸⁾
J.F. Labs,⁽²⁶⁾ M. Langston,⁽²⁰⁾ A. Lath,⁽¹⁶⁾ J.A. Lauber,⁽¹⁰⁾ D.W.G. Leith,⁽²⁶⁾
X. Liu,⁽⁷⁾ M. Loretti,⁽²¹⁾ A. Lu,⁽⁶⁾ H.L. Lynch,⁽²⁶⁾ J. Ma,⁽³⁰⁾ G. Mancinelli,⁽²²⁾
S. Manly,⁽³²⁾ G. Mantovani,⁽²²⁾ T.W. Markiewicz,⁽²⁶⁾ T. Maruyama,⁽²⁶⁾
R. Massetti,⁽²²⁾ H. Masuda,⁽²⁶⁾ E. Mazzucato,⁽¹²⁾ A.K. McKemey,⁽⁴⁾
B.T. Meadows,⁽⁸⁾ R. Messner,⁽²⁶⁾ P.M. Mockett,⁽³⁰⁾ K.C. Moffit,⁽²⁶⁾
B. Mours,⁽²⁶⁾ G. Müller,⁽²⁶⁾ D. Müller,⁽²⁶⁾ T. Nagamine,⁽²⁶⁾ U. Nauenberg,⁽¹⁰⁾
H. Neal,⁽²⁶⁾ M. Nussbaum,⁽⁸⁾ Y. Ohnishi,⁽¹⁹⁾ L.S. Osborne,⁽¹⁶⁾
R.S. Panvini,⁽²⁹⁾ H. Park,⁽²⁰⁾ T.J. Pavel,⁽²⁶⁾ I. Peruzzi,⁽¹³⁾ L. Pescara,⁽²¹⁾
M. Piccolo,⁽¹³⁾ L. Piemontese,⁽¹²⁾ E. Pieroni,⁽²³⁾ K.T. Pitts,⁽²⁰⁾ R.J. Piano,⁽²⁴⁾
R. Prepost,⁽³¹⁾ C.Y. Prescott,⁽²⁶⁾ G.D. Punkar,⁽²⁶⁾ J. Quigley,⁽¹⁶⁾
B.N. Ratcliff,⁽²⁶⁾ T.W. Reeves,⁽²⁹⁾ P.E. Rensing,⁽²⁶⁾ L.S. Rochester,⁽²⁶⁾
J.E. Rothberg,⁽³⁰⁾ P.C. Rowson,⁽¹¹⁾ J.J. Russell,⁽²⁶⁾ O.H. Saxton,⁽²⁶⁾
T. Schalk,⁽⁷⁾ R.H. Schindler,⁽²⁶⁾ U. Schneekloth,⁽¹⁶⁾ B.A. Schumm,⁽¹⁵⁾
A. Seiden,⁽⁷⁾ S. Sen,⁽³²⁾ M.H. Shaevitz,⁽¹¹⁾ J.T. Shank,⁽³⁾ G. Shapiro,⁽¹⁵⁾
S.L. Shapiro,⁽²⁶⁾ D.J. Sherden,⁽²⁶⁾ C. Simopoulos,⁽²⁶⁾ H.J. Simpson,⁽²⁶⁾
N.B. Sinev,⁽²⁰⁾ S.R. Smith,⁽²⁶⁾ J.A. Snyder,⁽³²⁾ M.D. Sokoloff,⁽⁸⁾ P. Stamer,⁽²⁴⁾

H. Steiner,⁽¹⁵⁾ R. Steiner,⁽¹⁾ M.G. Strauss,⁽¹⁷⁾ D. Su,⁽²⁶⁾ F. Suekane,⁽²⁸⁾
A. Sugiyama,⁽¹⁹⁾ S. Suzuki,⁽¹⁹⁾ M. Swartz,⁽²⁶⁾ A. Szumilo,⁽³⁰⁾ T. Takahashi,⁽²⁶⁾
F.E. Taylor,⁽¹⁶⁾ A. Tolstykh,⁽²⁶⁾ E. Torrence,⁽¹⁸⁾ J.D. Turk,⁽³²⁾ T. Usher,⁽²⁶⁾
J. Va'vra,⁽²⁶⁾ C. Vannini,⁽²³⁾ E. Vella,⁽²⁶⁾ J.P. Venuti,⁽²⁹⁾ P.G. Verdini,⁽²³⁾
S.R. Wagner,⁽²⁶⁾ A.P. Waite,⁽²⁶⁾ S.J. Watts,⁽⁴⁾ A.W. Weidemann,⁽²⁷⁾
J.S. Whitaker,⁽³⁾ S.L. White,⁽²⁷⁾ F.J. Wickens,⁽²⁵⁾ D.A. Williams,⁽⁷⁾
D.C. Williams,⁽¹⁶⁾ S.H. Williams,⁽²⁶⁾ S. Willocq,⁽³²⁾ R.J. Wilson,⁽⁹⁾
W.J. Wisniewski,⁽⁵⁾ M. Woods,⁽²⁶⁾ G.B. Word,⁽²⁴⁾ J. Wyss,⁽²¹⁾
R.K. Yamamoto,⁽¹⁶⁾ J.M. Yamartino,⁽¹⁶⁾ S.J. Yellin,⁽⁶⁾ C.C. Young,⁽²⁶⁾
H. Yuta,⁽²⁸⁾ G. Zapalac,⁽³¹⁾ R.W. Zdarko,⁽²⁶⁾ C. Zeitlin,⁽²⁰⁾ and J. Zhou⁽²⁰⁾

(1) *Adelphi University, Garden City, New York 11580*

(2) *INFN Sezione di Bologna, I-40126 Bologna, Italy*

(3) *Boston University, Boston, Massachusetts 02215*

(4) *Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom*

(5) *California Institute of Technology, Pasadena, California 91125*

(6) *University of California at Santa Barbara, Santa Barbara, California 93106*

(7) *University of California at Santa Cruz, Santa Cruz, California 95064*

(8) *University of Cincinnati, Cincinnati, Ohio 45221*

(9) *Colorado State University, Fort Collins, Colorado 80523*

(10) *University of Colorado, Boulder, Colorado 80309*

(11) *Columbia University, New York, New York 10027*

(12) *INFN Sezione di Ferrara and Università di Ferrara, I-44100 Ferrara, Italy*

(13) *INFN Lab. Nazionali di Frascati, I-00044 Frascati, Italy*

(14) *University of Illinois, Urbana, Illinois 61801*

(15) *Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720*

(16) *Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

(17) *University of Massachusetts, Amherst, Massachusetts 01003*

(18) *University of Mississippi, University, Mississippi 38677*

(19) *Nagoya University, Chikusa-ku, Nagoya 464 Japan*

(20) *University of Oregon, Eugene, Oregon 97403*

(21) *INFN Sezione di Padova and Università di Padova, I-35100 Padova, Italy*

(22) *INFN Sezione di Perugia and Università di Perugia, I-06100 Perugia, Italy*

(23) *INFN Sezione di Pisa and Università di Pisa, I-56100 Pisa, Italy*

(24) *Rutgers University, Piscataway, New Jersey 08855*

(25) *Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX United Kingdom*

(26) *Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309*

(27) *University of Tennessee, Knoxville, Tennessee 37998*

(28) *Tohoku University, Sendai 980 Japan*

(29) *Vanderbilt University, Nashville, Tennessee 37235*

(30) *University of Washington, Seattle, Washington 98195*

(31) *University of Wisconsin, Madison, Wisconsin 53706*

(32) *Yale University, New Haven, Connecticut 06511*