

Systems biology approach to bioremediation

Romy Chakraborty¹, Cindy H Wu¹ and Terry C Hazen^{1,2,3}

Bioremediation has historically been approached as a 'black box' in terms of our fundamental understanding. Thus it succeeds and fails, seldom without a complete understanding of why. Systems biology is an integrated research approach to study complex biological systems, by investigating interactions and networks at the molecular, cellular, community, and ecosystem level. The knowledge of these interactions within individual components is fundamental to understanding the dynamics of the ecosystem under investigation. Understanding and modeling functional microbial community structure and stress responses in environments at all levels have tremendous implications for our fundamental understanding of hydrobiogeochemical processes and the potential for making bioremediation breakthroughs and illuminating the 'black box'.

Addresses

¹ Ecology Department, Earth Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA.

² Department of Civil & Environmental Engineering and Center for Environmental Biotechnology, The University of Tennessee, 676 Dabney Hall, Knoxville, TN 37966-1605, USA

³ Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, USA

All authors are partially supported by the U.S. Department of Energy and LBNL under Contract No. DE-AC02-05CH11231.



Introduction

Bioremediation, a process mediated by microorganisms, is a sustainable way to degrade and detoxify environmental contaminants. Though bioremediation has been used to varying degrees for more than 60 years, for example petroleum land farming, it historically has been implemented as a very 'black box' engineering solution where amendments are added and the pollutants are degraded. This approach is often successful but all too often the results are less than desirable, that is, no degradation of the contaminant or even production of more toxic daughter products. The key to successful bioremediation is to harness the naturally occurring catabolic capability of microbes to catalyze transformations of environmental pollutants. Simulated experiments using defined microbial consortia in the laboratory is a great starting point in


providing crucial initial indication (within certain constraints) of the process. However, unlike bench-scale simulations, in situ bioremediation in reality is a complex phenomenon involving more than one contaminant and mediated by different strains of microbes involving different metabolic pathways, across geochemical gradients, geophysical and hydrological complexities.

Systems biology approach

Recently, modern tools of genomics, transcriptomics, proteomics, metabolomics, phenomics, and lipidomics have been applied to investigate systems biology of microbial communities in a myriad of environments (Figure 1). Systems biology is an integrated research approach to study complex biological systems, by investigating interactions and networks at the molecular, cellular, community, and ecosystem levels. Amalgamation of the results from the various 'omics' tools has provided crucial insights into the survival, metabolism and interaction of microbes in their native environments including groundwater and marine systems [1–4], extreme milieus [5], deep-sea sediments and vents [6,7], and animal microbiomes [8]. A systems biology approach is being adopted to unravel key processes to understand, optimize, predict and evaluate microbial function and survival strategies in the ecosystem of interest. However, successful application of this approach requires overcoming several challenges, including the high cost associated with sample processing equipment, materials and reagents, large amount of samples required, the need for skilled personnel to process the samples, massive amount of data generated, and the time consuming nature in integration and synthesis of the data. Currently, few bioremediation projects utilize the systems biology approach due to limitations in funding, expertise, and resources. This review will describe a compilation of research projects that would constitute a perfect study employing a systems biology approach for remediation of radionuclides, metals, hydrocarbons and chlorinated solvents, and suggest directions for future development.

To use a systems biology approach to bioremediation projects they must involve the characterization of microbial community composition, cellular and molecular activity and are complicated by the presence of toxic chemicals that alters the normal behavior of the microbial community. In addition, the ultimate objective of bioremediation projects is the elimination or detoxification of toxic compounds, which requires an understanding of all possible influence from environmental variables and cell–cell interactions. The selection process for which methods to use in a systems biology study includes consideration for cost, time frame, personnel, and the objectives of the project. If the

Figure 1



Systems biology from molecules to ecosystems. In general terms, an ecosystem consists of communities, populations, cells, protein, RNA, and DNA. The approaches use geochemical, ecological, genomic, proteomic, metabolomic, and computational techniques. Analyze DNA, RNA, and protein at the cellular levels to understand impacts on the cell in terms of how bioremediation functions, and analyze communities, and populations to understand impacts on structure/function relationships and finally interactomes at the ecosystem level in terms of bioremediation practices.

focus is to elucidate microbial community composition, DNA based ‘-omics’ tools such as 16S rRNA clone library, PhyloChip or sequencing should be used. If the interest is to understand cellular pathways and identify functional genes involved in microbially mediated reactions, tools that identify RNA, and proteins such as GeoChip, RNA-seq, and various mass spectrometry methods should be used. If the intention is to characterize small molecules produced by the microbes, matrix-assisted laser desorption/ionization (MALDI), desorption electrospray ionization (DESI), nuclear magnetic resonance (NMR) spectroscopy can be used. Concomitant monitoring of limiting nutrients, electron donors, electron acceptors, and hydrology is also crucial for a systems biology conceptual model to be useful.

To gain an understanding of complex *in situ* bioremediation processes, monitoring techniques that inventory and monitor terminal electron acceptors and electron donors, enzyme probes that measure functional activity in the environment, functional genomic microarrays, phylogenetic microarrays, metabolomics, proteomics, and quantitative PCR can provide unprecedented insights into the key microbial reactions employed (Figure 1). In general terms, an ecosystem consists of communities, populations, cells, protein, RNA, and DNA. We can analyze DNA, RNA, and protein at the cellular levels to understand the impacts on the cells, and analyze community and populations to understand effect of bioremediation on structure/function relationships (Figure 1). In some cases, a change in redox state is the simplest tool to bring about detoxification of hazardous metals and organic compounds. This is particularly true for metals and radionuclides like U(VI), Cr(VI), and Tc(VII). While these cannot be degraded, they can be biotransformed decreasing their bioavailability, mobility and thus toxicity [9–11]. Microbes can directly mediate such immobilization and detoxification by changing the valence states, utilizing them as electron acceptors [12,13] when appropriate electron donors are present. Measurement of enzyme activity during bioremediation is a reliable, inexpensive tool for measuring microbial respiration and metabolism. Dehydrogenase enzyme based assays such as INT (iodo nitro tetrazolium) [14] and TTC (triphenyl tetrazolium chloride) assays [15] have been used successfully in monitoring microbial respiration in the bioremediation of explosives [16], metals, PAHs [17], and oil [18]. Other enzymatic or DNA/RNA based probes from key microbial metabolic pathways can be effectively used as a tool in tracking bioremediation processes as used previously during degradation of trichloroethene (TCE) [19] and of petroleum hydrocarbons [20,21]. Proteogenomic analysis during U(VI) reduction field studies have been able to identify and track *Geobacter*-specific biomarker peptide citrate synthase [22] during the process. Using qPCR as a technique for detection of phylogenetic and catabolic genes as indication of microbially mediated remediation is a popular and successful approach for monitoring detoxification in metal as well as hydrocarbon contaminated sites. Examples include monitoring *Anaeromyxobacter* strains involved in reduction of U(VI) [23] and of *Dehalococcoides* spp. in bioremediation of chlorinated solvents [24]. In order to identify and track the entire microbial community during bioremediation processes, metagenomic analysis including 16S rRNA-based clone libraries has been broadly used for metals [25,26] as well as for hydrocarbons and chlorinated solvents [27,28,29] among others. Collectively, these techniques for metagenomics have reiterated that the microbial diversity existing in most environments is greater than expected [30]. Recently, high throughput microarrays like the PhyloChip and the GeoChip have been extensively used in metal and organics bioremediation studies

[31[•],32,33,34[•],35], in order to quickly characterize microbial community and function. PhyloChip, the 16S rRNA-phylogenetic microarray characterizes and monitors microbial community dynamics whereas GeoChip, the functional gene microarray tracks functional gene activity changes of microbes in the environment [34[•],36[•]]. Microbial community proteomics and metabolomics have been a major breakthrough in providing deeper insight into the microbial cellular function and gene products interplaying in the environment [37[•]]. A novel application of Immunomagnetic separation for targeting and monitoring specific microorganisms during *in situ* bioremediation [38] holds promise to enable transcriptomics, proteomics, or metabolomics-based studies directly on cells collected from the field. Integration of all of these techniques using the latest advances in bioinformatics and modeling will enable break-through science in environmental biotechnology. We discuss a review of these techniques as used in field studies and lab simulations from sites contaminated with metals, radionuclides, hydrocarbons, and chlorinated solvents (Table 1).

Case studies

The U.S. Department of Energy (DOE) is responsible for the remediation and long-term stewardship of a significant number of plumes containing various contaminants including radionuclides and metals, at sites spread across the United States (<http://www.em.doe.gov/Pages/siteslocations.aspx>). Several groups of researchers have been involved since 2004 in active implementation of basic research to understand the systems biology of contaminated sites, and predicting feasible remediation technologies.

Radionuclide biotransformation

Groundwater and soil at the Area 3 FRC site in Oak Ridge is not only contaminated with Uranium (up to 200 μ M), but poses an unique bioremediation problem due to its low pH ($\sim$ 3), high nitrate (200 mM), and high calcium concentrations along with presence of chlorinated organic solvents. Research at this site by various investigators exemplifies successful application of systems biology tools to reveal a deeper understanding of the microbiology at play in the subsurface. Previously, 16S clone library-based community analysis during an *in situ* biostimulation test at this site have identified *Desulfovibrio*, *Geobacter*, *Anaeromyxobacter*, *Desulfosporosinus*, *Acidovorax*, and *Geothrix* spp. present concomitant with U(VI) reduction [26]. Clone libraries of functional gene markers like *dsrAB*, *nirK*, *nirS*, *amoA*, and *pmoA* [39,40] showed high microbial diversity in functional genes. However, recent metagenomic analysis from well FW106 specifically using a random shotgun sequencing-based strategy revealed a highly enriched community dominated by denitrifying β -*Proteobacteria* and γ -*Proteobacteria* [2]. GeoChip analysis of several groundwater monitoring wells reported widespread diversity of *dsrAB* genes [34[•],41],

which showed that sulfate-reducing bacteria were key players in U(VI) reduction. During the U(VI) reoxidation phase as studied in a sediment column with samples from FRC, observed decrease in biomass, but increase in microbial activity [42]. Using the PhyloChip, the study showed no decline in *Geobacter* or *Geothrix* spp. during the reoxidation phase, but members of *Actinobacteria*, *Firmicutes*, *Acidobacteria*, and *Desulfovibrionaceae* exhibited increased abundance [42]. GeoChip analysis during the reoxidation phase from field samples showed a decline in *dsr* genes but reoxidation did not appear to effect microbial functional diversity [33] suggesting that the microbial community was able to recover and continue to reduce U(VI) in the post oxidation phase.

Metals bioimmobilization

The Hanford 100H area adjacent to the Columbia River in Washington is contaminated with Chromium (Cr) as a result of being a weapons production site. In 2004, Hydrogen Release Compound HRCtm was injected in an effort to mediate sustained bioimmobilization of Cr(VI) *in situ* by stimulating indigenous microbial flora [43]. Hubbard *et al.* used time-lapse seismic and radar tomographic geophysical monitoring to determine spatio-temporal distribution of the injected HRC and biogeochemical transformations associated with Cr(VI) bioremediation post injection of HRC [44]. Direct cell counts revealed that while cell numbers reached 10^8 cells/ml [43], Cr(VI) levels decreased from 100 ppb to below background levels within a year. PhyloChip analysis showed enrichment of sulfate reducers along with nitrate reducers, iron reducers, and methanogenic populations during this time [43]. Targeted enrichments resulted in isolation of sulfate-reducing *Desulfovibrio vulgaris* like strain RCH1, nitrate reducing strain *Pseudomonas stutzeri* strain RCH2, and iron-reducing strain *Geobacter metallireducens* strain RCH3 [45], all capable of Cr(VI) reduction [45]. μ FlowFISH (integrated fluorescence *in situ* hybridization and flow cytometry) analysis was able to detect and sort *Pseudomonads* similar to strain RCH2 directly from Hanford 100H field water samples collected in 2009 and 2010 [46].

Hydrocarbon bioremediation

The dependence of petroleum-based energy source has fueled industrial growth and prosperity. However, it also brought dispersal of hydrocarbons into different environments. Fortunately, the organic nature of hydrocarbons enables microbes to metabolize these petroleum compounds as substrates. Notable reviews on a systems biology approach to bioremediation a Atlas and Hazen [47[•]], Harayama *et al.* [48], Zhou *et al.* [49], Fredrickson *et al.* [50], and de Lorenzo [51].

The MC252 oil spill in the Gulf of Mexico in 2010 was the largest in US history. Many environmental factors

Table 1

Fundamental systems biology parameters measured at the bioremediation sites

Site	Contaminant	Key parameters measured	References
<u>Field Research Center, Oak Ridge, TN</u>	Uranium (VI), nitrate	(a) 16S clone libraries	[26]
		(b) Metagenomics	[2]
		(c) PhyloChip	[42]
		(d) Functional gene clone libraries	[39,40]
		(e) GeoChip	[33]
<u>Hanford 100H, Hanford, WA</u>	Chromium (VI)	(a) Seismic and radar tomographic monitoring	[43]
		(b) Microbial cell counts	[44]
		(c) PhyloChip	[44]
		(d) Microbial isolation	[45]
		(e) FISH	[46]
<u>Gulf of Mexico</u>	Oil	(a) Dissolved oxygen	[52]
		(b) Enzyme activities	[53]
		(c) Microbial counts	[27*,53]
		(d) Hydrocarbon analyses	[54,27*,52,53]
		(e) 16S clone libraries	[27*]
		(f) PhyloChip	[27*]
		(g) GeoChip	[27*,56*]
		(h) PLFA	[27*,53]
		(i) Isotope chemistry	[27*,55]
<u>Savannah River Site, SC</u>	TCE, PCE	(a) Microbial cell counts	[62]
		(b) Fluorescent antibody	[62]
		(c) PLFA	[62]
		(d) Functional gene analysis	[62,66]
		(e) Isotope chemistry	[67]
<u>Test Area North, ID</u>	TCE	(a) Microbial cell counts	[68]
		(b) PLFA	[68]
		(c) Phenotypic microarray	[68]
		(d) DGGE	[68]
		(e) qPCR	[69]
		(f) RFLP	[69,70]
		(g) Functional gene analysis	[70]
		(h) PhyloChip	[31*]
		(i) Isotope chemistry	[63]

distinguished this spill from previous ones, including hydrocarbon composition, environmental variables, depth of the spill, and the availability of systems biology tools. Information on chemical analyses is crucial in support of a system's biology approach for oil bioremediation in the MC252 spill. While Camilli *et al.* [52] concluded that microbial respiration rates within the deep plume were extremely low based on dissolved oxygen concentration, measurement of microbial respiration rates, enzyme activity, phosphate concentration, and polar membrane lipid concentration in surface water affected by the oil spill. Edwards *et al.* concluded that enzyme activities and respiration rates were found to be higher inside the oil slick [53]. Valentine *et al.* [54] investigated the fate of methane, propane, and ethane gases of the deep hydrocarbon plume at depth greater than 799 m, and found that propane and ethane were degraded faster than methane. ^{13}C -labeled substrates, as well as ^{13}C and ^3H tracers, were used to measure $\delta^{13}\text{C}$ -DIC. In another study, methane was found to be the most abundant hydrocarbon released during the MC252 spill, and that there was a rapid response of methanotrophic bacteria rapidly respiring

the released methane [55]. PhyloChip, clone library, GeoChip, phospholipid fatty acid (PLFA), and isotope chemistry were used to compare microbial communities inside and outside the deep plume [27*]. The results identified *Oceanospirillales*, which were found to degrade hydrocarbons at 5°C inside the plume. The GeoChip demonstrated genes that were significantly correlated to concentration of oil contaminants, such as *phdC1* (naphthalene degradation), and *alkB* (oxidation of alkanes), as well as a shift in C, N, P, S cycling processes in the deep plume samples [56*]. The involvement of federal agencies and pending lawsuits is the impetus for a concerted effort in collating all data collected resulting in a comprehensive database useful for researchers. By integrating chemical analyses with studies utilizing a systems biology approach, there was an unprecedented near real-time understanding of chemical and biological reactions involved in the hydrocarbon degradation. In order to gain a more comprehensive understanding of the microbiological processes, data from transcriptomics studies will provide information on whether the cultivatable dominant microbes are the *in situ* active ones, and

proteomics studies will identify enzymes central to hydrocarbon degradation.

Chlorinated solvents bioremediation

Chlorinated solvents, such as TCE and dichloroethene (DCE), are recalcitrant carcinogenic compounds that persist in the environment once released. Microbes, such as *Dehalococcoides*, are capable of using the chlorinated solvents as electron acceptors anaerobically and dechlorinating the compounds to ethene [31[•],57]. Another biodegradation pathway is the aerobic co-metabolism of the chlorinated compounds to carbon dioxide and chloride by microbes such as methane-oxidizers with methane monooxygenases (MMOs) [31[•]]. Descriptions of techniques that monitor mass loss, geochemical fingerprints, isotope fractionation associated with biodegradation, microbial communities in biostimulation and natural attenuation studies, quantitative real-time PCR methods targeting reductive dehalogenase genes are included in several reviews [58,59,57].

Between 1955 and 1972, low-level radioactive isotopes, sewage and chlorinated solvents were injected into the aquifer through a 95 m deep well at Test Area North (TAN) in Idaho National Laboratory. The plume contained TCE concentrations ranging from 5 ppb to 300 ppm extending for more than 2 km. An enhanced *in situ* bioremediation pilot study started in 1999 to treat the chlorinated solvents contaminated groundwater by injecting the electron donor Lactate to stimulate *in situ* reductive dechlorination. A comparison of microbial communities in the core and groundwater samples was assessed by characterizing total biomass, PLFA analysis, culturing and community-level physiological profiling (CLPP) using Biolog GN microplates [60]. DGGE analysis indicated that wells with high concentrations of chlorinated solvents had different microbial communities from wells with minimal concentrations of the contaminants, and that attached and the free-living microbes had different functional and composition profile [60]. Additionally, qPCR of the *Dehalococcoides* sp. 16S rRNA genes provided the most convincing result in quantifying dechlorinating potential of a community compared to community analysis by terminal restriction fragment length polymorphism (T-RFLP), and RFLP analysis with clone sequencing [61]. Erwin *et al.* [62] demonstrated the presence of bacteria harboring MMOs and potential of TCE co-metabolism at TAN from a pristine area using PCR amplification to generate a function gene fragment library and sequencing. Stable carbon isotope ratios of groundwater samples taken in 2000 confirmed the complete conversion of TCE to ethene, and minimal biodegradation of t-DCE [63]. Using the PhyloChip for bacterial composition characterization, a decrease in reductive dechlorinating organisms and an increase in methane-oxidizing microbes capable of

aerobic co-metabolism of TCE was observed [31[•]]. Further studies that would complement the investigation at the TAN site would be to employ a shotgun proteomics approach as reported by Werner *et al.* [64[•]]. Their method allowed for detection of peptides, such as *FdhA*, *TceA*, *PceA*, and *HupL* that could potentially be used as bioindicators of chlorinated ethene dehalorespiration.

Conclusion

The combination of the systems biology techniques as demonstrated in the case studies above allowed for enhanced understanding of complex bioremediation processes. Investigation of the MC252 spill is the most comprehensive bioremediation study using a systems biology approach to date as a result of available funding, resources, expertise, as well as, interests from the scientific communities and regulating agencies. Future projects can benefit from the experiences obtained from the MC252 spill investigation. However, while significant advances have been made in rapid generation and availability of 'omics'-based data in key microbial processes in the environment, a key bottleneck lies in the ability to quickly analyze the output using appropriate, user friendly, simplified bioinformatic tools to make meaningful conclusions. Currently, user-friendly bioinformatics pipelines available for analysis of sequencing and microarray data, include Qiime (qiime.sourceforge.net) [65] and PhyloTrac (www.phylotrac.org/), respectively. In order to fully utilize the data generated from the various 'omics' tools, better annotation of the genes, pathways, and metabolites are needed. A comprehensive database of all available genomics, proteomics, and metabolomics information from bioremediation research will provide a platform for scientist to exchange information including data obtained, and analysis methods and pipeline. This will require coordination from scientists to share data, and database managers to update, maintain, and provide quality control. Taken together, these tools will allow for accurate interpretation of the 'omics' data, leading to generation of judicious predictive models and strategies for successful implementation of bioremediation applications in the future.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Benndorf D, Balcke GU, Harms H, von Bergen M: **Functional metaproteome analysis of protein extracts from contaminated soil and groundwater.** *ISME J* 2007, **1**:224-234.
2. Hemme CL, Deng Y, Gentry TJ, Fields MW, Wu L, Barua S, Barry K, Tringe SG, Watson DB, He Z *et al.*: **Metagenomic insights into evolution of a heavy metal-contaminated groundwater microbial community.** *ISME J* 2010, **4**:660-672.
3. Edward FD: **Marine microbial diversity: the tip of the iceberg.** *Trends Biotechnol* 1997, **15**:203-207.

4. DeLong EE: **Microbial community genomics in the ocean.** *Nat Rev Microbiol* 2005, **3**:459-469.
 5. Baker BJ, Banfield JF: **Microbial communities in acid mine drainage.** *FEMS Microbiol Ecol* 2003, **44**:139-152.
 6. Harmsen HJM, Prieur D, Jeanthou C: **Distribution of microorganisms in deep-sea hydrothermal vent chimneys investigated by whole-cell hybridization and enrichment culture of thermophilic subpopulations.** *Appl Environ Microbiol* 1997, **63**:2876-2883.
 7. Hu Y, Fu C, Yin Y, Cheng G, Lei F, Yang X, Li J, Ashforth E, Zhang L, Zhu B: **Construction and preliminary analysis of a deep-sea sediment metagenomic fosmid library from Qiongdongnan Basin, South China Sea.** *Mar Biotechnol* 2010, **12**:719-727.
 8. Gill SR, Pop M, DeBoy RT, Eckburg PB, Turnbaugh PJ, Samuel BS, Gordon JL, Relman DA, Fraser-Liggett CM, Nelson KE: **Metagenomic analysis of the human distal gut microbiome.** *Science* 2006, **312**:1355-1359.
 9. Desjardins V, Bayard R, Huck N, Manceau A, Gourdon R: **Effect of microbial activity on the mobility of chromium in soils.** *Waste Manag* 2002, **22**:195-200.
 10. Suzuki Y, Kelly SD, Kemner KM, Banfield JF: **Direct microbial reduction and subsequent preservation of uranium in natural near-surface sediment.** *Appl Environ Microbiol* 2005, **71**:1790-1797.
 11. Chen JM, Hao OJ: **Microbial chromium (VI) reduction.** *Crit Rev Environ Sci Technol* 1998, **28**:219-251.
 12. Elias DA, Krumholz LR, Wong D, Long PE, Suflita JM: **Characterization of microbial activities and U reduction in a shallow aquifer contaminated by uranium mill tailings.** *Microb Ecol* 2003, **46**:83-91.
 13. Finneran KT, Anderson RT, Nevin KP, Lovley DR: **Potential for bioremediation of uranium-contaminated aquifers with microbial U(VI) reduction.** *Soil Sediment Contam* 2002, **11**:339-357.
 14. Mathew M, Obbard JP: **Optimisation of the dehydrogenase assay for measurement of indigenous microbial activity in beach sediments contaminated with petroleum.** *Biotechnol Lett* 2001, **23**:227-230.
 15. Guochen Z, Feng Z, Hang Y, Xue L, Jianzhong L: **Comparison between INT and TTC assay to determine the dehydrogenase activity of flocs.** *Water Resource and Environmental Protection (ISWREP), 2011 International Symposium on 20-22 May 2011.* 2011:1690-1693.
 16. Beltz LA, Neira DR, Axtell CA, Iverson S, Deaton W, Waldschmidt TJ, Bumpus JA, Johnston CG: **Immunotoxicity of explosives-contaminated soil before and after bioremediation.** *Arch Environ Contam Toxicol* 2001, **40**:311-317.
 17. Mosher JJ, Levison BS, Johnston CG: **A simplified dehydrogenase enzyme assay in contaminated sediment using 2-(p-iodophenyl)-3-(p-nitrophenyl)-5-phenyl tetrazolium chloride.** *J Microbiol Methods* 2003, **53**:411-415.
 18. Bento FM, Camargo FAO, Okeke BC, Frankenberger WT: **Comparative bioremediation of soils contaminated with diesel oil by natural attenuation, biostimulation and bioaugmentation.** *Bioresour Technol* 2005, **96**:1049-1055.
 19. Hazen T, Chakraborty R, Fleming J, Gregory I, Bowman J, Jimenez L, Zhang D, Pfiffner S, Brockman F, Saylor G: **Use of gene probes to assess the impact and effectiveness of aerobic *in situ* bioremediation of TCE.** *Arch Microbiol* 2009, **191**:221-232.
 20. Bell TH, Yergeau E, Martineau C, Juck D, Whyte LG, Greer CW: **Identification of nitrogen-incorporating bacteria in petroleum-contaminated arctic soils by using [15N]DNA-based stable isotope probing and pyrosequencing.** *Appl Environ Microbiol* 2011, **77**:4163-4171.
 21. Beller HR, Kane SR, Legler TC, Alvarez PJJ: **A real-time polymerase chain reaction method for monitoring anaerobic, hydrocarbon-degrading bacteria based on a catabolic gene.** *Environ Sci Technol* 2002, **36**:3977-3984.
 22. Wilkins MJ, Callister SJ, Miletto M, Williams KH, Nicora CD, Lovley DR, Long PE, Lipton MS: **Development of a biomarker for *Geobacter* activity and strain composition; Proteogenomic analysis of the citrate synthase protein during bioremediation of U(VI).** *Microb Biotechnol* 2011, **4**:55-63.
 23. Thomas SH, Padilla-Crespo E, Jardine PM, Sanford RA, Löffler FE: **Diversity and distribution of *Anaeromyxobacter* strains in a uranium-contaminated subsurface environment with a nonuniform groundwater flow.** *Appl Environ Microbiol* 2009, **75**:3679-3687.
 24. Ritalahti KM, Amos BK, Sung Y, Wu Q, Koenigsberg SS, Löffler FE: **Quantitative PCR targeting 16S rRNA and reductive dehalogenase genes simultaneously monitors multiple *Dehalococcoides* strains.** *Appl Environ Microbiol* 2006, **72**:2765-2774.
 25. Vronis HA, Anderson RT, Ortiz-Bernad I, O'Neill KR, Resch CT, Peacock AD, Dayvault R, White DC, Long PE, Lovley DR: **Microbiological and geochemical heterogeneity in an *in situ* uranium bioremediation field site.** *Appl Environ Microbiol* 2005, **71**:6308-6318.
 26. Cardenas E, Wu W-M, Leigh MB, Carley J, Carroll S, Gentry T, Luo J, Watson D, Gu B, Ginder-Vogel M *et al.*: **Microbial communities in contaminated sediments, associated with bioremediation of uranium to submicromolar levels.** *Appl Environ Microbiol* 2008, **74**:3718-3729.
 27. Hazen TC, Dubinsky EA, DeSantis TZ, Andersen GL, Piceno YM, Singh N, Jansson JK, Probst A, Borglin SE, Fortney JL *et al.*: **Deep-sea oil plume enriches indigenous oil-degrading bacteria.** *Science* 2010, **330**:204-208.
- This is one of the first research to employ all aspects of the systems biology approach (metagenomics with phylogenetic and functional microarrays, isotopic characterization, SRFTIR, and *in situ* geochemical monitoring) to understand microbial response in an unprecedented environmental disaster.
28. Dojka MA, Hugenholtz P, Haack SK, Pace NR: **Microbial diversity in a hydrocarbon- and chlorinated-solvent-contaminated aquifer undergoing intrinsic bioremediation.** *Appl Environ Microbiol* 1998, **64**:3869-3877.
 29. Militon C, Boucher D, Vachelard C, Perchet G, Barra V, Troquet J, Peyretailade E, Peyret P: **Bacterial community changes during bioremediation of aliphatic hydrocarbon-contaminated soil.** *FEMS Microbiol Ecol* 2010, **74**:669-681.
 30. Schloss PD, Handelsman J: **Status of the microbial census.** *Microbiol Mol Biol Rev* 2004, **68**:686-691.
 31. Conrad ME, Brodie EL, Radtke CW, Bill M, Delwiche ME, Lee MH, Swift DL, Colwell FS: **Field evidence for co-metabolism of trichloroethene stimulated by addition of electron donor to groundwater.** *Environ Sci Technol* 2010, **44**:4697-4704.
- This research demonstrates how the use of a number of systems biology parameters can reveal mechanisms of bioremediation that were previously not obvious.
32. Rastogi G, Osman S, Vaishampayan P, Andersen G, Stetler L, Sani R: **Microbial diversity in uranium mining-impacted soils as revealed by high-density 16S microarray and clone library.** *Microb Ecol* 2010, **59**:94-108.
 33. Van Nostrand JD, Wu W-M, Wu L, Deng Y, Carley J, Carroll S, He Z, Gu B, Luo J, Criddle CS *et al.*: **GeoChip-based analysis of functional microbial communities during the reoxidation of a bioreduced uranium-contaminated aquifer.** *Environ Microbiol* 2009, **11**:2611-2626.
 34. He Z, Gentry TJ, Schadt CW, Wu L, Liebich J, Chong SC, Huang Z, Wu W, Gu B, Jardine P *et al.*: **GeoChip: a comprehensive microarray for investigating biogeochemical, ecological and environmental processes.** *ISME J* 2007, **1**:67-77.
- The first description of GeoChip-functional gene microarray providing direct linkage of microbial functional gene to ecosystem function and processes.
35. Liang Y, Li G, Van Nostrand JD, He Z, Wu L, Deng Y, Zhang X, Zhou J: **Microarray-based analysis of microbial functional diversity along an oil contamination gradient in oil field.** *FEMS Microbiol Ecol* 2009, **70**:324-333.
 36. DeSantis T, Brodie E, Moberg J, Zubieta I, Piceno Y, Andersen G: **High-density universal 16S rRNA microarray analysis reveals**

broader diversity than typical clone library when sampling the environment. *Microb Ecol* 2007, **53**:371-383.

See annotation to [34*].

37. VerBerkmoes NC, Denev VJ, Hettich RL, Banfield JF: **Systems Biology: functional analysis of natural microbial consortia using community proteomics.** *Nat Rev Microbiol* 2009, **7**:196-205.

Excellent review on microbial community proteogenomic approaches being developed for systems biology studies.

38. Chakraborty R, Hazen TC, Joyner DC, Küsel K, Singer ME, Sitté J, Torok T: **Use of immunomagnetic separation for the detection of *Desulfovibrio vulgaris* from environmental samples.** *J Microbiol Methods* 2011, **86**:204-209.
 39. Fields MW, Bagwell CE, Carroll SL, Yan T, Liu X, Watson DB, Jardine PM, Criddle CS, Hazen TC, Zhou J: **Phylogenetic and functional biomarkers as indicators of bacterial community responses to mixed-waste contamination.** *Environ Sci Technol* 2006, **40**:2601-2607.
 40. Palumbo AV, Schryver JC, Fields MW, Bagwell CE, Zhou JZ, Yan T, Liu X, Brandt CC: **Coupling of functional gene diversity and geochemical data from environmental samples.** *Appl Environ Microbiol* 2004, **70**:6525-6534.
 41. Waldron PJ, Wu L, Nostrand JDV, Schadt CW, He Z, Watson DB, Jardine PM, Palumbo AV, Hazen TC, Zhou J: **Functional gene array-based analysis of microbial community structure in groundwaters with a gradient of contaminant levels.** *Environ Sci Technol* 2009, **43**:3529-3534.
 42. Brodie EL, DeSantis TZ, Joyner DC, Baek SM, Larsen JT, Andersen GL, Hazen TC, Richardson PM, Herman DJ, Tokunaga TK *et al.*: **Application of a high-density oligonucleotide microarray approach to study bacterial population dynamics during uranium reduction and reoxidation.** *Appl Environ Microbiol* 2006, **72**:6288-6298.
 43. Faybishenko B, Hazen TC, Long PE, Brodie EL, Conrad ME, Hubbard SS, Christensen JN, Joyner D, Borglin SE, Chakraborty R *et al.*: **In situ long-term reductive bioimmobilization of Cr(VI) in groundwater using hydrogen release compound.** *Environ Sci Technol* 2008, **42**:8478-8485.
 44. Hubbard SS, Williams K, Conrad ME, Faybishenko B, Peterson J, Chen JS, Long P, Hazen T: **Geophysical monitoring of hydrological and biogeochemical transformations associated with Cr(VI) bioremediation.** *Environ Sci Technol* 2008, **42**:3757-3765.
 45. Han RY, Geller JT, Yang L, Brodie EL, Chakraborty R, Larsen JT, Beller HR: **Physiological and transcriptional studies of Cr(VI) reduction under aerobic and denitrifying conditions by an aquifer-derived pseudomonad.** *Environ Sci Technol* 2010, **44**:7491-7497.
 46. Liu P, Meagher RJ, Light YK, Yilmaz S, Chakraborty R, Arkin AP, Hazen TC, Singh AK: **Microfluidic fluorescence in situ hybridization and flow cytometry (μ FlowFISH).** *Lab on a Chip* 2011, **11**:2673-2679.
 47. Atlas RM, Hazen TC: **Oil biodegradation and bioremediation: a tale of the two worst spills in US history.** *Environ Sci Technol* 2011, **45**:6709-6715.
- This feature article provides an interesting and informative comparison of the Exxon Valdez and the BP Deepwater Horizon spills and analyzed the bioremediation strategies employed in each of the spill.
48. Harayama S, Kasai Y, Hara A: **Microbial communities in oil-contaminated seawater.** *Curr Opin Biotechnol* 2004, **15**:205-214.
 49. Zhou JZ, He Q, Hemme CL, Mukhopadhyay A, Hillesland K, Zhou AF, He ZL, Van Nostrand JD, Hazen TC, Stahl DA *et al.*: **How sulphate-reducing microorganisms cope with stress: lessons from systems biology.** *Nat Rev Microbiol* 2011, **9**:452-466.
 50. Fredrickson JK, Romine MF, Beliaev AS, Auchtung JM, Driscoll ME, Gardner TS, Nealson KH, Osterman AL, Pinchuk G, Reed JL *et al.*: **Towards environmental systems biology of *Shewanella*.** *Nat Rev Microbiol* 2008, **6**:592-603.
 51. de Lorenzo V: **Systems biology approaches to bioremediation.** *Curr Opin Biotechnol* 2008, **19**:579-589.
 52. Camilli R, Reddy CM, Yoerger DR, Van Mooy BAS, Jakuba MV, Kinsey JC, McIntyre CP, Sylva SP, Maloney JV: **Tracking hydrocarbon plume transport and biodegradation at Deepwater Horizon.** *Science* 2010, **330**:201-204.
 53. Edwards BR, Reddy CM, Camilli R, Carmichael CA, Longnecker K, Van Mooy BAS: **Rapid microbial respiration of oil from the Deepwater Horizon spill in offshore surface waters of the Gulf of Mexico.** *Environ Res Lett* 2011, **6**:035301.
 54. Valentine DL, Kessler JD, Redmond MC, Mendes SD, Heintz MB, Farwell C, Hu L, Kinnaman FS, Yvon-Lewis S, Du MR *et al.*: **Propane respiration jump-starts microbial response to a deep oil spill.** *Science* 2010, **330**:208-211.
 55. Kessler JD, Valentine DL, Redmond MC, Du MR, Chan EW, Mendes SD, Quiroz EW, Villanueva CJ, Shusta SS, Werra LM *et al.*: **A persistent oxygen anomaly reveals the fate of spilled methane in the deep Gulf of Mexico.** *Science* 2011, **331**:312-315.
 56. Lu Z, Deng Y, Van Nostrand JD, He Z, Voordeckers J, Zhou A, Lee Y.-J., Mason OU, Dubinsky EA, Chavarria KL *et al.*: **Microbial gene functions enriched in the Deepwater Horizon deep-sea oil plume.** *ISME J*, doi:10.1038/ismej.2011.91.
- This is the companion paper to the Hazen *et al.* [27*] research that utilized the systems biology approach to understand microbial response to the Gulf of Mexico MC252 spill in 2010. This paper reports in more detail the results from the functional microarray (GeoChip) of microbial gene response to an oil spill.
57. Cupples AM: **Real-time PCR quantification of *Dehalococcoides* populations: methods and applications.** *J Microbiol Methods* 2008, **72**:1-11.
 58. Illman WA, Alvarez PJ: **Performance assessment of bioremediation and natural attenuation.** *Crit Rev Environ Sci Technol* 2009, **39**:209-270.
 59. Scow KM, Hicks KA: **Natural attenuation and enhanced bioremediation of organic contaminants in groundwater.** *Curr Opin Biotechnol* 2005, **16**:246-253.
 60. Lehman RM, O'Connell SP, Banta A, Fredrickson JK, Reysenbach AL, Kieft TL, Colwell FS: **Microbiological comparison of core and groundwater samples collected from a fractured basalt aquifer with that of dialysis chambers incubated in situ.** *Geomicrobiol J* 2004, **21**:169-182.
 61. Rahm BG, Chauhan S, Holmes VF, Macbeth TW, Sorenson KSJ, Alvarez-Cohen L: **Molecular characterization of microbial populations at two sites with differing reductive dechlorination abilities.** *Biodegradation* 2006, **17**:523-534.
 62. Erwin DP, Erickson IK, Delwiche ME, Colwell FS, Strap JL, Crawford RL: **Diversity of oxygenase genes from methane- and ammonia-oxidizing bacteria in the Eastern Snake River Plain aquifer.** *Appl Environ Microbiol* 2005, **71**:2016-2025.
 63. Song DL, Conrad ME, Sorenson KS, Alvarez-Cohen L: **Stable carbon isotope fractionation during enhanced in situ bioremediation of trichloroethene.** *Environ Sci Technol* 2002, **36**:2262-2268.
 64. Werner JJ, Ptak AC, Rahm BG, Zhang S, Richardson RE: **Absolute quantification of *Dehalococcoides* proteins: enzyme bioindicators of chlorinated ethene dehalorespiration.** *Environ Microbiol* 2009, **11**:2687-2697.
- This research utilized a shotgun proteomics approach to characterize *Dehalococcoides* proteins in complex samples and identifies peptides as bioindicators of dehalospiration.
65. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Pena AG, Goodrich JK, Gordon JI *et al.*: **QIIME allows analysis of high-throughput community sequencing data.** *Nat Methods* 2010, **7**:335-336.
 66. Hazen T, Chakraborty R, Fleming J, Gregory IR, Bowman JP, Jimenez L, Zhang D, Pfiffner SM, Brockman FJ, Sayler GS: **Use of gene probes to assess the impact and effectiveness of aerobic in situ bioremediation of TCE.** *Arch Microbiol* 2009, **191**:221-232.
 67. Seaman JC, Looney BB, Harris MK: **Research in support of remediation activities at the Savannah River Site.** *Vadose Zone J* 2007, **6**:316-326.

68. Conrad ME, Brodie EL, Radtke CW, Bill M, Delwiche ME, Hope Lee M, Swift DL, Colwell FS: **Field evidence for co-metabolism of trichloroethene stimulated by addition of electron donor to groundwater.** *Environ Sci Technol* 2010, **44**:4697-4704.
69. Rahm BG, Chauhan S, Holmes VF, Macbeth TW, Sorenson KS Jr, Alvarez-Cohen L: **Molecular characterization of microbial populations at two sites with differing reductive dechlorination abilities.** *Biodegradation* 2006, **17**:523-534.
70. Ritalahti KM, Amos BK, Sung Y, Wu Q, Koenigsberg SS, Löffler FE: **Quantitative PCR targeting 16S rRNA and reductive dehalogenase genes simultaneously monitors multiple dehalococcoides strains.** *Appl Environ Microbiol* 2006, **72**:2765-2774.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California.

Ernest Orlando Lawrence Berkeley National Laboratory is an equal opportunity employer.