



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Room-Temperature, Ignition-Scale Hohlraum Experiments on NIF

D. J. Strozzi, J. E. Ralph, T. Ma, D. E. Hinkel, D. A.
Callahan, J. L. Kline, J. D. Moody, J. R. Rygg, G. D.
Kerbel, M. M. Marinak, S. H. Glenzer

November 4, 2013

APS DPP 2013
Denver, CO, United States
November 11, 2013 through November 15, 2013

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.



Room-temperature, ignition-scale hohlraum experiments on NIF

**D. J. Strozzi, J. E. Ralph, T. Ma, D. E. Hinkel, D. A. Callahan,
J. L. Kline, J. D. Moody, O. Jones, J. R. Rygg, G. D. Kerbel,
M. M. Marinak**

Lawrence Livermore National Laboratory

S. H. Glenzer

SLAC

NIF room-temperature (“warm”) ignition-scale hohlraum experiments with higher-Z gas fill

Does higher Z gas fill improve inner-beam propagation = round implosion with less cross-beam transfer to inners ($\Delta\lambda$)? Jury still out.

Motivation:

- Hotter hohlraum plasma could reduce inner-beam SRS (more Landau damping)
- Higher Z hohlraum gas fill absorbs more laser via inverse bremsstrahlung
- Warm shots allow gases that freeze at cryo, e.g. hydrocarbons
- Warm shots easier to field, cheaper (no cryo hardware)

Shape: in-flight shell vs. stagnated hotspot


- Warm hotspots close to round with less $\Delta\lambda$ than cryos
- But warm shell pancaked – need more $\Delta\lambda$ to make round
- Opposite “swing” of cryos – shell round but hotspot pancaked

Backscatter: Warms have less inner-beam SRS, more outer-beam SBS

- Pure $\Delta\lambda$ scaling or intrinsic effect of gas fill?

Warm shots use hydrocarbon gas fills, retuned laser picket and trough

	Cryo	Warm	Comments
Hohlraum fill	0.96 mg/cc He ⁴	0.82 mg/cc C ₅ H ₁₂	higher Z, same n _e
Capsule fill	D-He ³	propane: C ₃ {H or D} ₈	D, He leak warm



19ns: shown on next slide


Higher Z hohlraum gas fill increases inverse bremsstrahlung laser absorption, gives hotter plasma

$$\kappa_{IB} \propto \left[1 - \frac{n_e}{n_{cr}}\right]^{-1/2} \frac{n_e^2}{T_e^{3/2}} \ln \Lambda \frac{\sum_i f_i Z_i^2}{\sum_i f_i Z_i}$$


$$\frac{\text{warm } \kappa_{IB}(\text{C}_5\text{H}_{12})}{\text{cryo } \kappa_{IB}(\text{He})} = \frac{4.57}{2} = 2.29$$

Directly depends on gas fill


Hydra modeling: electron temperature at mid peak power (19 ns):




Implosion must be symmetric for ignition



Warm shot history



Hotspot shape: less transfer to be round warm vs. cryo





- After shots 2 and 4 we chose $\Delta\lambda = 3.5$ Å as giving round hotspot

Suggests higher Z improves inner beam propagation

But warm 2D ConA's (last two shots) revealed pancaked in-flight shell

Cryo


warm




Hydra playbook [O. Jones]: $\Delta\lambda = 4.7 \text{ \AA}$ to make warm shell round

Best shape shots

	E_{laser} [kJ]	P_{peak} [TW]	Hohlraum length	$\Delta\lambda_{23,30}$ [Å]
Cryo N130314	1340	360	+700 um: low shell P_4	6.0, 4.5
Warm N130627	same	1.06x	same	3.5, 3.5






Tent talks (already happened):
Nagel, Haan, Town

Warm hotspot also complicated
by propane capsule fill: cooler
hotspot, significant shell
emission

Plot courtesy R. Town

¹R. H. H. Scott et al., PRL 2013

Warms have less inner SRS, more outer SBS than cryos. Pure $\Delta\lambda$ scaling, or gas fill effect?



Jury out on whether higher Z gas fill improves inner beam propagation

- Warm and cryo shots have different symmetry and backscatter
- Purely due to $\Delta\lambda$ or gas Z effect?
- Need to compare implosions with good symmetry throughout time
- Not achieved yet warm, maybe cryo (stalk vs. tent mount)
- Time-varying $\Delta\lambda$ or cone fraction may be needed
- Should check early time symmetry of warm shots – keyhole, re-emit

- **BACKUP SLIDES**

Jury out on whether higher Z gas fill improves inner beam propagation

Cryo shape:

- 2D ConA's: in-flight P_2 sausage and P_4
- Reduce $\Delta\lambda$ to remove P_2 , +700 um hohlraum to remove P_4
- Round shell swings to pancaked hotspot – tent?

Warm shape:

- Opposite P_2 swing from cryo: shell pancake but hotspot round!
- More $\Delta\lambda$ needed to remove shell P_2
- Warm hotspot complicated by cool hotspot (propane), shell emission

To settle if high-Z gas improves inner beam propagation, need to compare implosions with good symmetry throughout time

Time-varying $\Delta\lambda$ or cone fraction may be needed - warm and cryo

Should check symmetry and strength of warm first 3 shocks - keyhole or re-emit shot

Other relevant presentations – all before this one!

Tent and shape:

O. Jones, IFSA 2013

S. Nagel, Wed. AM, NO4.00014

S. Haan Tues. PM, JO7.00001

Low-mode asymmetries:


A. Kritcher Wed AM NO4.00004

R. Town Wed PM QI3.00002


Inline SRS Hydra modeling:

M. Marinak, Wed. AM, NP8.00090

Overall reflectivity slightly lower on warms. Pure $\Delta\lambda$ scaling, or gas fill effect?




P2 shape: positive for sausage (prolate), negative for pancake (oblate)




Equatorial
X-ray
Detector:

In-flight shell images
shot N130509



Hotspot images
shot N130405



-
- **Nuclear:** Deuterated propane C_3D_8 -> up to $2.6\text{E}11$ neutrons, T_{ion} up to 1.7 keV

But: 2D ConA shots show warms have large in-flight pancake, cryos had in-flight sausage

Red: warm

Blue: cryo

□: $L_{\text{hohl}} -300 \mu\text{m}$

⊕: $L_{\text{hohl}} +700 \mu\text{m}$


□: N121219: cryo $-300 \mu\text{m}$

⊕: N130211: cryo $+700 \mu\text{m}$


□: N130509: warm $-300 \mu\text{m}$

⊕: N130627: warm $+700 \mu\text{m}$

$P_0 = \text{radius}$




P_2




Outer beam SBS: DrD sensors show more on cone 50 than 44, and give power scaling

N130125:
970 kJ shot





“Tang curve:” avg. nonzero 50’s



- DrD = drive diagnostic sensor - at least one beam in each quad
 - 3ω power history - forward and backward (separated in time)
- N130125: one quad on each cone had 18% higher power: power scaling on one shot!
- Why more SBS on 50's than 44's?
 - 50 focal spot smaller -> higher intensity
 - Cross-beam energy transfer calculations: post-transfer power on 50's > 44's
 - Could be pure intensity scaling; plasma conditions may also play role


SBS on 50° outer cone

N130405:
1.3 MJ shot



- Cryo shots show some outer SBS late in time, esp. for longer pulses or high power
- Warm platform good for studying outer SBS and mitigation – cheaper, reproducible

**Warm radius slightly larger:
cooler hotspot, shell emission?**



NIF

