



Exceptional service in the national interest



ZAPP: The Z Astrophysical Plasma Properties Collaboration

Gregory A. Rochau

Presented at the 41st European Physical Society Conference on Plasma Physics – June 23-27, 2013



Sandia National Laboratories is a multi-program laboratory managed and operated by Sandia Corporation, a wholly owned subsidiary of Lockheed Martin Corporation, for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-AC04-94AL85000.

ZAPP represents a large collaboration between the NNSA labs and the academic community



Jim Bailey, Taisuke Nagayama,
Guillaume Loisel, Stephanie Hansen,
Dave Bliss, Tom Nash
Sandia National Laboratories



Roberto Mancini, Iain Hall, Tom
Lockard, Dan Mayes
University of Nevada – Reno



Don Winget, Mike Montgomery, Ross
Falcon, Thomas Gomez, Alan Wooton,
Jennifer Ellis, Sean Moorhead, Roger
Bengtson
University of Texas – Austin



Anhil Pradhan, C. Orban, Mark
Pinsonneault, and S.N. Nahar
Ohio State University



Mark Koepke
West Virginia University



Duane Leidahl, Carlos Iglesias, Brian Wilson
Lawrence Livermore National Laboratory



Manolo Sherrill, Heidi Tierney ,Chris Fontes,
James Colgan, Dave Kilcrease
Los Alamos National Laboratory






C. Blancard, Ph. Cosse, G. Faussurier, F.
Gilleron, J.C. Pain
**French Alternative Energies and Atomic
Energy Commission (CEA)**




Joe MacFarlane, Igor Golovkin
Prism Computational Sciences

Summary: ZAPP experiments measure fundamental properties of atoms in plasmas to solve important astrophysical puzzles.


- Why can't we predict the location of the convection zone boundary in the Sun?
 - Opacity of Fe at ~ 200 eV
- How does ionization and line formation occur in accreting objects and warm absorbers?
 - Ionization distribution and spectral properties of photoionized Ne and Si
- Why doesn't spectral fitting provide the correct properties for White Dwarfs?
 - Stark-broadened H-Balmer line profiles




Experiments on Z access a broad range of the energy-density phase space



Experiments on Z access a broad range of the energy-density phase space



The Z facility at Sandia National Laboratories is the most powerful pulsed power machine in the world.




23 MJ
Electrical

80 TW
Electrical

> 26 MA
Current
>300 TW,
> 2 MJ
X-ray


Wire Array Z pinch




$E_{\text{x-ray}} \sim 10\% E_{\text{Electric}}$

The z-pinch dynamic hohlraum (ZPDH) produces record currents of 25.8 MA with 1.5% reproducibility

Load Currents (20 shot average)



Z-pinch Dynamic Hohlraum



Standard ZPDH Characteristics

360 W wires – 11.4 μ m diameter

$m = 8.5$ mg W total

$V_{\text{marx}} = 85$ kV (20.3 MJ)

$I_p = 25.8 \pm 0.4$ MA [20 shots]

Sanford et al., POP 9 (2002)

Lemke et al., POP 12 (2004)


Bailey et al., POP 13 (2006)

Slutz et al., POP 13 (2006)


Rochau et al., PRL 100 (2008)

The ZPDH x-ray emission is reproducible to $\pm 10\%$ in peak power and $\pm 7\%$ in energy

Radial X-ray Power and Energy (20 shot average)




Z-pinch Dynamic Hohlraum




	ZR >2011	Z <2007
Marx Energy	20.3 MJ	11.4 MJ
Ipeak	25.8 MA (1.5%)	21.7 MA* (2.1%)
Mass	8.5 mg	3.8 mg
Peak Power	220 TW (10%)	120 TW (14%)
Radiated Energy	1.6 MJ (7%)	0.82 MJ (17%)


The ZPDH can also radiatively heat samples placed above the z-pinch to $T_e \sim 200$ eV.



Framing Pinhole Camera Images




Axial Fe Foil Temperature





The ZPDH simultaneously drives four independent experiments on a single ZAPP shot

1 Axial Experiment




3 Radial Experiments






The x-ray drive on radial samples needs to be corrected for the view-factor


Monochromatic Images (277 eV)



Calibrated VISRAD* view-factor model



Absolute Power Measurements




Assumption:


Each 'pixel' emits a Planckian spectrum at its characteristic T_r

The calibrated VISRAD model is used to infer the radiation drive spectrum at each sample

**r-θ Peak Brightness
Temperature Contours**




Peak Drive Flux on a Sample



ZAPP campaigns simultaneously study multiple issues spanning 200x in temperature and 10^6 x in density



Solar Opacity



Question:


Why can't we predict the location of the convection zone boundary in the Sun?

Achieved Conditions:

$T_e \sim 200$ eV, $n_e \sim 10^{23}$ cm $^{-3}$



Photoionized Plasmas



Question:

How does ionization and line formation occur in accreting objects?

Achieved Conditions:

$T_e \sim 20$ eV, $n_e \sim 10^{18}$ cm $^{-3}$



White Dwarf Line-Shapes



Question:

Why doesn't spectral fitting provide the correct properties for White Dwarfs?

Acheived Conditions:


$T_e \sim 1$ eV, $n_e \sim 10^{17}$ cm $^{-3}$



ZAPP campaigns simultaneously study multiple issues spanning 200x in temperature and 10^6 x in density



Solar Opacity



Question:

Why can't we predict the location of the convection zone boundary in the Sun?

Achieved Conditions:

$T_e \sim 200 \text{ eV}$, $n_e \sim 10^{23} \text{ cm}^{-3}$



Photoionized Plasmas



Question:

How does ionization and line formation occur in accreting objects?

Achieved Conditions:

$T_e \sim 20 \text{ eV}$, $n_e \sim 10^{18} \text{ cm}^{-3}$



White Dwarf Line-Shapes



Question:


Why doesn't spectral fitting provide the correct properties for White Dwarfs?

Achieved Conditions:

$T_e \sim 1 \text{ eV}$, $n_e \sim 10^{17} \text{ cm}^{-3}$



Models for solar interior structure disagree with helioseismology observations.



Convection-Zone (CZ) Boundary
Models are off by $10\text{-}30 \sigma$

Models depend on:

- Composition (revised in 2005*)
- EOS as a function of radius
- The solar matter *opacity*
- Nuclear cross sections


Question: Is opacity uncertainty the cause of the disagreement?

Objective: Measure Fe opacity at CZ base conditions.

The ZPDH radiating shock is used to both heat and backlight samples to stellar interior conditions.






Foil is heated during the ZPDH implosion

Thin Foil




Foil is backlit at shock stagnation

Thin Foil




The achieved temperature and density depend on the target design.

Thin Tamper




Thick Tamper




Modern computations of Fe opacity show large disagreements with data at CZ base conditions

Thin Tamper* (156 eV, $6.9 \times 10^{21} \text{ cm}^{-3}$)



Thick Tamper (182 eV, $31 \times 10^{21} \text{ cm}^{-3}$)



Present Status

- Agreement between data and computation becomes worse at increasing temp. and dens.
- Disagreements at CZ base conditions can partially explain the CZ boundary problem.
- The differences are probably not unique to Fe... more scrutiny of the data is prudent.

ZAPP campaigns simultaneously study multiple issues spanning 200x in temperature and 10^6 x in density



Solar Opacity



Question:


Why can't we predict the location of the convection zone boundary in the Sun?

Achieved Conditions:

$T_e \sim 200 \text{ eV}$, $n_e \sim 10^{23} \text{ cm}^{-3}$



Photoionized Plasmas



Question:

How does ionization and line formation occur in accreting objects?

Achieved Conditions:

$T_e \sim 20 \text{ eV}$, $n_e \sim 10^{18} \text{ cm}^{-3}$



White Dwarf Line-Shapes



Question:

Why doesn't spectral fitting provide the correct properties for White Dwarfs?

Achieved Conditions:


$T_e \sim 1 \text{ eV}$, $n_e \sim 10^{17} \text{ cm}^{-3}$



We learn about black holes from the matter falling into them – these are photoionized plasmas

Conceptual Picture of a Black-Hole Accretion Disk

$$\xi \sim 10 - 10,000 \text{ erg.cm.s}^{-1}$$



Photoionization parameter

$$\xi \equiv \frac{4\pi F}{n_e} \text{ [erg.cm.s}^{-1}\text{]}$$

Laboratory Plasmas

$$n_e \sim 10^{19} \text{ cm}^{-3}$$


$$F > 1 \text{ TW/cm}^2 \text{ for } \xi > 10$$

- Can we model the ionization?
- Can we model the line emission?

We learn about black holes from the matter falling into them – these are photoionized plasmas

Conceptual Picture of a Black-Hole Accretion Disk

$$\xi \sim 10 - 10,000 \text{ erg.cm.s}^{-1}$$



Photoionization parameter

$$\xi \equiv \frac{4\pi F}{n_e} \text{ [erg.cm.s}^{-1}\text{]}$$

Laboratory Plasmas


$$n_e \sim 10^{19} \text{ cm}^{-3}$$

$$F > 1 \text{ TW/cm}^2 \text{ for } \xi > 10$$

- Can we model the ionization?
- Can we model the line emission?

A Specific Problem: Emission from L-shell ions is not seen in some prominent black-hole accretion disks.

Measured Fe Emission from MCG 6-30-15




No observed emission from Fe ionized to the L-shell


Resonant Auger Destruction (RAD) was accepted as the reason*

- 2 competing processes for the de-excitation of L-shell ions:

Radiative Decay




Auger Decay



- **Thin Plasma:** high probability of observing the photon
- **Thick Plasma:** high probability of the photon being resonantly absorbed
→ Higher probability of Auger Decay for the ensemble


New models* suggest that RAD may not be as efficient as previously thought.



Question: Is Resonant Auger Destruction the reason we don't see emission from L-shell ions in some black-hole accretion disks?

Objective: Measure spectra in a highly photo-ionized lab plasma.

ZAPP experiments achieve $\xi \sim 20$ at the correct column depths to study the RAD question.



Measured Quantities

Flux..... $F = 1.3 \text{ TW/cm}^2$

Electron

Density..... $n_e = 8.5 \cdot 10^{18} e^-/\text{cm}^3$



Photoionization

Parameter..... $\xi = 19 \text{ erg.cm.s}^{-1}$



Column Density..... $N_i = 5 \cdot 10^{17} \text{ at/cm}^2$

Electron Temp..... $T_e = 19.5 \text{ eV}$

Si Absorption Measurements



Recent emission measurements demonstrate that L-shell emission is not 100% quenched by RAD.



Present Status

- Z Data demonstrates that L-shell emission does escape at column depths $>1E17$ at/cm².
- Present data can discriminate between models of the ionization distribution AND relative line strengths.
- Absolute intensity is needed to determine efficiency of RAD process.

ZAPP campaigns simultaneously study multiple issues spanning 200x in temperature and 10^6 x in density



Solar Opacity



Question:

Why can't we predict the location of the convection zone boundary in the Sun?

Achieved Conditions:

$T_e \sim 200$ eV, $n_e \sim 10^{23}$ cm $^{-3}$



Photoionized Plasmas



Question:

How does ionization and line formation occur in accreting objects?

Achieved Conditions:

$T_e \sim 20$ eV, $n_e \sim 10^{18}$ cm $^{-3}$



White Dwarf Line-Shapes



Question:

Why doesn't spectral fitting provide the correct properties for White Dwarfs?

Achieved Conditions:


$T_e \sim 1$ eV, $n_e \sim 10^{17}$ cm $^{-3}$




The properties of White Dwarfs are determined by spectral fitting, but disagrees with other methods

- White Dwarfs are fundamentally important
 - Evolutionary endpoint for ~98% of stars
 - Simple in structure and evolution
 - Cosmic laboratories (cosmochronology)
- WD surface temperature and total mass are usually determined by fitting the observed spectra
- The spectroscopic method and gravitational redshift disagree by >10% in the stellar mass

Spectral fit of WD J1916+3938*



New Stark broadened line-shape calculations* partially fix the problem – are they right?




Question: Are inaccurate H-Balmer line shapes responsible for the inaccurate determination of WD mass?

Objective: Measure H-Balmer line shapes at relevant temperature and density.


ZAPP experiments utilize radiatively heated gas cells to provide benchmark data for the WD problem

- Gas cells provide a precisely known atom density
- Large cell size provides optical depths needed for high-n lines
- Large cell minimizes the effect of boundary layers
- Long fielding distance provides uniform heating flux

Gas Cell Model





Gas Cell





Simultaneous streaked absorption and emission in absolute units provide a unique capability


Emission



Absorption




Measured H-Balmer line shapes can discriminate between theories



Present Status

- Measurement of relative line-shapes up to $n=7$ provides a strong constraint on models
- Additional measurements at higher density may be required to fully address the WD problem
- Continued scrutiny on the data is prudent:
 - Reproducibility of the result
 - Plasma uniformity

ZAPP experiments inform and challenge the interpretation of spectral data from the world's multi-billion dollar x-ray observatories.



NATURE Editorial: November, 2013

THIS WEEK

EDITORIALS

MENTORING The heavy responsibility to the next generation p.438

WORLD VIEW Beware the real risk of World Cup fever p.439



POISON Strawberry-frog parents give protection to kids p.441


Nailing fingerprints in the stars

Laboratory-based experiments are sorely needed to complement the rapidly proliferating spectral data originating from observations by the latest space telescopes.

What are stars made of? After astronomers detected a bright-yellow, unknown spectral line in sunlight in 1868, they named the new element helium after the Greek Sun god Helios. But it was some 30 years before physicists on Earth managed to detect — and so confirm the discovery of — helium in a laboratory.

It is a pattern that has been repeated many times since: the indirect detection of elements and molecules through spectral signatures in

quantum mechanics. But heavier elements have many electrons that can participate in transitions — iron has 26, making the probabilities of possible transitions between levels too complex to calculate accurately. Measuring emissions in the lab is the only alternative. Physicists can use tunable lasers to excite electrons into more levels and measure further transitions. This information can then feed back to the astronomical observations. Extra funds would significantly improve this



Chandra
\$2.8B



XMM-Newton
\$1.1B



Suzaku
\$0.5B

“Laboratory-based experiments are sorely needed to complement the rapidly proliferating spectral data originating from the latest space telescopes”

Extra Slides