

Mining the Agave Microbiome for adaptations to arid environments

Devin Coleman-Derr^{*1}, Tanja Wojke¹, Gretchen North², Laila Partida-Martinez³, Kristen DeAngeli⁴, Scott Clingenpeel¹, Stephen Gross¹, Susannah Tringe¹, Axel Visel¹

¹ Department of Energy Joint Genome Institute // LBNL - Walnut Creek, CA

² Department of Biology, Occidental College, Los Angeles, CA

³ Laboratory of Microbial Interactions, CINVESTAV, Irapuato, Mexico

⁴ Department of Microbiology, University of Massachusetts, Amherst, MA

^aTo whom correspondence may be addressed. E-mail: dacoleman-derr@lbl.gov

March 25, 2013

ACKNOWLEDGMENTS:

The work conducted by the US Department of Energy (DOE) Joint Genome Institute is supported by the Office of Science of the DOE under Contract Number DE-AC02-05CH11231. The views and opinions of the authors expressed herein do not necessarily state or reflect those of the United States Government, or any agency thereof, or the Regents of the University of California.

DISCLAIMER:

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California.

Introduction: A major challenge facing the biofuels industry is the identification of high-yield plant feedstocks that can be cultivated with minimal resource inputs without competing for land and water supplies with existing food crops. Recent research has demonstrated that the *Agave* plant, cultivated in Mexico and Southwestern United States for the production of fiber and alcohol, meets these criteria¹. Agaves grow on non-arable rocky soils in regions characterized by prolonged drought and extreme temperatures, due in part to physiological adaptations that prevent excess water-loss in arid environments². Plant-microbial symbioses can play a role in helping plants adapt to heat and drought stress, increasing the accessibility of soil nutrients, or compete with plant pathogens³. Whether agaves have similar beneficial microbe interactions in their native environment is unknown. We aim to provide a comprehensive characterization of the *Agave* microbiome, with the goal of identifying specific community members that may contribute to *Agave* biotic and abiotic stress tolerance.

Sampling Plan: We are investigating the microbial communities of both wild and cultivated *Agave* species. *Agave* specimens and associated soil samples were collected from three sites in Southern California and from four sites in Central Mexico. In California, all samples were collected from the wild species *Agave deserti*. In Mexico, samples of cultivated *Agave tequiliana* were collected from two *Agave* plantations, while the species *Agave salmiana* was collected from its native habitat. For comparison, a smaller number of samples were collected from two species of native cacti (*Myrtillocactus geometrizans* and *Opuntia robusta*).



Figure 1. Collection sites in Mexico and the United States

Sample Types:




Figure 2. The six sample types collected from all plant specimens.

For each *Agave* plant, samples were collected from the leaf episphere (leaf surface), rhizosphere (root surface) and leaf and root interiors (endospheres), as well as from surrounding soils (root zone and bulk soil). Additionally, sampling was repeated over two time points in 2012 (spring and summer), corresponding to the beginning and ends of the rainy season.

Sample Times:




Figure 3. Two sample collection points, in late spring and summer.

iTag Analysis Pipeline: A 16S iTag survey of all samples is being used to select specific microbiome compartments for further characterization, including shotgun metagenomics and single-cell genomics.




Figure 4. Flow chart for the pipeline of analysis tools used in processing the 16S iTag amplicon data. Initial processing of the MiSeq generated FASTQ reads is handled by a custom PERL script (itagger.pl) and subsequent statistical analyses are performed in the R platform.




Figure 5. Alpha and beta diversity plots by Sample type, Sampling Site, and Species. Each row of panels (from top to bottom) represent filtered readcounts, Shannon Diversity, and Phylum-level relative abundance.




Figure 6. Heatmap of log-abundance of readcounts in all *Agave deserti* samples for OTUs with the top two percent of variance across samples. Samples are ordered by Ward clustering of the Bray Curtis distances, and false colored at left by sample type. Indicator species analysis will be used to determine OTUs correlating with each compartment.




Figure 7. Principle Components Analysis of *Agave deserti* samples colored by sample type (top left), Bulk soil samples colored by location (top right), and Rhizosphere samples colored by species (bottom left). Plots were generated from the Bray Curtis distances for the measureable rarefied OTU table. The respective Analysis of similarity statistic is found in a corner of each plot.

Conclusion: We are investigating and comparing the microbial communities of native and cultivated *Agave* species in California and Mexico under a panel of different environmental conditions. Our project will expand our understanding of microbial diversity in desert soils, catalog and characterize the microbial factors that contribute to *Agave*'s successful adaptation to the extreme environments of its endemic range. Ultimately, we aim to enable microbiome manipulation aimed at improving the suitability of *Agave* for use in the rapidly growing biofuels industry.

References:

- Davis, S., et al. The global potential for *Agave* as a biofuel feedstock. *GCB BioEnergy*, 68-78, (2011).
- Nunez, H. et al. *Agave* for tequila and biofuels: an economic assessment and potential opportunities. *GCB BioEnergy*, 43-47, (2011).
- Rodriguez, R. et al. Stress Tolerance in plants via habitat-adapted symbiosis. *ISMEJ*, 404-416, (2008).

Contact:

Devin Coleman-Derr, Postdoctoral Researcher, Microbial Ecology Group, Joint Genome Institute, Walnut Creek, CA, dcoleman-derr@lbl.gov