

Transcriptomic and Metagenomic Exploration of Agave Drought and Stress Tolerance

Stephen Gross ^{1,2}, Jeffrey Martin ^{1,2}, June Simpson ³, Laila Partida-Martínez ³, Gretchen North ⁴,
Yuri Peña-Ramirez ⁵, Aidee Orozco-Hernandez ⁵, Kristen DeAngelis ⁶, Tanja Woyke ^{1,2}, Scott
Clingenpeel ^{1,2}, Susannah Tringe ^{1,2}, Zhong Wang ^{1,2}, and Axel Visel ^{1,2}

1 Department of Energy Joint Genome Institute, Walnut Creek, CA

2 Genomics Division, Lawrence Berkeley National Lab, Berkeley, CA

3 CINVESTAV, Irapuato, Guanajuato, Mexico

4 Occidental College, Los Angeles, CA

5 Brown-Forman, Inc., Amatitán, Jalisco, Mexico

6 University of Massachusetts, Amherst, MA

^aTo whom correspondence may be addressed. E-mail: smgross@lbl.gov

March 25, 2012

ACKNOWLEDGMENTS:

The work conducted by the US Department of Energy (DOE) Joint Genome Institute is supported by the Office of Science of the DOE under Contract Number DE-AC02-05CH11231. Agave microbiome work is supported in part by the 2012 DOE JGI Community Sequencing Program. The views and opinions of the authors expressed herein do not necessarily state or reflect those of the United States Government, or any agency thereof, or the Regents of the University of California.

DISCLAIMER:

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California

Transcriptomic and Metagenomic Exploration of Agave Drought and Stress Tolerance



Stephen Gross ^{1,2}, Jeffrey Martin ^{1,2}, June Simpson ³, Laila Partida-Martínez ³, Gretchen North ⁴, Yuri Peña-Ramirez ⁵, Aidee Orozco-Hernandez ⁵, Kristen DeAngelis ⁶, Tanja Woyke ^{1,2}, Scott Clingenpeel ^{1,2}, Susannah Tringe ^{1,2}, Zhong Wang ^{1,2}, and Axel Visel ^{1,2}

¹ Department of Energy Joint Genome Institute, Walnut Creek, CA
² Genomics Division, Lawrence Berkeley National Lab, Berkeley, CA
³ CINVESTAV, Irapuato, Guanajuato, Mexico
⁴ Occidental College, Los Angeles, CA
⁵ Brown-Forman, Inc., Amatitlán, Jalisco, Mexico
⁶ University of Massachusetts, Amherst, MA




Abstract

Understanding drought and heat tolerance in plants will be critical for sustained agriculture and bioenergy production in a changing global climate. Because of their exceptional ability to thrive in arid, hot environments with minimal soil nitrogen, Agave species have been identified as a candidate bioenergy feedstock and provide a model for studies of adaptations to drought and heat. While some physiological mechanisms underlying these traits have been studied in detail, the molecular basis of the extreme heat and drought tolerance in these plants remains unclear. We have constructed *de novo* reference transcriptomes for *Agave tequilana*, an economically important species cultivated in Mexico for spirit distillation, and *A. deserti*, an extremely thermo- and drought-tolerant species native to the Colorado Desert, using 368 and 185 gigabasepairs, respectively, of Illumina short-read sequence data. These data sets enable cross-species comparative analysis of protein-coding sequences and tissue-specific gene expression levels. Based on these reference transcriptomes, we also perform quantitative expression profiling of agaves subjected to heat and drought stress in controlled greenhouse experiments, enabling a transcriptome-wide understanding of Agave stress responses. Complementary to plant-endogenous mechanisms, Agave plants are associated with microbes that may play important roles in mitigating abiotic stress in arid resource-limited environments. Using targeted sequence-based microbial community profiling, we study the microbiomes of cultivated *A. tequilana*, and wild *A. deserti* and *A. salmiana*. We also use deep metagenomic sequencing to gain further insight into Agave rhizosphere and phyllosphere microbe communities, and perform isolation and single-cell genome sequencing of individual microbial cells residing within Agave tissues. These studies aim to identify microbial species, genes, and pathways conferring additional stress resistance to agaves. Taken together, our work builds a robust platform to accelerate discovery of plant and plant-associated microbial adaptations to major abiotic stresses.

Overview

I. Agave can supplement other bioenergy feedstocks

Agave species, adapted to their native habitat in arid regions of Mexico and the United States, hold promise as a biofuel feedstock [1], capable of growing on marginal lands where other bioenergy plants cannot. The ability of agaves to withstand hot and arid conditions relies upon Crassulacean Acid Metabolism (CAM)—a specialized form of photosynthesis allowing agaves to keep leaf stomata (pores) closed during the day, minimizing water loss through evapotranspiration.



II. Agaves are productive with minimal resources

Agaves are capable of producing lignocellulosic biomass with little water and nitrogen inputs. Species *A. salmiana* and *A. mapisaga* have been reported to produce up to 40 metric tonnes (Mg) of dry biomass per hectare per year [2].


Feedstock	Inputs			Outputs	
	Water (cm yr ⁻¹)	Drought tolerance	Nitrogen (kg ha ⁻¹ yr ⁻¹)	Dry biomass (Mg ha ⁻¹ yr ⁻¹)	Ethanol (liters yr ⁻¹)
Corn grain	50–80	low	90–120	7–10	2900
Corn stover				3–6	900
<i>Miscanthus</i>	75–120	low	0–15	15–40	4600–12,400
Poplar coppice	70–105	moderate	0–50	5–11	1500–3400
Agave spp.	30–80	high	0–12	10–34	3000–10,500

Comparison of inputs (water and nitrogen) and outputs (biomass and ethanol) of agaves and other biofuel feedstock species. Though agaves are harvested at several years of age, their annualized growth rate is comparable to *Miscanthus*. Table is modified from reference [4].

Building Agave reference transcriptomes

De novo assembly of Agave transcriptomes

Without sequence information, molecular studies of agaves are difficult. To address this need, we chose *A. tequilana*, which is currently cultivated for tequila production, and *A. deserti*, an extremely drought and heat tolerant agave, as our reference species. Agaves have large genomes (~4–7 Gb) [5, 6], so we focused on sequencing the protein-coding transcriptome using Illumina RNA-seq and *de novo* transcriptome assembly with Rnnotator [7], a *de novo* transcriptome assembly pipeline developed at JGI.




Robust transcriptome assemblies

A resource for Agave molecular biology

Transcriptome assemblies were filtered for artifacts and sequences from associated microorganisms. The resulting Agave transcriptomes, while not a 100% complete representation of Agave genes, contain a significant number of loci and encode the majority of proteins within other plant species.

A	Species dataset	Raw sequence	No. of loci	No. of transcripts	N50 transcript length	Assembled transcriptome size
	<i>A. tequilana</i>	368 Gbp	139,525	204,530	1387 nt	204.8 Mbp
	<i>A. deserti</i>	185 Gbp	88,718	128,869	1323 nt	125.0 Mbp

A. Summary of transcriptome assemblies for *A. tequilana* and *A. deserti*.



B. Comparing the *A. deserti* and *A. tequilana* transcriptomes by BLAT alignment. The majority of transcript in one species aligns to a transcript in the other.


C. Alignment of 82 *A. tequilana* sequences in GenBank to the *A. tequilana* transcriptome assembly by BLAT. 81 of 82 sequences were found.

D. Shared OrthoMCL v5 [8] protein clusters between *Arabidopsis*, rice, *A. tequilana* and *A. deserti*.

Expression profiling of agaves

Adding functional data to annotation

With reference transcriptomes in place, expression profiling experiments can be initiated. Using data from the transcriptome assemblies, tissue-specific expression profiles can be studied. Additional experiments are in progress to understand transcriptome responses to heat and drought stress.



A. Heatmap of Photosystem I and II genes expressed in *A. deserti* roots, folded leaves, and sections along the proximal-distal axis of an unfolded leaf. Data indicates photosystem genes are most highly expressed in the distal half of the leaf, while the proximal end, lacking chlorophyll pigmentation, has relatively low expression of photosystem genes.


B. Heatmap of KNOX gene expression along the proximal-distal axis of an unfolded leaf. Arrow indicates the ortholog of *Zea mays* Knotted1 (*Kn1*), expressed higher in proximal leaf sections than distal leaf sections.

C. Overview of ongoing expression profiling experiments, testing *A. tequilana* responses to prolonged drought and heat.

Agave microbiomes and adaptations to stress

Discovering the Agave microbial community

We have initiated sequence-based studies of the *A. tequilana*, *A. salmiana* and *A. deserti* microbiomes. Ultimately, we aim to identify microbes conferring stress and disease resistance to agaves. With both transcriptome and microbe sequences in hand, we will have a strong foundation for powerful plant-microbe interaction studies.



A. Microscopic examination of epiphytic microbes associated with the Agave phyllosphere (leaf surface). Red-stained plant cell walls, blue—epiphytic microbes.

B. Microbial communities of Agave to be examined by sequence-based techniques.

C. Fast UniFrac [9] PCoA analysis of *A. deserti* microbial communities collected during Summer 2011 in Southern California. Data demonstrate distinct microbial communities associated with different parts of the plant.

Acknowledgements and References

This work performed at the U.S. Department of Energy Joint Genome Institute was supported in part by the Office of Science of the U.S. Department of Energy under contract DE-AC02-05CH112. Agave microbiome work is supported in part by the 2012 DOE JGI Community Sequencing Program.

- [1] Davis, A. S. et al. The global potential for Agave as a biofuel feedstock. *GCB Bioenergy* 3, 68–78, (2011).
- [2] Nobel, P. S. et al. High annual productivity of certain agaves and cacti under cultivation. *Plant Cell Environ* 15, 329–35, (1992).
- [3] Simpson, J. et al. Genomic resources and transcriptome mining in *Agave tequilana*. *GCB Bioenergy* 3, 25–36, (2011).
- [4] Somerville, C. et al. Feedstocks for lignocellulosic biofuels. *Science* 329, 790–2, (2010).
- [5] Palomino, G. et al. Nuclear genome size analysis of *Agave tequilana* Weber. *Caryologia* 56, 37–46, (2003).
- [6] Robert, M. L. et al. Wild and agronomically important *Agave* species (Asparagaceae) show proportional increases in chromosome number, genome size, and genetic markers with increasing ploidy. *Bot J Linn Soc* 158, 215–22, (2008).
- [7] Martin, J. et al. Rnnotator: an automated *de novo* transcriptome assembly pipeline from stranded RNA-seq reads. *BMC Genomics* 11, 663, (2010).
- [8] Chen, F. et al. OrthoMCL-DB: querying a comprehensive multi-species collection of ortholog groups. *Nucleic Acids Res* 34, D363–8, (2006).
- [9] Hamady, M. et al. Fast UniFrac: facilitating high-throughput phylogenetic analyses of microbial communities including analysis of pyrosequencing and PhyloChip data. *ISME J* 4, 17–27, (2010).