

Microbial diversity in restored wetlands of San Francisco Bay

Susanna Theroux^{1*}, Wyatt Hartman¹, Shaomei He^{1,2}, and Susannah Tringe¹

¹ LBNL Department of Energy Joint Genome Institute, 2800 Mitchell Drive, Walnut Creek, USA

² University of Wisconsin-Madison - Madison, WI, USA

**To whom correspondence should be addressed:* Email: stheroux@lbl.gov

December 9, 2013

ACKNOWLEDGMENTS:

The work conducted by the U.S. Department of Energy Joint Genome Institute is supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. The project was funded by the DOE Early Career Research Program, grant number KP/CH57/1 to Susannah Green Tringe and was also supported by DOE JGI Community Sequencing Program.

DISCLAIMER:

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California.

Microbial diversity in restored wetlands of San Francisco Bay

Susanna Theroux^{1*}, Wyatt Hartman¹, Shaomei He^{1,2}, Susannah Tringe¹

1. DOE Joint Genome Institute, Walnut Creek, CA 2. University of Wisconsin-Madison, Madison, WI

*Contact: stheroux@lbl.gov

ABSTRACT

Wetland ecosystems may serve as either a source or a sink for atmospheric carbon and greenhouse gases. This delicate carbon balance is influenced by the activity of below-ground microbial communities that return carbon dioxide and methane to the atmosphere. Wetland restoration efforts in the San Francisco Bay-Delta region may help to reverse land subsidence and possibly increase carbon storage in soils. However, the effects of wetland restoration on microbial communities, which mediate soil metabolic activity and carbon cycling, are poorly studied. In an effort to better understand the underlying factors which shape the balance of carbon flux in wetland soils, we targeted the microbial communities in a suite of restored and historic wetlands in the San Francisco Bay-Delta region. Using DNA and RNA sequencing, coupled with greenhouse gas monitoring, we profiled the diversity and metabolic potential of the wetland soil microbial communities along biogeochemical and wetland age gradients. Our results show relationships among geochemical gradients, availability of electron acceptors, and microbial community composition. Our study provides the first genomic glimpse into microbial populations in natural and restored wetlands of the San Francisco Bay-Delta region and provides a valuable benchmark for future studies.

OBJECTIVES

The aims of our study are to:

- ☐ Sample microbial diversity across wetland age and salinity gradients
- ☐ Detect microbial response to geochemical fluctuation and the influence of above-ground plant species
- ☐ Link microbial diversity to carbon flux measurements from wetland soils
- ☐ Evaluate wetland carbon source/sink potential and associated indicator microbial species

SITE DESCRIPTIONS

Twitchell Island wetland (121.65°W, 38.11°N) is a pilot-scale wetland restoration project established in 1997 to assess carbon storage land subsidence-reversal potential of converted agricultural land in the Sacramento-San Joaquin River Delta (**Figure 1a**). The wetland consists of two adjacent ponds, each about 3 ha (**Figure 1b**) with artificially maintained water depths of 25cm and 55cm. We sampled the West Pond along a transect from the water source inlet (A) to points furthest from the inflow (C/L, **Figure 1b**) in February and August 2011.




Figure 1. Twitchell wetland (a) and sampling locations (b).

In Phase II of the project, we sampled a suite of wetlands in San Pablo Bay, Suisun Bay and in the Delta region of San Francisco Bay (**Figure 2**). These wetlands encompassed a salinity gradient ranging from 0ppt (Sandmound Slough) to 40ppt (China Camp). Five wetlands sampled were historical wetlands (blue markers, **Figure 2**) and six wetlands were restored wetlands (red markers, **Figure 2**).



Figure 2. Wetland sampling locations in San Pablo Bay, Suisun Bay and the SF Delta. Blue markers are historical wetlands, red markers are restored wetlands.

Phase I Results: Twitchell Wetland

Increasing methane production rates were observed at Twitchell wetland along the transect from site A→C/L (**Figure 1**). Methane emissions were highest at the Twitchell wetland sites C/L which were furthest from the water inlet (**Figure 3**). These higher CH₄ emissions corresponded to sites with lowest oxygen and nitrate availability (He et al., in review).




Figure 3. Methane production at Twitchell sites in February and August.

Microbial communities were strongly associated with sampling site (A, B, C/L) and plant type (Cattail, Tule, Bulk soil, **Figure 4**). High methane-production sites were enriched for phyla typically observed in anoxic environments (Riviere et al., 2009), consistent with the expectation that after inundation, soil environments become more anaerobic and methanogenic.




Figure 4: Nonmetric multi-dimensional scaling (NMDS) of relative Operational Taxonomic Unit (OTU) abundance.

MATERIALS AND METHODS




Figure 5. Sampling schematic for San Pablo Bay, Suisun Bay, and Delta sites.

Twitchell wetland: Soils were dissected into bulk, tule rhizome and cattail rhizome samples. A subsample was taken from February 2011 samples for 12-day lab incubation studies to quantify greenhouse gas (GHG) flux. For August 2011 samples, stationary whole-plant chambers were used to measure GHG flux. Soil samples were extracted for DNA and microbial community profiles were generated by pyrotag sequencing targeting the 16S rRNA gene.

San Pablo/Suisun/Delta wetlands: Soil cores were collected using an AMS split core sampler. Adjacent DNA/RNA and GHG cores were collected in triplicate for two plant types at each site (**Figure 5**). Intact soil cores were analyzed for CO₂/CH₄ flux on a Los Gatos Research GHG Analyzer.



Bulk Decomposed



Cattail Rhizome



Tule Rhizome

Phase II Results: San Pablo Bay, Suisun Bay, Delta Wetlands




Figure 6. Methane production versus salinity for historic and restored wetlands.

Methane production rates clustered along salinity and plant type gradients (**Figure 7a**). Tule (*Schoenoplectus acutus*) and cattail (*Typha* spp.) were associated with the highest rates of methane production. Net methane consumption was measured at the oligohaline, historic wetland Sandmound Slough in the tule and three-square bulrush soils.




Figure 7. Methane production versus salinity for historic and restored wetlands separated by plant type (a) and by sampling location for individual plant types (b-d).

The influence of sampling location (**7b-d**), salinity, and plant type on methane production were all statistically significant (ANOVA, $p > 0.0001$), although these factors co-vary. Site was a stronger predictor of methane flux ($R^2 = 0.62$) than salinity ($R^2 = 0.144$) or plant type ($R^2 = 0.40$), although only methane flux from freshwater restored wetlands was significantly greater than the mean.

CONCLUSIONS

- ☐ At Twitchell wetland, microbial community composition is primarily governed by sampling location and sample types (bulk soil, tule rhizome, cattail rhizome).
- ☐ Low methane production samples at Twitchell wetland were associated with the site closest to the inflow, correlated to higher availabilities of sulfate, nitrate and oxygen. Lower numbers of methanogen OTUs were detected at these sites.
- ☐ Soil methane production clustered according to sampling location, plant type and salinity along the San Pablo/Suisun/Delta wetland sampling survey. Sampling location had the strongest influence on methane production.
- ☐ Restored wetlands had higher average methane production rates than historic wetlands possibly due to inundation regime, larger labile carbon pools or decreased availability of terminal electron acceptors.

REFERENCES

- He S, Malfatti S, McFarland JW, Anderson FE, Pati A, Huntemann M, Tremblay J, Glavina del Rio T, Waldrop M, Windham-Myers L, Tringe SG. In review. Patterns in wetland microbial community composition and functional gene repertoire associated with methane emissions. ISME J.
- Poffenberger H, Needelman B, Megonigal J. (2011). Salinity influence on methane emissions from tidal marshes. Wetlands 31:831-842.
- Riviere D, Desvignes V, Pelletier E, Chaussonnerie S, Guermazi S, Weissenbach J, Li T, Camacho P, Sghir A. (2009). Towards the definition of a core of microorganisms involved in anaerobic digestion of sludge. ISME J 3: 700-714.