

Genome Editing in Escherichia coli with Cas9 and synthetic CRISPRs

Ze Peng*, Sarah Richardson, David Robinson, Samuel Deutsch and Jan-Fang Cheng

US Department of Energy Joint Genome Institute, Walnut Creek, CA 94598

*Email Address: zpeng@lbl.gov

March 2014

The work conducted by the U.S. Department of Energy Joint Genome Institute is supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231


DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California.

Ze Peng*, Sarah Richardson, David Robinson, Samuel Deutsch and Jan-Fang Cheng
US Department of Energy Joint Genome Institute, Walnut Creek, CA 94598
*Email address: zpeng@lbl.gov


CRISPR systems are prokaryotic adaptive immune systems which integrate short fragments of foreign DNA into the host chromosome at one end of repetitive element known as a CRISPR (clustered regularly interspaced short palindromic repeat). The CRISPR serves as memory, which associated proteins (e.g. Cas9) then read in order to recognize and destroy invasive DNA.

Recently, the Cas9-CRISPR system has proven to be a useful tool for genome editing in eukaryotes, which repair the double stranded breaks made by Cas9 with non-homologous end joining or homologous recombination. *Escherichia coli* lacks non-homologous end joining and has a very low homologous recombination rate, effectively rendering targeted Cas9 activity lethal. We have developed a heat curable, serializable, plasmid based system for selectionless Cas9 editing in arbitrary *E. coli* strains that uses synthetic CRISPRs for targeting and λ -red to effect repairs of double stranded breaks. We have demonstrated insertions, substitutions, and multi-target deletions with our system, which we have tested in several strains.




Genome Editing in *E. coli* HME63 cells (one & two targets)

We have tested several editing conditions of *rpsL* and *Amp* in HME63 host cells which have genome integrated λ -red. We have found that: *rpsL* is easier to edit than *amp*; oligos causing deletions have the highest success rate; and the first spacer in CRISPR has a higher editing rate than the second.



mdfA gene deletion in DH1 cells


We made *mdfA* gene deletions in DH1 cells transformed with vector pKD46(λ -red) and vector pBHK-Cas9. The basic procedure, gene and arms structures before and after deletion and results are shown below:



Ten cell membrane transport related genes have been deleted in *E. coli* DH1 cells


We constructed a new pKD46 derived vector which contains both λ -red and Cas9. We created ten knock out strains for cell membrane transport related protein genes using this new vector, which retains the temperature sensitivity of the original plasmid.

#	gene	deletions	gene deleted clone
1	mdfA	(8/19)	EC112
2	basB	(2/11)	EC096
3	exbB	(2/24)	EC097
4	glpT	(1/3)	EC098
5	mdlB	(2/7)	EC099
6	tehA	(3/12)	EC100
7	ycjF	(5/12)	EC101
8	yhjD	(1/13)	EC102
9	yhjG	(4/20)	EC104
10	yhjX	(1/14)	EC103



A long gene cluster has been deleted in *E. coli* DH1 cells

We knocked out the 4kb thiamine synthesis gene cluster in DH1 cells, with single and double spacer CRISPRs.



Cas9 Editing Also Works in Other *E. coli* strains

Besides HME63 and DH1, we have attempted CRISPR-based editing with the strains EPI300 (top), S17-1 (middle) and JM109 (bottom).

Given our success with *E. coli*, we hope to move into other useful prokaryotic model systems.

