


Dirac Composite Fermions at the Half-Filled Landau Level

Wei Pan


Sandia National Laboratories
Albuquerque, NM, USA

Main result – CFs are probably Dirac fermions

resistivity scaling



linear density dependence




samples and measurements

Outline:

- Introduction
- Resistivity scaling at $\nu=1/2$
- Density dependence of the CF conductivity

Quantum Hall effects





$$R_{xy} = (h/e^2)/\nu$$

IQHE – single particle
 $\nu = 1, 2, 3 \dots$

FQHE – many-body
 $\nu = 1/3, 2/5, 3/7 \dots$
 $2/3, 3/5, 4/7 \dots$

similarity between IQHE and FQHE: R_{xx}






similarity between IQHE and FQHE: R_{xy}

Composite Fermion (CF) Model

Jain, PRL 1989


Halperin, Lee, and Read (HLR), PRB 1993

one composite fermion = one electron + 2 flux quanta



$$B_{\text{eff}} = B - 2\Phi_0 \times n = B - 2nh/e = B - B_{1/2}$$

Classical transport features around $\nu=1/2$



$$\nu = 1/2$$

$$B_{\text{eff}} = B - B_{1/2} = 0$$

CFs form fermi sea

IQHE of CF FQHE of e^-

p

$$\nu = p/(2p+1)$$

1

$1/3$


2

$2/5$

3

$3/7$

....



Unclarified aspects of CF theory

.....

The most fundamental problem is the lack of particle-hole symmetry

(copied from Son's presentation at the workshop on Strongly Interacting Topological Phases)


Kivelson, Lee, Krotov, and Gan, PRB (1997).
Lee, PRL (1998).
Kamburov et al. PRL (2014).
Barkeshli, Mulligan, and Fisher, PRB (2015).
Balram, Toke, and Jain, PRL (2015).
Murthy and Shankar, arXiv: 1508.06974.

Dirac composite fermions


Son, *Phys. Rev. X* **5**, 031027 (2015); Metlitski and Vishwanath, arXiv:1505.05142;
Wang and Senthil, arxiv:1507.08290; Geraedts et al, arXiv:1508.04140; ...

$$\frac{\rho_{xy} - 2}{\rho_{xx}^2} = \tilde{\sigma}_{xy} - \frac{1}{2}$$


HLR CFs: $(\rho_{xy}-2)/\rho_{xx}^2 = 0$



Dirac CFs: $(\rho_{xy}-2)/\rho_{xx}^2 = -1/2$




Sample: high quality quantum well




PRL (2003)
PRB (2015)

Density: $n = 1.190 \times 10^{11} \text{ cm}^{-2}$
Mobility: $\mu \sim 13 \times 10^6 \text{ cm}^2/\text{Vs}$

Magneto-transport measurements around $\nu=1/2$





Highly uniform 2DEG






Accurate Hall resistivity

Quantization better than 0.1%












$$B_{1/2} = 9.814\text{T}$$





$$\frac{\rho_{xy} - 2}{\rho_{xx}^2} = -0.7$$

- Finite thickness
- Landau level mixing



$$\kappa = e^2/\epsilon l_B/\hbar\omega_c = 0.87 \text{ at } \nu = 1/2$$

Details ...


- Mixing of ρ_{xx} and ρ_{xy}

$$\rho_{xy} = \rho_{xy}^0 + \alpha\rho_{xx}$$


$$\rho_{xy}(+B) = \rho_{xy}^0(B) + \alpha\rho_{xx}(B)$$

$$\rho_{xy}(-B) = -\rho_{xy}^0(B) + \alpha\rho_{xx}(B)$$

$$\rho_{xy}^0 = (\rho_{xy}(+B) - \rho_{xy}(-B))/2$$





The position of the 1/2 state



straight line, through origin

$$B_{\nu=1/2} = 9.843\text{T}$$







- Landau level mixing $\kappa = e^2/\epsilon l_B/\hbar\omega_c = 0.87$ at $\nu = 1/2$

DENSITY DEPENDENCE OF CF CONDUCTIVITY



HIGFET

(Heterojunction-Insulated Gate Field-Effect Transistor)

high mobility down to very low densities









HIGFET



Kane, Pfeiffer, West, and Harnett, APL, 1993


Straight sidewall is important







device works!

Linear I-V at very low densities




SdH oscillations at $B \sim 0.2\text{T}$







Composite fermions in HIGFET




CF conductivity *versus* density




CF conductivity





Electron conductivity







Linear dependence in graphene



Tan *et al.* PRL (2007).

CFs

- Resistivity scaling
- Linear density dependence of conductivity



Collaborators:


W. Kang

University of Chicago

K.W. Baldwin, K.W. West, L.N. Pfeiffer, and D.C. Tsui

Princeton University

Thank you



Very big density tunable range

$\sim 1 \times 10^9$ to $\sim 7.5 \times 10^{11} \text{ cm}^{-2}$

