

LA-UR-18-20421

Approved for public release; distribution is unlimited.

Title: Fission modeling at LANL

Author(s): Stetcu, Ionel

Intended for: Talk for visiting students from MSU

Issued: 2018-01-19

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



Fission Modeling at LANL

Ionel Stetcu

Theoretical Division, Los Alamos National Laboratory



A Comprehensive Understanding of Fission?

Fission Fragment Yields






[SPIDER]

Fission Cross Sections & Fission Fragment Angular Distributions




[TPC]

Prompt Fission Neutrons & Photons



[Chi-nu]



[DANCE]

T-2 Nuclear Physics Group, Theoretical Division, LANL

T-2 group
(working on nuclear fission)

Selected Fission Topics


- Toshihiko Kawano
- Patrick Jaffke (postdoc)
- Skip Kahler*
- Eric J. Lynn (consultant)
- Peter Möller*
- Denise Neudecker (XCP-3)
- Arnie Sierk*
- Ionel Stetcu
- Patrick Talou

- ◆ (n,f) , (γ,f) , (t,pf) , etc. cross sections
- ◆ Fission fragment yields $Y(A,Z,TKE)$
- ◆ Fragment angular distributions
- ◆ The $(n,\gamma f)$ process
- ◆ Prompt fission neutron spectrum
- ◆ Correlations in prompt fission data
- ◆ Prompt fission γ rays
- ◆ Benchmarking (not just fission) and evaluations

Fission Fragment yields

Fission Fragment Yields

(slide borrowed from Peter Möller & J. Randrup)



Brownian shape motion

Nuclear deformation energy: $E_{\text{def}}(i, j, k, l, m)$


Bias potential: $V_{\text{bias}}(i) = V_0 (Q_0/Q_2)^2$

Level density parameter: $a_A = A/(8 \text{ MeV})$

Temperature T : $E^* - E_{\text{def}} = a_A T^2$

$$\Rightarrow V(\chi) = E_{\text{def}} + V_{\text{bias}}$$

χ {
 i
 j
 k
 l
 m



P. Möller *et al*, Nature 409 (2001) 785

Metropolis walk:





N. Metropolis *et al*, J Chem Phys 26 (1953) 1087

Change shape: $\chi \rightarrow \chi'$?

$$\begin{cases} V(\chi') < V(\chi): \text{ move with } P = 1 \\ V(\chi') > V(\chi): \text{ move with } P = \exp(-\Delta V/T) \end{cases}$$

Scission: Critical neck radius $c_0 \approx 2.5 \text{ fm}$

Selected results



P. Möller



Fission Fragment Yields

- Results are very promising
P. Möller and C. Schmitt, EPJA 53, 7 (2017)
- E^* -dependent yields parameterized by 5 Gaussians – *Patrick Jaffke* (postdoc)

$$Y_i(A, E^*) = \frac{W_i(E^*)}{\sqrt{2\pi\sigma_i^2(E^*)}} \exp \left[-\frac{(A - \mu_i(E^*))^2}{2\sigma_i^2(E^*)} \right]$$

Fermi-function fits

Quadratic fits



→ Working on Pu, U and Np series

Langevin model calculations

- Classical dissipative system with fluctuating damping forces

Nuclear shape coordinates

$$\begin{aligned} \frac{dq_j}{dt} &= \frac{\partial H}{\partial p_j} = \frac{\partial(K + V)}{\partial p_j} = \frac{\frac{1}{2} M_{jk}^{-1} p_j p_k}{\partial p_j} = M_{jk}^{-1} p_k \\ \frac{dp_j}{dt} &= \frac{\partial H}{\partial q_j} = -\frac{\partial V}{\partial q_j} \dot{q}_k \dot{q}_l - \eta_{jk} \dot{q}_k + \sqrt{\frac{2T}{\Delta t}} \gamma_{jk} \Theta_k(t) \end{aligned}$$


Inertia tensor

Stochastic function


Dissipation tensor Nuclear temperature

A. J. Sierk



Some results with the Langevin model



^{235}U



^{239}Pu



Fission Fragment properties within TD-SLDA

Highlight from first microscopic calculation of ^{240}Pu fission:

- **Fully unrestricted 3D calculation with full nuclear energy density functional**
- **Long evolution times from outer fission saddle to full scission**
- **TKE predicted within 3% of expected values**
- **Light fission fragment heavily deformed, heavy fission fragment spherical (as expected)**
- **Demonstrated the essential role played by pairing correlations in nuclear shape evolution**

Bulgac et. al, PRL **116**, 122504 (2016)


Time-Dependent Density Functional Theory with Nuclear Energy Density Functional a la Skyrme

Construct constrained initial state near the outer fission barrier

Evolve in time beyond the scission point


Full characterization of fission fragment properties as a function of initial excitation energy of the fissioning nucleus (neutron incident energy):

- Average mass, charge, excitation energy, and angular momentum
- TKE of fission fragments



I. Stetcu

Average fission fragment properties



- Significant amount of collective shape and pairing oscillations
- Time from saddle to fission is sensitive to the functional




TABLE I. The simulation number, the pairing parameter η , the excitation energy (E^*) of $^{240}\text{Pu}_{146}$ and of the fission fragments [$E_{H,L}^* = E_{H,L}(t_{\text{SS}}) - E_{\text{gs}}(N_{H,L}, Z_{H,L})$], the equivalent neutron incident energy (E_n), the scaled initial mass moments $q_{20}(0)$ and $q_{30}(0)$, the “saddle-to-scission” time t_{SS} , TKE evaluated as in Ref. [71], TKE, atomic (A_L^{syst}), neutron (N_L^{syst}), and proton (Z_L^{syst}) extracted from data [72] using Wahl’s charge systematics [73] and the corresponding numbers obtained in simulations, and the number of postscission neutrons for the heavy and light fragments ($\nu_{H,L}$), estimated using a Hauser-Feshbach approach and experimental neutron separation energies [8,74,75]. Units are in MeV, fm², fm³, fm/c as appropriate.


S no.	η	E^*	E_n	q_{zz}	q_{zzz}	t_{SS}	TKE^{syst}	TKE	A_L^{syst}	A_L	N_L^{syst}	N_L	Z_L^{syst}	Z_L	E_H^*	E_L^*	ν_H	ν_L
S1	0.75	8.05	1.52	1.78	-0.742	14 419	177.27	182	100.55	104.0	61.10	62.8	39.45	41.2	5.26	17.78	0	1.9
S2	0.5	7.91	1.38	1.78	-0.737	4360	177.32	183	100.56	106.3	60.78	64.0	39.78	42.3	9.94	11.57	1	1
S3	0	8.08	1.55	1.78	-0.737	14 010	177.26	180	100.55	105.5	60.69	63.6	39.81	41.9	3.35	29.73	0	2.9
S4	0	6.17	-0.36	2.05	-0.956	12 751	177.92	181		103.9		62.6		41.3	7.85	9.59	1	1

Bulgac et. al, PRL **116**, 122504 (2016)


Fission Dynamics

Fission of ^{240}Pu at excitation energy $E_x = 8.08 \text{ MeV}$


Neutron pairing gap (MeV)




Proton pairing gap (MeV)



Neutron density (fm⁻³)



Proton density (fm⁻³)



Time= 0.000000 fm/c

Modeling of prompt fission neutron and gamma properties

CGMF

❑ CGM


- Monte-Carlo implementation of the de-excitation of compound nuclei using the Hauser-Feshbach model
- Full treatment of neutron-gamma competition
- Phenomenological approach, many parameters from experiment or systematics

❑ F

- Monte-Carlo sampling of fission fragment yields
- (Monte-Carlo sampling of pre-fission neutrons)
- Parameterization of
 1. fission yields (mass, charge, TKE)
 2. TXE sharing between FFs
 3. FF angular momenta

$$Y(A, Z, TKE, J, \pi; E_n) \approx Y(A; E_n)Y(Z|A)Y(TKE|A; E_n)P(J)P(\pi)$$

Fission simulation



□ Assumptions:

- ◆ prompt fission products emitted from the fully accelerated fragments
- ◆ no emission occurs during the evolution from saddle to scission
- ◆ no emission at the neck rupture

□ Model:

- Consider fission fragments compound nuclei
- Model neutron and gamma emissions in Hauser-Feshbach formalism
- Monte-Carlo: access to more quantities (e.g. average multiplicities), correlations

PHYSICAL REVIEW C 87, 014617 (2013)



Monte Carlo Hauser-Feshbach predictions of prompt fission γ rays:
 Application to $n_{th} + {}^{235}\text{U}$, $n_{th} + {}^{239}\text{Pu}$, and ${}^{252}\text{Cf}$ (sf)



B. Becker,¹ P. Talou,^{2,*} T. Kawano,² Y. Danon,¹ and I. Stetcu²

¹Gaertner LINAC Laboratory, Rensselaer Polytechnic Institute, New York 12180, USA

²T-2 Nuclear Theory Group, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

Hauser-Feshbach formalism

initial conditions play an important role



Neutron emission probability:

$$P(\epsilon_n)dE \propto T_n(\epsilon_n)\rho(Z, A-1, E - \epsilon_n - S_n)$$


- ✓ Transmission coefficients computed using an optical model
- ✓ Density of states

Gamma emission probability:




$$P(\epsilon_\gamma)dE \propto T_\gamma(\epsilon_\gamma)\rho(Z, A, E - \epsilon_\gamma)$$

- ✓ Transmission coefficients calculated from the gamma strength function (Brink-Axel hypothesis)
- ✓ Density of states
- ✓ Discrete levels
- ✓ Branching ratios

Neutron-induced fission schematics



Multi-chance fission






- Double-hump fission barrier penetrability calculations
- Fission probabilities/barrier not observables
- Other fission observables sensitive to the fission probabilities?

TKE distribution

For each FF mass, assume a Gaussian distribution* given by $TKE(A)$ and $\sigma_{TKE}(A)$

$$Y(A|TKE) = \frac{1}{\sqrt{2\pi}\sigma_{TKE}(A)} \exp\left(-\frac{(TKE - \overline{TKE}(A))^2}{2\sigma_{TKE}^2(A)}\right)$$



- ❖ use $TKE(A)$ at thermal, rescaled to obtain $TKE(E_n)$ from Madland
- ❖ assume the slope fit by Madland holds for all fissioning systems (for a particular initial isotope)
- ❖ $TKE(n_{th})$ fitted within reasonable values to reproduce $\bar{\nu}$; for other systems use systematics
- ❖ use $\sigma_{TKE}(A)$ at thermal for all fissioning systems

* Needs corrections because of the Q value


Mass and charge yields

Vogt et. al., PRC **85**, 024608 (2012)

exponential energy dependence

$$Y(A; E_n) = S_L(A; E_n) + \sum_{i=1,2} S_i(A; E_n)$$


$$S_i(A; E_n) = \frac{N_i(E_n)}{\sqrt{2\pi}\sigma_i(E_n)} \left[\exp\left(-\frac{(A - \bar{A} - D_i(E_n))^2}{2\sigma_i^2(E_n)}\right) + \exp\left(-\frac{(A - \bar{A} + D_i(E_n))^2}{2\sigma_i^2(E_n)}\right) \right] \quad S_L(A; E_n) = \frac{N_L(E_n)}{\sqrt{2\pi}\sigma_L} \exp\left(-\frac{(A - \bar{A})^2}{2\sigma_L^2}\right)$$



Quadratic energy dependence

$$2N_1 + 2N_2 + N_L = 2$$

In the future we plan to use data from model simulations (Moller/Sierk/Bulgac)



Charge distribution:

$$Z_p = A_h \frac{Z_c}{A_c} + \Delta Z \quad (\text{Wahl Systematics})$$

TXE sharing and angular momenta


$$TXE = Q_f(A_l, Z_l; A_h, Z_h; A_c, Z_c) - TKE$$

$$= M_l + M_h - M_c + E_{inc} + B_n(A_c, Z_c) - TKE$$



$$R_T = \frac{T_l}{T_h}$$

$$P(J) \propto (2J + 1) \exp(-J(J + 1)/(2B^2))$$

$$B^2 = \frac{\mathcal{I}T}{\hbar^2} \quad \mathcal{I} = \alpha \mathcal{I}_{rig}^0(Z, A, \beta)$$



$R_T(A)$ kept constant at all energies



- α fitted to fine tune nubar below 2nd chance fission
- energy dependence extended to 20 MeV

Forward propagation of detector response

$$\begin{pmatrix} \tilde{O}_1 \\ \tilde{O}_2 \\ \vdots \\ \tilde{O}_m \end{pmatrix} = \hat{R} \begin{pmatrix} O_1 \\ O_2 \\ \vdots \\ O_m \end{pmatrix}$$

Bypass the complicated problem of inverting the detector response function







FIG. 16. PFG energy spectra E'_γ for the different multiplicities M'_γ , starting from 2 (top) to 16 (bottom). The measured spectra at DANCE (black markers/lines) are compared to CGMF results (blue solid lines) with $\alpha_\gamma=1.3$.




Important considerations for γ rays



Experimental time coincidence window





Energy detection threshold

Selected gamma observables




Position of the peaks is determined by available nuclear structure information

Entrance channel dependence




High-energy γ rays (> 8 MeV)

- Impact of GDR in $T_\gamma(\epsilon_\gamma)$ in FF
- Constraints on CGMF input parameters




Dietrich, Browne et al, PRC **10**, 795 (1974)


- Similar trend as in experiment between 3-8 MeV
- Sensitivity to γ strength function
- Sensitivity to level densities







Makii, Nishio et al, ND2016





Multiplicity-dependent spectra



Selected observables vs. incident energy




Neutron multiplicity probability







Sensitivity to select parameters


Sensitivity to multi-chance fission probabilities



Sensitivity to $\sigma_{\text{TKE}}(\text{A})$





Sensitivity to the deformation



$n_{th} + \text{U-235}$

PFNS (A)

$\Omega_{n-LF}(A)$





Exp. data by Gök, Hamsch, PRC **90**, 064611 (2014)

Correlated fission data



$\langle v \rangle(\text{TKE})$

- How much energy does it cost the fragment to emit a neutron?



Neutron Angular Distributions

- Neutrons are focused along the fission



Summary

- Theoretical approaches to fission fragment yields:
 - Brownian motion
 - Langevin model
 - TD-SLDA

- Emission of prompt fission neutrons and gammas/ correlations (CGMF):
 - De-excitation of fission fragments described within the Monte-Carlo Hauser-Feshbach formalism
 - Spontaneous (^{252}Cf , ^{240}Pu , ^{242}Pu) and neutron-induced fission up to 20 MeV (^{235}U , ^{238}U , ^{239}Pu , ^{241}Pu)
 - Simple model for TKE and spin as a function of incident energy
 - Reasonable description of experimental data, but not everything
 - Incorporate input from models (Moller, Sierk, Bulgac)
 - Incorporated into transport simulations (part of MCNP)