

LA-UR-18-24646

Approved for public release; distribution is unlimited.

Title: Direct Search for Dark Photons with the SeaQuest Spectrometer

Author(s):
Uemura, Sho
Liu, Kun
Liu, Ming Xiong

Intended for: Conference on the Intersections of Particle and Nuclear Physics,
2018-05-29/2018-06-03 (Palm Springs, California, United States)

Issued: 2018-05-30

Disclaimer:


Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Direct Search for Dark Photons with the SeaQuest Spectrometer

Sho Uemura, Kun Liu, Ming Liu
Los Alamos National Laboratory
SeaQuest Collaboration

The SeaQuest facility



- Fixed-target muon spectrometer, Fermilab 120 GeV proton beam
- Primary program: Drell-Yan measurements of sea quark distributions
 - ▶ E906 (unpolarized targets, 2012–2017), E1039 (polarized targets, 2019–2020)
- Parasitic searches for dark photons approved 2015 (E1067)



arXiv:1706.09990


Production and signatures at SeaQuest

- Three dominant production mechanisms: meson decay, proton bremsstrahlung, Drell-Yan (yields from arXiv:1804.00661)
- Prompt $A' \rightarrow \mu^+ \mu^-$: bump-hunt
- Displaced $A' \rightarrow \mu^+ \mu^-$: background suppressed by vertexing
- Displaced $A' \rightarrow e^+ e^-$: background absorbed in dump




SeaQuest searches for dark photons

- Dimuons in main SeaQuest dataset
 - ▶ Bump-hunt at high mass (ongoing effort)
- Dimuon displaced-vertex trigger
 - ▶ Commissioned 2017
- Dielectron trigger
 - ▶ EMCal for electron PID (proposals in preparation)





E906 preliminary



Displaced vertex trigger

- Two new fine-grained scintillator hodoscopes measure track Y
- Existing station 4 paddles are used for muon ID
- FPGA trigger extrapolates tracks to the beam plane and fires on pairs of tracks with large Z





Trigger hodoscopes

- Extruded scintillator bars detect charged particles, wavelength-shifting fibers collect light and transport it to SiPMs
- Each station is split into quadrants; active area has a beam gap of $|y| > 7.5$ cm but minimal gap in x
- Station 1: $z = 8$ m, 1 cm segmentation, 80 bars/quadrant
- Station 2: $z = 15$ m, 2 cm segmentation, 50 bars/quadrant





Readout and services

- Power supplies provide independent control of every SiPM bias voltage
- Postage-stamp preamps read out the SiPMs and send signals to discriminators
- Discriminators fan out to TDCs (readout) and CAEN V1495 FPGA boards (trigger)



Trigger logic

- Two levels: identify displaced tracks, trigger on pairs
- L1: three-way coincidence within each quadrant
 - ▶ Identify displaced tracks ($z_0 \in [400, 650] \text{ cm}$) in each quadrant using hit patterns (“roads”)
- L2: two-out-of-four coincidence between opposite-sign quadrants






Installation and commissioning

- Trigger hodoscopes installed on the SeaQuest beamline spring 2017
- Displaced vertex trigger rate is $\sim 5\%$ of the SeaQuest Drell-Yan trigger, acceptable for parasitic running
- 5 days of good data taken with the displaced vertex trigger before accelerator shutdown: 8×10^{15} live protons on target
- Expect $O(10^{18})$ POT over the next two years, parasitic with new SeaQuest polarized target run




Detector performance

- Efficiencies $\sim 95\%$ excepting $\sim 10\%$ bad channels
- Inefficiency: bar gaps
- Bad channels: inconsistent optical coupling (will fix)




Trigger performance in 2017 data

- Displaced-vertex trigger is 150 ns late relative to the event (cabling mistake during commissioning)
- Damage report: lost all hits in station 1 drift chamber (120 ns drift time) and ~50% of hits in other drift chambers
- We can make straight tracks in stations 2–4 and try to assess backgrounds in the non-bend view
 - ▶ Problem: very bad (and angle-dependent) tracking efficiency
- We can look at hodoscope hit information (not in time relative to events that fired the displaced-vertex trigger) to understand trigger backgrounds
 - ▶ May allow us to optimize the trigger criteria



Schedule and prospects

- Updated reach estimates soon
 - ▶ Realistic trigger geometry and acceptance
- SeaQuest will run with polarized target (E1039) for two years: displaced-vertex dark photon search will run parasitically
 - ▶ Displaced-vertex trigger will be ready to go when beam is available
- Possible PID upgrade (using recycled PHENIX EMCal) will add sensitivity to dielectron decay channel and other new physics (SIMPs, iDM)



arXiv:1804.00661

EMCal upgrade

- One PHENIX EMCal sector: $2 \times 4 \text{ m}^2$ wall of Pb-scintillator shashlyks
- Refurbish the readout: replace PMTs with SiPMs, new readout electronics
- Simple energy threshold can trigger on non-MIP particles
- Track matching enables electron ID

