

Molecular Dynamics Simulations of Ion-Containing Polymers

Amalie L. Frischknecht

LAMMPS Workshop and Symposium
August 2, 2017



*Exceptional
service
in the
national
interest*



Sandia National Laboratories is a multi-mission laboratory managed and operated by National Technology and Engineering Solutions of Sandia, LLC., a wholly owned subsidiary of Honeywell International, Inc., for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA-0003525.

Acknowledgments

Sandia National Labs

Lisa Hall
Dan Bolintineanu
Christina Ting
Mark Stevens

Joe Cordaro
Dale Huber

Lauren Abbott
Eric Sorte
Cy Fujimoto
Todd Alam

Funding: Sandia LDRD
Center for Integrated Nanotechnologies

University of Pennsylvania

Francisco Buitrago
Robert Middleton
Phil Griffin
Karen Winey



NIST



Jacob Tarver
Madhu Tyagi
Christopher Soles



Ion-Containing Polymers

polymers with covalently-bonded ionic groups

ionomers




two main classes of ionomers



- dry (melt): no solvent
- hydrated: with water

Nanoscale Phase Segregation

ionomer melts




hydrated PEMs



STEM showing
aggregates

PEPAA_{9.5}-Zn56



Seitz et al., *J Am Chem Soc* 132, 8165 (2010)


Li, N. & Guiver, M. D. *Macromolecules* 47, 2175 (2014)

Ionic Aggregates in Ionic Polymers

“ionomer peak”


- ubiquitous
- low wavevector peak in scattering
- from inter-aggregate scattering

PSS with Zn^{+2}



Yarusso & Cooper, *Macromolecules*, 1983

Nafion in water





Schmidt-Rohr, K. & Chen, Q. *Nat Mater* **7**, 75–83 (2007)


Proton Exchange Membranes

ion-selective membranes
typically in water

- water purification
- fuel cells



Nafion™





Applications of Ionomers: Batteries?

Issues with current electrolytes in Li-ion batteries:

- organic solvents
- PEO + lithium salts + solvent
 - need containment
 - flammable!

- solvent free PEO + salt
 - conductivity dominated by anions
 - salt concentration at electrodes
 - extra heating





ionomers as next generation electrolytes?

- safer: no solvent
- serve as electrolyte & separator
- less packaging
- improved electrochemical stability
- higher efficiency: single ion conductors

Questions

- What do ionic aggregates look like, exactly?
- What's the dependence on polymer architecture?
- How do ion association and aggregate morphology affect dynamics?




Difficult to measure local structure/dynamics experimentally.


Can simulations resolve issue?

Rest of the Talk

- melts of precise poly(ethylene-co-acrylic acid) (pAA)
 - morphology and comparison to X-ray
 - CG model dynamics
 - dynamics and comparison to QENS






- hydrated, sulfonated Diels-Alder polyphenylenes (SDAPP)
 - morphology and cluster analysis



Model Materials: Precise Ionomers

PE backbone with **precisely** spaced carboxylic acid functional groups




Wagener group, University of Florida

M. E. Seitz et al., *J. Am. Chem. Soc.* **2010**, 132, 8165-8174.


Atomistic MD Simulations

- Variations in:
 - cation type; today: H or Li⁺
 - neutralization level = % COO-M⁺ vs COOH
 - spacing length: p9, p15, p21
- All atom L-OPLS force-field
- 80-200 polymers, 81, 90, or 84 backbone carbons/polymer
 - ~ 64 Å box, total of ~25,000 atoms
- NVT ensemble, **150°C → well above T_g**
- LAMMPS



Morphology: Li-neutralized pAA


p9AA-100%Li



Bolintineanu et al, *ACS Macro Lett*, **2013**


Morphology: Li-neutralized pAA

p9AA-10%Li




TYPE II (stringy)

p9AA-43%Li




TYPE II (stringy)

p9AA-100%Li



TYPE III
(fully percolated)

p21AA-43%Li




TYPE I
(compact, isolated)

coloring by cluster

Bolintineanu et al, *ACS Macro Lett*, 2013

Closer look at aggregates








Two mechanisms of aggregate formation:

1. Counterion-oxygen association → dominant at **moderate to high** neutralization
2. Hydrogen-bonded networks → dominant at **low** neutralization

Direct Comparison to X-ray Scattering

$$S(q) = \sum_i c_i f_i^2 + 4\pi\rho \int_0^\infty \frac{\sin(qr)}{qr} r^2 \sum_{i,j} c_i c_j f_i f_j (g_{ij}(r) - 1) dr$$



- 120°C, same neutralization
- $S(q)$ scaled to match amorphous halo height


¹ Buitrago et al, MM 48,1210 (2015)

² Seitz et al, JACS 132, 8165 (2010)

- excellent agreement in peak positions
- good agreement in peak shapes


X-ray Scattering Doesn't Determine Morphology

need simulations or imaging!



Buitrago, C. F. *et al. Macromolecules* **48**, 1210–1220 (2015).

Quasi-Elastic Neutron Scattering




incoherent, inelastic: sensitive to self-motion of hydrogens





$G_s(r, t)$: given an atom was at $r=0$ at time $t=0$, the probability that the atom is at r at time t

from MD:
$$S(Q, t) = \int G_s(r, t) \frac{r \sin(Qr)}{Q} dr$$

from QENS:
$$S(Q, t) = \frac{\int S_{exp}(Q, \omega) e^{i\omega t} d\omega}{\int R(Q, \omega) e^{i\omega t} d\omega}$$



Acid Copolymers



excellent agreement between QENS and MD


relevant length scales:

amorphous halo: $Q \approx 1.35 \text{ \AA}^{-1}$

ionomer peak: $Q \approx 0.3 - 0.6 \text{ \AA}^{-1}$


Fit $S(Q,t)$ to Two KWW Functions

$$S(Q, t) = \phi_{slow} e^{-\left(\frac{t}{\tau_{slow}}\right)^{\beta_{slow}}} + (1 - \phi_{slow}) e^{-\left(\frac{t}{\tau_{fast}}\right)^{\beta_{fast}}}$$




- One stretched exponential relaxation cannot fit data, two needed
- Extract time scales τ and stretching parameters β

Dynamics from KWW



Fast: local dynamics that are insensitive to composition (vibrations, librations, etc.)

Dynamics from KWW



Slower: structural dynamics of the chain

Highly composition sensitive

Length scale dependent


Fast: local dynamics that are insensitive to composition (vibrations, librations, etc.)

increasing acid content slows dynamics


Acid aggregates rearrange

p21AA: one chain


$t = 0$




$t = 0.1$ ns



$t = 1$ ns




next: Li ionomer dynamics




Rest of the Talk


- melts of precise poly(ethylene-co-acrylic acid) (pAA)
 - atomistic simulations
 - morphology and comparison to X-ray
 - dynamics and comparison to QENS
 - coarse-grained simulations




- hydrated, sulfonated Diels-Alder polyphenylenes (SDAPP)
 - atomistic simulations



SDAPP Membranes




- high T_g
- high modulus
- high thermomechanical stability
- high conductivity



SDAPP IEC = 3.4 meq/g
Nafion IEC = 0.9 meq/g

Fujimoto, C., Hickner, M., Cornelius, C. & Loy, D. *Macromolecules* **38**, 5010–5016 (2005); Tang, Z. *et al.*, *J Electrochem Soc* **161**, A1860–A1868 (2014)

SDAPP Simulations



short SDAPP chain

yellow = sulfur
red = oxygen
cyan = carbon
white = hydrogen


70 chains
3 monomers/chain

sulfonic acids/monomer = $S = 1, 2, 4$
waters/sulfonic acid = $\lambda = 3, 5, 10$


LOPLS-AA, improvements for aromatics
TIP4P/2005 water model

Cluster Morphologies


$S = 1$ sulfonic acid/monomer



$\lambda = 3$



$\lambda = 5$




$\lambda = 10$




increasing water

Cluster Morphologies


$$\lambda = 3$$



S = 1



S = 2



S = 4



increasing sulfonation level



How do we better characterize percolated morphologies?

New Clustering Algorithm


A. Rodriguez and A Laio, *Science* 344, 1492 (2014)

base on density as well as distance

- identify cluster centers
- atoms are in the same cluster as their nearest neighbor with a higher density





SDAPP Clusters




density-based algorithm resolves differences in percolated systems

Cluster Size and Shape

$$R_g^2 = \lambda_1 + \lambda_2 + \lambda_3 \quad \kappa^2 = 1 - 3(\lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3)/R_g^4$$






$$S = 1, \lambda = 3, \kappa^2 = 0.4$$



$$S = 4, \lambda = 10, \kappa^2 = 0.05$$

Summary

- simulation reveals ionic aggregate morphologies
 - MD agrees with x-ray
- atomistic MD in agreement with QENS for acid copolymers/ionomers
 - chains dynamics slowed by aggregates
- MD for SDAPP
 - density-based algorithm for more information on percolated clusters
 - ion transport will depend on cluster shape



Future work: continued correlation of dynamics with morphology