

Anisotropic heavy-hole g-factors & relevance to photon-to-spin conversion schemes in semiconductor quantum dot circuits

A. Bogan,^{1,2} S. A. Studenikin,¹ M. Korkusinski,¹ G. C. Aers,¹ L. Gaudreau,¹
P. Zawadzki,¹ A. Kam,¹ A. S. Sachrajda,¹ D. G. Austing,¹
L. A. Tracy,³ J. L. Reno,³ and T. W. Hargett³

¹Security and Disruptive Technologies Portfolio, National Research Council of Canada,
Ottawa K1A 0R6, Canada

²Department of Physics and Astronomy, University of Waterloo, Ontario NL2 3G1, Canada

³Sandia National Laboratories, Albuquerque, New Mexico 87185, USA






Why holes in GaAs quantum dots?

- Holes:
 - Strongly anisotropic g^* -factor (heavy-hole g^* -factor tunable to zero in appropriately oriented external magnetic field).
 - Expect reduced coupling to nuclei compared to electrons.
 - Strong spin-orbit interaction.
- GaAs:
 - Traditional workhorse for spin qubits in quantum dots.
 - Direct bandgap semiconductor that can host inter-band excitations necessary for a photon-to-spin interface.
 - Clean and readily available.

Few-hole double quantum dot device in an undoped GaAs/AlGaAs hetero-structure

L. A. Tracy et al., Appl. Phys. Lett. **104**, 123101 (2014)
L. A. Tracy et al., Sandia Report SAND2015-8132 (2015)






GaAs dots for holes:
Klochan Appl. Phys. Lett. (2010)
Mak Appl. Phys. Lett. (2013)
Komijani Phys. Rev. B (2013)
Wang Nanoletters (2016)
Wang Nanotechnology (2016)
...

Resonant tunneling spectroscopy of two hole system

See Bogan et al., Phys. Rev. Lett. (2017)

Spin-flip tunneling (via spin-orbit)



Upper triangle:
 $(1,0) \rightarrow (1,1) \rightarrow (2,0) \rightarrow (1,0)$
 Lower triangle:
 $(2,1) \rightarrow (1,1) \rightarrow (2,0) \rightarrow (2,1)$




(2,0) state is a singlet

Similar situation with electrons:
 Pfund Phys. Rev. B (2007) [InAs nano-wire DQD]
 ...

Resonant tunneling spectroscopy of two hole system

See Bogan et al., Phys. Rev. Lett. (2017)

Spin-conserving tunneling




(2,0) state is a triplet



Resonant tunneling spectroscopy of two hole system

See Bogan et al., Phys. Rev. Lett. (2017)

Spin-flip tunneling






(2,0) state is a triplet



Resonant tunneling spectroscopy of two hole system

See Bogan et al., Phys. Rev. Lett. (2017)



More complex spin-flip tunneling




(2,0) state is a triplet

Effective heavy-hole g^* -factor in tilted magnetic field: g^* -factor anisotropy

See Bogan et al., Phys. Rev. Lett. (2017)




Two-Hole Transport Magneto-Spectroscopy



$g_{\min} = 0.04 \pm 0.04$ (for in-plane B-field)

Minimal heavy-hole light-hole mixing

Problem of long distance entanglement distribution



Attractive Aspects:

1. Photo-generated spins can be heralded by non-destructive charge detection techniques.
2. Long spin coherence time (aka native quantum memory), e.g., 200 μs [Bluhm Nat. Phys. (2011)].
3. BSM with 100% success probability can in principle be achieved [Engel & Loss Science (2005); Yokoshi Phys. Rev. B (2010)].
4. Spin-to-photon conversion not required.

BSM node with for example atomic ensemble quantum memory...

Challenges:

1. Non-destructive signaling of successful photon storage challenging.
2. Stored photons must be regenerated before BSM which is an extra step.
3. BSM with linear optics succeeds with probability of 50% at most.


CD=Charge detector

on-to-spin
on schemes
D circuits

with Physica E (2001)
c IEEE (2003)
Phys. (2011)
5)
Jpn (2017)

Polarization schemes

FARADAY CONFIG.



Vrijen & Yablonovitch, Physica E (2001)
 Yablonovitch et al, Proc. IEEE (2003)
 Kosaka, J. Appl. Phys. (2011)
 Oiwa, J. Phys. Soc. Jpn (2017)

See also time-bin schemes: Kosaka Phys. Rev. A (2012)

For transfer
 of polarization-encoded state of photon
 to spin state of QD

A (near) zero g-factor is an important property to realize the necessary V- or Λ -shaped three-level system for a coherent photon-to-spin interface

Transfer
 state of photon
 polarization
 $\rightarrow$ state of heavy-hole

Natural anisotropy gives
 heavy-hole g-factor ~ 0
 for in-plane B-field
 [Bogan PRL (2017)].



Transfer
 state of photon
 polarization
 $\rightarrow$ state of electron

Need to engineer
 electron g-factor ~ 0
 [e.g., see Allison PRB
 (2014)]
 & strain...

Throughput scaling model

To demonstrate potential performance advantage of full BSM capability in spin-based QD scheme over optical scheme with linear optics and atomic ensemble QMem

See Gaudreau et al., Semicond. Sci. Technol. (2017)



$$P_h = 10^{-(2L/L_0)/10}$$

Fiber losses

$$R_t = E_M^{2N}$$

Memory losses (in)

$$\cdot E_M^{2N}$$



Memory losses (out)

$$\cdot (1 - (1 - P_h)^K)^N$$

Heralded photon losses

$$\cdot E_{BSM}^{N-1}$$

Incomplete BSM



$$P_h = 10^{-(2L/L_0)/10}$$

Fiber losses

$$\cdot E_M^2$$

Memory losses (in)

$$R_t = (1 - (1 - P_h)^K)^N$$



Heralded photon losses

$$\cdot E_{BSM}^{N-1}$$


Incomplete BSM

Coherent hybrid interplay between phonon & spin in piezoelectric material such as GaAs during manipulation & measurement of QDs


See Korkusinski et al., Phys. Rev. Lett. (2017)



For surface phonon velocity 2700 m/s,
 $\lambda = 2\ell = 8.6 \mu\text{m}$ gives $\epsilon_{\text{LOC}} = 1.3 \mu\text{eV}$




LZS
 spectroscopy
 of S- T_+ qubit



Features related to coherent and incoherent driving by phonon bath

See Korkusinski et al., Phys. Rev. Lett. (2017)



Landau-Zener-Stückelberg
Petta Science (2010)
Studenikin Phys. Rev. Lett. (2012)
Nichol Nat Comm (2015)
...