

Clay Swelling in Dry Supercritical Carbon Dioxide: Effects of Interlayer Cations on the Structure, Dynamics, and Energetics of CO Intercalation Probed by XRD, NMR and GCMD Simulations

Narasimhan Loganathan, Geoffrey M Bowers, A. Ozgur Yazaydin, Herbert Todd Schaef, John Loring, Andrey G. Kalinichev, and R. James Kirkpatrick

J. Phys. Chem. C, **Just Accepted Manuscript** • DOI: 10.1021/acs.jpcc.7b12270 • Publication Date (Web): 08 Feb 2018

Downloaded from <http://pubs.acs.org> on February 10, 2018

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



1
2
3 Clay Swelling in Dry Supercritical Carbon Dioxide: Effects of
4
5
6 Interlayer Cations on the Structure, Dynamics, and Energetics of CO₂
7
8
9 Intercalation Probed by XRD, NMR and GCMD Simulations
10
11
12

13
14 *Narasimhan Loganathan*^{1, *}, *Geoffrey M. Bowers*², *A. Ozgur Yazaydin*^{1,3}, *H. Todd Schaefer*⁴,
15
16 *John S. Loring*⁴, *Andrey G. Kalinichev*⁵ and *R. James Kirkpatrick*⁶
17
18

19
20 ¹ Department of Chemistry, Michigan State University, East Lansing, Michigan 48824,
21
22 United States
23
24

25
26 ² Department of Chemistry and Biochemistry, St. Mary's College of Maryland, St. Mary's
27
28 City, Maryland 20686, United States
29

30
31 ³ Department of Chemical Engineering, University College London, London, WC1E7JE,
32
33 United Kingdom
34
35

36
37 ⁴ Pacific Northwest National Laboratory, Richland, WA, 99352, United States
38

39
40 ⁵ Laboratoire SUBATECH (UMR 6457 - Institut Mines-Télécom Atlantique, Université de
41
42 Nantes, CNRS/IN2P3), 44307, Nantes, France
43
44

45
46 ⁶ College of Natural Science, Michigan State University, East Lansing, Michigan 48824,
47
48 United States
49

50
51 * **Corresponding author e-mail:** naresh20@msu.edu
52

53
54 **Telephone:** (+1)517-353-1106
55
56
57
58
59
60

Abstract

In situ XRD and NMR experiments combined with molecular dynamics simulations using the grand canonical ensemble (GCMD) show that cation size, charge and solvation energy play critical roles in determining the interlayer expansion of smectite clay minerals when exposed to dry *sc*CO₂ under conditions relevant to petroleum reservoirs and geological CO₂ sequestration conditions. The GCMD results show that the smectite mineral, hectorite, containing interlayer alkali and alkaline earth cations with small ionic radii and high solvation energies (e.g., Na⁺ and Ca²⁺) does not intercalate CO₂ and that the fully collapsed interlayer structure is the energetically most stable configuration. With Cs⁺ and Ba²⁺ the monolayer structure is the stable configuration, and CO₂ should spontaneously enter the interlayer. With Cs⁺ there is not even an energy barrier for CO₂ intercalation, in agreement with the XRD and NMR results. ¹³C NMR and simulations show that the average orientation of the intercalated CO₂ is with their O-C-O axes parallel to the basal clay surface and that undergo a rapid rotation about an axis perpendicular to the main molecular axis. The simulations show that the strength of the interaction between the exchangeable cation and the clay structure dominates the intercalation energetics in dry *sc*CO₂.

Introduction

Injection of CO₂ into deep geological formations and depleted oil and gas reservoirs is an effective method for enhancing oil recovery and a potential pathway for reducing anthropogenic CO₂ in the atmosphere. Target geological storage sites are primarily composed of porous rock formations in which the CO₂ is confined by overlying caprocks that act as low permeability barriers to upward fluid flow.¹⁻⁸ Long-term confinement of CO₂ depends strongly on its interactions with these caprocks, which are typically shales containing significant amounts of clay minerals. Intercalation and nano-confinement of fluids in the 2-dimensional interlayer galleries of expandable clays (principally smectites and smectite layers in mixed layer illite-smectites) may affect the properties and porosity of the caprocks by expanding or shrinking the interlayer pores (tracked via the basal spacings)⁹⁻²⁴ and can lead to substantial changes in the structural and dynamical behavior of the intercalated fluid molecules with respect to the bulk fluid phase. Numerous studies²⁵⁻³⁹ provide critical molecular-scale information about the role of the clay composition, location of its structural charge, and the properties of the charge compensating cation in determining the structure, dynamics, energetics and reactivity of H₂O in the interlayer galleries and inter-particle pore space.

Recent experimental and molecular simulation studies at conditions relevant to subsurface injection of supercritical CO₂^{9-23,40-48} have shown that the intercalation and retention of CO₂ in smectite interlayers is possible but depends strongly on the amount and thermodynamic properties of water in the system. For example, Giesting et al.^{11,12} and Schaefer et al.^{17,19} have shown that the H₂O saturation state (relative humidity, R.H.) of supercritical CO₂ (scCO₂) controls CO₂ intercalation in Na-, Mg- and Ca-montmorillonite at 323 K and 90 bar. The results show that no CO₂ enters the dry and collapsed interlayers of these smectites and that CO₂ intercalation only occurs in the presence of at least a sub-monolayer of H₂O. This humidity-dependent intercalation was further validated by Loring et al.,²⁰⁻²² who also showed

1
2
3 that the maximum CO₂ intercalation at 90 bar and 323 K occurs when the interlayer H₂O
4 content is equivalent to that of a monolayer hydrate structure. *In situ* experimental NMR, IR,
5 and XRD of the similar smectite mineral, hectorite, at 90 bar CO₂ and 323 K by Bowers et al.⁹
6 showed that the properties of the charge balancing cation can greatly change the CO₂
7 intercalation behavior. These results show that for the cations with relatively high hydration
8 energies and small ionic radii (Na⁺, Ca²⁺), CO₂ intercalation requires the presence of H₂O
9 molecules to prop open the interlayers. In contrast, Cs⁺ with a larger ionic radius and smaller
10 hydration energy allows Cs-hectorite to intercalate CO₂ readily even in the complete absence
11 of water (R.H. of the scCO₂ = 0). With all three cations, increasing H₂O saturation of the scCO₂-
12 rich fluid phase causes a decreasing interlayer molar CO₂/(CO₂+H₂O) ratio, with Cs-hectorite
13 maintaining a higher ratio at all conditions. Computational molecular simulations of nano-
14 confined CO₂-H₂O mixtures in smectites in equilibrium with water-saturated CO₂ at 348 K and
15 125 bar using the grand canonical Monte Carlo (GCMC) ensemble by Botan et al.⁴⁰ have
16 suggested that the stable state of Na-montmorillonite exposed to supercritical CO₂ is the bilayer
17 hydrate (basal spacing ~15.0 Å) but that maximum CO₂ intercalation occurs at the monolayer
18 hydrate state. Recently GCMC simulations by Kadoura et al.⁴¹ show that for montmorillonite
19 at high R.H.s (≥ 60%) at 323 K and 90 bar there is no significant difference in the interlayer
20 CO₂ content with cations having different hydration energies (Na⁺, Ca²⁺, Mg²⁺). Simulation
21 studies using GCMC and Gibbs ensemble Monte Carlo (GEMC) have indicated that for Na-
22 smectite under similar conditions CO₂ intercalation can vary depending on the clay layer charge
23 and its structural location.⁴²⁻⁴⁵ Recently, we reported grand canonical molecular dynamics
24 (GCMD) simulations that show that varying temperature (323–368 K) and pressure (90-
25 150 bar) have little impact on the structural environments and dynamics of intercalated CO₂
26 molecules in Na-hectorite in equilibrium with H₂O-saturated scCO₂.⁴⁸
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Intercalation of dry CO₂ in smectite interlayers has been less extensively studied but is
4 important as a limiting energetic and structural case and may have significant application to
5 situations when dry CO₂ is injected into the subsurface. Mineralogical studies using X-ray
6 computed tomography before and after CO₂ injection have suggested that scCO₂ can laterally
7 displace the brine in that occupies the pore volumes of the host sedimentary rocks causing
8 either anhydrous or nearly anhydrous conditions, especially during the early stages of
9 injection.^{1,6,49,50} Under these circumstances, the dry CO₂ could potentially enter and dehydrate
10 the smectite interlayers, cause swelling or shrinking of the clay, and thus generate mechanical
11 stresses and alter the overall pore structure of shale caprocks. It could also suppress hydration
12 reactions involving the mineral phases due to a lack of residual H₂O in the pore fluid. Our
13 previous experimental studies have shown that collapsed Na-montmorillonite does not
14 intercalate CO₂ or expand when exposed to dry CO₂ under reservoir conditions (323 K and
15 90 bar), whereas Cs- and NH₄-montmorillonite do.⁵¹ Parallel GCMD calculations are
16 consistent with these results, showing a large energy barrier for CO₂ intercalation in Na-
17 montmorillonite but no such barrier Cs- and NH₄-montmorillonites.⁵¹ These results suggest
18 that the exchangeable cation has a significant effect on the intercalation behavior under dry
19 CO₂ conditions, although the energetic origin of these effects and the associated structural and
20 dynamical behavior have not been explored in detail.

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45 In the present work, we combine *in situ* experimental nuclear magnetic resonance
46 spectroscopy (NMR) and X-ray diffraction (XRD) studies at a CO₂ pressure of 90 bar and a
47 temperature of 323K with GCMD calculations to investigate the structure, dynamics and
48 energetics of the intercalation of dry CO₂ in the smectite mineral, hectorite, containing
49 monovalent and divalent cations spanning a range of hydration and solvation properties. We
50 used hectorite as our host material for the experiments and modeling studies, because it has a
51 very low Fe content that improves NMR spectral resolution by minimizing paramagnetic
52
53
54
55
56
57
58
59
60

1
2
3 effects affecting most natural montmorillonites. Like montmorillonite, hectorite develops most
4 of its permanent structural charge by cation substitution in the octahedral layer, and it has been
5 used for many NMR studies to examine smectite behavior.^{9,24,26,28,29,31,52}
6
7
8
9

10 **Experimental Methods**

11 *In Situ XRD*

12
13
14 Na- and Cs-exchanged hectorite samples were characterized by XRD at 323 K first
15 under vacuum (10^{-3} Torr) and then during exposure to dry CO₂ at a pressure of 90 bar using
16 high pressure and temperature X-ray diffraction techniques previously described.^{9,17,19,51} The
17 Cs-hectorite was also reexamined under vacuum conditions at 323K after the CO₂ was
18 removed. The base hectorite was SHCa-1, which is available at the Clay Source Repository of
19 the Clay Minerals Society.
20
21
22
23
24
25
26
27

28
29 Hectorite is a 2:1 trioctahedral smectite clay mineral that develops negative structural
30 charge due to Li⁺ for Mg²⁺ substitution in the octahedral layer. The structural formula of this
31 sample, that was processed to remove carbonate and quartz impurities, is
32 $[M_{0.36}(Mg_{2.65}Li_{0.35})(Si_{3.99}Al_{0.01})O_{10}(F_{1.1}OH_{0.9})]$.^{28,29} For these experiments, measured aliquots
33 of dilute clay suspensions in water were pipetted onto a beryllium post, with each sample
34 occupying a circular area with an approximate diameter of 2-3 mm. Each clay sample was
35 allowed to air dry to produce a thin preferentially oriented film. The beryllium post was then
36 placed on an XYZ stage and the samples were aligned with the aid of a laser system, allowing
37 exposure of only one clay film at a time to the XRD beam. Once the sample coordinates were
38 established, the pressurized reactor was assembled, mounted into the instrument, and heated to
39 323 K. Subsequent exposure to vacuum for 30 minutes removed most of the water from the
40 clay samples. Pressurization with CO₂ was accomplished through an ISCO pump connected
41 directly to the reactor.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 XRD patterns were collected using a Bruker D8 Discover instrument with a rotating Cu
4 anode (CuK alpha = 1.5418 Å), a programmable XYZ customized stage, and a Vantec 500
5 detector set at a sample-to-detector distance of 15 cm. This instrument, operated at 50 kV and
6 24 mA, is capable of producing an intensely focused 0.5 mm beam. XRD patterns for each
7 sample were acquired for 300 seconds, integrated over a 2θ range of 4-45°, and subsequently
8 analyzed with the MDI JADE® XRD software package to obtain peak positions and peak
9 profiles.

19 *In Situ MAS NMR*

21 *In situ* ^{13}C MAS NMR spectra were obtained at 323 K and 90 bars P_{CO_2} for the same
22 Na- and Cs-exchanged hectorite samples examined by XRD using the high pressure apparatus
23 that has been described previously.^{9,24} For these experiments, clay filled rotors were dried
24 overnight in a vacuum oven at 323K and 10^{-3} Torr. The rotors were sealed after backfilling the
25 vacuum oven with dry N_2 gas. Hectorite samples prepared in a similar way and analyzed by IR
26 spectroscopy show no detectable H_2O content with Cs^+ and ≤ 0.5 H_2O per cation with Na^+ .
27 99% isotopically enriched $^{13}\text{CO}_2$ was then introduced into the rotor using a one-way flow
28 system.^{53,54} The spectra were collected using a 7.0 T Varian DDR 2 console with a 5 mm HXY
29 probe at a spin frequency of 3 kHz using a $\pi/2$ pulse width of 5 μs , a spectral width of 50 kHz,
30 and a 30 s inter-pulse delay to obtain 3000 transients. The ^{13}C spectra were referenced to
31 tetramethylsilane (TMS) through a secondary standard of adamantane using the high-frequency
32 peak. All NMR data were processed using iNMR, a commercial program by MestReC, and all
33 NMR results were iteratively fit using Abscissa, a freeware app written by Rudiger Bruhl. For
34 ^{13}C , the first two data points were removed due to acoustic ringing before zero filling to 32k
35 points and applying 20 Hz of exponential apodization. In almost all cases a standard baseline
36 correction was performed to flatten the baseline. All peak fits were performed by iteratively
37 fitting the data set to Lorentzian lineshapes.

Simulation Methods

The model hectorite used in our simulations has a structural formula of $M^+(\text{Mg}_5\text{Li})\text{Si}_8\text{O}_{20}(\text{OH})_4$. This model has no tetrahedral substitutions, in accordance with the experimental sample, which contains a negligible fraction (0.25%) of tetrahedral Al^{3+} for Si^{4+} substitution.^{9,26,28,29,31} The simulation model, however, differs from the experimental sample in two important respects. First, the model has a 30% higher layer charge. Second, the OH^- site in the model contains only OH^- groups, whereas the experimental sample has 55% F^- for OH^- substitution. The hydration behavior of synthetic fluoro-hectorite has been discussed previously.^{27,55,56} Previous simulation studies employing this model have shown good agreement with experimental data justifying its use here.^{30,33,35,47,48}

The simulation supercell consists of 16 crystallographic unit cells of hectorite ($4 \times 2 \times 2$) encompassing two interlayer galleries with surface areas of 360.23 \AA^2 . This supercell size is sufficient to eliminate any finite size effects. Isomorphic substitutions of $\text{Li}^+/\text{Mg}^{2+}$ in the supercell were performed in a quasi-disordered pattern following Lowenstein's rule to prevent $\text{Li}^+-\text{O}-\text{Li}^+$ next nearest neighbor arrangements.⁵⁷ The structural details of the model are described elsewhere.^{30,33,35,48} Overall charge balance for the model is obtained by cation substitution in the interlayer galleries. Nine different compositions were simulated, each with the interlayer occupied by a single type of alkali or alkaline earth cation (Li^+ , Na^+ , K^+ , Rb^+ , Cs^+ , Mg^{2+} , Ca^{2+} , Sr^{2+} , or Ba^{2+}).

Molecular dynamics simulations were performed in the grand canonical ensemble using the RASPA software package.^{58,59} Basal spacings were varied from 9.0 \AA to 18.0 \AA at steps of 0.2 \AA and were constrained at the initial value throughout each simulation run. All of the charge compensating cations were initially placed randomly in the interlayer region and were allowed to move during the GCMC runs. Importantly, during the simulations we permit lateral movements of the T-O-T layers parallel to each other (in the x and y directions) without

1
2
3 disrupting the structure. These translational movements are essential, because the minimum
4 energy interlayer structure depends on the relative positions of the adjacent T-O-T layers and
5 varies with the type of the interlayer cation and the number of intercalated fluid molecules, as
6 reported in previous studies.^{40,48,51,60}
7
8
9

10
11
12 All the GCMD simulations were performed as described in our previous study of H₂O
13 and CO₂ exchange in Na-hectorite.⁴⁸ The GCMD approach is a hybrid technique which
14 incorporates both Monte Carlo and molecular dynamics procedures by sampling the insertion
15 and deletion of fluid molecules with equal probability at every MD time step. Therefore,
16 temperature, volume and the chemical potential (μ) of the fluid molecules are fixed in our
17 simulations, but the number of fluid molecules absorbed in the interlayer can fluctuate. The
18 interatomic interactions for the T-O-T clay structure and the metal cations were described using
19 the ClayFF force field,⁶¹ which is widely used in clay interfacial simulations. The CO₂
20 molecules were represented using the rigid EPM2 model.⁶² Three-dimensional periodic
21 boundary conditions were employed with a cutoff of 9.0 Å for short range non-electrostatic
22 interactions, and the long-range electrostatic interactions were computed using Ewald
23 summation⁶³ with an accuracy of 10⁻⁶. Fugacities are required to determine the acceptance of
24 insertion and deletion moves for the fluid species (here CO₂) and were calculated using the
25 Peng-Robinson equation of state.⁶⁴
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 All simulations were performed at $T=323$ K and $P=90$ bar (the experimental conditions
45 described above) for 15 ns to reach equilibrium followed by another 10 ns of data production
46 with a time step of 1 fs. The reported adsorption profile and related energetics, interlayer
47 structure and dynamics were calculated using the data from the last 5 ns of each equilibrium
48 simulation run. The methods of analysis can be found in our previous papers.^{33,35,48,65} The
49 computed energetics of each system is evaluated using the immersion energies, which were
50 computed by considering the largest basal spacings (18.0 Å) as the reference state because its
51
52
53
54
55
56
57
58
59
60

1
2
3 value is the closest to that of bulk CO₂ than any other basal spacing. The use of immersion
4
5 energies to investigate the equilibrium states of smectites have been discussed previously
6
7 mostly for water adsorption^{33,35,37,48,51,65} and is extended here to CO₂.
8
9

10 **Results and Discussion**

11 *XRD*

12
13
14 The *in situ* XRD results (Figure 1) show that the basal spacing (d_{001} basal reflection)
15
16 of Cs-hectorite increases reversibly by about 1 Å on exposure to dry CO₂, but that the basal
17
18 spacing of Na-hectorite does not change. We interpret these results to indicate that under our
19
20 experimental conditions, CO₂ readily enters the interlayer galleries of Cs-hectorite but not Na-
21
22 hectorite. The specific basal spacings of the Cs-hectorite before and after exposure to dry CO₂
23
24 (11.06 Å and 12.05 Å, respectively) are the same as previously reported.⁹ They are also
25
26 consistent with values reported in the literature for similar samples and conditions.^{33,37,51,57,65,66}
27
28
29 After a few minutes of exposure to vacuum conditions following equilibration with dry scCO₂,
30
31 the basal spacing of Cs-hectorite returns to its original value, suggesting that there is not a
32
33 significant quantity of recalcitrant CO₂ in the interlayers.⁹ For Na-hectorite, previous *in situ* IR
34
35 data for this sample⁹ show nearly complete removal of interlayer water and a lack of expansion
36
37 when exposed to dry scCO₂. Fully dehydrated Na-SHCa-1 sample has a basal spacing of
38
39 9.7 Å,²⁶ and the value of 9.81 Å observed here indicates removal of all or most of the interlayer
40
41 water. In addition, XRD did not detect formation of a carbonate phase in either the Cs- or Na-
42
43 hectorite.
44
45
46
47
48

49 *¹³C MAS NMR*

50
51 As for the XRD results, the ¹³C MAS NMR results indicate sorption of CO₂ in the
52
53 interlayers of Cs-hectorite but not Na-hectorite. As discussed by Bowers, et al.,⁹ the spectrum
54
55 of CO₂ in an empty rotor and with Na-SHCa-1 are very similar (Figure 2). Both contain a center
56
57 band at ~124 ppm with a full width at half height (FWHH) of ~0.11 ppm that represents CO₂
58
59 molecules that are undergoing rapid, isotropic tumbling in the supercritical fluid phase and that
60

1
2
3 experience little interaction with the clay surface.⁹ Both these samples also show a pair of small
4 spinning sidebands (SSBs). For the *scCO*₂ without clay present, these SSBs must be associated
5 with CO₂ either adsorbed on the rotor wall or being compressed and densified against the wall
6 due to the centrifugal forces caused by the rotor rotation. For the Na-SHCA-1 sample, the SSBs
7 are likely to arise from similar effects, possibly including weak adsorption on or compressed
8 against the exterior surfaces of the clay particles.⁹ Note that the spectrum for the Na-SHCA-1
9 sample in Figure 2 is greatly expanded vertically to show the SSBs. In contrast, for the Cs-
10 hectorite, the center band is much broader, with a FWHH of 0.67 ppm, and there is a broader,
11 asymmetric spinning sideband manifold. The sideband manifold for this sample is very similar
12 to that reported previously for Cs-hectorite dried over P₂O₅ or equilibrated at 43% R.H. before
13 exposure to dry *scCO*₂.⁹ It indicates a residual chemical shift anisotropy (CSA) associated with
14 adsorption of CO₂ molecules in an environment where they undergo anisotropic rotation. We
15 previously assigned this pattern to adsorbed CO₂ molecules in a monolayer hydrate-type
16 (~12.5 Å) interlayer arrangement with their long (O=C=O) molecular axes having a time-
17 averaged orientation parallel to the basal clay surface and experiencing rapid (>10⁵ Hz)
18 reorientation about an axis normal to the clay basal surface.⁹

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40 The ¹³C MAS NMR spectra for Cs-hectorite here are also similar to those published
41 previously for Cs-exchanged Wyoming montmorillonite (Cs-SWy-2) with dry CO₂ at 90 bar
42 and 323 K.⁵¹ Most importantly, both show similar SSB patterns indicating CO₂ intercalation
43 into the interlayer galleries where confinement induces restricted molecular reorientation. For
44 the Cs-hectorite, however, the sidebands have on average a FWHH of ~2.92 ppm, compared
45 to ~1.37 ppm for the Cs-montmorillonite. The FWHH of the center band for the Cs-hectorite
46 is also about twice that of the center band of the Cs-montmorillonite. This difference is
47 surprising, because SWy-2 contains substantially more structural Fe than SHCA-1,^{9,29,31,51} and
48 we expected that its ¹³C NMR spectrum would show significant paramagnetic broadening as a
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 result. Thus, it appears that in Cs-montmorillonite the motion of the CO₂ molecules is
4 effectively decoupling the ¹³C and electron spin systems. The origin of the differences in peak
5 widths for the Cs-hectorite and montmorillonite are uncertain at this time, but may be due to
6 dynamical disorder, small differences in the interlayer *z* dimension, or differences in the NMR
7 powder averaging behavior related to the mean particle size, crystallite size, and packing
8 density/geometry in the rotors. The spectrum for Cs-hectorite was obtained with a 5 mm rotor,
9 whereas that for the Cs-montmorillonite was obtained with a 7.5 mm rotor, and the Cs-hectorite
10 samples are ground and sieved to provide a controlled particle size in the rotors. In general,
11 the *sc*CO₂ fluid pockets in the center of the rotor cavity are larger with 7.5 mm than with 5 mm
12 rotors, which may influence the exchange of CO₂ between interlayer and bulk fluid
13 environments.
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 **Computational Results**

29 *Adsorption Profile and Energetics*

30
31
32
33 The computational modeling results show that the properties of the exchangeable
34 cations have significant effects on the structural and energetic behavior of smectite-CO₂
35 systems and are in good agreement with the experimental results discussed above. For all the
36 simulated systems, CO₂ intercalation begins at a basal spacing of ~11.0 Å except with Mg²⁺,
37 for which it begins at ~10.7 Å. The amount of intercalated CO₂ then increases stepwise with
38 increasing basal spacing (Figures 3a and 3b). Except with Cs⁺, the basal spacing at which
39 intercalation begins is greater than the basal spacings of the fully collapsed structures.
40
41
42
43
44
45
46
47
48
49

50 In parallel, the computed immersion energies show an energy barrier to CO₂
51 intercalation that decreases with increasing cation size for both the alkali and alkaline earth
52 series (Figures 3c and 3d). This energy barrier is the difference between the maximum energy
53 of the system (typically near 10.5 – 11.0 Å) and the value for the fully collapsed structure with
54 the same cation. With Cs⁺, there is no energy barrier, and the immersion energy decreases
55
56
57
58
59
60

1
2
3 continuously from that of the collapsed structure to the first energy minimum near 12.5 Å. With
4
5 K^+ , Rb^+ , Sr^{2+} , and Ba^{2+} , the barrier is quite small, but it is much larger with Li^+ , Na^+ , Mg^{2+} and
6
7 Ca^{2+} . With Li^+ , Na^+ , Mg^{2+} and Ca^{2+} , the global minimum energy structure is the collapsed state,
8
9 whereas it is the monolayer structure near 12.5 Å with K^+ , Rb^+ , Cs^+ , Sr^{2+} and Ba^{2+} .

10
11
12
13 We interpret these results to show that CO_2 should spontaneously enter the interlayer
14
15 galleries of fully dehydrated and collapsed K^+ , Rb^+ , Cs^+ , Sr^{2+} , and Ba^{2+} hectorite, but not Li^+ ,
16
17 Na^+ , Mg^{2+} and Ca^{2+} hectorite. This conclusion is in full agreement with the experimental XRD
18
19 results above that show CO_2 intercalation in Cs-hectorite but not Na-hectorite. Note that the
20
21 computed immersion energies here are not free energies and do not include entropic effects.
22
23 This could make a difference in interpreting relatively small immersion energy differences
24
25 between the monolayer and bilayer structures of the K^+ , Rb^+ , Cs^+ , Sr^{2+} , and Ba^{2+} systems, but
26
27 are unlikely to be significant for interpreting the results for the Li^+ , Na^+ , Mg^{2+} and Ca^{2+} systems.
28
29 The computed intercalation energies are consistent with the results of comparable GCMD
30
31 calculations for Na-, NH_4 -, and Cs-montmorillonite, which show a large barrier to CO_2
32
33 intercalation with Na^+ but essentially no barrier with NH_4^+ and Cs^+ .⁵¹ This conclusion is also
34
35 consistent with the free energy calculations by Makaremi et al.⁴³, which show a large energy
36
37 barrier for CO_2 intercalation into Na-montmorillonite.
38
39
40
41
42

43
44 The relationship between the basal spacings and computed energies that highlight the
45
46 need to prop open the interlayers of smectites that contain relatively small charge-balancing
47
48 cations to allow CO_2 intercalation, as observed in previous experimental and simulation
49
50 studies.^{9-14,16-23,40-45,48} These studies have indicated that the small cations with high hydration
51
52 energies, the interlayers needed to be propped open by some H_2O molecules to incorporate
53
54 CO_2 , and the maximum CO_2 intercalation (maximum CO_2/CO_2+H_2O ratio) occurs near the
55
56 monolayer hydrate interlayer structure.^{9-14,40-45,48} To our knowledge, there have been no studies
57
58 of CO_2 intercalation in pillared clays, which are propped open by interlayer precipitates.
59
60

1
2
3 For all the simulated systems here, the increase in interlayer CO₂ content with
4 increasing basal spacing shows 2 broad plateaus between 12.4-13.5 Å and 15.5-16.3 Å (Figures
5 3a and 3b), corresponding to monolayer and bilayer CO₂ structures, respectively. In parallel,
6 the immersion energies have broad minima at the same basal spacing ranges (Figures 3c and
7 3d). Basal spacings associated with monolayer CO₂ adsorption were previously reported for
8 Na-fluorohectorite exposed to gaseous CO₂ at much lower temperature and pressure conditions
9 (253 K and 15 bar) using XRD.⁶⁷ For our hectorite simulations with Li⁺, Na⁺, Mg²⁺ and Ca²⁺,
10 these monolayer and bilayer structures are metastable with respect to the collapsed structures,
11 and for Na-hectorite interlayer adsorption of CO₂ is not experimentally accessible. We expect
12 fully dry Li-, Mg-, and Ca-hectorites to behave in the same way. As for the Cs-hectorite XRD
13 experiments and simulations, the computational results suggest that dry hectorite exchanged
14 with K⁺, Rb⁺, Sr²⁺, and Ba²⁺ should intercalate CO₂ and expand. For the Cs-hectorite
15 simulations reported here, the monolayer structure is the experimentally observed stable state,
16 in agreement with the global energy minimum in the computed immersion energy at the
17 monolayer structure compared to the bilayer structure. This is also the case with Rb⁺, K⁺, Sr²⁺
18 and Ba²⁺. However, with divalent cations, the immersion energy differences between the
19 monolayer and bilayer structures are small.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 For our systems with alkali cations, the computed interlayer CO₂ content decreases as
44 the size of the cation increases from Li⁺ to Cs⁺. For the monolayer structures this decrease is
45 from 2.7 to 1.8 CO₂ per unit cell, and for the bilayer structures, it is from 5.0 to 3.8 molecules
46 per unit cell. In contrast, with the alkaline earth cations cation size has no effect on the amount
47 of intercalated CO₂, with all systems showing ~2.6 and ~5.3 CO₂ per unit cell at the monolayer
48 and bilayer spacings, respectively. The decreasing interlayer CO₂ content with increasing ionic
49 radii with the alkali metal cations correlates well with experimental and simulation studies of
50 montmorillonite exposed to dry CO₂ under identical thermodynamic conditions.⁵¹ In contrast,
51
52
53
54
55
56
57
58
59
60

1
2
3 recent experimental study by Bowers et al.⁹ have reported that the interlayer CO₂ content of
4 Cs-hectorite exposed to scCO₂ with increasing water content is greater than that of Na-
5 hectorite. This difference is probably due to the larger hydration energy of Na⁺ causing more
6 interlayer H₂O adsorption leaving less open interlayer space for CO₂ intercalation in the studies
7 by Bowers et al.⁹ However, under dry scCO₂ exposure, the smaller the size of the cation, the
8 larger the interlayer space available for CO₂ intercalation thus leading to higher CO₂ content
9 for Na- than Cs-hectorite.

10
11
12 In the current study, at any given basal spacing, the number of intercalated CO₂
13 molecules per cation is always greater with the alkaline earth cations than with the alkali
14 cations. As discussed in our previous experimental and simulation studies,^{9,19} this difference
15 can be attributed to the smaller number of divalent cations required to compensate the hectorite
16 layer charge, thereby increasing the interlayer space accessible for CO₂ compared to smectites
17 with monovalent cations. With alkali cations the plateau regions also become increasingly flat
18 with increase in ionic size, whereas the plateau shapes are similar for all the alkaline earth
19 cations. This is likely because the change in ion size is substantially greater for the alkali cation
20 series (Li⁺ vs Cs⁺) than the alkaline earth cation series (Mg²⁺ vs Ba²⁺). The absence of
21 substantial differences with the different alkaline earth cations despite the ~160 kJ/mol
22 difference in the CO₂ solvation energies between Mg²⁺ and Ba²⁺ supports the geometrical
23 origin of this effect.

24
25
26 The energetic contributions from the cation-clay layer, cation-CO₂, and CO₂-clay layer
27 interactions provide deeper insight into the origin of the dependence of the CO₂ intercalation
28 on the properties of the cation and demonstrate that the magnitude of the interaction between
29 the cation and the clay T-O-T layers dominates this behavior (Figure 4). For both the alkali and
30 alkaline earth cation series, the magnitude of the cation-clay layer interaction for the fully
31 collapsed structures decreases (becomes less negative) with increasing cation size (Figures 4a
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 and 4b). The energies at the collapsed basal spacings are the relevant comparators, since for
4
5 each system expansion starts from this value. With the alkali cations, this energy varies
6
7 from -3675.3 kJ/mol with Li^+ to -3400.1 kJ/mol with Cs^+ , and with the alkaline earth cations it
8
9 varies -3764.9 kJ/mol with Mg^{2+} to -3487.9 kJ/mol with Ba^{2+} . Importantly, the difference in
10
11 the clay-cation interaction energies between the collapsed structures and the basal spacings at
12
13 which intercalation begins ($\sim 11.0\text{\AA}$) is large for cations with high hydration energies and
14
15 decreases with increasing cation size. Hence, there is a cost of $>\sim 150$ kJ/mol in the clay-cation
16
17 interaction energies with Li^+ , Na^+ , Mg^{2+} and Ca^{2+} for any CO_2 to intercalate, in contrast to
18
19 $<\sim 60$ kJ/mol with the other cations. At monolayer spacings the clay-cation interaction energies
20
21 are very similar for both the alkali and alkaline earth cations, and the differences at larger basal
22
23 spacings are very small. Despite a need for only half the number of cations to compensate the
24
25 layer charge with the alkaline earth cations vs. the alkali cations, the magnitude of the
26
27 interaction energies for both types of cations with the clay surface are very similar. This
28
29 similarity indicates that the divalent cations have a greater interaction energy per cation with
30
31 the clay structure than the monovalent cations, as expected from the charge differences. For
32
33 the cation- CO_2 interactions at a given basal spacing, the interaction energies are negative and
34
35 decrease (becomes less negative) with increasing cation size in each cation series. They are
36
37 larger in general for the alkaline earth series (Figures 4c and 4d). Both these trends are expected
38
39 based upon recently computed cation- CO_2 solvation energies in scCO_2 fluid.^{51,68} The
40
41 interaction energies of CO_2 molecules with the clay structure are slightly negative but do not
42
43 vary greatly with the different alkaline and alkaline earth cations (Figures 4e and 4f). Together,
44
45 these results show that the increase in the cation- CO_2 interaction energy with decreasing cation
46
47 size cannot overcome the decreasing cation-clay interaction energy in the absence of H_2O .
48
49
50
51
52
53
54

55 56 57 *Computed Interlayer Structure* 58 59 60

1
2
3 It is normally not possible to determine the detailed arrangement of the cations and fluid
4 molecules in smectite interlayer galleries by diffraction methods due to static and dynamic
5 structural disorder, and computed structures provide insight into the interlayer fluid structures.
6
7 For the computed systems here, the atomic density profiles (ADPs) normal to the clay basal
8 surface show that the average positions of the different species vary somewhat with the cation.
9
10 For the monolayer structures here, the ADPs of C_{CO_2} , O_{CO_2} , and the cation are all centered at
11 the middle of the interlayer at ~ 2.8 Å to ~ 3.0 Å from both basal surfaces and become narrower
12 with increasing cation size in each series (Figures 5a-5i). The ADPs of Li^+ , Na^+ and Mg^{2+} are
13 significantly broader than those of the other cations, reflecting their relatively small size and
14 ability to move perpendicular to the basal surfaces. The ADPs of the intercalated CO_2
15 molecules do not vary significantly with cation size, and both O_{CO_2} and C_{CO_2} share the same
16 plane as the cations, centered at ~ 2.9 Å from the two basal surfaces. With the alkali cations,
17 the O_{CO_2} and C_{CO_2} intensities decrease with increasing cation size due to the decreasing amount
18 of intercalated CO_2 (Figures 5a-5e). They do not vary with the different alkaline earth cations,
19 reflecting the independence of the amount of intercalated CO_2 on the cation for this series. This
20 result is consistent with much of the CO_2 being located in free volume in the interlayer rather
21 than being coordinated to the cations, as the energetic results above suggest (Figures 5f-5i).
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 The average orientation of the O-C-O axis of the intercalated CO_2 molecules lies
43 parallel to the basal surfaces with all the cations (Figures 6a and 6b), as expected from the
44 ADPs of O_{CO_2} and C_{CO_2} and in agreement with the ^{13}C NMR results above. However, the range
45 of angles probed by the O-C-O vector is somewhat larger with the alkali cations (45° - 135°)
46 than with the alkaline earth cations (55° - 125°), indicating slightly more dynamically restricted
47 local structural environment with the alkaline earth cations. This difference is probably due to
48 the larger interlayer CO_2 contents with the alkaline earth cations leading to less interlayer space.
49
50
51
52
53
54
55
56
57
58
59
60
60

1
2
3 time oriented parallel to the clay layers and part of the time wobbling. The fraction of the time
4 spent wobbling decreases with increasing cation size with the alkali cations but does not change
5 significantly with the alkaline earth cations.
6
7
8

9
10 As expected, the computed mean interatomic distance between the cations and CO₂
11 molecules increases with increasing cation size in the order Cs⁺ > Ba²⁺ > Na⁺ > Mg²⁺ (Figures
12 7a -7d). Similarly, cations with small ionic radii are coordinated to the basal hectorite surface
13 at shorter interatomic M-O_b distances (<2.5 Å for Na⁺ and Mg²⁺) than Cs⁺ (~3.0 Å). The mean
14 interatomic distances between CO₂ molecules and surface oxygen atoms (O_b), however, do not
15 change with changing cation size. Paralleling the cation-oxygen interatomic distances, the
16 running coordination numbers (RCN) between all the interlayer species vary considerably with
17 the size of the cation. For the monolayer structures, the total coordination of CO₂ molecules by
18 cations, surface O_b atoms, and other CO₂ molecules is ~15.2 for Na⁺, Cs⁺, and Mg²⁺ but is
19 ~17.0 for Ba²⁺ ions. The RCN of cations with O_b atoms, however, increases from ~2 for Na⁺
20 and Mg²⁺ to ~6.5 and ~7.5 for Ba²⁺ and Cs⁺, respectively. The adsorption structure and RCN
21 values for Na- and Cs-hectorites are in excellent agreement with previous studies on hectorites
22 under ambient and elevated *T* and *P* conditions.^{14,17,47}
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 The two-dimensional density maps (also called planar atomic density distributions, or
42 PADDs) for the monolayer structures show that the differences in the RCNs for different
43 cations are the result of significantly different local adsorption environments (Figures 8a-8d).
44 The Na⁺ and Mg²⁺ ions are located predominately between two O_b atoms, one on each basal
45 surface giving rise to the O_b RCN of ~2. For Na-hectorite (Figure 8a), the sets of three
46 neighboring site maxima represent the same Na⁺ ion (blue) that is hopping between different
47 sites, illustrating a potential pathway for Na⁺ diffusion along these narrow channels unhindered
48 by intercalated CO₂. Likewise, the pairs of maxima for Mg²⁺ (Figure 8b) represent the same
49 ion, but in this case the site hopping is more hindered by the increased number of intercalated
50
51
52
53
54
55
56
57
58
59
60

1
2
3 CO₂ molecules. As a result, the arrangement of CO₂ molecules is more ordered in Mg-hectorite
4 than in Na-hectorite, where the CO₂ molecules adsorb in narrow channels formed by Na⁺ ions
5 and wobble between adjacent hectorite surfaces. This difference correlates well with the more
6 restricted orientation distribution for CO₂ molecules with Mg²⁺ than with Na⁺. Nonetheless,
7 there are on average four CO₂ molecules around both Na⁺ and Mg²⁺, consistent with the RCN
8 values (Figures 7a and 7b). For both Na- and Mg-hectorite, the intercalated CO₂ molecules
9 form small clusters, with their arrangement predominantly in a slipped parallel geometry. A
10 small fraction of CO₂ molecules in these clusters are in a distorted T-shaped arrangement,
11 particularly in Mg-hectorite because of its larger number of intercalated CO₂ molecules. A T-
12 shaped arrangement is highly unlikely for Na-hectorite because of the narrow channels formed
13 by the Na⁺ ions and CO₂ molecules. For both Na- and Mg-hectorite here, the intercalated CO₂
14 molecules are adsorbed with one O_{CO2} located near the center of a ditrigonal cavity with the
15 other above a neighboring Si tetrahedron.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 Surprisingly, the adsorption structures of CO₂ molecules in Cs- and Ba-hectorite are
35 generally similar to that in Na- and Mg-hectorite, despite the considerably different ionic sizes
36 and interlayer CO₂ contents. Importantly, the RCNs of intercalated CO₂ with surface O_b atoms
37 is ~7-8 with all cations because of the similar adsorption structures described above. The RCN
38 of CO₂ molecules with interlayer cations is ~4 for Na⁺-, Cs⁺- and Mg²⁺- and is ~5 with Ba²⁺, in
39 agreement with the PADDs in Figures 8a-8d. Similarly, the RCN between intercalated CO₂
40 molecules varies significantly between the structures with alkali cations (~3.5-4) and alkaline
41 earth cations (~5-6) and is due to higher interlayer CO₂ content in the latter case. The contour
42 patterns for CO₂ molecules in Cs-hectorite are more disordered than in the other models due to
43 the lower CO₂/cation ratio and subsequently larger free interlayer space available for
44 librational/wobbling motion. The coordination environments of the Cs⁺ and Ba²⁺ ions,
45 however, are quite different (Figures 8c and 8d). The large RCN of Cs⁺ with O_b atoms (~7.5)
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 is due to the fact that on average the Cs^+ ions are located above the center of a ditrigonal cavity
4 of one surface and above a Si tetrahedron on the opposite surface. The Cs^+ ions oscillate
5 between the centers of different ditrigonal cavities and above different Si tetrahedra the,
6 resulting in the disordered contours in Figure 8c. In contrast, the Ba^{2+} ions are adsorbed
7 between Si tetrahedra from both basal surfaces, leading to the average RCN value of ~ 6.0 .
8
9
10
11
12
13
14

15 The computed CO_2 orientations and locations in the PADDs are in excellent agreement
16 with the ^{13}C NMR results for Cs-hectorite, which show a restricted environment for the CO_2
17 molecules in which the time averaged O-C-O vector is parallel to the basal surfaces and the
18 molecules undergo rapid rotation or libration (frequencies $> \sim 10^5$ Hz) about an axis
19 perpendicular to their O-C-O axes. The PADD results show that on the 10 ns time scale of the
20 simulations the motion is best thought of as hindered libration rather than free rotation. The
21 tight contours in the PADDs show that the positions of the O_{CO_2} are typically quite restricted,
22 even with Cs^+ . On the \sim ms time scale of probed by the ^{13}C NMR line shapes, however, this
23 may not be true. As shown in our previous experimental study of variably humidified
24 CO_2 /hectorite interactions,⁹ the absence of CO_2 molecules with O-C-O vectors near 0° and 180°
25 (Figure 6a and 6b) shows that they never experience isotropic reorientation because they are
26 never oriented perpendicular to the basal surfaces.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 *Residence times*

44
45
46 The dynamic interaction among the interlayer species and O_b atoms of the basal
47 surfaces can be quantitatively characterized by the computed mean residence times, which for
48 our systems decrease with increasing cation size for all interatomic pairs (Table 1). Overall,
49 the mean residence times are at least twice as long with smaller cations (Li^+ , Na^+ , Mg^{2+} , Ca^{2+})
50 than with larger cations (K^+ , Rb^+ , Cs^+ and Ba^{2+}). This relationship holds for both the
51 intermittent residence times, $c(t)$, which allow for the possibility that individual atomic pairs
52
53
54
55
56
57
58
59
60

1
2
3 can re-coordinate after becoming separated during the simulation, and the continuous residence
4
5 times, $C(t)$, which do not allow such re-coordination.^{33,35,48} Here, the intermittent residence
6
7 times are typically about an order of magnitude longer than the continuous ones, as also
8
9 reported in our previous studies of hydrated interlayers and hectorite exposed to H₂O saturated
10
11 scCO₂.^{33,35,48} The $c(t)$ and $C(t)$ values for cation-O_b pairs are consistently longer than those for
12
13 cation-O_{CO2} and O_b-O_{CO2} coordination, indicating that interactions between cations and basal
14
15 surfaces are the principal controls of the interlayer structure and dynamics. This conclusion is
16
17 consistent with the energetic results above, which show that the extent of CO₂ intercalation is
18
19 strongly dependent on cation-clay interactions (Figures 4a-4f). As the size of interlayer cations
20
21 increases, the residence times for cation-O_b pairs decreases by an order of magnitude
22
23 paralleling the expected stronger coordination of smaller cations with clay surfaces, which are
24
25 dominated by electrostatic interactions. This result is in good agreement with the atomic
26
27 density maps for especially the alkali cations, which show much more ordered contours for Na⁺
28
29 than for Cs⁺ (Figures 8a and 8c). This difference is less pronounced with the divalent cations,
30
31 because their higher charge leads to overall stronger coordination with the surface. The cation-
32
33 O_b residence times are in reasonable agreement with previous simulation studies of hectorite
34
35 under ambient conditions.^{33,35} The cation-CO₂ residence times are longer than O_b-CO₂
36
37 residence times, particularly for smaller ions, consistent with their CO₂ solvation energies.^{51,68}
38
39 The difference in cation-CO₂ and O_b-CO₂ residence times decreases with increasing cation size,
40
41 especially with the alkali cations.
42
43
44
45
46
47
48

49 **Comparison of Na- and Cs-hectorite**

50
51 The effects of the exchangeable cations on the structure and energetics of CO₂
52
53 molecules intercalated in smectites and the connections to the experimental results are well
54
55 illustrated by comparing the specific details of the Na- and Cs-hectorite systems. The collapsed
56
57 interlayer distances for Na- (9.5 Å) and Cs-hectorite (10.7 Å) computed from our simulations
58
59
60

1
2
3 are in good agreement with the values obtained in XRD measurements under vacuum
4 conditions (9.8 Å and Cs-11.08 Å, respectively) and also in good agreement with published
5
6 computed values, which vary slightly depending on the interatomic potentials used in the
7
8 simulations.^{12,14,16,17,20,69,70,71}
9

10
11
12 Upon exposure to dry scCO₂ at 323 K and 90 bar, the XRD data indicate interlayer
13 expansion for Cs-hectorite but not for Na-hectorite. This is in excellent agreement with our
14
15 simulations that show that the collapsed structure is the most energetically favorable state for
16
17 Na-hectorite (Figure 3c). In contrast, there is no energy barrier for Cs-hectorite, and this system
18
19 favors interlayer expansion when in contact with dry scCO₂. Our simulations also indicate that
20
21 despite the stronger interaction energy between Na⁺ and CO₂ molecules (-95 kJ/mol) than
22
23 between Cs⁺ and CO₂ molecules (-30 kJ/mol), CO₂ intercalates in Cs-hectorite because the
24
25 stronger cation-clay interactions with Na⁺ result in a much larger energy cost for expansion
26
27 from the collapsed to the monolayer structure (~400 kJ/mol) than with Cs⁺ (~190 kJ/mol).
28
29 Thus, the energy difference for the cation-clay interactions between the collapsed state and the
30
31 basal spacing at which CO₂ intercalation starts is ~175 kJ/mol with Na⁺ and only ~35 kJ/mol
32
33 with Cs⁺ (see Figure 4a). Because of the small energy barrier, CO₂ intercalation is favourable
34
35 in Cs-hectorite, and not in Na-hectorite. These results indicate that the cation-clay interactions
36
37 play a critical role during CO₂ intercalation.
38
39
40
41
42
43

44 **Conclusions**

45
46
47 The intercalation of CO₂ in the interlayers of the smectite clay hectorite with a range of
48
49 mono- and divalent cations in contact with dry scCO₂ at 323 K and 90 bar was quantitatively
50
51 analyzed by GCMD simulations and *in situ* XRD and NMR experiments. The GCMD results
52
53 show that clays containing small charge compensating cations, do not spontaneously
54
55 incorporate CO₂ in their interlayers when exposed to dry scCO₂, but that those with larger
56
57 exchangeable cations do. The thermodynamic energy barriers between the collapsed state of
58
59
60

1
2
3 the clay structure and the onset of CO₂ intercalation are large with small cations and decrease
4
5 with increasing cation size. With Li⁺, Na⁺, Mg²⁺, and Ca²⁺, the fully collapsed state has the
6
7 global energy minimum, but with K⁺, Rb⁺, Cs⁺, Sr²⁺, and Ba²⁺ the monolayer state has the
8
9 global energy minimum, suggesting that hectorite with these cations should spontaneously
10
11 intercalate CO₂ and expand but that with the smaller cations it should not. This conclusion is
12
13 in good agreement with the experimental observation that Na-hectorite does not incorporate
14
15 CO₂ under these conditions but that Cs-hectorite does. These results are also in good agreement
16
17 with experimental and GCMD results showing that the similar smectite mineral
18
19 montmorillonite incorporates dry CO₂ and expands with interlayer NH₄⁺ and Cs⁺, but not with
20
21 Na⁺. A detailed quantitative analysis of different contributions to the total interaction energies
22
23 in the investigated systems demonstrates that the structural and energetic characteristics of
24
25 interlayer expansion with dry scCO₂ are largely controlled by the strength of the interactions
26
27 between the cations and the clay T-O-T structure. This is unlike the case of H₂O intercalation,
28
29 where the hydration energy of cations dominates the interlayer expansion.^{33,35}
30
31
32
33
34

35 Structurally, irrespective of the cation size, charge and CO₂ solvation energy, at
36
37 monolayer basal spacings the computed average positions of the intercalated CO₂ molecules
38
39 and the cations are all at the mid-plane of the interlayer. Both the ¹³C NMR spectra and GCMD
40
41 simulations show that the intercalated CO₂ molecules are oriented on average with their O-C-O
42
43 axes parallel to the basal hectorite surfaces. These NMR results for Cs-hectorite show
44
45 restricted dynamics for the interlayer CO₂ in which they undergo rapid (frequency > ~10⁵ Hz)
46
47 rotation or libration around an axis perpendicular to their long molecular axes. The GCMD
48
49 calculations suggest that on the time scale of our simulations (~ 10 ns) this is principally
50
51 librational hopping among reasonably well-defined sites. The GCMD results suggest that
52
53 aggregation of the interlayer CO₂ molecules results in both slipped parallel and T-shaped
54
55 arrangements with slipped parallel dominant. The number of intercalated CO₂ molecules is
56
57
58
59
60

1
2
3 larger for interlayers with higher charge cations due to the larger free interlayer space available
4
5 for CO₂ intercalation due to the smaller number of charge compensating interlayer cations
6
7 present. Since the cation-clay interactions dominate the thermodynamics of interlayer
8
9 expansion with dry CO₂, the expansion behavior of smectites with different layer charges or
10
11 positions of isomorphic substitution (e.g., tetrahedral Al for Si substitution vs. dominantly the
12
13 octahedral substitution in hectorite and montmorillonite) could be quite different than that
14
15 discussed here.
16
17

18 19 **Acknowledgements**

20
21
22 All the calculations in this work were performed using computational resources at the
23
24 National Energy Research Scientific Computing Center, which is supported by the Office of
25
26 Science of the U.S. Department of Energy under ECARP No. m1649. The authors acknowledge
27
28 iCER computational facility at Michigan State University for additional computational
29
30 resources. The work was supported by the United States Department of Energy, Office of
31
32 Science, Office of Basic Energy Sciences, Chemical Sciences, Biosciences, and Geosciences
33
34 Division through the sister grants DE-FG02-10ER16128 (Bowers, P.I.) and DE-FG02-
35
36 08ER15929 (Kirkpatrick, P.I.), as well as through its Geosciences program at Pacific
37
38 Northwest National Laboratory (J.S.L.). H.T.S. was supported by the DOE Office of Fossil
39
40 Energy at PNNL through the National Energy Technology Laboratory, Morgantown, West
41
42 Virginia. A.G.K. acknowledges the support of the industrial chair “Storage and Disposal of
43
44 Radioactive Waste” at the IMT-Atlantique, funded by ANDRA, Areva, and EDF, and of the
45
46 European Union’s Horizon 2020 research and innovation program under grant agreement
47
48 No. 640979.
49
50
51
52
53

54 55 **Notes**

56
57
58 The authors declare no competing financial interest.
59
60

Cation	Cation-O _b	Cation-O _{CO2}	O _b -O _{CO2}
$c(t) : C(t)$			
Li ⁺	10.4 : 0.81	6.3 : 0.65	3.1 : 0.29
Na ⁺	7.7 : 0.54	3.8 : 0.42	2.4 : 0.21
K ⁺	2.1 : 0.31	1.9 : 0.29	1.1 : 0.13
Rb ⁺	0.9 : 0.14	1.1 : 0.09	0.9 : 0.08
Cs ⁺	0.7 : 0.08	0.8 : 0.02	0.8 : 0.05
Mg ²⁺	12.1 : 1.0	8.4 : 0.64	5.5 : 0.31
Ca ²⁺	8.8 : 0.79	6.2 : 0.48	3.8 : 0.24
Sr ²⁺	5.2 : 0.51	4.1 : 0.34	2.1 : 0.15
Ba ²⁺	3.1 : 0.35	2.5 : 0.25	1.3 : 0.12

Table 1. Calculated intermittent $c(t)$ and continuous $C(t)$ residence times (ns) for different atomic pairs for monolayer basal spacing (12.4 Å) in the interlayers of hectorite with different charge compensating cations at 323 K and 90 bar.

FIGURES




Figure 1. X-ray diffraction tracings of the Cs-SHCa-1 (left) and the Na-SHCa-1 (right) clay under vacuum (10^{-3} Torr, 323 K) and during exposure to 90 bar scCO₂ at 323 K. Note that the smaller peak intensities in the samples exposed to scCO₂ are due to signal attenuation by the dense CO₂ fluid.




Figure 2. ^{13}C MAS NMR spectra Cs-SHCa-1 in scCO_2 (top), Na-SHCa-1 in scCO_2 (middle), and a rotor only filled with scCO_2 , all at 90 bar pressure and 323 K. The broad peak near 110 ppm is the ^{13}C background of the NMR probe, and the narrow singularity near 100 ppm reflects a DC offset in the receiver channels.




Figure 3. a) and b) are the average number of intercalated CO_2 molecules (per unit cell); c) and d) are the computed immersion energies for hectorite models with different charge compensating cations as functions of interlayer basal spacing at 323 K and 90 bar. Dotted vertical lines indicate the collapsed interlayer distances for every cations used in our study. The reference state for computing the immersion energies is the 18.0 Å basal spacing.




Figure 4. Computed variation in interaction energies between 3 different pairs (a) and b) are Hectorite–Cation (top), c) and d) are Cation–CO₂ (middle), e) and f) are Hectorite–CO₂ (bottom) as functions of interlayer spacing using hectorite with different of mono- and di-valent cations at 323 K and 90 bar. Dotted vertical lines indicate the collapsed interlayer distances for every cations used in our study.




Figure 5. Computed atomic density profiles of O_b (dark blue vertical lines), interlayer mono- and divalent cations (orange), O_{CO_2} (green) and C_{CO_2} (violet) of hectorite as functions of distance from the basal clay surface at monolayer distances (12.4 Å) under conditions of 323 K and 90 bar.




Figure 6. Computed orientational distributions of intercalated CO₂ molecules in the interlayers of hectorite at monolayer distances under 323 K and 90 bar conditions with different charge compensating cations. θ is the angle between the O-O vector of a CO₂ molecule and the normal to the hectorite basal surface. a) monovalent cations. b) divalent cations.




Figure 7. Radial distribution functions (RDFs, solid lines) and corresponding running coordination numbers (RCNs, dashed lines) for different atomic pairs in the interlayers of a) Na- b) Mg- c) Cs- d) Ba-hectorite at monolayer distances under 323 K and 90 bar conditions.




Figure 8. Computed PADDs of interlayer metal cations and intercalated CO₂ molecules in hectorite interlayers at monolayer distances and 323 K and 90 bar. a) Na⁺ and CO₂ in Na-hectorite; b) Mg²⁺ and CO₂ in Mg-hectorite; c) Cs⁺ and CO₂ in Cs-hectorite; d) Ba²⁺ and CO₂ in Ba-hectorite. Color code: O_b – gray (dark and light), Si – yellow (dark and light), Cation (Na⁺, Cs⁺, Mg²⁺, Ba²⁺) – blue, O_{CO2} – green, C_{CO2} – pink, Dark colors correspond to the O_b and Si of one hectorite basal surface, and light colors correspond to those atoms of the opposite basal surface.

References

1. Gauss, I. Role and Impact of CO₂-Rock Interactions During CO₂ Storage in Sedimentary Rocks. *Int. J. Green. Gas Cont.* **2010**, *4*, 73-89.
2. Lackner, K. S. A Guide to CO₂ Sequestration. *Science* **2003**, *300*, 1677-1678.
3. Benson, S. M.; Cole, D. R. CO₂ Sequestration in Deep Sedimentary Formations. *Elements* **2008**, *4*, 325-331.
4. Bachu, S.; Bonijoly, D.; Bradshaw, J.; Burruss, R.; Holloway, S.; Christensen, N. P.; Mathiassen, O. M. CO₂ Storage Capacity Estimation: Methodology and Gaps. *Int. J. Green. Gas Cont.* **2007**, *1*, 430-443.
5. Wilkin, R. T.; Digiulio, D. Geochemical Impacts to Groundwater from Geologic Carbon Sequestration: Controls on pH and Inorganic Carbon Concentrations from Reaction Path and Kinetic Modeling. *Environ. Sci. Technol.* **2010**, *44*, 4821-4827.
6. Berrezueta, E.; Menendez, G.-L.; Breitner, D.; Luquot, L. Pore System Changes During Experimental CO₂ Injection into Detritic Rocks. Studies of Potential Storage Rocks from Some Sedimentary Basins of Spain. *Int. J. Green. Gas Cont.* **2013**, *4*, 73-89
7. Meakin, P.; Tartakovsky, A. M. Modeling and Simulation of Pore-Scale Multiphase Fluid Flow and Reactive Transport in Fractured and Porous Media. *Rev. Geophys.* **2009**, *47*, RG3002.
8. Davis, A. J.; Kent, B. D. Surface Complexation Modeling in Aqueous Geochemistry. *Rev. Mineral.* **1990**, *23*, 177-260.
9. Bowers, G. M.; Schaef, H. T.; Loring, J. S.; Hoyt, D. W.; Burton, S. D.; Walter, E. D.; Kirkpatrick, R. J. Role of Cations in CO₂ Adsorption, Dynamics and Hydration in Smectite Clays under In Situ Supercritical CO₂ Conditions. *J. Phys. Chem. C* **2017**, *121*, 577-592.
10. Wang, Z.; Felmy, A. R.; Thompson, C. J.; Loring, J. S.; Joly, A. G.; Rosso, K. M.; Schaef, H. T.; Dixon, D. A. Near-Infrared Spectroscopic Investigation of Water in Supercritical CO₂ and the effect of CaCl₂. *Fluid Phase Equil.* **2013**, *338*, 155-163.
11. Giesting, P.; Guggenheim, S.; van Groos, A. F. K.; Busch, A. Interaction of Carbon Dioxide with Na-Exchanged Montmorillonite at Pressures to 640 bar: Implications for CO₂ Sequestration. *Int. J. Green. Gas Cont.* **2012**, *8*, 73-81.
12. Giesting, P.; Guggenheim, S.; van Groos, A. F. K.; Busch, A. X-ray Diffraction Study of K- and Ca-Exchanged Montmorillonites in CO₂ Atmospheres. *Environ. Sci. Technol.* **2012**, *46*, 5623-5630.
13. Lee, M.-S.; McGrail, B. P.; Glezakou, V.-A. Microstructural Response of Variably Hydrated Ca-rich Montmorillonite to Supercritical CO₂. *Environ. Sci. Technol.* **2014**, *48*, 8612-8619.
14. Ilton, E. S.; Schaef, H. T.; Qafoku, O.; Rosso, K. M.; Felmy, A. R. In Situ X-ray Diffraction Study of Na⁺ Saturated Montmorillonite Exposed to Variably Wet Supercritical CO₂. *Environ. Sci. Technol.* **2012**, *46*, 4241-4248.
15. Guggenheim, S.; van Groos, A. F. K. An Integrated Experimental System for Solid-Gas-Liquid Environmental Cells. *Clays Clay Miner.* **2014**, *62*, 470-476.

16. Rother, G.; Ilton, E. S.; Wallacher, D.; Hauss, T.; Schaef, H. T.; Qafoku, O.; Rosso, K. M.; Felmy, A. R.; Krukowski, E. G.; Stack, A. G. et al. CO₂ Sorption to Subsingle Hydration Layer Montmorillonite Clay Studies by Excess Sorption and Neutron Diffraction Measurements. *Environ. Sci. Technol.* **2013**, *47*, 205-211.
17. Schaef, H. T.; Ilton, E. S.; Qafoku, O.; Martin, P. F.; Felmy, A. R.; Rosso, K. M. In Situ XRD study of Ca²⁺ Saturated Montmorillonite (STx-1) Exposed to Anhydrous and Wet Supercritical Carbon Dioxide. *Int. J. Green. Gas Cont.* **2012**, *6*, 220-229.
18. Romanov, V. N. Evidence of Irreversible CO₂ Intercalation in Montmorillonite. *Int. J. Green. Gas Cont.* **2013**, *14*, 220-226.
19. Schaef, H. T.; Loring, J. S.; Glezakou, V. A.; Miller, Q. R. S.; Chen, J.; Owen, A. T.; Lee, M.-S.; Ilton, E. S.; Felmy, A. R.; McGrail, B. P. et al. Competitive Sorption of CO₂ and H₂O in 2:1 Layer Phyllosilicates. *Geochim. Cosmochim. Acta* **2015**, *161*, 248-257.
20. Loring, J. S.; Schaef, H. T.; Turcu, R. V. F.; Thompson, C. J.; Miller, Q. R.; Martin, P. F.; Hu, J.; Hoyt, D. W.; Qafoku, O.; Ilton, E. S. et al. In Situ Molecular Spectroscopic Evidence for CO₂ Intercalation into Montmorillonite in Supercritical Carbon Dioxide. *Langmuir* **2012**, *28*, 7125-7128.
21. Loring, J. S.; Ilton, E. S.; Chen, J.; Thompson, C. J.; Martin, P. F.; Benezeth, P.; Rosso, K. M.; Felmy, A. R.; Schaef, H. T. In Situ Study of CO₂ and H₂O Partitioning Between Na-Montmorillonite and Variably Wet Supercritical Carbon Dioxide. *Langmuir* **2014**, *30*, 6120-6128.
22. Loring, J. S.; Schaef, H. T.; Thompson, C. J.; Turcu, R. V. F.; Miller, Q. R.; Chen, J.; Hu, J.; Hoyt, D. W.; Martin, P. F.; Ilton, E. S. et al. Clay Hydration/Dehydration in Dry to Water-Saturated Supercritical CO₂: Implications for Caprock Integrity. *Energy Procedia* **2013**, *37*, 5443-5448.
23. Krukowski, E. G.; Goodman, A.; Rother, G.; Ilton, E. S.; Guthrie, G.; Bodnar, R. J. FT-IR Study of CO₂ Interaction with Na⁺ Exchanged Montmorillonite. *Appl. Clay Sci.* **2015**, *114*, 61-68.
24. Bowers, G. M.; Hoyt, D. W.; Burton, S. D.; Ferguson, B. O.; Varga, R.; Kirkpatrick, R. J. In Situ ¹³C and ²³Na Magic Angle Spinning NMR Investigation of Supercritical CO₂ Incorporation in Smectite-Natural Organic Matter Composites. *J. Phys. Chem. C* **2014**, *118*, 3564-3573.
25. Bowers, G. M.; Bish, D. L.; Kirkpatrick, R. J. Cation Exchange at the Mineral-Water Interface: H₃O⁺/K⁺ Competition at the Surface of Nano-Muscovite. *Langmuir* **2008**, *24*, 10240-10244.
26. Bowers, G. M.; Singer, J. W.; Bish, D. L.; Kirkpatrick, R. J. Alkali Metal and H₂O Dynamics at the Clay/Water Interface. *J. Phys. Chem. C* **2011**, *115*, 23395-23407.
27. Marry, V.; Dubois, E.; Malikova, N.; Durand-Vidal, S.; Longeville, S.; Brey, J. Water Dynamics in Hectorite Clays: Influence of Temperature Studied by Coupling Neutron Spin Echo and Molecular Dynamics. *Environ. Sci. Technol.* **2011**, *45*, 2850-2855.
28. Weiss, C. A.; Kirkpatrick, R. J.; Altaner, S. P. Variations in Interlayer Cation Sites of Clay Minerals as Studied by ¹³³Cs MAS Nuclear Magnetic Resonance Spectroscopy. *Amer. Mineral.* **1990**, *75*, 970-982.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
29. Weiss, C. A.; Kirkpatrick, R. J.; Altaner, S. P. The Structural Environments of Cations Adsorbed onto Clays – ^{133}Cs Variable-Temperature MAS NMR Spectroscopic Study of Hectorite. *Geochim. Cosmochim. Acta* **1990**, *54*, 1655-1669.
30. Morrow, C. P.; Yazaydin, A. O.; Krishnan, M.; Bowers, G. M.; Kalinichev, A. G.; Kirkpatrick, R. J. Structure, Energetics and Dynamics of Smectite Clay Interlayer Hydration: Molecular Dynamics and Metadynamics Investigation of Na-Hectorite. *J. Phys. Chem. C* **2013**, *117*, 5172-5187.
31. Reddy, U. V.; Bowers, G. M.; Loganathan, N.; Bowden, M.; Yazaydin, A. O.; Kirkpatrick, R. J. Water Structure and Dynamics in Smectites: ^2H NMR Spectroscopy of Mg, Ca, Sr, Cs and Pb-Hectorite. *J. Phys. Chem. C* **2016**, *120*, 8863-8876.
32. Marry, V.; Malikova, N.; Cadene, A.; Dubois, E.; Durand-Vidal, S.; Turq, P.; Breu, J.; Longeville, S.; Zanotti, J. M. Water Diffusion in a Synthetic Hectorite by Neutron Scattering – Beyond the Isotropic Translational Model. *J. Phys.: Cond. Matter* **2008**, *20*, 104205.
33. Loganathan, N.; Yazaydin, A. O.; Bowers, G. M.; Kalinichev, A. G.; Kirkpatrick, R. J. Structure, Energetics and Dynamics of Cs^+ and H_2O in Hectorite: Molecular Dynamics Simulations with Unconstrained Substrate Surface. *J. Phys. Chem. C* **2016**, *120*, 10298-10310.
34. Porion, P.; Faugere, A. M.; Delville, A. Multiscale Water Dynamics within Dense Clay Sediments Probed by ^2H Multiquantum NMR Relaxometry and Two-Time Stimulated Echo NMR Spectroscopy. *J. Phys. Chem. C* **2013**, *117*, 26119-26134.
35. Loganathan, N.; Yazaydin, A. O.; Bowers, G. M.; Kalinichev, A. G.; Kirkpatrick, R. J. Cation and Water Structure, Dynamics and Energetics in Smectite Clays: A Molecular Dynamics Study of Ca-Hectorite. *J. Phys. Chem. C* **2016**, *120*, 12429-12439.
36. Greathouse, J. A.; Hart, D. B.; Bowers, G. M.; Kirkpatrick, R. J.; Cygan, R. T. Molecular Simulation of Structure and Diffusion at Smectite-Water Interfaces: Using Expanded Clay Interlayers as Model Nanopores. *J. Phys. Chem. C* **2015**, *119*, 17126-17136.
37. Ngouana-Wakou, B. F.; Kalinichev, A. G. Structural Arrangements of Isomorphic Substitutions in Smectites: Molecular Simulation of the Swelling Properties, Interlayer Structure and Dynamics of Hydrated Cs-Montmorillonite Revisited with New Clay Models. *J. Phys. Chem. C* **2014**, *118*, 12758-12773.
38. Sato, T.; Watanabe, T.; Otuka, R. Effects of Layer Charge, Charge Location, and Energy Change on Expansion Properties of Dioctahedral Smectites. *Clays Clay Miner.* **1992**, *29*, 873-882.
39. Wang, J.; Kalinichev, A. G.; Kirkpatrick, R. J. Effects of Substrate Structure and Composition on the Structure, Dynamics and Energetics of Water at Mineral Surfaces: A Molecular Dynamics Modeling Study. *Geochim. Cosmochim. Acta* **2006**, *70*, 562-582.
40. Botan, A.; Rotenberg, R.; Marry, V.; Turq, P.; Noetinger, B. Carbon Dioxide in Montmorillonite Clay Hydrates: Thermodynamics, Structure and Transport from Molecular Simulation. *J. Phys. Chem. C* **2010**, *114*, 14962-14969.
41. Kadoura, A.; Nair, A. K. N.; Sun, S. Molecular Simulation Study of Montmorillonite in Contact with Variably Wet Supercritical Carbon Dioxide. *J. Phys. Chem. C* **2017**, *121*, 6199-6208.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
42. Myshakin, E. M.; Saidi, W. A.; Romanov, V. N.; Cygan, R. T.; Jordan, K. D. Molecular Dynamics Simulations of Carbon Dioxide Intercalation in Hydrated Na-Montmorillonite. *J. Phys. Chem. C* **2013**, *117*, 11028-11039.
 43. Makaremi, M.; Jordan, K. D.; Guthrie, G. D.; Myshakin, E. M. Multiphase Monte Carlo and Molecular Dynamics Simulations of Water and CO₂ Intercalation in Montmorillonite and Beidellite. *J. Phys. Chem. C* **2015**, *119*, 15112-15124.
 44. Rao, A.; Leng, Y. Molecular Understanding of CO₂ and H₂O in a Montmorillonite Clay Interlayer Under CO₂ Geological Sequestration Conditions. *J. Phys. Chem. C* **2016**, *120*, 2642-2654.
 45. Rao, A.; Leng, Y. Effect of Layer Charge on CO₂ and H₂O Intercalations in Swelling Clays. *Langmuir* **2016**, *32*, 11366-11374.
 46. Sena, M. M.; Morrow, C. P.; Kirkpatrick, R. J.; Krishnan, M. Structure, Energetics, and Dynamics of Supercritical Carbon Dioxide at Smectite Mineral-Water Interfaces: Molecular Dynamics Modeling and Adaptive Force Investigation of CO₂/H₂O Mixtures Confined in Na-Montmorillonite. *Chem. Mater.* **2015**, *27*, 6946-6959.
 47. Yazaydin, A. O.; Bowers, G. M.; Kirkpatrick, R. J. Molecular Dynamics Modeling of Carbon Dioxide, Water and Natural Organic Matter in Na-Hectorite. *Phys. Chem. Chem. Phys.* **2015**, *17*, 23356-23367.
 48. Loganathan, N.; Yazaydin, A. O.; Bowers, G. M.; Kalinichev, A. G.; Kirkpatrick, R. J. Molecular Dynamics Study of CO₂ and H₂O Intercalation in Smectite Clays: Effect of Temperature and Pressure on Interlayer Structure and Dynamics in Hectorite. *J. Phys. Chem. C* **2017**, *121*, 24527-24540.
 49. Luquot, L.; Gouze, P. Experimental Determination of Porosity and Permeability Changes Induced Injection of CO₂ into Carbonate Rocks. *Chem. Geol.* **2009**, *265*, 148-159.
 50. Vickerd, M. A.; Thring, R. W.; Arocena, J. M.; Li, J. B. Changes in Porosity due to Acid Gas Injection as Determined by X-ray Computed Tomography. *J. Can. Petrol. Technol.* **2006**, *45*(8), 17-22.
 51. Schaefer, H. T.; Loganathan, N.; Bowers, G. M.; Kirkpatrick, R. J.; Yazaydin, A. O.; Burton, S. D.; Hoyt, D. W.; Ilton, E. S.; Thanthiriwatte, K. S.; Dixon, D. A. et al. Tipping Point for Expansion of Layered Aluminosilicates in Weakly Polar Solvents: Supercritical CO₂. *Appl. Mater. Interf.* **2017**, *9*, 36783-36791.
 52. Bowers, G. M.; Singer, J. W.; Bish, D. L.; Kirkpatrick, R. J. Structure and Dynamical Relationships of Ca²⁺ and H₂O in Smectite/²H₂O Systems. *Amer. Mineral.* **2014**, *99*, 318-331.
 53. Hoyt, D. W.; Turcu, R. V. F.; Sears, J. A.; Rosso, K. M.; Burton, S. D.; Felmy, A.; Hu, J. Z. High-Pressure Magic Angle Spinning Nuclear Magnetic Resonance. *J. Magn. Reson.* **2011**, *212*, 378-385.
 54. Turcu, R. V. F.; Hoyt, D. W.; Rosso, K. M.; Sears, J. A.; Loring, J. S.; Felmy, A.; Hu, J. Z. Rotor Design for High Pressure Magic Angle Spinning Nuclear Magnetic Resonance. *J. Magn. Reson.* **2013**, *226*, 64-69.
 55. Tenorio, R. P.; Alme, L. R.; Engelsberg, M.; Fossum, J. O.; Hallwass, F. Geometry and Dynamics of Intercalated Water in Na-Fluorohectorite Clay Hydrates. *J. Phys. Chem. C* **2008**, *112*, 575-580.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
56. Dazas, B.; Lanson, B.; Breu, J.; Robert, J-L.; Pelletier, M.; Ferrage, E. Smectite Fluorination and its Impact on Interlayer Water Content and Structure: A way to Fine Tune the Hydrophobicity of Clay Surfaces? *Micro. Meso. Mater.* **2013**, *181*, 233-247.
 57. Lowenstein, W. The Distribution of Aluminium in the Tetrahedra of Silicates and Aluminates. *Amer. Mineral.* **1954**, *39*, 92-96.
 58. Boinepalli, S.; Attard, P. Grand Canonical Molecular Dynamics. *J. Chem. Phys.* **2003**, *119*, 12769-12775.
 59. Dubbeldam, D.; Calero, S.; Ellis, D. E.; Snurr, R. Q. RASPA: Molecular Simulation Software for Adsorption and Diffusion in Flexible Nanoporous Materials. *Mol. Sim.* **2016**, *42*, 81-101.
 60. Young, D. A.; Smith, D. E. Simulations of Clay Mineral Swelling and Hydration: Dependence upon Interlayer Ion Size and Charge. *J. Phys. Chem. B* **2000**, *104*, 9163-9170.
 61. Cygan, R. T.; Liang, J.-J.; Kalinichev, A. G. Molecular Models of Hydroxide, Oxyhydroxide and Clay Phases and the Development of a General Force Field. *J. Phys. Chem. B* **2004**, *108*, 1255-1266.
 62. Cygan, R. T.; Romanov, V. N.; Myshakin, E. M. Molecular Simulation of Carbon Dioxide Capture by Montmorillonite Using an Accurate and Flexible Force Field. *J. Phys. Chem. C* **2012**, *116*, 13079-13091.
 63. Allen, M. P.; Tildesley, D. J. Computer Simulations of Liquids; Clarendon Press: Oxford, U.K., 1987.
 64. Peng, D. Y.; Robinson, D. B. A New Two-Constant Equation of State. *Ind. Eng. Chem. Fundam.* **1976**, *15*, 59-64.
 65. Smith, D. E. Molecular Computer Simulations of the Swelling Properties and Interlayer Structure of Cesium Montmorillonite. *Langmuir* **1998**, *14*, 5959-5967.
 66. Berend, I.; Cases, J. M.; Francois, M.; Uriot, J. P.; Michot, L.; Masion, A.; Thomas, F., Mechanism of Adsorption and Desorption of Water-Vapor by Homoionic Montmorillonites 2. The Li⁺, Na⁺, K⁺, Rb⁺, Cs⁺-Exchanged Forms. *Clays Clay Miner.* **1995**, *43*, 324-336.
 67. Hemmen, H.; Rolseth, E. G.; Fonseca, D. M.; Hansen, E. L.; Fossum, J. O.; Plivelic, R. X-ray Studies of Carbon Dioxide Intercalation in Na-Fluorohectorite Clay at Near-Ambient Conditions. *Langmuir* **2012**, *28*, 1678-1682.
 68. Criscenti, L. J.; Cygan, R. T. Molecular Simulations of Carbon Dioxide and Water: Cation Solvation. *Environ. Sci. Technol.* **2013**, *47*, 87-94.
 69. Sutton, R.; Sposito, G. Molecular Simulation of Interlayer Structure and Dynamics in 12.4Å Cs-Smectite Hydrates. *J. Coll. Interf. Sci.* **2001**, *237*, 174-184.
 70. Seidl, W.; Breu, J. Single Crystal Structure Refinement of Tetramethylammonium-Hectorite. *Z. Kristallogr.* **2005**, *220*, 169-176.
 71. Kosakowski, G.; Churakov, S. V.; Thoenen, T. Diffusion of Na and Cs in Montmorillonite. *Clays Clay Miner.* **2008**, *56*, 190-206.

TOC Graphic

