



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Mid infrared observations of Van Maanen 2: no substellar companion.

J. Farihi, E.E. Becklin, B.A. Macintosh

November 5, 2004

Astrophysical Journal

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

MID INFRARED OBSERVATIONS OF VAN MAANEN 2: NO SUBSTELLAR COMPANION

J. Farihi and E. E. Becklin

*Department of Physics & Astronomy, 8371 Math Sciences Building, University of
California, Los Angeles, CA 90095-1562*

`jfarihi@astro.ucla.edu, becklin@astro.ucla.edu`

and

B. A. Macintosh

*Institute of Geophysics & Planetary Physics, Lawrence Livermore National Laboratory,
7000 East Avenue L-413, Livermore, CA 94551*

`bmac@igpp.llnl.gov`

ABSTRACT

The results of a comprehensive infrared imaging search for the putative $0.06M_{\odot}$ astrometric companion to the 4.4 pc white dwarf van Maanen 2 are reported. Adaptive optics images acquired at $3.8\mu\text{m}$ reveal a diffraction limited core of $0.09''$ and no direct evidence of a secondary. Models predict that at 5 Gyr, a $50M_{\text{J}}$ brown dwarf would be only 1 magnitude fainter than van Maanen 2 at this wavelength and the astrometric analysis suggested a separation of $0.2''$. In the case of a chance alignment along the line of sight, a 0.4 mag excess should be measured. An independent photometric observation at the same wavelength reveals no excess. In addition, there exist published *ISO* observations of van Maanen 2 at $6.8\mu\text{m}$ and $15.0\mu\text{m}$ which are consistent with photospheric flux of a 6750 K white dwarf. If recent brown dwarf models are correct, there is no substellar companion with $T_{\text{eff}} \gtrsim 500$ K.

Subject headings: binaries: general—stars: low-mass, brown dwarfs—stars: individual(van Maanen 2)—white dwarfs

1. INTRODUCTION

Van Maanen’s star, also known as van Maanen 2 was discovered in 1917 by Adrian van Maanen (van Maanen 1917). Its discovery was quite accidental, as van Maanen was searching for common proper motion companions to HD4628 (Lalande 1299) and noticed the even larger proper motion of a nearby star. At 4.4 pc, van Maanen 2 is the nearest single white dwarf (Holberg, Oswalt, & Sion 2002).

Makarov (2004) reported the astrometric detection of a substellar companion to van Maanen 2 through analysis of *Hipparcos* data. The data suggested an orbital solution with a lower bound companion mass of $0.06 \pm 0.02 M_{\odot}$, a period of 1.57 yr, and a maximum separation on the sky of $0.3''$. Based on the published parameters, a binary orbital calculator (A. Ghez 2004, private communication) indicated a separation of $0.19''$ at position angle 274° near the middle of January 2004. This paper presents the results of a direct imaging search carried out during that time frame.

2. OBSERVATIONS, DATA REDUCTION, & PHOTOMETRY

2.1. Keck Adaptive Optics Observations

Van Maanen 2 was observed using the facility adaptive optics system (Wizinowich et al. 2000a,b) and the NIRC2 camera at Keck Observatory on 12 January 2004. Five dithered images were obtained through an L' ($3.4\text{--}4.1\mu\text{m}$) filter. To reduce thermal backgrounds and provide a more symmetric PSF, a circular pupil stop was used which describes an inscribed circle on the hexagonal Keck primary, with a diameter of ~ 9 meters. Each image consisted of 100 coadds of 0.4 seconds each. Van Maanen 2 itself was used as the guide star, with the adaptive optics system running at a rate of 80 Hz. The correction was quite good, with a Strehl ratio (measured relative to the PSF of the 9 meter pupil) of 0.75 and a full width at half maximum of $0.089''$, essentially diffraction limited. The plate scale was $0.01''$ per pixel.

The adaptive optics images were reduced using standard programs in the IRAF environment. For a given raw image, the sky was extracted by taking the median of the four remaining dithered images with rejection of hot and cold pixels. A flat frame was created by averaging all five sky frames and subsequently normalizing. Each dithered frame was then sky subtracted, flat fielded, and the resulting five frames were registered and averaged.

Due to the uncertainties in the wings of the adaptive optics PSF and weather that was not photometric, it was difficult to perform photometry on the Keck data. Therefore photometric observations of van Maanen 2 were acquired at the IRTF.

2.2. *IRTF and ISO Observations*

$3.8\mu\text{m}$ data on van Maanen 2 and two standard stars were acquired on 1 February 2004 by Alan Stockton at the 3 meter NASA Infrared Telescope Facility with NSFCAM (Rayner et al. 1993). For the standards, a 9 point dither pattern was used with each frame consisting of 0.15 second exposure times 100 coadds – yielding a total integration time of 135 seconds. The UKIRT faint standard SA92-342 ($L' = 10.44$ mag) was observed immediately before van Maanen 2 and the Elias standard star HD22686 ($L' = 7.19$ mag) was observed immediately afterward (Leggett et al. 2003). Two sets of images, acquired in the same manner as the standards, were obtained for van Maanen 2 for a total of 18 frames and 270 seconds integration. Conditions during the observations were reported as photometric.

The data on the two standards were reduced using standard programs in the IRAF environment. For a given raw image, the sky was extracted by taking the median of the eight remaining dithered images with rejection of hot and cold pixels. A flat frame was created by averaging all nine sky frames and subsequently normalizing. Each dithered frame was then sky subtracted, flat fielded, and the resulting nine frames were registered and averaged. Reducing the two sets of images on van Maanen 2 was more difficult due to the faintness of the target in each raw frame. It was found that the target was more easily seen in pairwise subtracted images rather than in sky subtracted images. All 18 pairwise subtracted frames were flat fielded, registered and averaged, creating one final image for photometric measurements.

The flux of both standards was measured with aperture radii of $1.2''$ and $2.4''$ and corrected for extinction. The error in determining the zero point was 0.03 mag. Van Maanen 2 was measured in the smaller aperture to minimize noise, corrected to the larger aperture (with an error of 0.04 mag) and extinction corrected. This yielded a signal to noise of 48 and a measurement error of 0.03 mag. The photometric measurement including all errors was $L' = 11.40 \pm 0.06$ mag for van Maanen 2.

ISOCAM observations of van Maanen 2 were carried out in 1997 in an effort to provide observational constraints on the origin of metals in the photospheres of cool white dwarfs (Chary, Zuckerman, & Becklin 1999). Data were taken at $6.8\mu\text{m}$ and $15.0\mu\text{m}$ on van Maanen 2 and several other white dwarfs as well as an A0V calibrator star. The data are listed in Table 1.

3. RESULTS & DISCUSSION

3.1. *Adaptive Optics - No Direct Evidence*

In order to estimate the brightness of the reported companion, a model for a $50M_J$ brown dwarf at 5 Gyr was chosen. This mass lies conservatively in the lower range of possible values and required no interpolation within available models. The model age is likely to be greater than the 3.67 Gyr cooling age (Bergeron, Leggett, & Ruiz 2001) of van Maanen 2 plus the ~ 0.5 Gyr main sequence lifetime (Maeder 1989) for a $\sim 4M_\odot$ progenitor of the $0.83M_\odot$ white dwarf (Weidemann 1987, 1990, 2000; Bragaglia, Renzini & Bergeron 1995; Bergeron et al. 2001). A substellar companion of this mass and age would be a late T dwarf ($T_{\text{eff}} \sim 800$ K) and have $M_{L'} = 14.3$ mag which is $L' = 12.5$ mag at 4.4 pc (Burrows et al. 1997). Photometric L' band data on known brown dwarfs do exist and the measurements agree with the models used here to within 0.3 mag for spectral type T6 (Leggett et al. 2002). Van Maanen 2 is predicted to have $L' = 11.43$ mag based on the model predicted $V - K = 0.84$ color of a 6750 K helium white dwarf (Bergeron, Wesemael, & Beauchamp 1995) and the $K - L' = 0.12$ color of a 6750 K blackbody. $V - K = 0.89$ is the measured color – the extrapolation was done from V in case of any contamination by a companion at K .

The reduced L' image (Figure 1) shows no indication whatsoever of a companion with the brightness expected from a brown dwarf of the type reported by Makarov (2004). From the published orbital parameters, the companion should have been at a separation of $0.19''$ and position angle of 274° on the date of the observation. The full width at half maximum of van Maanen 2 in the reduced image is $0.089''$ and the distance to the first Airy ring is $\approx 0.14''$. There are two extremely faint features within the Airy ring at position angles of 284 and 295° . Small aperture flux measurements, relative to the primary, at eight different evenly spaced locations around the Airy ring indicate that these features are unlikely to be real. The flux at 284 and 295° , $f/f_0 = 0.069$ and 0.073 respectively, are both within 2σ (0.018) of the average flux ($f/f_0 = 0.056$) in the ring and are almost certainly artifacts due to imperfect optical guide star corrections. If the feature at 295° were real, its brightness after subtracting the flux of the primary in the Airy ring implies $L' = 15.9$ mag and a mass of $\sim 15M_J$, assuming an age of 5 Gyr (Burrows et al. 1997). This is simply not massive enough to cause the reported astrometric wobble.

An artificial star was planted at $0.19''$ from the primary in order to test the ability to detect faint companions at this separation. The adaptive optics PSF, extracted from the reduced image of van Maanen 2, was used for the stellar profile of the planted star. To be conservative, this artificial star is a full 2.0 magnitudes fainter than the white dwarf – this was confirmed by placing the star at many positions on the image and measuring its flux

without contamination by the primary. The simulated star is readily seen in Figure 2.

Large errors in the reported orbital parameters allow for the possibility that the companion remained unresolved due to chance alignment during the observation. The values and computed errors in separation and position angle, derived from the binary orbital calculator, are $0.19'' \begin{smallmatrix} +0.07 \\ -0.18 \end{smallmatrix}$ and $274^\circ \begin{smallmatrix} +180 \\ -2 \end{smallmatrix}$ respectively. Hence the putative companion could have been located at almost any position in its orbit by the epoch of the observation. However, there is further evidence against this possibility.

3.2. Photometry - No Indirect Evidence

In the case of a chance alignment, any substellar companion would still cause excess emission at mid infrared wavelengths. This is primarily due to the size difference between brown dwarfs and white dwarfs; a ratio of 10:1 in radius. It is not possible for a significant occultation to occur. At most, a white dwarf could block only $\sim 1\%$ of the light from an orbiting brown dwarf.

As mentioned above, a value of $L' = 11.40 \pm 0.06$ mag was measured for van Maanen 2. This is to be compared with the value predicted for a 6750 K white dwarf, $L' = 11.43$ mag. The combined flux of the white dwarf plus a $50M_J$ brown dwarf at 5 Gyr would have a magnitude of $L' = 11.09$ mag. Hence it is concluded there is no excess emission at this wavelength.

Furthermore, the published *ISO* observations place even stronger constraints on the absence of flux from a brown dwarf around van Maanen 2. A recent model (Burrows, Sudarsky, & Lunine 2003) of the flux from a 5 Gyr, $25M_J$ brown dwarf was integrated over the *ISO* $15.0\mu\text{m}$ filter LW3. This results in a flux of about 1.0 mJy in this filter. The reported measurement taken at van Maanen 2 in 1997 was 0.5 ± 0.2 mJy. This measurement is consistent with photospheric flux from the white dwarf and an *excess* of 5σ most likely would have been detected. If the *ISO* results are correct and the models are right, any companion with $T_{\text{eff}} \gtrsim 500$ K is ruled out – this includes 10 Gyr old brown dwarfs with $M \geq 35M_J$ (Burrows et al. 1997, 2003).

Table 1 summarizes all existing photometric data on van Maanen 2 and the corresponding fluxes are plotted in Figure 3. It should be clear from the figure that the measured fluxes are all consistent with a single white dwarf with a temperature around 6800 K.

4. CONCLUSION

Three compelling infrared observations of van Maanen 2 are presented which rule out the presence of a substellar companion warmer than ~ 500 K. First, $3.8\mu\text{m}$ adaptive optics images reveal no significant flux at the predicted magnitude and position of the putative companion reported by Makarov (2004). Second, $3.8\mu\text{m}$ photometric observations are consistent with no excess emission at this wavelength as would be expected from a $50M_J$ brown dwarf at 5 Gyr. And third, $15.0\mu\text{m}$ data also are consistent with no excess emission, ruling out any possible companion massive enough to induce the reported astrometric wobble.

Part of the data presented herein were obtained at Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology (CIT), the University of California and the National Aeronautics and Space Administration (NASA). This publication makes use of data acquired at the Infrared Telescope Facility, which is operated by the University of Hawaii under Cooperative Agreement no. NCC 5-538 with NASA, Office of Space Science, Planetary Astronomy Program. Some data used in this paper are part of the Two Micron All Sky Survey (2MASS), a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center (IPAC)/CIT, funded by NASA and the National Science Foundation (NSF). 2MASS data were retrieved from the NASA/IPAC Infrared Science Archive, which is operated by the Jet Propulsion Laboratory, CIT, under contract with NASA. The authors wish to express their gratitude to A. Stockton, J. Rayner and A. Tokunaga for their assistance in acquiring the IRTF observations of van Maanen 2. Sincere thanks go to J. Larkin and M. Barczys for donating some of their Keck adaptive optics time for this project. Acknowledgement also goes to B. Hansen for bringing this system to our attention, to B. Zuckerman for a careful reading of this manuscript and many discussions, to A. Ghez for the binary orbit calculator, and to B. Vacca for help in the data reduction. Part of this work was performed under the auspices of the U.S. Department of Energy, National Nuclear Security Administration by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48. This research has been supported in part by grants from NASA to UCLA.

Facilities: Keck, IRTF

REFERENCES

- Bergeron, P., Leggett, S. K., & Ruiz, M. T. 2001, *ApJS*, 133, 413
- Bergeron, P., Wesemael, F., & Beauchamp, A. 1995, *PASP*, 107, 1047

- Bessell, M. S., & Brett, J. M. 1988, PASP, 100, 1134
- Bragaglia, A., Renzini, A., & Bergeron, P. 1995, ApJ, 443, 735
- Burrows, A. 1997, ApJ, 491, 856
- Burrows, A., Sudarsky, D., & Lunine, J. I. 2003, ApJ, 596, 587
- Chary, R., Zuckerman, B., & Becklin, E.E. 1999, Conf. Proceedings: the Universe as Seen by ISO, ed. P. Cox & M. F. Kessler (Noordwijk: ESA/ESTEC), 289
- Holberg, J. B., Oswalt, T. D., & Sion, E. M. 2002, ApJ, 571, 512
- Leggett, S. K. et al. 2002 ApJ, 564, 452
- Leggett, S. K. et al. 2003 MNRAS, 345, 144
- Maeder, A. 1989, Proceedings of the 5th IAP Workshop: Astrophysical Ages and Dating Methods, eds. E. Vangioni-Flam, M. Cassé, J. Audouze, J. Tran Thanh Van, 71
- Makarov, V. V. 2004, ApJ, 600, L71
- Rayner, J. T., et al. 1993, SPIE, 1946, 490
- van Maanen, A. 1917, PASP, 29, 258
- Weidemann, V. 1987, A&A, 188, 74
- Weidemann, V. 1990, ARA&A, 28, 103
- Weidemann, V. 2000, A&A, 363, 647
- Wizinowich, P., et al. 2000, PASP, 112, 315
- Wizinowich, P., Acton, D. S., Lai, O., Gathright, J., Lupton, W., & Stomski, P., 2000, SPIE 4007, 64




Fig. 1.— $3.8\mu\text{m}$ adaptive optics image of van Maanen 2. There is no direct evidence for a companion one magnitude fainter than the primary.




Fig. 2.— Adaptive optics image of van Maanen 2 with an artificial star placed at $0.19''$ from the white dwarf. The magnitude of the planted star is $L' \approx 13.4$ mag and is clearly seen.




Fig. 3.— Spectral energy distribution of van Maanen 2 from the data in Table 1. Overplotted is a 6800 K blackbody demonstrating that the data reveal no excess emission in the infrared and are consistent with a single star.

Table 1. Photometric Data on van Maanen 2

Filter	$\lambda_0(\mu\text{m})$	Magnitude	$F_\nu(\text{mJy})$
<i>B</i>	0.44	12.91	27.9 ± 1.4
<i>V</i>	0.55	12.39	40.2 ± 1.2
<i>R</i>	0.64	12.13	43.1 ± 1.3
<i>I</i>	0.80	11.90	42.0 ± 1.3
<i>J</i>	1.22	11.69	34.4 ± 0.6
<i>H</i>	1.65	11.57	24.7 ± 0.5
<i>K_s</i>	2.16	11.50	16.7 ± 0.5
<i>L'</i>	3.76	11.40	6.8 ± 0.4
LW2	6.75	...	2.0 ± 0.4
LW3	15.0	...	0.5 ± 0.2

Note. — *BVRI* photometry is from Bergeron et al. (2001) and is on the Johnson-Cousins system. *JHK_s* data are from 2MASS and agree with the near infrared data reported by Bergeron et al. (2001). The *L'* data are from the present work and the LW2 & LW3 fluxes are from *ISO* (Chary, Zuckerman, & Becklin 1999).