

# Ab-initio shock states and thermodynamic properties of liquid silicates

Joshua P Townsend<sup>1</sup>, Luke Shulenburger

<sup>1</sup>*High Energy Density Physics Theory, Sandia National Laboratories, Albuquerque, New Mexico, USA*

We present *ab-initio* calculations of the principal Hugoniot states, entropy, sound velocity, as well as transport properties for liquid silicate  $Mg_2SiO_4$ . Our investigations heavily employ density functional theory-based molecular dynamics (QMD) calculations to self-consistently compute thermodynamic quantities. In this presentation, we discuss the Hugoniot states and possible de-mixing of liquid  $Mg_2SiO_4$ . We then turn to new calculations of entropy on the Hugoniot of  $Mg_2SiO_4$  using the 2PT method [1] and compare to results from more traditional thermodynamic integration. Wherever possible, we compare our results to experiment. The results highlight the utility and predictive power of QMD methods for high energy density physics.

Sandia National Laboratories is a multimission laboratory managed and operated by National Technology and Engineering Solutions of Sandia, LLC., a wholly owned subsidiary of Honeywell International, Inc., for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA0003525.

## REFERENCES

[1] M.P. Desjarlais, "First-principles calculation of entropy for liquid metals", PRE **88**, 062145 (2013).