



Computational Geochemistry at Sandia National Labs: Gas Hydrates and Other Applications

Jeffery A. Greathouse
Geochemistry Department
Sandia National Laboratories

People

Randy Cygan

Blake Simmons

Bob Bradshaw

Eric Majzoub

Funding

Laboratory Directed Research & Development

DOE Basic Energy Sciences




Outline

- **Atomistic simulations**
- Methane Hydrate
- HCFC Hydrates
- Other Molecular Simulation Projects



Time and Length Scales



Molecular Modeling Methods

Experiment
Ab initio

Forcefield

+

Structure

Structure

Molecular Mechanics

$$F = ma$$

Energy Minimization


Conformational Analysis

Lattice Dynamics


Molecular Dynamics

Monte Carlo

Molecular Model



Cluster



Periodic

Quantum Mechanics

$$H\psi = E\psi$$

Hartree-Fock and DFT



EM, CA, MD

Structure
Physical properties
Thermodynamics
Kinetics
Spectroscopy



Potential Parameters

Intermolecular Terms (rigid or flexible models)




$$U_{ij} = \frac{q_i q_j}{r} + 4\epsilon \left[\left(\frac{\sigma_{ij}}{r} \right)^{12} - \left(\frac{\sigma_{ij}}{r} \right)^6 \right]$$


Long-range electrostatics handled in reciprocal space by **Ewald sum** or **particle mesh** techniques

Potential Parameters

Intramolecular Terms (flexible models)




Bond Stretch $E = K(r - r_0)^2$




Angle Bend $E = K(\theta - \theta_0)^2$

$$E = k(1 + \cos(n\phi - \phi_o))$$



Torsion

$$E = k(1 + \cos(n\chi - \chi_o))$$



Out-of-Plane Stretch

Simulation vs Experiment

Property	Simulation	Experiment
Unit cell data	Lattice constants (constant P simulation)	Diffraction
Local atomic coordination	Radial distribution functions	EXAFS
Interfacial structure	Atomic density profiles	X-Ray Scattering
Vibrational motion	Power spectra	IR/Raman



Outline


- Atomistic simulations
- **Methane Hydrate**
- HCFC Hydrates
- Other Molecular Simulation Projects




Natural Gas Hydrates

Components

- Water in various cage sizes (host)
- Gas-phase guest molecules



5^{12}



$5^{12}6^2$





$5^{12}6^4$



Natural Gas Hydrates


Components

- Water in various cage sizes (host)
- Gas-phase guest molecules



46 H_2O

Structure I



136 H_2O

Structure II




34 H_2O

Structure H


Hydrate Crystal Structure	I (12.0 Å)		II (17.1 Å)		H (a=12.3 Å, c=10.2 Å)		
Cavity	Small 5^{12}	Large $5^{12}6^2$	Small 5^{12}	Large $5^{12}6^4$	Small 5^{12}	Large $4^3 5^6 6^3$ $5^{12}6^8$	
# per unit cell	2	6	16	8	3	2	1
Coordination #	20	24	20	28	20	20	36





Effect of Water Potential

Methane Hydrate (sI), 200 K, 20 bar



	Water Dipole Moment (D)	Bulk Density (kg·m ³)
SPC	2.274	945 ± 2
SPC/E	2.351	966 ± 2
TIP4P	2.177	961 ± 2
Polarizable TIP4P	3.15 ± 0.14	999 ± 2
Flexible SPC	2.44 ± 0.10	955 ± 2



Bulk Water

1.85

997

English and MacElroy, *J. Comp. Chem.* **2003**


g = Radial Distribution Function




Sandia National Laboratories



MD Simulations of H₂ Hydrate



Drastic change in structure
when # H₂/unit cell > 48




1 H₂/small cage
4 H₂/large cage


Alavi et al, *J. Chem. Phys.*, 2005

Methane Hydrate Formation

Methane Hydrate (sI), 300 K, 30 bar, rigid SPC water





Hydrate cluster formation at 6.9 ns




Face-sharing 5^{12} dimer (sII)

Moon et al, *J. Am. Chem. Soc.*, **2003**

Methane sl, Cage Structure




Methane-Methane
Coordination # = 13.5



Methane-Water
Coordination # = 23 (first peak)


Methane Hydrate Thermal Expansion



Greathouse et al, *J. Phys.Chem. B* **2006**, 110, 6428-6431




Methane Hydrate Power Spectrum



Greathouse et al, *J. Phys.Chem. B* **2006**, *110*, 6428-6431



Methane Hydrate Thermal Expansion



Normal modes from methane in large cage similar to methane gas




Outline

- Atomistic simulations
- Methane Hydrate
- **HCFC Hydrates**
- Other Molecular Simulation Projects



Desalination of Brackish Water




<http://wrrri.nmsu.edu/>

All wet

Texas is rich in brackish groundwater, with more than an estimated 2.7 billion acre-feet spread throughout the state's 16 regional water planning areas. Desalination makes millions of gallons of that water drinkable.

Water planning area

- A Panhandle
- B Region B
- C Region C
- D North East Texas
- E Far West Texas
- F Region F
- G Brazos
- H Region H
- I East Texas
- J Plateau
- K Lower Colorado
- L South Central Texas
- M Rio Grande
- N Coastal Bend
- O Llano Estacado
- P Lavaca



SOURCE: Texas Water Development Board.

NOTE: An acre-foot of water is equivalent to the amount of water needed to cover an acre of land with a foot of water.

Drinking water.....< 1k mg/L

Brackish water.....1k-10k mg/L

Seawater.....> 30k mg/L




R141b-Hydrate, Structure II

(R141b = 1,1-dichloro-1-fluoroethane, 265 K)


Small cages are empty.

R141b molecules freely rotate within large cages.




Density Functional Theory (DFT)

Cluster model of R141b in **LARGE** cage



$$\Delta E_{\text{formation}} = -49.9 \text{ kcal/mol}$$

Cluster model of R141b in **SMALL** cage




$$\Delta E_{\text{formation}} = +38.6 \text{ kcal/mol}$$

- Strong interactions between guest molecule and cage waters lead to destabilization of the hydrate; R141b is not expected to occupy the small cage


R141b Isomer Instability

Cluster model of R141b in **LARGE** cage



$$\Delta E_{\text{formation}} = -49.9 \text{ kcal/mol}$$


Isomer (1,2-dichloro-1-fluoroethane)



$$\Delta E_{\text{formation}} = -43.9 \text{ kcal/mol}$$

- Electron density map (through C-C-Cl plane) for isomer in large cage exhibits strong interaction between isomer and water molecules as indicated by sharing of electrons

R141b Hydrate Thermal Expansion



Experimental forming conditions:


R141b 280 K, 1 atm pressure

Isomer No formation seen at 273 K, 1 atm

Greathouse et al, *J. Phys.Chem. C* **2007**, submitted.




R141b Hydrate Power Spectrum




R141b Peak Comparison

Slight shift (5-10 cm^{-1}) in C-Cl and C-F stretch modes between pure gas and hydrate phases



Experimental Raman Spectrum



Gas Hydrate Summary and Conclusions

➤ Conclusions

- Flexible atomistic forcefield (CVFF) is able to predict hydrate structure and aid in interpretation of Raman spectra
- Quantum calculations are useful in refining guest/cage diameters

➤ Future work

- Monte Carlo simulations to determine cage occupancies
- Kinetic Monte Carlo simulations to study hydrate nucleation and growth
- Effect of mineral surfaces on hydrate nucleation



Outline

- Atomistic simulations
- Methane Hydrate
- HCFC Hydrates
- **Other Molecular Simulation Projects**





UO₂-Clay Simulations

➤ Montmorillonite (**octahedral charge**), Beidellite (**tetrahedral charge**), and Pyrophyllite (**no charge**)



➤ **Simulation specifics:**

- Monte Carlo and Molecular Dynamics Simulations
- Nonbonded interactions only (rigid-body)
- Constant pressure interlayer simulations
- Constant volume adsorption simulations (external surface)


➤ **Goal:** examine equilibrium water content and layer spacing, sorption tendencies of:



Large-Scale Simulations of Uranyl Adsorption




Integrate density peaks
to obtain percent sorbed




$$K = \frac{\% \text{U}_{\text{sorbed}}}{\% \text{U}_{\text{aqueous}}} = \frac{\text{area sorbed}}{\text{area aqueous}}$$

Sorption Equilibria (K_D)



UO_2^{2+} sorption is influenced by solution concentration






Na^+ sorption is unaffected by solution concentration

Greathouse and Cygan, *ES&T*, in press



Carbonato Complexes and Oligomers




High carbonate concentration hinders uranyl sorption
Curtis et al, *App. Geochem.* **2004**; Fuller et al, *ES&T* **2003**.


Zinc-based metal organic frameworks (MOFs): nanoporous materials with exceptional properties

- MOFs are inorganic complexes linked by coordinated organic ligands
- Crystalline nanoporous materials
 - **Homogeneous, periodic pores**
 - **$\sim 4 - 19 \text{ \AA}$ aperture diameter; $13 - 29 \text{ \AA}$ Van der Waals diameter (calculated)**
 - **Extremely high surface areas (up to $4,500 \text{ m}^2/\text{g}$)**
 - **Very low density**
- ➔ Tunable pore environment for applications
- ➔ Rational design possible because crystallinity enables first-principles modeling

MOF5 (IRMOF-1): Zn_4O tetrahedra linked by benzene dicarboxylate linkers





IRMOF-1: C_4 axis (above); C_3 axis (below)





MD simulations reveal substantial molecular motion by the solvent-free MOF5 lattice

0.0% Water, lattice parameter = 25.61 Å, 300 K



Increasing the amount of H₂O results in chains of hydrogen-bonded water molecules




2.3% Water (9 H₂O/unit cell), 25.37 Å, 300 K



Water molecules gradually replace framework O atoms. Zn-O_{water} distance $\approx$ 1.9 Å.

Close examination indicates that H_2O disrupts the tetrahedral geometry of the Zn_4O cluster

Equilibrium structure, no H_2O



MOF5 structural collapse

9.5% Water (36 H₂O/unit cell), 20.14 Å, 300

