

LA-UR-20-29477

Approved for public release; distribution is unlimited.

Title: Neutron-Induced Fission and Scattering Measurements at LANSCE

Author(s): Kelly, Keegan John

Intended for: Discussion of LANL with CENTAUR students, for advertising purposes and informing them about work/life at LANL.

Issued: 2020-11-18

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Neutron-Induced Fission and Scattering Measurements at LANSCE


CENTAUR Meets LANL
11/10/2020

Keegan J. Kelly,
on behalf of P-27 and the Chi-Nu collaboration

Outline

- The Los Alamos Neutron Science Center (LANSCE)
- The Chi-Nu Experiment: Prompt Fission Neutron Spectra
- Neutron Scattering Measurements
- Collaborations with T and X Divisions at LANL
- Future Work


LANSCCE Layout and Experimental Areas



- Simultaneous H⁺ and H⁻ beams
- >15 experiments at once
- Chi-Nu, DANCE, SPIDER, TPC, TKE, ICEHouse, Proton Radiography, Materials Science, UCN, IPF, and more

Chi-Nu Goal: Measure PFNS χ Matrix for Actinides


Fig: Neudecker et al. NDS 131 (2016) 289



Ultimate Goal: Stand the Test of Time

- Repeated mistakes in experiments with regard to uncertainties:
 - Frequently no covariance matrices, many uncertainty sources missing or ignored
 - Sometimes not possible to know how to accurately correct old data
- Potentially unknown correlations within and between experiments

Chi-Nu is a single experiment to measure the major actinide PFNS for all incident and outgoing energies of interest with a thorough analysis of systematic uncertainties and covariances




The Chi-Nu Arrays



The Chi-Nu Experiment

Devlin *et al.* NDS 148 322 (2018)


Haight *et al.* NDS 123 130 (2015)



The Chi-Nu Experiment

Devlin *et al.* NDS 148 322 (2018)

Haight *et al.* NDS 123 130 (2015)




Chi-Nu is a double time-of-flight experiment

- $t_f - t_0 \rightarrow E_n^{inc}$
- $t_n - t_f \rightarrow E_n^{inc}$
- PPAC fission counter
- Two n -detector arrays used separately
- Combined to form single PFNS shape

Correlations Between All E_{out} and All E_{inc} Points


- Li-glass and liquid data combined to form single shape result for each E_{inc}
- Correlations between PFNS points within each detector array
- Correlations between PFNS points across detector arrays
- Correlations between PFNS points from each E_{inc} bin and E_{out}

Required for accurate representation of Chi-Nu data in evaluations



Mean PFNS Energy, $\langle E \rangle$; Observing New Physics


- Never been calculated for a PFNS measurement
- $\langle E \rangle$ points are *highly* correlated
- Implies that the relative value of the $\langle E \rangle$ points is fairly well known
 - More well known than the total uncertainty would suggest




Repurposing Liq. Scints. for Dual n - γ Analysis

- Able to leverage data collected from other experiments using the Chi-Nu liquid scintillator array
 - High statistics, to better guide data analysis
 - Difficult to obtain high statistics with trial detector purchases
- Start with easy case: natural carbon

Begin by simply looking for n - γ coincidence in post-processing analysis




Operated by Triad National Security, LLC for the U.S. Department of Energy's NNSA



Random Coincidence Background Technique

- Random coincidence rates derived from Poisson probabilities for *uncorrelated* detection rates [†]
 - true coincidence rate must be low
- Calculate the total probability for:
 1. Detecting a γ at time t_γ
 2. Not detecting n over coinc. time $t_n - t_\gamma$
 3. Detecting n at time t_n


$$\begin{aligned} \text{Coinc. Rate} &= r_b = r_\gamma r_n \Delta_t \\ &\Rightarrow b = \frac{\gamma n}{N_{t_0}} \\ &\text{with } \gamma, n = \text{counts} \end{aligned}$$




Correlated n - γ Distributions Collected as Well

Extract n angular distributions from elastic scattering and correlated n - γ distributions from inelastic scattering

Elastic Scattering



Inelastic Scattering




Very few experiments have measured n - γ distributions

→ Measured at limited energies and angles


Even “Simple” Angular Dists. Can Be Complicated!

Extract n angular distributions from elastic scattering and correlated n - γ distributions from inelastic scattering

Elastic Scattering



Inelastic Scattering





Distributions change with changing angular momentum distributions

\Rightarrow **Need to measure the entire distribution**

Fe Levels Can Be Resolved

- Fe levels are reasonably dense
- The liquid scint. time resolution allows for low-lying state separation




$(n,2n)$ and $(n,3n)$ Reactions Observed in Pt Data

- Fe levels are reasonably dense
- The liquid scint. time resolution allows for low-lying state separation

- Natural Pt contains 5 isotopes

- ^{190}Pt : 0.01%
- ^{192}Pt : 0.79%
- ^{194}Pt : 32.97%
- ^{195}Pt : 33.83%
- ^{196}Pt : 25.24%
- ^{198}Pt : 7.16%

- Regions of high counts can be understood as inelastic scattering, $(n,2n)$, and $(n,3n)$ reactions



Collaboration Across the Wide World of LANL

- Direct communication with evaluators (T and X division)
 - Work with ENDF evaluators to get data correctly incorporated into evaluations
 - Assist with interpretation of other data including in evaluation
 - Extremely informative and rewarding process
- MCNP developers up the road (X Division)
 - Direct inclusion of models and theories into calculations
 - Able to influence the direction of development for MCNP
- Opportunities for collaboration and experiment hosting from ISR and NEN
 - LANSCE is a key experimental resource for LANL, and divisions all over LANL utilize it
 - Cross-group and cross-division mobility is a feature of LANL
- Working with experimenters from all over the world
 - When you're an experimental hub, you get the opportunity to be involved in many different experiments