


Reaction Pathways of Cyclopentanone Oxidation Intermediates

Sarah Khanniche, William H. Green

Wednesday, June 26 2019


ORLÉANS,
JUNE
23 - 27 2019

Cyclopentanone(CPO) as a Biofuel




Aim of the project

Combustion of CPO




What are the elementary reactions?
How fast are they?

Initiation reactions




Chain propagation reactions



To identify reaction pathways of cyclopentanone oxidation intermediates using *ab initio* calculations and RMG simulations

RMG simulations

Rate-based Model Enlarging Algorithm (flux-based algorithm)




$$0.001 < \epsilon < 0.1$$

R_{char} = reaction system's characteristic rate

Outline of the talk

Combustion of CPO



1. *Ab initio* calculations

QOO: H-migration vs HO₂ elimination


QOOH: Ring opening vs cyclic ether formation

2. RMG simulations

Evolution of the system with time


Sensitivity analysis

Computational methods



RMG simulations


Rate-based Model Enlarging Algorithm (flux-based algorithm)




$$0.001 < \epsilon < 0.1$$

R_{char} = reaction system's characteristic rate

Oxidation of 2-oxo-cyclopentyl





Oxidation of 3-oxo-cyclopentyl





Oxidation of 2-oxo- & 3-oxo-cyclopentyl

ΔH_{0K} in kcal mol⁻¹





Oxidation of 2-oxo- & 3-oxo-cyclopentyl

ΔH_{0K} in kcal mol⁻¹



Oxidation of 2-oxo- & 3-oxo-cyclopentyl



ΔH_{0K} in kcal mol⁻¹



Oxidation of 2-oxo- & 3-oxo-cyclopentyl



ΔH_{0K} in kcal mol⁻¹

1. Shallow αRO_2 well vs deep βRO_2 well





Oxidation of 2-oxo- & 3-oxo-cyclopentyl

ΔH_{0K} in kcal mol⁻¹





Oxidation of 2-oxo- & 3-oxo-cyclopentyl

ΔH_{0K} in kcal mol⁻¹



Oxidation of 2-oxo- & 3-oxo-cyclopentyl



ΔH_{0K} in kcal mol⁻¹



Oxidation of 2-oxo- & 3-oxo-cyclopentyl



ΔH_{0K} in kcal mol⁻¹

2. There is a 2 kcal mol⁻¹ gap between TS α' Habs and TS β' Habs





Oxidation of 2-oxo- & 3-oxo-cyclopentyl

ΔH_{0K} in kcal mol⁻¹





Oxidation of 2-oxo- & 3-oxo-cyclopentyl

ΔH_{0K} in kcal mol⁻¹



Oxidation of 2-oxo- & 3-oxo-cyclopentyl



ΔH_{0K} in kcal mol⁻¹



Oxidation of 2-oxo- & 3-oxo-cyclopentyl



ΔH_{0K} in kcal mol⁻¹

3. QOOH α' and QOOH α are more stable than QOOH β' and QOOH β





Oxidation of 2-oxo- & 3-oxo-cyclopentyl

ΔH_{0K} in kcal mol⁻¹





HO₂ elimination vs H-abstraction

ΔH_{0K} in kcal mol⁻¹





HO₂ elimination vs H-abstraction

ΔH_{0K} in kcal mol⁻¹





HO₂ elimination vs H-abstraction

ΔH_{0K} in kcal mol⁻¹





HO₂ elimination vs H-abstraction


ΔH_{0K} in kcal mol⁻¹



Kinetics, $k_{\infty}(T)$



Kinetics, $k_{\infty}(T)$




$A = 1.9 \times 10^3 \text{ s}^{-1}$
 $n = 2.45$
 $E_a = 21.9 \text{ kcal mol}^{-1}$

$A = 8.9 \times 10^4 \text{ s}^{-1}$
 $n = 1.98$
 $E_a = 21.3 \text{ kcal mol}^{-1}$

$A = 4.0 \times 10^8 \text{ s}^{-1}$
 $n = 1.27$
 $E_a = 22.7 \text{ kcal mol}^{-1}$

$$k(T) = AT^n \exp\left(\frac{E_a}{RT}\right)$$



$A = 4.1 \times 10^4 \text{ s}^{-1}$
 $n = 2.08$
 $E_a = 20.4 \text{ kcal mol}^{-1}$

$A = 1.0 \times 10^5 \text{ s}^{-1}$
 $n = 2.51$
 $E_a = 27.1 \text{ kcal mol}^{-1}$


$A = 6.3 \times 10^8 \text{ s}^{-1}$
 $n = 1.21$
 $E_a = 23.7 \text{ kcal mol}^{-1}$

$A = 7.9 \times 10^8 \text{ s}^{-1}$
 $n = 1.29$
 $E_a = 29.1 \text{ kcal mol}^{-1}$

Oxidation of 2-oxo-cyclopentyl

10


ΔH_{0K} in kcal mol⁻¹




Oxidation of 3-oxo-cyclopentyl

11


ΔH_{0K} in kcal mol⁻¹




Acyclic radicals



Cyclic Ether




Enol



Outline of the talk

Combustion of CPO



1. *Ab initio* calculations

QOO: H-migration vs HO₂ elimination

QOOH: Ring opening vs cyclic ether formation

2. RMG simulations


Evolution of the system with time

Sensitivity analysis

Oxidation of CPO

isothermal isobaric
batch reactor

$T = 700\text{ K}$
 $P = 10\text{ bar}$




- O2(2)
- H2O2(13)
- cyc5one(1)
- H2O(9)
- S(51697)
- prodAB(52)
- CO(14)
- CO2(15)
- S(290627)
- S(2001)

2.6% CPO, 19% O_2 , 76% N_2


451 species & 18188 reactions

$\varepsilon = 0.01$







Sensitivity Analysis

$T = 700 \text{ K}$
 $P = 10 \text{ bar}$




Sensitivity of the concentration of CPO (c) to the kinetics (k_i) of the RMG-generated model




$$T = 700 \text{ K}$$

$$P = 10 \text{ bar}$$

Sensitivity Analysis



Sensitivity of the concentration of CPO (c) to the thermo (G_i) of the RMG-generated model



1. *Ab initio* calculations

- ❑ PES for $\alpha R\bullet + O_2$ and $\beta R\bullet + O_2$ @ CBS-QB3 level of theory

QOO species: Competition between HO_2 elimination (termination) and α' -Habs (propagation)

QOOH species: enol formation for $\alpha QOOH\alpha'$, cyclic ether formation (3-membered rings)

Mechanism of 20 species, 21 rate constants and 15 thermodynamic data

2. RMG simulations

- ❑ Mechanism of 450 species and 18800 reactions

Olefin obtained from HO_2 elimination

- Compute thermo and kinetics for the species/reactions of the sensitivity analysis
- Ignition delay = $f(T)$

Acknowledgments



US Department of Energy (DOE)
Co-optimization of Fuels & Engines project

Pr. William H. Green
Matt Johnson
Xiaorui Dong

Green Group members
RMG developers

Green Group @ MIT



